

## Thermodynamic and structural studies of the solid deposit formation in paraffinic petroleum cuts

Michel Dirand, Dominique Petitjean, Mohammed Bouroukba

► **To cite this version:**

Michel Dirand, Dominique Petitjean, Mohammed Bouroukba. Thermodynamic and structural studies of the solid deposit formation in paraffinic petroleum cuts. David, N and Jaubert, JN and Privat, R. XXXIX JEEP - 39TH EDITION OF THE JOINT EUROPEAN DAYS ON EQUILIBRIUM BETWEEN PHASES, 2013, Nancy, France. 3, 2013, MATEC Web of Conferences. <10.1051/matecconf/20130301006>. <hal-01292159>

**HAL Id: hal-01292159**

**<https://hal.univ-lorraine.fr/hal-01292159>**

Submitted on 10 May 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Thermodynamic and structural studies of the solid deposit formation in paraffinic petroleum cuts

M. Dirand, D. Petitjean and M. Bouroukba

LRGP (UMR CNRS 7274), ENSIC, Université de Lorraine, 1 rue Grandville F54001 NANCY Cedex

A general review [1, 2] is presented concerning jointly the crystallographic structures and the transition physico-chemical data ( $T_{\text{fus}}$ ,  $\Delta_{\text{fus}}H$ ,  $\Delta_{\text{fus}}S$ ,  $\Delta_{\text{eb}}H$ ,  $\Delta_{\text{sub}}H$ ,  $C_p$ ) in the pure normal alkanes, as well as the structural and thermodynamic behavior of their synthetic binary, ternary and multinary model mixtures and of real petroleum waxes, particularly the solubility and mixture properties [3-6]. A major part of the structural and thermodynamic data of the literature and our experimental results are listed from methane up to the alkane with carbon atom number equal to 390 [1, 2] as well as their variations versus the atom carbon number.

The experimental results [3] and those of literature concerning the pure n-alkane dissolution enthalpies allow to determine the enthalpy of the formation of one or several crystallized solid solutions of the mixtures [4] and to highlight the influence of the characteristic parameters of the paraffinic distribution (theoretical average chain length, n-alkanes number, monophasic or polyphasic state) on the excess properties in the solid state. This data bank permits to test the predictive capacities of the UNIQUAC thermodynamic model. After modifying the expression of the interaction energy to take into account the effect of the size difference between the n-alkanes of the mixtures and the internal disorder induced by the n-alkane length distribution, this model give very good results for the prediction of the excess enthalpy of complex multialkane samples [4].

### References

1. M. Dirand, M. Bouroukba, V. Chevallier, D. Petitjean, E. Behar, V. Ruffier-Meray, Review of J. Chem. Eng. Data **47** 115 (2002)
2. M. Dirand, M. Bouroukba, A-J. Briard, V. Chevallier, D. Petitjean, J-P. Corriou, J. Chemical Thermodynamics **34** 1255 (2002)
3. A-J. Briard, M. Bouroukba, D. Petitjean, M. Dirand, J.Chem. and Eng. Data **48** 1574 (2003)
4. A-J. Briard, M. Bouroukba, M. Dirand, J. Chem. and Eng. Data **49** 899 (2004)
5. A-J. Briard, M. Bouroukba, D. Petitjean, M. Dirand, J. Chem. Eng. Data **49** 1201 (2004)
6. A-J. Briard, M. Bouroukba, D. Petitjean, N. Hubert, J.C. Moïse, M. Dirand, Fuel **85** 764 (2006)