

Mobile Insertion Cassette Elements Found in Small Non-Transmissible Plasmids in Proteeae May Explain qnrD Mobilization

Thomas Guillard, Antoine Grillon, Christophe de Champs, Céline Cartier, Janick Madoux, Béatrice Berçot, Anne-Laure Lebreil, Alain Lozniewski,

Jacques Riahi, Véronique Vernet-Garnier, et al.

▶ To cite this version:

Thomas Guillard, Antoine Grillon, Christophe de Champs, Céline Cartier, Janick Madoux, et al.. Mobile Insertion Cassette Elements Found in Small Non-Transmissible Plasmids in Proteeae May Explain qnrD Mobilization. PLoS ONE, 2014, 9 (2), pp.e87801. 10.1371/journal.pone.0087801. hal-01481324

HAL Id: hal-01481324 https://hal.univ-lorraine.fr/hal-01481324

Submitted on 2 Mar 2017 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mobile Insertion Cassette Elements Found in Small Non-Transmissible Plasmids in *Proteeae* May Explain *qnrD* Mobilization

Thomas Guillard^{1,2,3}, Antoine Grillon³, Christophe de Champs^{1,3}, Céline Cartier³, Janick Madoux³, Béatrice Berçot^{2,5}, Anne-Laure Lebreil¹, Alain Lozniewski⁴, Jacques Riahi⁵, Véronique Vernet-Garnier^{1,3}, Emmanuelle Cambau^{2,5}*

1 EA 4687, UFR Médecine SFR CAP-Santé Université de Reims Champagne-Ardenne, Reims, France, 2 EA3964, PRES Sorbonne Paris Cité Université Paris Diderot-Paris 7, Paris, France, 3 Laboratoire de Bactériologie-Virologie-Hygiène, CHU Reims - Hôpital Robert Debré, Reims, France, 4 Laboratoire de Bactériologie, CHU Nancy - Hôpital Central, Nancy, France, 5 Laboratoire de Bactériologie, AP-HP Groupe Hospitalier Lariboisière - Saint Louis, Paris, France

Abstract

qnrD is a plasmid mediated quinolone resistance gene from unknown origin, recently described in Enterobacteriaceae. It encodes a pentapeptide repeat protein 36–60% different from the other Qnr (A, B, C, S and VC). Since most qnrD-positive strains were described as strains belonging to Proteus or Providencia genera, we hypothesized that *anrD* originated in Proteeae before disseminating to other enterobacterial species. We screened 317 strains of Proteeae for qnrD and its genetic support by PCR. For all the seven qnrD-positive strains (4 Proteus mirabilis, 1 Proteus vulgaris and 2 Providencia rettgeri) the gene was carried onto a small non-transmissible plasmid, contrarily to other qnr genes that are usually carried onto large multi-resistant plasmids. Nucleotide sequences of the *gnrD*-bearing plasmids were 96% identical. Plasmids contained 3 ORFs apart from *qnrD* and belonged to an undescribed incompatibility group. Only one plasmid, in *P. vulgaris*, was slightly different with a 1,568-bp insertion between qnrD and its promoter, leading to absence of quinolone resistance. We sought for similar plasmids in 15 reference strains of Proteeae, but which were tested negative for qnrD, and found a 48% identical plasmid (pVERM) in Providencia vermicola. In order to explain how qnrD could have been inserted into such native plasmid. we sought for gene mobilization structures. *qnrD* was found to be located within a mobile insertion cassette (mic) element which sequences are similar to one mic also found in pVERM. Our conclusions are that (i) the small non-transmissible qnrDplasmids described here may result from the recombination between an as-yet-unknown progenitor of qnrD and pVERM, (ii) these plasmids are maintained in Proteeae being a *qnrD* reservoir (iii) the mic element may explain *qnrD* mobilization from non-transmissible plasmids to mobilizable or conjugative plasmids from other Enterobacteriaceae, (iv) they can recombined with larger multiresistant plasmids conjugated in Proteeae.

Citation: Guillard T, Grillon A, de Champs C, Cartier C, Madoux J, et al. (2014) Mobile Insertion Cassette Elements Found in Small Non-Transmissible Plasmids in *Proteeae* May Explain *qnrD* Mobilization. PLoS ONE 9(2): e87801. doi:10.1371/journal.pone.0087801

Editor: Ulrike Gertrud Munderloh, University of Minnesota, United States of America

Received August 29, 2013; Accepted December 30, 2013; Published February 4, 2014

Copyright: © 2014 Guillard et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: TG was partially funded by an annual grant from Université de Reims Champagne-Ardenne (EA 4687) and the study was found by an annual grant from Université Paris Diderot (EA 3964). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Emmanuelle.cambau@lrb.aphp.fr

Introduction

Quinolones inhibit replication and transcription by inhibiting the bacterial type II topoisomerases, DNA gyrase and topoisomerase IV [1]. Quinolones attach to the DNA-topoisomerase complex, which becomes irreversible, leading to immobilization of the enzymes resulting in bacteriostasis and to the release of DNA double-strand breaks that activate the SOS system and produce the "poison" effect responsible for the intense bactericidal action of quinolones [2]. Fluoroquinolones (the main subgroup of quinolones) are currently among the most heavily prescribed antimicrobials in the world because of their pharmacodynamic and pharmacokinetic properties [1]. They are very potent, especially for treatment of urinary tract infections due to *Enterobacteriaceae* [1,2], and as a consequence of their intense use, quinolone resistance rate has increased much for the last years [1,3,4]. Quinolone resistance mechanisms are multiple such as those reducing permeability of the bacterial wall, increasing efflux, reducing target affinity, producing inactivating enzymes and target protection proteins. Clinical resistance mostly results from the combination of several mechanisms [1]. Most of these mechanisms are chromosome-mediated [1,2] but plasmid-mediated genes have been described for a decade [5]. *qnr* genes were the first plasmid-mediated quinolone resistance genes described in 1998 [6].

Qnr proteins are pentapeptide repeat proteins that protect DNA gyrase and topoisomerase IV from quinolone binding. Six families of plasmid-mediated *qnr* gene were described: *qnrA*, *qnrB*, *qnrC*, *qnrD* and *qnrS* [7–10]. Several alleles have been reported for each gene: 7 alleles of *qnrA*, 73 of *qnrB*, 1 of *qnrC*, 2 of *qnrD*, 9 of *qnrS* and 6 *qnrVC*. Alleles are numbered consecutively in a *qnr* library (http://lahey.org/qnrStudies, last update December 08, 2013). Most of *qnr* genes were detected in *Enterobacteriaceae*, where they located on large conjugative multi-resistance plasmids, such as pMG252 described in the original *Klebsiella pneumoniae qnrA1*-

positive isolate [6]. The qnrB gene is the most prevalent in clinical bacterial strains, whereas qnrS gene is prevalent in environmental strains. Since qnrA alleles were found also as chromosome-borne in Shewanella spp. [11], this species was proposed to be a qnrA progenitor. Similarly, a *qnrB* allele was found in the chromosome of Citrobacter spp. [12], a qnrS and qnrVC allele in Vibrionaceae [10,13]. qnr genetic environment usually showed mobile elements that can explain their spread from their progenitor, although they confer only a low level resistance to quinolones. Most of qnrA and qmB genes were reported on large conjugative multi-drug resistant plasmids including aminoglycoside and beta-lactam resistance genes such as those encoding extended spectrum beta-lactamases, and in the vicinity of complex type 1 integrons [14,15]. On the contrary, qnrS and qnrD genes were described on small nonconjugative plasmids, that may harbor mob genes allowing their mobilization [16,17].

qnrD was first reported on a 4,270-bp (p2007057) plasmid in four Salmonella enterica isolates [7]. We subsequently described two Providencia rettgeri clinical isolates carrying qnrD on a 2,683-bp plasmid [18]. We hypothesized that *qnrD* can be originated from Proteeae or present in these bacteria as a reservoir. Our study determined how much *qnrD* is prevalent in recent clinical isolates of Proteeae, and investigated how qnrD could have disseminated to other bacterial species. We found seven nearly identical small qnrD-bearing plasmids, where qnrD is surrounded by sequences of mobile insertion cassette (mic) that may explain qnrD dissemination. Since the half part of the plasmid sequence excluding qnrD, was found in a native plasmid of Providencia vermicola, we think that and might be present on the chromosome of a close, as yet unknown, Proteeae species and that its DNA recombination with P. vermicola native plasmid led to the small non-transmissible gnrpositive plasmid found in clinical isolates.

Materials and Methods

qnrD Detection in Proteeae Clinical Isolates

A total of 317 clinical isolates including 190 Proteus mirabilis, 59 Morganella morganii, 33 Proteus vulgaris, 20 Providencia stuartii, 14 Providencia rettgeri and 1 Proteus penneri were screened. They were isolated between 2008 and 2011 at five University hospitals (CHU Dijon, CHU Lariboisière Paris, CHU Nancy, CHU Pitié-Salpêtrière Paris, CHU Reims) from clinical specimens regardless antibiotic susceptibility. Fifteen reference strains of the tribe of Proteeae were purchased at the Collection de l'Institut Pasteur (CIP) (n = 13) and at the Leibniz-Institut DSMZ-German Collection of Microorganisms and Cells Cultures (Braunschweig, Germany) (n = 2) (Table 1).

Detection of *qnr* genes was performed by multiplex real time PCR as previously described [19]. Briefly, after DNA extraction using the NucliSens easyMAG (bioMérieux, Marcy l'Etoile, France), multiplex real-time PCR was carried out using Light-Cycler[®] 480 (LC480) with HRM Supermix (Roche Molecular Diagnostics, Germany). The melt-curve analysis showed characteristic curves for each *qnr* gene family. All strains detected were confirmed to be *qnr*-positives by conventional PCR-sequencing.

The *qnrD*-positive isolates were compared for genome fingerprinting by Random Amplification of Polymorphic DNA (RAPD) using the Ready to Go RAPD analysis kit according to the manufacturer's instructions (Amersham Biosciences).

Characterization of *qnrD*-bearing Plasmids

Plasmid extraction was performed by the Kieser method from the *qnrD*-positive strains [20]. Plasmids harboring *qnrD* were sequenced on both strands using a Walk DS strategy [18]. Nucleotide sequences were aligned using the program BioEdit version 7.09.0 and BLAST searches using the National Center for Biotechnology Information website (http://www.ncbi.nlm.nih. gov) and the blastn algorithms. Putative promoters, ribosome binding site, and transcription start site were searched using BPROM (http://linux1.softberry.com) and Promoter Prediction by Neural Network [21].

Incompatibility groups of the plasmids were determined using the PCR-based replicon typing (PBRT) as described elsewhere [22]. Briefly, four multiplex PCR were used for the detection of A/ C, T, FIIAs, W, N, FIB, L/M, I1-Iγ, X, HI2, FIA, and Y replicons. Replicons P, R, U, F, FIC, and K were detected by simplex PCR, as previously described, and replicons, FII1K, FII2K, NewXXX or also named ZK, LVPK and Amet (Table S1) as described by D. Decré and G. Arlet. The primer design was based on pGSH500 and pKNP4 (AJ009980 and CP000649 respectively), pLVPK (AY378100) and pK245 (DQ449578), and pMET-1 (EU383016) sequences.

Search for *qnrD* and Small Non-transmissible Plasmids in Reference Strains of *Proteeae*

Plasmid extraction was performed by the Kieser method from the 15 reference strains and from 50 *qnrD*-negative strains randomly selected [20]. Based on the nucleotide sequences of the *qnrD*-plasmids characterized in the clinical isolates, six pairs of primers were designed corresponding to overlapping fragments (Table S1). Amplification products were visualized on 2% agar gel after electrophoresis and sequenced.

Investigation of *qnrD* Transfer

Conjugation experiments were carried out using liquid and filter mating methods. Liquid mating experiments were performed in brain-heart infusion (BHI) broth with E. coli I53 azide-resistant [15], P. mirabilis ATCC 29906 Rif^R [23] and E. coli C600 Rif^R [23] as recipient strains. Donor and recipient cells in the logarithmic phase of growth were mixed (1 ml each) and incubated at 37°C for 3 hours without shaking. For filter conjugation, cultures (logarithmic phase of growth) of the donor and the different recipient cells were collected on a MF-type filter (Millipore, Molsheim, France). The hydrophobic edge membrane was incubated at 37°C on the surface of a BHI agar plate overnight. Transconjugants were selected on BHI agar plates containing sodium azide (100 µg/ml) or rifampin (250 μ g/ml), with regard to the recipient strain, and nalidixic acid (50 µg/ml) or ciprofloxacin (0.015 µg/ml, 0.03 µg/ ml, $0.06 \,\mu\text{g/ml}$ and $0.12 \,\mu\text{g/ml}$ or ofloxacin (0.03 $\mu\text{g/ml}$, $0.06 \ \mu g/ml$, $0.12 \ \mu g/ml$, and $0.25 \ \mu g/ml$).

The main capture systems (*intl1*, ISCR1, IS26, IS10, and ISEcp1) that have been associated with plasmid-mediated quinolone resistance determinants were detected using specific PCR (Table S1). In silico analysis was conducted for searching insertion sequence (IS) using IS Finder (http://www-is.biotoul.fr/is.html) and inverted repeats using REPFIND [24].

Quinolone Resistance Testing

Since the *qnrD*-positive clinical isolates may have additional mechanisms conferring quinolone resistance, including chromosomal mutations, we determined the quinolone resistance of both the original *Proteeae* strains and the transformants obtained after transfer of *qnrD*-bearing plasmids into competent *E. coli* DH10B cells (Invitrogen, Cergy-Pontoise, France). Plasmid DNA was extracted from the *qnrD*-positive strains using the QIamp DNA mini Kit (Qiagen, Courtaboeuf, France) and introduced by electroporation (Gene Pulser, Biorad, Marnes la Coquette, Table 1. Reference strains of Proteeae harbored native plasmids but no qnr genes.

Organism	Strain collection designation	Plasmid content ^a	
Proteus mirabilis	CIP 103181, ATCC 29906	No plasmid	
Proteus hauseri	CIP 106868, ATCC 700826	No plasmid	
Proteus myxofaciens	CIP 106872, ATCC 19692	No plasmid	
Proteus penneri	CIP 103030, ATCC 33519	No plasmid	
Proteus vulgaris	CIP 104989, ATCC 29905	No plasmid	
Providencia alcalifaciens	CIP 82.90, ATCC 9886	14 kb (pPALC1, [35])	
Providencia rettgeri	CIP 103182, ATCC 29944	5.5 kb (pPRET1, [35])	
Providencia rustigianii	CIP 103032	150 Kb, 12 Kb	
Providencia stuartii	CIP 104687, ATCC 29914	No plasmid	
Providencia vermicola	CIP 108829, DSM 17385	3.6 kb (pVERM)	
Providencia heimbachae	CIP 103031, ATCC 35613	No plasmid	
Providencia burhodogranariea	DSM 19968, ATCC-BAA1590	-	
Providencia sneebia	DSM 19967, ATCC BAA-1589	10.8 Kb (pPSN1), 7.6 Kb (pPSN2), 4.3 Kb (pPSN3) [35]	
Morganella morganii subsp. morganii	CIP A231, ATCC 25830	No plasmid	
Morganella morganii subsp. sibonii	CIP 103648, ATCC 49948	No plasmid	

^aDetermined by agarose gel electrophoresis of plasmids extracted from the reference strains of *Proteeae* using the Kieser method. Control plasmids for size were pIP55 (130 kb), pIP135 (70.4 kb) and pGHS09-09 (2.6 kb).;

–, not done.

doi:10.1371/journal.pone.0087801.t001

France). Transformants were selected on BHI agar plates containing ciprofloxacin (0.015 μ g/ml, 0.03 μ g/ml, 0.06 μ g/ml and 0.12 μ g/ml) or ofloxacin (0.03 μ g/ml, 0.06 μ g/ml, 0.12 μ g/ml, and 0.25 μ g/ml). In order to distinguish transformants and quinolone resistant mutants growing on quinolone containing media, we sequenced the QRDRs in *gyrA* and *parC*, in addition to *qnrD* detection by PCR.

Minimal inhibitory concentrations (MICs) of nalidixic acid, ciprofloxacin, levofloxacin, moxifloxacin, norfloxacin, and ofloxacin were determined by the agar dilution method using Mueller–Hinton agar plates containing serially twofold-diluted antibiotics. Plates were inoculated with a Steers-type multiprong device with $ca. 10^4$ CFU per spot, and were read after incubation for 18 hours at 37° C.

Results

Small Non-transmissible Plasmids are the Genetic Support of *qnrD* in *Proteeae* Isolates

qnr gene screening of the 317 clinical isolates resulted in eight (2.6%) qnr-positive strains with seven strains (87.5% of qnr-positive isolates) carrying qnrD: four P. mirabilis (DPROT11, DPROT104, DPROT189, and DPROT304), two P. rettgeri (DIJ09-518 and GHS09-09) and one P. vulgaris (DPROT78). The remaining qnr-positive isolate was a P. mirabilis isolate harboring qnrA1 (DPROT289). The seven qnrD genes were 100% identical to the original gene described [7]. On the basis of epidemiological findings (Table 2) and RAPD profiles (data not shown), the isolates belonging to the same species (4 P. mirabilis, 2 P. rettger) were distinct strains.

Conjugation assays, performed extensively, failed in transferring *qnrD* from clinical isolates to any of the recipient strains. From the seven *qnrD*-positive isolates, DNA extraction by the Kieser method showed several plasmids but a similar small plasmid of *ca.* 2.6 Kb in six isolates and one *ca.* 4 Kb in the remaining isolate. *qnrD*-plasmids were negative for known incompatibility groups.

Purified *qnrD*-plasmids were successfully transferred by electroporation into *E. coli* DH10B. Plasmid sequencing showed that six of the seven plasmids were of *ca.* 2.6-Kb long: pDIJ09–518a, pGHS09–09, pRS12–11, pRS12–104, pRS12–189, and pLRB12– 304 being 2,683-bp, 2,683-bp, 2,863-bp, 2,683-bp, 2,656-bp and 2,658-bp long, respectively. They were carried, respectively, by two *P. rettgeri* clinical isolates previously described [18] and 4 clinical strains of *P. mirabilis*. The remaining plasmid (pRS12–78) was 4,236-bp long and was carried by a *P. vulgaris* isolate.

Quinolone MICs are presented in Table 3. Quinolone resistance conferred by the *qnrD*-bearing plasmids was similar for all plasmids, with a 60-fold increase in ciprofloxacin MIC. For the plasmid pRS12–78, transformation experiments on ofloxacin or ciprofloxacin, even at low concentration, selected S83L *gyrA* chromosomal mutants harboring pRS12–78. Consequently we cannot measure the exact level of resistance conferred by the plasmid. However, since the ciprofloxacin MIC was similar to that on a *gyrA* mutant [25], we concluded that the plasmid did not confer quinolone resistance, which is explained by the insertion between the putative promoter and the *qnrD* start codon (see below). No mutation in the QRDR of *gyrA* and *parC* were found in the other transformants.

Nucleotide sequences (GenBank accession numbers: pDIJ09– 518a, HQ834472; pGHS09-09, HQ834473; pRS12-11, KF364953; pRS12–104, KF364955; pRS12–189, KF364956; pLRB12–304, KF364957) of the six *ca.* 2.6-kb plasmids were 96% identical. These plasmids contain four putative open reading frames (ORFs) including *qnrD*, which we arbitrarily named ORF1 to ORF4. Two ORFs were similar to those found in p2007057 found in *S. enterica* [7] : the *qnrD* ORF1 (100% identity) and the ORF2 (97% identity). ORF2 encodes a hypothetical protein with 88% similarity to a hypothetical protein previously described (accession number AF448250). ORF3 and ORF4 are also hypothetical proteins with no significant matches.

Plasmid pRS12-78 (GenBank accession number KF364954) was identical to the 2.6-Kb plasmids but with an additional

Table 2. Epidemiological findings of the 7 qnrD-positive strains detected among 317 clinical isolates of Proteeae.

Strains	Hospital	Ward	Date of isolation	Specimen
P. mirabilis DPROT 11	CHU Reims	Nephrology	11/10/2010	Urine
P. mirabilis DPROT 104	CHU Reims	Diabetology	04/09/2010	Sputum
P. mirabilis DPROT 189	CHU Reims	Nephrology	02/18/2011	Urine
P. mirabilis DPROT 304	CHU Lariboisière-Paris	Long term care	02/02/2010	Urine
P. vulgaris DPROT 78	CHU Reims	Long-term care	12/12/2010	Urine
P. rettgeri DIJ09-518	CHU Dijon	Hematology	12/08/2008	Stool
P. rettgeri GHS09-09	CHU Pitié-Salpêtrière-Paris	Emergency room	10/09/2009	Urine

doi:10.1371/journal.pone.0087801.t002

1,568 bp insert between *qnrD* and its promoter (Figure 1A). BLAST comparison highlighted a putative ORF5 exhibiting 76% identity with a hypothetical protein (GenBank accession number

Table 3. Resistance conferred by *qnrD*-bearing plasmids as measured by determination of minimal inhibitory concentrations (MIC) of quinolones on transformants with regard to parental strains of *Proteeae*.

Strains	MIC (µg/ml)					
	NAL	NOR	OFX	CIP	MXF	LVX
P. mirabilis DPROT11	>256	12	>32	>32	>32	16
P. mirabilis DPROT104	8	0.5	1	0.25	1	0.25
P. mirabilis DPROT189	>256	12	>32	>32	>32	16
P. mirabilis DPROT304	>256	8	>32	>32	>32	16
P. rettgeri GHS09-09	4	0.25	0.5	0.125	0.5	0.25
P. rettgeri DIJ09–518	>256	8	8	4	12	4
P. vulgaris DPROT78	>256	8	8	4	>32	4
E. coli DH10B	1	0.03	0.015	0.006	0.03	0.006
<i>E. coli</i> DH10B/ pRS12-11	4	0.5	0.25	0.125	0.125	0.125
<i>E. coli</i> DH10B/ pRS12–104	4	0.5	0.25	0.125	0.125	0.125
<i>E. coli</i> DH10B/ pRS12–189	2	0.5	0.25	0.125	0.125	0.125
<i>E. coli</i> DH10B/ pLRB12–304	4	0.5	0.25	0.125	0.125	0.125
<i>E. coli</i> DH10B/ pGHS09-09	2	0.5	0.25	0.125	0.125	0.125
<i>E. coli</i> DH10B/ pDIJ09–518a	2	0.5	0.25	0.125	0.125	0.125
<i>E. coli</i> DH10B/ pRS78	16	0.125	0.125	0.03	0.06	0.006
<i>E. coli</i> DH10B/ p2007057ª	2	0.125	0.25	0.125	ND	ND

NAL, Nalidixic acid; NOR, Norfloxacin; CIP, Ciprofloxacin; OFX, Ofloxacin; LVX, Levenfloxacin; MXF, Moxifloxacin; ND, no data.

^aMICs reported by Cavaco et al. [7].

doi:10.1371/journal.pone.0087801.t003

ADR29581) close to the ATPase domain of the *E. coli* ParA protein. Moreover, at 147 nucleotides downstream the stop codon of ORF5, the presence of inverted repeats followed by a short string of A's suggested a factor-independent termination site.

Interestingly, among the randomly selected *qnrD*-negative strains for plasmid characterization using Kieser extraction, none of them harbored small plasmids (data not shown).

qnrD is Included in a Mobile Insertion Cassette

In silico analysis did not show any gene involved in mobilization and specific PCRs for intl1, ISCR1, IS26 and ISEcp1 were negative. We also looked for mobile insertion cassette (mic) elements that are similar to transposons. A mic element is bracketed by two inverted repeats without any transposaseencoding gene found nearby and duplication of the target site is usually observed indicating the transposition. In the 2,683-bp long plasmids, qnrD was localized within a 1,519-bp "mic" element, bracketed by two 24-bp imperfect inverted repeats (IR) with six mismatches (Figure 1B). No transposase-encoding gene was found. Acquisition of the mic-qnrD element was likely the result of a transposition process because it is bracketed by a 2-bp duplication of the target site (CA). The mic-qnrD element described here carries the necessary elements for expression of the qnrD gene: (i) promoter sequences including a -35 box (TTTTATA) and a -10 box (TCTTAAAAT) separated by 11-bps, (ii) and a Shine-Dalgarno sequence (AAAAGGG) upstream on the qnrD initiation codon (ATG).

In the 4,686-bp long plasmid pRS12–78, *qnrD* is a part of a 3,072-bp "mic" element given that the ORF5 described above was inserted between *qnrD* and a putative promoter (Figure 1A). Although this second mic-*qnrD* element is slightly different in length, it has the same characteristics.

Reference Strains of *Proteeae* Carried Native Plasmids but no *qnrD* Gene

qnrD was not detected by multiplex or simplex real-time PCR in the fifteen reference strains of *Proteeae* but plasmids were found in most of the reference strains (Table 1). These plasmids did not belong to any known incompatibility group detected using the PBRT method [22].

Overlapping PCRs designed on the sequence of pDIJ09–518a and performed using template DNA of the *Proteeae* reference strains showed that one small plasmid of *P. vermicola*, pVERM, carried a fragment exhibiting 89% identity with the sequence of pDIJ09–518a. pVERM was linearized using *AclI* restriction enzyme. After gel extraction, the sequence was determined using a walk DS strategy on the amplification product obtained with primers pDIJ-2558F and orf2–3'R. pVERM is a 3,682-bp long



Figure 1. *qnrD* is located within a mobile insertion cassette (mic) element. A. Sequence analysis of the mic element including *qnrD* from the sequence of the plasmid pRS12–78, which contains a 1,603 bp insert (ORF5) upstream from *qnrD*. The nucleotide numbering is shown on the left-hand side. Putative -35, -10 promoter regions as well as the ribosome binding site (RBS) and the putative transcription start sites (+1) of *qnrD* are in bold. The start and the stop codons of *qnrD* (in blue) and of the ORF5 (in magenta) are underlined. For the termination site of ORF5, two black arrows indicate inverted repeats, and the string of "A's" is boxed. The inverted repeats (IRL and IRR) bracketing the mic element are boxed as arrows shaded in grey. In pRS12–78, *qnrD* is far from its own promoter due to the insertion of ORF5. B. Comparison of the mic elements found in the plasmid, pDIJ09–518, pRS12–78, pVERM and in p2007057. Open reading frames (ORFs) are indicated by horizontal arrows. White arrows are identical ORFs harbored by the different plasmids beyond the mic. Red arrows are identical ORFs harbored by the different plasmid subtin the mic. The right and left inverted repeats (IRR and IRL) are indicated as black triangles with the duplication sites (CA). Nucleotide sequences of the IRL and IRR are outlined and the nucleotides which differed between them are in bold type and underlined. Asterisks show nucleotide differences of IRs with regard to those in pDIJ09–518a.

doi:10.1371/journal.pone.0087801.g001

plasmid containing a 1,794-bp region sharing more than 98% identity with the qnrD-plasmids. In this region, we identified three ORFs similar to ORF2, ORF3 and ORF4 described in pDIJ09-518a. BLAST of the 1,888-bp remaining region showed a 267-bp ORF1 (named pVERM_ORF1 in the following) covering 76% of a putative protein described in Vibrio brasiliensis (accession number ZP_080099857.1) with 31% similarities. Two 24-bp long imperfect IR, with eight mismatches, and CA duplications were found to bracket pVERM_ORF1 and pVERM_ORF2 leading to identify another mic element. The 2,664-bp long mic element harbored by pVERM is different from mic-qnrD for 1,145 bps containing pVERM_ORF1 and lacking qnrD (Figure 1B). Nonetheless, IRR and IRL are similar to those bracketing mic-qnrD on the qnrD plasmids such as pDIJ09-518a with 2 and 1 single nucleotide differences for IRLs and IRRs respectively (Figure 1B). No known mobilization structure such as IS were identified in this 2,664-bp long mic element.

Discussion

We investigated the emergence of the new plasmid-borne quinolone resistance gene, qnrD in Proteeae clinical bacterial isolates. The qnr genes are encoding a group of pentapeptide repeat proteins recently discovered but certainly present on bacterial chromosomes long before quinolones were used clinically. They emerged 10 years ago, following the 1990-2010 worldwide increase in fluoroquinolone usage [3]. qnrD was initially described in Salmonella enterica Kentucky and Bovismorbificans isolates [7], our investigation showed that qnrD was the qnr gene mainly present in isolates belonging to the Proteeae tribe (Proteus, Providencia and Morganella genera) since the screening of all qnr genes in 317 Proteeae clinical isolates resulted in 8 qnr-positive strains but seven (87.5%) positive for qnrD. This unexpected result converges with data in GenBank where among 38 sequences of qnrD half were reported in Proteeae (Table S2, last update October 30, 2013). We consequently hypothesized that Proteeae are the reservoir of qnrD genes.



Figure 2. *qnrD* **may have emerged and disseminated from** *Proteeae* **to** *Enterobacteriaceae* **using mic elements and a recombination event with the native PVERM plasmid.** Comparison of the plasmid structures of pDIJ09-518a from *P. rettgeri*, pVERM from *P. vermicola*, p2007057 from *S. enterica* Bovismorbificans, pIGC156 from *E. coli* and pSC101 from *S. enterica* Typhimurium. The open reading frames are shown as arrows with the arrowhead indicating the direction of transcription. Red color is used to shade arrows in the *qnrD*-mic element. Grey shading shows the areas of homology between the plasmids.

doi:10.1371/journal.pone.0087801.g002

In Proteeae clinical isolates, we found qnrD on small ca. 2.6-kb non-transmissible plasmids, in contrast with what was mainly described for other qnr genes carried by large conjugative plasmids [5]. Although qnrD was also reported to on such large conjugative plasmids, none of these plasmids was well characterized [26-29]. All the 15 qnrD-bearing plasmids found in Proteeae (the 7 from our study and 8 plasmids from GenBank sequences, Table S2) have the same size (2,683-bp) and have nearly identical sequences, which is in favor of a common origin or ancestor [30–32]. The only exception found during our study is the plasmid pRS12-78 that presented a plasmid scaffold identical to the other characterized plasmids but with a 1,568-bp insertion corresponding to a putative additional ORF. By contrast, the plasmids bearing qnrD found in strains of Enterobacteriaceae other than Proteeae are 4,270 bp long with sequences nearly identical to p2007057 described in S. enterica. These latter plasmids usually harbor a mob gene that may explain their transfer. The two types of plasmids carrying qnrD show a highly conserved backbone, but the smallest are not transmissible. This was previously described for qnrS, as it was described on a 10,066-bp non-conjugative but mobilizable plasmid in S. enterica Typhimurium [16]. It was suggested that this plasmid results from the recombination of a gnrS1 conjugative plasmid from S. enterica Infantis (pINF5) and a small mobilizable plasmid from E. coli (pEC278). The qnrD-2,683-bp plasmids were stable in E. coli DH10B after electroporation, indicating a host range including other species of Enterobacteriaceae. However, the gene encoding the replicase is probably different to what is usually found since the incompatibility group cannot be assessed. Non classic replication systems were recently reported for some small plasmids in Proteeae [18,33] and we share the view of Galac et al. that cryptic plasmids in *Proteeae*, including the small *qnrD*-plasmids, may use a not yet characterized replication method [33]. Since reference type strains of *Proteeae* did not harbor *qnrD*, we sought for similar plasmid backbone in their constitutive DNA. We found, in a reference strain of *P. vermicola*, a native plasmid that shares 1,794 bps with more than 98% identity with our *qnrD*-bearing plasmids.

We described here *qnrD* as located in a mobile insertion cassette (mic) element. Mic elements were first described in Bacillus cereus (MIC231A1) as a DNA cassette able to be mobilized in *trans* by the IS231A transposase [34]. MIC231A1 contains an active Dstereospecific endopeptidase (adp) gene instead of a transposase. Further studies conducted on B. cereus showed the presence of antibiotic resistance determinant such as a fosfomycin resistance gene in MIC231D, described as a novel active composite cassette [35]. It was then proposed mic element could be also a vehicle for antibiotic resistance genes beside class 1 integrons where multiple antibiotic cassettes can be inserted. With regard to plasmid mediated quinolone resistance genes, qnrS2 was described in a mic element in a strain of Aeromonas punctata isolated in water sampled from the River Seine in Paris [36]. Unfortunately, as reported by Cattoir et al., no transposases were found, on the basis of the IR nucleotide sequences, suitable for demonstrating trans-transposition in an in vitro assay [36]. qnrA genes were mainly associated with ISCR1 at the vicinity of complex class 1 integrons which include various other resistance determinants (e.g. bla genes or aac(6')-Ib-cr). qnrB genes were also described beside class 1 integrons, but associated with ISEcp1 and ORF1005, another type of inserted sequences [37,38]. Unlike qnrA and qnrB, qnrS genes were mainly associated with structure such as ISEcl2, IS26 and mic structure [16,36,39], and were not reported as part of complex integrons. These gene-capture systems explain the mobilization of qnr genes from their progenitors to self-transmissible plasmids leading to their widespread in Enterobacteriaceae.

We think that *qnrD* was imported from a *Proteeae* species and inserted in pVERM through mic mobilization (Figure 2). Further

recombination, insertion or deletion events may have occurred after the mic-qnrD mobilization including qnrD deletion, resulting in the plasmids pVERM and pDIJ09-518a reported here. Indeed, the IRs found in the mic-qnrD plasmids from Proteeae clinical strains are more similar to those found in pVERM than those found in plasmids in S. enterica. pVERM differed from qnrD-plasmids by 1,888 bps located inside the very same lacking qnrD mic. This qnrD-lacking-mic element found in pVERM is 1,145 bp longer than mic-qnrD and carried a pVERM_ORF1 present neither on pDIJ09-518 nor on p2007057. While mic-qnrD IRR and IRL showed seven unmatched nucleotides for pVERM vs six for mic*anrD*, more importantly a comparison of IRLs between mic-*anrD* and pVERM found only 2 SNPs and only 1 SNPs was found comparing both IRRs. Finally, IRs in both gnrD-plasmid and pVERM were 24-bp long. (Figure 1A). Moreover, none of the 50 *anr*-negative clinical strains, randomly selected, harbored such small plasmids. Therefore, our current hypothesis is that pVERM could be the ancestor plasmid where *qnrD* was later deleted. Alternatively, pVERM could be a *qnrD*-plasmid resulting from another mic-qnrD harbored by an unknown qnrD progenitor, from which qnrD was subsequently lost.

The qnrD progenitor is not yet identified. For the other qnr genes, progenitors are mainly bacteria living in the environment (Shewanella spp., Citrobacter spp., and Vibrio spp) where Proteeae are also present [13,36,40,41]. The Proteeae could have acquired qnrD from its progenitor by recombination between an as-yet-unknown bacterial source of qmD and the cryptic plasmid pVERM harbored by P. vermicola leading to small non-mobilizable 2,683-bp long plasmids carrying *qnrD* embedded in a mic-*qnrD* element (Figure 2). Given that all plasmids of about 4,270 bps such as p2007057 (e.g. P. mirabilis isolates CGP180 and CGH40 described by Zhang et al.) also harbored a mic-qnrD with a conserved 24-bp IRL and a modified 26-bp IRR showing 1 and 7 nucleotide differences, respectively, with regard to 2,683-bp qnrD-plasmids, we think that the mic-qnrD element could have been later inserted by trans transposition in a mobilizable plasmid such as the 4,270-bp qnrDplasmid p2007057. It is also possible that the mic-qnrD was transferred from a common progenitor to both Proteeae and Salmonella spp. in parallel onto two different types of plasmids in

References

- Hooper D (2000) Mechanisms of action and resistance of older and newer fluoroquinolones. Clin Infect Dis. Suppl 2: S24–8.
- Drlica K, Malik M, Kerns RJ, Zhao X (2008) Quinolone-Mediated Bacterial Death. Antimicrob Agents Chemother 52: 385–392.
- Goossens H (2009) Antibiotic consumption and link to resistance. Clin Microbiol Infect 15 Suppl 3: 12–15.
- Garau J, Xercavins M, Rodríguez-Carballeira M, Gómez-Vera JR, Coll I, et al. (1999) Emergence and dissemination of quinolone-resistant *Escherichia coli* in the community. Antimicrob Agents Chemother 43: 2736–2741.
- Strahilevitz J, Jacoby GA, Hooper DC, Robicsek A (2009) Plasmid-mediated quinolone resistance: a multifaceted threat. Clin Microbiol Rev 22: 664–689.
- Martínez-Martínez L, Pascual A, Jacoby GA (1998) Quinolone resistance from a transferable plasmid. Lancet 351: 797–799.
- Jacoby G, Cattoir V, Hooper D, Martínez-Martínez L, Nordmann P, et al. (2008) qnr Gene nomenclature. Antimicrob Agents Chemother 52: 2297–2299.
- Wang M, Guo Q, Xu X, Wang X, Ye X, et al. (2009) New plasmid-mediated quinolone resistance gene, qmrC, found in a clinical isolate of Proteus mirabilis. Antimicrob Agents Chemother 53: 1892–1897.
- Cavaco LM, Hasman H, Xia S, Aarestrup FM (2009) *qnrD*, a novel gene conferring transferable quinolone resistance in *Salmonella enterica* serovar Kentucky and Bovismorbificans strains of human origin. Antimicrob Agents Chemother 53: 603–608.
- Fonseca EL, Vicente ACP (2013) Epidemiology of *qnrVC* alleles and emergence out of the *Vibrionaceae* family. J Med Microbiol 62: 1628–1630.
- Poirel L, Rodriguez-Martinez J-M, Mammeri H, Liard A, Nordmann P (2005) Origin of plasmid-mediated quinolone resistance determinant QnrA. Antimicrob Agents Chemother 49: 3523–3525.
- Jacoby GA, Griffin CM, Hooper DC (2011) Citrobacter spp. as a Source of qnrB Alleles. Antimicrob Agents Chemother 55: 4979–4984.

which the IRs sequences evolved and diverged especially for the IRR.

Finally, we investigate the quinolone resistance conferred by the *qnrD*-bearing plasmids. The MICs of fluoroquinolones were similar and comparable to MICs previously reported for p2007057, except for the plasmid pRS12–78. In this plasmid, the insertion of a 1,568-bp long sequence, can hamper *qnrD* expression from a putative promoter then located far from *qnrD*. Although the level of resistance is rather low for classical quinolones such as nalidixic acid, it can reach 60-fold increase in the MIC of ciprofloxacin. Such an increase can cause failure in experimental model of pyelonephritis and in pulmonary infections [42–44]. Further studies are needed to investigate the fitness cost of *qnrD* harbored by these small plasmids, but as described for *qmA3* [45], *qnrD* could contribute to a fitness gain leading to the emergence of *qnrD*-bearing plasmids even in the absence of quinolones.

Supporting Information

Table S1List of primers used in this study.(DOC)

Table S2 List of *qnrD* deposited in GenBank.

 (DOC)

Acknowledgments

We are grateful to Prof G. Arlet and Dr D. Decré for providing primers sequences in order to characterize incompatibility groups NewXXX, FII1K, FII2K, U, R, LVPK and Amet.

Part of this work was presented at the 53th Interscience Conference on Antimicrobial Agents and Chemotherapy, Denver, U.S.A., 2013.

Author Contributions

Conceived and designed the experiments: TG CdC EC. Performed the experiments: TG AG CC JM. Analyzed the data: TG CdC EC. Contributed reagents/materials/analysis tools: TG EC CdC VVG BB AL JR ALL. Wrote the paper: TG EC.

- Cattoir V, Poirel L, Mazel D, Soussy C-J, Nordmann P (2007) Vibrio splendidus as the source of plasmid-mediated QnrS-like quinolone resistance determinants. Antimicrob Agents Chemother 51: 2650–2651.
- Cambau E, Lascols C, Sougakoff W, Bébéar C, Bonnet R, et al. (2006) Occurrence of *qnrA*-positive clinical isolates in French teaching hospitals during 2002–2005. Clin Microbiol Infect 12: 1013–1020.
- Lascols C, Podglajen I, Verdet C, Gautier V, Gutmann L, et al. (2008) A plasmid-borne Shewanella algae Gene, *qnrA3*, and its possible transfer in vivo between *Kluyvera ascorbata* and *Klebsiella pneumoniae*. J Bacteriol 190: 5217–5223.
- Kehrenberg C, Hopkins KL, Threlfall EJ, Schwarz S (2007) Complete nucleotide sequence of a small *qnrS1*-carrying plasmid from *Salmonella enterica* subsp. enterica Typhimurium DT193. J Antimicrob Chemother 60: 903–905.
- Gay K, Robicsek A, Strahilevitz J, Park CH, Jacoby G, et al. (2006) Plasmidmediated quinolone resistance in non-Typhi serotypes of *Salmonella enterica*. Clin Infect Dis 43: 297–304.
- Guillard T, Cambau E, Neuwirth C, Nenninger T, Mbadi A, et al. (2012) Description of a 2,683-base-pair plasmid containing *qnrD* in two *Providencia rettgeri* isolates. Antimicrob Agents Chemother 56: 565–568.
- Guillard T, Moret H, Brasme L, Carlier A, Vernet-Garnier V, et al. (2011) Rapid detection of *qnr* and *qepA* plasmid-mediated quinolone resistance genes using real-time PCR. Diagn Microbiol Infect Dis 70: 253–259.
- Kieser T (1984) Factors affecting the isolation of CCC DNA from Streptomyces lividans and Escherichia coli. Plasmid 12: 19–36.
- Reese MG (2001) Application of a time-delay neural network to promoter annotation in the Drosophila melanogaster genome. Comput Chem 26: 51–56.
- Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, et al. (2005) Identification of plasmids by PCR-based replicon typing. J Microbiol Methods 63: 219–228.

- De Champs C, Monne C, Bonnet R, Sougakoff W, Sirot D, et al. (2001) New TEM variant (TEM-92) produced by *Proteus mirabilis* and *Providencia stuartii* isolates. Antimicrob Agents Chemother 45: 1278–1280.
- Betley JN, Frith MC, Graber JH, Choo S, Deshler JO (2002) A ubiquitous and conserved signal for RNA localization in chordates. Curr Biol 12: 1756–1761.
- Cesaro A, Roth Dit Bettoni R, Lascols C, Mérens A, Soussy CJ, et al. (2008) Low selection of topoisomerase mutants from strains of *Escherichia coli* harbouring plasmid-borne *qnr* genes. J Antimicrob Chemother 61: 1007–1015.
- Zhao J, Chen Z, Chen S, Deng Y, Liu Y, et al. (2010) Prevalence and dissemination of *oqxAB* in *Escherichia coli* isolates from animals, farmworkers, and the environment. Antimicrob Agents Chemother 54: 4219–4224.
- Ogbolu DO, Daini OA, Ogunledun A, Alli AO, Webber MA (2011) High levels of multidrug resistance in clinical isolates of Gram-negative pathogens from Nigeria. Int J Antimicrob Agents 37: 62–66.
- 28. Li J, Wang T, Shao B, Shen J, Wang S, et al. (2012) Plasmid-Mediated Quinolone Resistance Genes and Antibiotic Residues in Wastewater and Soil Adjacent to Swine Feedlots: Potential Transfer to Agricultural Lands. Environ Health Perspect.
- Zhao J-Y, Dang H (2012) Coastal Seawater Bacteria Harbor a Large Reservoir of Plasmid-Mediated Quinolone Resistance Determinants in Jiaozhou Bay, China. Microb Ecol.
- Kristiansson E, Fick J, Janzon A, Grabic R, Rutgersson C, et al. (2011) Pyrosequencing of Antibiotic-Contaminated River Sediments Reveals High Levels of Resistance and Gene Transfer Elements. PLoS ONE 6: e17038.
- Mazzariol A, Kocsis B, Koncan R, Kocsis E, Lanzafame P, et al. (2012) Description and plasmid characterization of qnrD determinants in Proteus mirabilis and Morganella morganii. Clin Microbiol Infect 18: E46–E48.
- Zhang S, Sun J, Liao X-P, Hu Q-J, Liu B-T, et al. (2013) Prevalence and plasmid characterization of the *qnrD* determinant in *Enterobacteriaceae* isolated from animals, retail meat products, and humans. Microbial Drug Resistance 19: 331–335.
- Galac MR, Lazzaro BP (2011) Comparative pathology of bacteria in the genus Providencia to a natural host, *Drosophila melanogaster*. Microbes and Infection 13: 673–683.
- Chen Y, Braathen P, Léonard C, Mahillon J (1999) MIC231, a naturally occurring mobile insertion cassette from *Bacillus cereus*. Mol Microbiol 32: 657– 668.

- De Palmenaer D, Vermeiren C, Mahillon J (2004) IS231-MIC231 elements from *Bacillus cereus* sensu lato are modular. Mol Microbiol 53: 457–467.
- Cattoir V, Poirel L, Aubert C, Soussy C-J, Nordmann P (2008) Unexpected occurrence of plasmid-mediated quinolone resistance determinants in environmental *Aeromonas* spp. Emerging Infect Dis 14: 231–237.
- Cattoir V, Nordmann P, Silva-Sanchez J, Espinal P, Poirel L (2008) ISEcp1mediated transposition of *qnrB*-like gene in *Escherichia coli*. Antimicrob Agents Chemother 52: 2929–2932.
- Jacoby GA, Walsh KE, Mills DM, Walker VJ, Oh H, et al. (2006) *qurB*, another plasmid-mediated gene for quinolone resistance. Antimicrob Agents Chemother 50: 1178–1182.
- Poirel L, Cattoir V, Soares A, Soussy C-J, Nordmann P (2007) Novel Ambler class A beta-lactamase LAP-1 and its association with the plasmid-mediated quinolone resistance determinant QnrS1. Antimicrob Agents Chemother 51: 631–637.
- Poirel L, Liard A, Rodriguez-Martinez J-M, Nordmann P (2005) Vibrionaceae as a possible source of Qnr-like quinolone resistance determinants. J Antimicrob Chemother 56: 1118–1121.
- 41. Saga T, Kaku M, Onodera Y, Yamachika S, Sato K, et al. (2005) Vibrio parahaemolyticus chromosomal qnr homologue VPA0095: demonstration by transformation with a mutated gene of its potential to reduce quinolone susceptibility in Escherichia coli. Antimicrob Agents Chemother 49: 2144–2145.
- 42. Allou N, Cambau E, Massias L, Chau F, Fantin B (2009) Impact of low-level resistance to fluoroquinolones due to *qnrA1* and *qnrS1* genes or a gyrA mutation on ciprofloxacin bactericidal activity in a murine model of *Escherichia coli* urinary tract infection. Antimicrob Agents Chemother 53: 4292–4297.
- Rodríguez-Martínez JM, Pichardo C, García I, Pachón-Ibañez ME, Docobo-Pérez F, et al. (2008) Activity of ciprofloxacin and levofloxacin in experimental pneumonia caused by *Klebsiella pneumoniae* deficient in porins, expressing active efflux and producing QnrA1. Clin Microbiol Infect 14: 691-697.
 Jakobsen I, Cattoir V, Jensen KS, Hammerum AM, Nordmann P, et al. (2012)
- 44. Jakobsen L, Cattoir V, Jensen KS, Hammerum AM, Nordmann P, et al. (2012) Impact of low-level fluoroquinolone resistance genes qtrA1, qtrB19 and qtrS1 on ciprofloxacin treatment of isogenic Escherichia coli strains in a murine urinary tract infection model. J Antimicrob Chemother. 67: 2438–44.
- Michon A, Allou N, Chau F, Podglajen I, Fantin B, et al. (2011) Plasmidic qurA3 Enhances Escherichia coli Fitness in Absence of Antibiotic Exposure. PLoS ONE 6: e24552.