



Chemical reproductive traits of diploid *Bombus terrestris* males: Consequences on bumblebee conservation

Thomas Lecocq, Maxence Gérard, Kevin Maebe, Nicolas Brasero, Lauren Dehon, Guy Smagghe, Irena Valterová, Thibaut de Meulemeester, Pierre Rasmont, Denis Michez

► To cite this version:

Thomas Lecocq, Maxence Gérard, Kevin Maebe, Nicolas Brasero, Lauren Dehon, et al.. Chemical reproductive traits of diploid *Bombus terrestris* males: Consequences on bumblebee conservation. *Insect Science*, 2017, 24 (4), pp.623 - 630. 10.1111/1744-7917.12332 . hal-01575608

HAL Id: hal-01575608

<https://hal.univ-lorraine.fr/hal-01575608>

Submitted on 4 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chemical reproductive traits of diploid *Bombus terrestris* males: consequences on bumblebee conservation

Thomas Lecocq¹, Maxence Gérard¹, Kevin Maebe², Nicolas Braserio¹, Lauren Dehon¹, Guy Smagghe², Irena Valterová³, Thibaut De Meulemeester⁴, Pierre Rasmont¹, Denis Michez¹

¹Laboratory of Zoology, Research institute of Biosciences, University of Mons, Place du Parc 20, 7000 Mons, Belgium

²Department of Crop Protection, Faculty of Bioscience Engineering, Ghent University, Coupure links 653, B-9000 Ghent, Belgium

³Academy of Sciences of the Czech Republic, Institute of Organic Chemistry and Biochemistry, Flamingovo nám 2, CZ-166 10 Prague, Czech Republic

⁴Naturalis Biodiversity Center, Darwinweg 2, Postbus 9517, 2300 RA Leiden, The Netherlands

Correspondence: Thomas Lecocq, Laboratory of Zoology, Research institute of Biosciences, University of Mons, B-7000, Mons, Belgium.

Tel: +3265373405. E-mail: thomas.lecocq@umons.ac.be

Running title: Diploid male chemical reproductive traits

Abstract The current bumblebee decline leads to inbreeding in populations that fosters a loss of allelic diversity and diploid male production. As diploid males are viable and their offspring are sterile, bumblebee populations can quickly fall in a vortex of extinction. In this paper, we investigate for the first time a potential pre-mating mechanism through a major chemical reproductive trait (male cephalic labial gland secretions) that could prevent monandrous virgin queens from mating with diploid males. We focus our study on the cephalic labial gland secretions of diploid and haploid males of *Bombus terrestris* (L.). Contrary to initial expectations, our results do not show any significant differentiation of cephalic labial gland secretions between diploid and haploid specimens. Queens seem therefore to be unable to avoid mating with diploid males based on their compositions of cephalic labial gland secretions. This suggests that the vortex of extinction of diploid males could not be stopped through pre-mating avoidance based on the cephalic labial gland secretions but other mechanisms could avoid mating between diploid males and queens.

Key Words

Bee decline; Bumblebees; Conservation; Diploid males; Pre-mating recognition; Reproductive traits.

Introduction

About 90% of the world's plant species are pollinated by animals, and mainly bees (Ollerton *et al.*, 2011). This makes bees a critically important functional group for ecosystem services in wild environments as well as in

crops (Free, 1993; Klein *et al.*, 2007). Among bees, females of the social genus *Bombus* (i.e. bumblebees) are among the most important pollinators of the northern temperate zone (Williams, 1985). However, some bumblebee species, as well as other wild bee species, are obviously experiencing a strong decline in their populations (Carvalho *et al.*, 2013; Rasmont *et al.*, 2015). Several factors were proposed to explain this decline: landscape modifications/fragmentations, intensive use of agrochemicals, pathogen infections, competition with alien species, climate change, and interactions between all of these factors (review in Winfree 2010; Potts *et al.* 2011). Ultimate consequences are a decrease of population size and an increase of population fragmentation that leads to a loss of genetic diversity through inbreeding and genetic drift (Zayed, 2009; Frankham *et al.*, 2010).

Loss of genetic diversity is particularly problematic in haplo-diploid species such as bees (Zayed & Packer, 2005; Zayed, 2009). Like most Hymenoptera, bees have a single-locus complementary sex determiner (sl-CSD): males are hemizygote while females are heterozygote for this locus (Crozier, 1977; Heimpel & de Boer, 2008). Inbreeding in haplo-diploid species with sl-CSD brings a heavy load through the production of homozygotes which show male phenotype (Duchateau *et al.*, 1994; Duchateau & Marien, 1995). Diploid male larvae can be recognized and killed by workers in social bee species like *Apis mellifera*, (Woyke, 1963) but they can develop into viable adults in bumblebee species (e.g. *Bombus terrestris* (L.); Duchateau *et al.* 1994). The production of diploid bumblebee males can be substantial: half of the diploid eggs, which normally develop into females, result in males after brother/sister mating (Duchateau *et al.*, 1994; Duchateau & Marien, 1995). These diploid males decrease the fitness of the local populations (Whitehorn *et al.*, 2009). First, the production of diploid males in place of workers implies that the number of the workers decreases up to 50% resulting in a small colony size and a low production of new queens (Duchateau *et al.*, 1994; Duchateau & Marien, 1995). Second, as diploid males produce diploid sperm (Ayabe *et al.*, 2004; Darvill *et al.*, 2012), the mating between monandrous virgin queens and diploid males produces 100% of triploid (i.e. sterile) offspring (Payne *et al.*, 2003; Takahashi *et al.*, 2008). Therefore, this process leads to a vortex of extinction: populations decreasing based on extrinsic factors (e.g. habitat fragmentation) show higher inbreeding, lower allelic diversity and production of diploid males, this production leading to an additional decrease of the population (Zayed & Packer, 2005). Consequently populations can be driven to quick extinction (Zayed & Packer, 2005) as currently observed in small and fragmented bumblebee populations (Darvill *et al.*, 2006; Takahashi *et al.*, 2008).

This vortex of extinction could be reduced if females are able to recognize and avoid mating with diploid males. In monandrous species, like most bumblebee species, the female choice requires pre-mating indicators of male

genetic equipment (Herzner *et al.*, 2006). Therefore, pre-mating mechanisms of avoidance of diploid males could be selected in monandrous bumblebee queens in view of the deleterious consequences of mating with diploid males. First bioassays show that *Bombus* diploid males have the same mating success as haploid males in experimental conditions (Duchateau & Marien, 1995). However, such experiments could lead to unnatural mating because queens have no choice between different sexual partners. For example, sibling individuals mate in experimental condition even if *B. terrestris* queens seem to be less willing to mate with their siblings than with an unrelated partner when they have the choice (Whitehorn *et al.*, 2009). Therefore, we expect that pre-mating cues produced by males could allow female pre-mating discrimination to avoid diploid males in natural conditions. As diploid males show differences in body size and physiology (Duchateau & Marien, 1995; Gerloff *et al.*, 2003), we hypothesize that they could display original chemical reproductive traits.

The present study investigates the cephalic labial gland secretions (CLGS) of haploid *versus* diploid males of *B. terrestris*. Courtship signals of bumblebees include both behavioral and chemical features (see Baer 2003 for a review). Among the features, the CLGS act as a key trait for the pre-mating recognition (Bergström *et al.*, 1981; Baer, 2003; Ayasse & Jarau, 2014): Most bumblebee males patrol along paths where they scent-mark objects with their CLGS that attract conspecific virgin females (see experimental demonstration that CLGS of bumblebee males attract virgin queens in Coppée *et al.*, 2011; Lecocq *et al.*, 2015b). This chemical reproductive trait consists of a species-specific complex mixture of (mainly aliphatic) compounds synthesized *de novo* (Calam, 1969; Lecocq *et al.*, 2011, 2013) with a interindividual variation (Coppée *et al.*, 2008; Žáček *et al.*, 2009; Lecocq *et al.*, 2013) which leads to differential female preferences for different kinds of males (e.g. young *versus* old males, sibling *versus* unrelated males; (Coppée, 2010; Coppée *et al.*, 2011; Lecocq *et al.*, 2015c). Therefore variations in this trait could allow the virgin queen to detect diploids and avoid mating with them.

Material and Methods

Collection and Rearing of Specimens

In order to produce diploid males (diploid male samples were also used in Gérard *et al.*, 2015), we obtained three colonies of *Bombus terrestris terrestris* (L.) from bee breeder Biobest N.V. We were reared these colonies at the laboratory of Zoology in Mons (UMons, Belgium) in a controlled climate room (see rearing protocol in Lhomme *et al.*, 2012, 2013; Gérard *et al.*, 2015): we obtained queens (n=11) and males (n=21) among the first generation from only one colony. We forced brother-sister mating by placing the newly emerged sibling queens and males in a flight cage exposed to natural light (see mating protocol in Lhomme *et al.*, 2013; Gérard *et al.*, 2015). We

used these mated queens to produce a second generation of inbred colonies. These queens overwintered during two months prior to a activation by CO₂ narcosis (see protocol in Lhomme *et al.*, 2013). Then, we placed the queens in small boxes with two workers and a cocoon to (i.e. to maximize the probability of colony initiation see protocol in Lhomme *et al.*, 2012, 2013). All 11 mated queens of the first generation initiated a colony, producing a total of 62 workers and 11 males of second generation (Table S1).

In order to produce haploid males, we reared eight queenless micro-colonies with five workers from one colony from bee breeder Biobest N.V. Each micro-colony was maintained in a plastic box in a controlled climate room (same rearing conditions than for colonies). The micro-colonies produced 41 haploid males (Table S1).

We increased the diversity in our dataset by sampling 38 wild males of *B. terrestris terrestris* in Belgium *in natura* with a net (Table S1). As wild males of *B. terrestris* could be confused with the closely related species *B. cryptarum*, the species identification was checked by CLGS analyses: the two species have species-specific CLGS composition used as diagnostic traits (e.g. Bertsch et al 2005; see CLGS analyses protocol below).

All wild (n=38) and reared (n=52) males (n total=90) were killed by freezing at -20°C. For the reared males, we used five to ten days old males when the *B. terrestris* males are the most attractive to females (Coppée *et al.*, 2011). For the wild males, we examined the wing condition (Tkalců, 1969) in order to avoid the sampling of old specimens that have a CLGS modification composition and are less attractive to females (Žáček *et al.*, 2009; Coppée *et al.*, 2011).

Determination of the Degree of Ploidy

We assessed the degree of ploidy by genotyped males with 13 microsatellite markers developed for *B. terrestris*: B11, B100, B121, B126 and B132 (Estoup *et al.*, 1993) and 0007_47n22, 0078_59o4, 0222_63d21, 0583_22I4, 0636_34m4, 0646_83e8, 0930_40o1, 0939_33h17 (Stolle *et al.*, 2009). We extracted sample DNA following the method described by Maebe *et al.* (2013) with some small modifications: 200 µl of 5% Chelex (InstaGeneTM matrix, BioRad) and 10 µl of proteinase K (20 mg/ml) were added to one middle leg and incubated at 56°C for 2h and 97°C for 15 min. The microsatellite loci: B11, B100, B121, B126 and B132 were amplified in multiplex by PCR as described in Maebe *et al.* (2013). For allele detection of the primers originated from Stolle *et al.* (2011), we used a tailed-primer approach (Schuelke, 2000). We used the PCR protocol described in Maebe *et al.* (2015): simplex PCR reactions were performed, containing 1.5 µl template DNA, 1 µl of 10x PCR buffer (Qiagen), 0.2 µl of 10 mM dNTP's (Qiagen), 0.1 µl of 10 µM forward primer, 0.4 µl of 10 µM reverse primer, 0.4 µl of 10 µM labeled M13-primer and 0.05 µl of 2.5 units/reaction Hotstar Taq DNA Polymerase (Qiagen). Initial denaturing at 95°C for 15 min was followed by 30 cycles of denaturing at 94°C for 30 s, annealing at 48,

52 or 58°C for 30 s, and extension at 72°C for 30 s, and ended at 72°C for 10 min with a final extension step. Pooled samples were visualized on a ABI-3730xl sequencer (Applied Biosystems) and fragments were scored manually using Peak Scanner Software v 1.0 (Applied Biosystems).

Comparative Chemical Analyses of Cephalic Labial Gland Secretions

We extracted the CLGS of 90 males following the entire cut head protocol described in De Meulemeester *et al.*, 2011). Vials containing the solvent and sample were (i) kept for 24 h at room temperature (20°C) to fulfill the extraction and (ii) stored at -40°C for two months prior to the analyses. We (i) analyzed the CLGS samples and identified compounds using gas chromatography-mass spectrometry (GC/MS) and (ii) quantified them by gas chromatography with flame ionization detector (GC/FID) (see GC/MS and GC/FID specifications and temperature program in Lecocq *et al.*, 2015). We identified the CLGS in Xcalibur™ with their mass spectra compared to those in the National Institute of Standards and Technology Library (NIST, U.S.A) with NIST MS Search 2.0. We determined the double bond positions from (i) mass spectra of dimethyl disulphide adducts of unsaturated components (Francis, 1981) (reaction time: four hours) and (ii) by chemical ionization with acetonitrile as a reaction gas (Oldham & Svatoš, 1999) (see protocol details in Lecocq *et al.*, 2013). We quantified the peak areas of CLGS compounds in GCSolution Postrun (Shimadzu Corporation) with automatic peak detection and noise measurement. We elaborated the data matrix (Table S2) with the relative amounts of each compound for each individual (the relative amounts are calculated by dividing the peak areas of compounds by the total area of compounds in each sample). We based our data matrix on the alignment of each compound between all samples performed with GCAligner 1.0 (Dellicour & Lecocq, 2013a,b).

We compared the CLGS between haploid and diploid males with non-metric multidimensional scaling (nMDS) ordination using a Bray-Curtis similarity matrix, three dimensions and 50 runs (R-package ecodist, Goslee & Urban, 2007). We transformed ($\log(x+1)$) the data matrix to reduce the great difference of abundance between compounds in high and low concentration (De Meulemeester *et al.*, 2011). An hypothetic CLGS differentiation between haploids and diploids was assessed by performing permutation-based version of the multivariate analysis of variance (perMANOVA) using the Bray-Curtis similarity matrix and 10,000 permutations (R-package vegan, Oksanen *et al.*, 2011) (perMANOVA assumption; i.e. similar variance homogeneity between groups was checked by performing a distance-based test for multivariate homogeneity of group dispersions for a one-way ANOVA design; Anderson, 2006). We determine CLGS compounds that are potentially specific to and

regular to diploid or haploid males with the indicator value (IndVal) method (Dufrêne & Legendre, 1997; see details in Lecocq *et al.*, 2015b). All statistical analyses were performed in R (R Development Core Team, 2013).

Results

Genetic analyses showed that reared queen-right colonies produced nine diploid males (Table S1). One diploid male was also found in wild specimens (Table S1). Other males were unambiguously haploid (Table S1).

Fifty six compounds were detected in the CLGS of analyzed specimens (Table S2). The general pattern obtained for CLGS of diploid males corresponded to those of haploid males. No qualitative difference was detected between haploid and diploid males (including no differences in double bound positions). The main compounds (i.e. compounds that have the highest relative amount in one individual at least) were farnesol, 2,3-Dihydrofarnesol, geranylcitronellol, and 2,3-dihydrofarnesyl dodecanoate. The nMDS (stress value = 0.16; $R^2=0.83$) did not show any obvious CLGS differentiations between haploid and diploid males except for a slight trend observed for the reared diploid males in the nMDS (i.e. these specimens are more or less grouped together at the edge of the nMDS) (Fig. 1). The perMANOVA ($DF = 1$, $F = -9.17$, $P = 0.79$) analyses confirmed the lack of significant CLGS differentiation between the two groups. The IndVal method did not reveal indicator compounds with strong significance ($IndVal > 0.70$) in haploid or diploid males (Table S2).

Discussion

Diagnostic Features of Diploid Bumblebee Males

In view of the dramatic consequences of diploid male production on population (Zayed & Packer, 2005), we expected that *B. terrestris* would display pre-mating mechanisms avoiding virgin queen mating with diploid males as already observed in some Hymenoptera (Herzner *et al.*, 2006). Such pre-mating mechanisms are most likely based on differences between reproductive traits used in pre-mating recognition between diploid and haploid males. Therefore, we investigated one reproductive trait, the CLGS that acts as a key trait for the bumblebee pre-mating recognition. However, our comparative statistical analyses based on CLGS composition did not detect obvious differences between diploid and haploid males in their CLGS, a main key trait for mate attraction (Fig. 1). The slight trend observed for the reared diploid males in the nMDS (i.e. these specimens are more or less grouped together at the edge of the nMDS; Fig. 1) could reflect a "family-effect" since all queens that produced these males are sister mated with their brothers as already observed in other study (Lecocq *et al.*, 2011). Nevertheless, our statistical analyses did not show significant differentiation between CLGS of diploid

and haploid males (see perMANOVA results). Even if the components of the male CLGS that constitute the pheromones are not known (i.e. those components that release a female response or convey information for females), this does not invalidate the basic result that there is no significant differentiation in diploid male CLGS since there is no specific and regular chemical compounds to diploid or haploid males (see IndVal results in Table S2). Although we are aware of the limited power of our statistical analyses due to our small sample size (10 diploid males), our data clearly demonstrates that there is no strong difference in the CLGS composition between haploid and diploid males.

According to our results we speculate that the queens are not able to discriminate the diploid males through their CLGS. However, since the bumblebee male pre-mating signal (Baer, 2003; Ayasse & Jarau, 2014; Streinzer & Spaethe, 2014) includes behavioral and other chemical features such as tendon glands (Jarau *et al.*, 2012) that are mainly poorly known, we cannot rule out the fact that the whole pre-mating signal of diploid males are evaluated by females as being strictly similar to those of haploid males. Indeed other traits involved in the species communication (e.g. cuticular hydrocarbons) or in the reproduction could be different in diploid *B. terrestris* males such as observed for cuticular compounds and testis size of diploid drone in the close relative *Apis mellifera* L. (Herrmann *et al.*, 2005). Moreover, diploid *B. terrestris* males are distinct in other characters such as wing shape, body size, and physiology (Duchateau & Marien, 1995; Gerloff *et al.*, 2003; Gérard *et al.*, 2015). These features could allow the female to diagnose diploid males and/or decrease the fitness of diploid males (e.g. fitness of a bumblebee male is correlated with body size, Cueva del Castillo & Fairbairn, 2012). Further studies on other traits and ethological test allowing choices of virgin females could help to define whether the *B. terrestris* females are not able to recognize diploid males. Nevertheless, current evidence suggests that queens are not able to discriminate the diploid males during the pre-mating recognition because (i) our results show that there are no diploid specific cephalic labial gland secretions that presumably contain the main long-distance signal of the bumblebee pre-mating recognition (e.g. Ayasse & Jarau, 2014), and (ii) diploid males have the same mating success as haploid males in experimental conditions where short distance pre-mating recognition signals should act (Duchateau & Marien, 1995).

Diploid Males in the Wild

Diploid males have been previously detected, mainly in bumblebee species with small and fragmented distribution like *B. florilegus* in Japan and *B. muscorum* in Great Britain, with a considerable proportion of such males of up to 50% (Darvill *et al.*, 2006; Takahashi *et al.*, 2008). However, diploid males were detected in

populations of widespread species like *B. occidentalis*, *B. perplexus*, and *B. terricola* in about 5% frequency (Whidden & Owen 2011). Our results show a ratio ($\approx 2\%$) of wild diploid males in *B. terrestris* similar to those found in other widespread bumblebee species. This low ratio was expected since *B. terrestris* is one of the most common and widespread bumblebee species in Europe (Rasmont *et al.*, 2008). Indeed, the large distribution and population density of this species (Rasmont *et al.*, 2008, 2015) make unlikely a limited genetic flow (i.e. genetic drift) between populations and therefore a high diploid ratio. However, our current sampling is limited to a small geographic region and to a small sample. Further studies on the diploid male ratio on large *B. terrestris* sampling and other European and non-European species are needed to validate this hypothesis.

Consequences for Bumblebee Conservation

If future studies confirm current evidence (see our results and Duchateau & Marien 1995) that diploid males cannot be recognized through their reproductive traits by females, this rules out the potential role of pre-mating recognition in curbing an ongoing vortex of extinction of diploid males in bumblebee species. Zayed & Packer (2005) and Hein *et al.* (2009) underline four crucial conditions that must be met to initialize the diploid male vortex: (i) fixed sex ratio, (ii) low reproductive rate, (iii) no mate choice (i.e. females cannot detect if potential mates are related mates), and (iv) extreme population fragmentation. Conservation plans should act on the two last factors since the two first conditions depend on intrinsic features of bumblebee species. The decrease of population fragmentation through the establishment of corridors (Holzschuh *et al.*, 2009) between populations could increase the genetic diversity of populations and thus prevent the initialization of diploid male vortex. Alternatively, individual translocations between populations (Arrendal *et al.*, 2004) could neutralize the fragmentation and enable females to choose exotic (unrelated) mates. Nevertheless, the beneficial effects of such translocations would appear to be restricted to areas in the immediate vicinity of the release sites (Arrendal *et al.*, 2004). Moreover, potential translocations should consider the interpopulational differentiations to maximize the genetic distinctiveness of translocated individuals while preserving the species diversity (i.e. avoid genetic homogenization through translocation; Lecocq *et al.*, 2015b).

Acknowledgments

This research was supported by the Belgian Science Policy (project BR/132/A1/BELBEES) and the Institute of Organic Chemistry and Biochemistry of the Academy of Sciences of the Czech Republic (RVO61388963). TL was grant holders of the *Fonds pour la Recherche dans l'Industrie et l'Agriculture*. Authors acknowledge David Notton (the Natural History Museum, London, UK) for correcting the English.

231 **Disclosure**

232 The authors declare that they have no conflict of interest.

233 **References**

- 234 Anderson, M.J. (2006) Distance-based tests for homogeneity of multivariate dispersions. *Biometrics*, 62, 245–
235 253.
- 236 Arrendal, J., Walker, C.W., Sundqvist, A.-K., Hellborg, L. and Vilà, C. (2004) Genetic evaluation of an otter
237 translocation program. *Conservation Genetics*, 5, 79–88.
- 238 Ayabe, T., Hoshiba, H. and Ono, M. (2004) Cytological evidence for triploid males and females in the
239 bumblebee, *Bombus terrestris*. *Chromosome Research*, 12, 215–223.
- 240 Ayasse, M. and Jarau, S. (2014) Chemical ecology of bumble bees. *Annual Review of Entomology*, 59, 299–319.
- 241 Baer, B. (2003) Bumblebees as model organisms to study male sexual selection in social insects. *Behavioral*
242 *Ecology and Sociobiology*, 54, 521–533.
- 243 Bergström, G., Svensson, B.G., Appelgren, M. and Groth, I. (1981) Complexity of bumble bee marking
244 pheromones: biochemical, ecological and systematical interpretations. *Systematics Association*, special vo,
245 175–183.
- 246 Bertsch, A., Schweer, H., Titze, A. and Tanaka, H. (2005) Male labial gland secretions and mitochondrial DNA
247 markers support species status of *Bombus cryptarum* and *B. magnus* (Hymenoptera, Apidae). *Insects*
248 *Sociaux*, 52, 45–54.
- 249 Calam, D.H. (1969) Species and sex-specific compounds from the heads of male bumblebees (*Bombus* spp.).
250 *Nature*, 221, 856–857.
- 251 Carvalheiro, L.G., Kunin, W.E., Keil, P., Aguirre-Gutiérrez, J., Ellis, W.N., Fox, R., Groom, Q., Hennekens, S.,
252 Van Landuyt, W., Maes, D., Van de Meutter, F., Michez, D., Rasmont, P., Ode, B., Potts, S.G., Reemer,
253 M., Roberts, S.P.M., Schaminée, J., Wallisdevries, M.F. and Biesmeijer, J.C. (2013) Species richness
254 declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecology*
255 *Letters*, 16, 870–878.
- 256 Coppée, A. (2010) *Bombus terrestris* (L. 1758): A complex species or a species complex ? - Intraspecific
257 pheromonal and genetic variations of *Bombus terrestris* (L.), Impacts on the speciation. Ph-D Thesis,
258 University of Mons, Mons (Belgium).
- 259 Coppée, A., Mathy, T., Cammaerts, M.-C., Verheggen, F.J., Terzo, M., Iserbyt, S., Valterová, I. and Rasmont, P.
260 (2011) Age-dependent attractivity of males' sexual pheromones in *Bombus terrestris* (L.) [Hymenoptera,
261 Apidae]. *Chemoecology*, 21, 75–82.
- 262 Coppée, A., Terzo, M., Valterová, I. and Rasmont, P. (2008) Intraspecific variation of the cephalic labial gland
263 secretions in *Bombus terrestris* (L.) (Hymenoptera: Apidae). *Chemistry and Biodiversity*, 5, 2654–2661.
- 264 Crozier, R.H. (1977) Evolutionary genetics of the Hymenoptera. *Annual Review of Entomology*, 22, 263–288.
- 265 Cueva del Castillo, R. and Fairbairn, D.J. (2012) Macroevolutionary patterns of bumblebee body size: detecting
266 the interplay between natural and sexual selection. *Ecology and Evolution*, 2, 46–57.

- 267 Darvill, B., Ellis, J.S., Lye, G.C. and Goulson, D. (2006) Population structure and inbreeding in a rare and
268 declining bumblebee, *Bombus muscorum* (Hymenoptera: Apidae). *Molecular Ecology*, 15, 601–611.
- 269 Darvill, B., Lepais, O., Woodall, L.C. and Goulson, D. (2012) Triploid bumblebees indicate a direct cost of
270 inbreeding in fragmented populations. *Molecular Ecology*, 21, 3988–95.
- 271 De Meulemeester, T., Gerbaux, P., Boulvin, M., Coppée, A. and Rasmont, P. (2011) A simplified protocol for
272 bumble bee species identification by cephalic secretion analysis. *Insectes Sociaux*, 58, 227–236.
- 273 Dellicour, S. and Lecocq, T. (2013a) *GALIGNER 1.0 and GCKOVATS 1.0 - Manual of a software suite to*
274 *compute a multiple sample comparison data matrix from eco-chemical datasets obtained by gas*
275 *chromatography*, University of Mons, Mons (Belgium).
- 276 Dellicour, S. and Lecocq, T. (2013b) GALIGNER 1.0: an alignment program to compute a multiple sample
277 comparison data matrix from large eco-chemical datasets obtained by GC. *Journal of Separation Science*,
278 36, 3206–3209.
- 279 Duchateau, M.J. and Marien, J. (1995) Sexual biology of haploid and diploid males in the bumble bee *Bombus*
280 *terrestris*. *Insectes Sociaux*, 42, 255–266.
- 281 Duchateau, M.J., Hoshiba, H. and Velthuis, H.H.W. (1994) Diploid males in the bumble bee *Bombus terrestris*.
282 Sex determination, sex alleles and viability. *Entomologia Experimentalis et Applicata*, 71, 263–269.
- 283 Dufrêne, M. and Legendre, P. (1997) Species assemblages and indicator species: The need for a flexible
284 asymmetrical approach. *Ecological Monographs*, 67, 345–366.
- 285 Estoup, A., Solignac, M., Harry, M. and Cornuet, J.-M. (1993) Characterization of (GT)_n and (CT)_n
286 microsatellites in two insect species *Apis mellifera* and *Bombus terrestris*. *Nucleic Acid Research*, 21,
287 1427–1431.
- 288 Francis, G.W. (1981) Alkylthiolation for the determination of double-bond position in unsaturated fatty acid
289 esters. *Chemistry and Physics of Lipids*, 29, 369–374.
- 290 Frankham, R., Ballou, J.D. and Briscoe, D.A. (2010) *Introduction to Conservation Genetics*, 2nd edn.
291 Cambridge University Press, Cambridge, UK.
- 292 Free, J.B. (1993) *Insect Pollination of Crops*, 2nd edn. Academic Press, London, UK.
- 293 Gérard, M., Michez, D., Fournier, D., Maebe, K., Smagghe, G., Biesmeijer, J.C. and De Meulemeester, T.
294 (2015) Discrimination of haploid and diploid males of *Bombus terrestris* (Hymenoptera; Apidae) based on
295 wing shape. *Apidologie*, 46, 644–653.
- 296 Gerloff, C.U., Ottmer, B.K. and Schmid-Hempel, P. (2003) Effects of inbreeding on immune response and body
297 size in a social insect, *Bombus terrestris*. *Functional Ecology*, 17, 582–589.
- 298 Goslee, S.C. and Urban, D.L. (2007) The ecodist package for dissimilarity-based analysis of ecological data.
299 *Journal of Statistical Software*, 22, 1–19.
- 300 Heimpel, G.E. and de Boer, J.G. (2008) Sex determination in the Hymenoptera. *Annual Review of Entomology*,
301 53, 209–230.
- 302 Hein, S., Poethke, H.-J. and Dorn, S. (2009) What stops the “diploid male vortex”?—A simulation study for
303 species with single locus complementary sex determination. *Ecological Modelling*, 220, 1663–1669.
- 304 Herrmann, M., Trenzcek, T., Fahrenhorst, H. and Engels, W. (2005) Characters that differ between diploid and
305 haploid honey bee (*Apis mellifera*) drones. *Genetics and Molecular Research*, 4, 624–641.

306 Herzner, G., Schmitt, T., Heckel, F., Schreier, P. and Strohm, E. (2006) Brothers smell similar: variation in the
307 sex pheromone of male European Beewolves *Philanthus triangulum* F. (Hymenoptera: Crabronidae) and
308 its implications for inbreeding avoidance. *Biological Journal of the Linnean Society*, 89, 433–442.

309 Holzschuh, A., Steffan-Dewenter, I. and Tscharnkte, T. (2009) Grass strip corridors in agricultural landscapes
310 enhance nest-site colonization by solitary wasps. *Ecological Applications*, 19, 123–132.

311 Jarau, S., Záček, P., Sobotník, J., Vrkoslav, V., Hadravová, R., Coppée, A., Vašíčková, S., Jiroš, P. and
312 Valterová, I. (2012) Leg tendon glands in male bumblebees (*Bombus terrestris*): structure, secretion
313 chemistry, and possible functions. *Die Naturwissenschaften*, 99, 1039–49.

314 Klein, A.M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A., Kremen, C. and Tscharnkte,
315 T. (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings. Biological*
316 *sciences / The Royal Society*, 274, 303–313.

317 Lecocq, T., Brasero, N., Martinet, B., Valterová, I. and Rasmont, P. (2015a) Highly polytypic taxon complex:
318 interspecific and intraspecific integrative taxonomic assessment of the widespread pollinator *Bombus*
319 *pascuorum* Scopoli 1763 (Hymenoptera: Apidae). *Systematic Entomology*, 40, 881–888.

320 Lecocq, T., Brasero, N., Meulemeester, T. De, Michez, D., Dellicour, S., Lhomme, P., Jonghe, R. De, Valterová,
321 I., Urbanová, K. and Rasmont, P. (2015b) An integrative taxonomic approach to assess the status of
322 Corsican bumblebees : implications for conservation. *Animal Conservation*, 18, 236–248.

323 Lecocq, T., Coppée, A., Mathy, T., Lhomme, P., Cammaerts-Tricot, M.-C., Urbanová, K., Valterová, I. and
324 Rasmont, P. (2015c) Subspecific differentiation in male reproductive traits and virgin queen preferences,
325 in *Bombus terrestris*. *Apidologie*, 46, 595–605.

326 Lecocq, T., Dellicour, S., Michez, D., Lhomme, P., Vanderplanck, M., Valterová, I., Rasplus, J.-Y. and
327 Rasmont, P. (2013) Scent of a break-up: phylogeography and reproductive trait divergences in the red-
328 tailed bumblebee (*Bombus lapidarius*). *BMC Evolutionary Biology*, 13, 263.

329 Lecocq, T., Lhomme, P., Michez, D., Dellicour, S., Valterová, I. and Rasmont, P. (2011) Molecular and
330 chemical characters to evaluate species status of two cuckoo bumblebees: *Bombus barbutellus* and *Bombus*
331 *maxillosus* (Hymenoptera, Apidae, Bombini). *Systematic Entomology*, 36, 453–469.

332 Lhomme, P., Ayasse, M., Valterová, I., Lecocq, T. and Rasmont, P. (2012) Born in an alien nest: how do social
333 parasite male offspring escape from host aggression? *PloS One*, 7, e43053.

334 Lhomme, P., Sramkova, A., Kreuter, K., Lecocq, T., Rasmont, P. and Ayasse, M. (2013) A method for year-
335 round rearing of cuckoo bumblebees (Hymenoptera: Apoidea: *Bombus* subgenus *Psithyrus*). *Annales de la*
336 *Société Entomologique de France*, 49, 117–125.

337 Maebe, K., Meeus, I., De Riek, J. and Smagghe, G. (2015) Quantitative trait loci for light sensitivity, body
338 weight, body size, and morphological eye parameters in the bumblebee, *Bombus terrestris*. *PLoS One*, 10,
339 e0125011.

340 Maebe, K., Meeus, I., Maharramov, J., Grootaert, P., Michez, D., Rasmont, P. and Smagghe, G. (2013)
341 Microsatellite analysis in museum samples reveals inbreeding before the regression of *Bombus veteranus*.
342 *Apidologie*, 44, 188–197.

343 Oksanen, F.J., Blanchet, G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P.,
344 Stevens, M.H.H. and Wagner, H. (2011) Vegan: Community Ecology Package.

345 Oldham, N.J. and Svatoš, A. (1999) Determination of the double bond position in functionalized monoenes by
346 chemical ionization ion-trap mass spectrometry using acetonitrile as a reagent gas. *Rapid Communications*
347 *in Mass Spectrometry*, 13, 331–336.

- 348 Ollerton, J., Winfree, R. and Tarrant, S. (2011) How many flowering plants are pollinated by animals? *Oikos*,
349 120, 321–326.
- 350 Payne, C.M., Lavery, T.M. and Lachance, M.A. (2003) The frequency of multiple paternity in bumble bee
351 (*Bombus*) colonies based on microsatellite DNA at the B10 locus. *Insectes Sociaux*, 50, 375–378.
- 352 Potts, S.G., Biesmeijer, J.C., Bommarco, R., Felicioli, A., Fischer, M., Jokinen, P., Kleijn, D., Klein, A.-M.,
353 Kunin, W.E., Neumann, P., Penev, L.D., Petanidou, T., Rasmont, P., Roberts, S.P.M., Smith, H.G.,
354 Sørensen, P.B., Steffan-Dewenter, I., Vaissière, B.E., Vila, M., Vujić, A., Woyciechowski, M., Zobel, M.,
355 Settele, J. and Schweiger, O. (2011) Developing European conservation and mitigation tools for
356 pollination services: Approaches of the STEP (Status and Trends of European Pollinators) project. *Journal*
357 *of Apicultural Research*, 50, 152–164.
- 358 R Development Core Team (2013) *R: A language and environment for statistical computing*, R Foundation for
359 Statistical Computing, Vienna, Austria.
- 360 Rasmont, P., Coppée, A., Michez, D. and De Meulemeester, T. (2008) An overview of the *Bombus terrestris* (L.
361 1758) subspecies (Hymenoptera: Apidae). *Annales de la Société entomologique de France (N.S.)*, 44, 243–
362 250.
- 363 Rasmont, P., Franzen, M., Lecocq, T., Harpke, A., Roberts, S., Biesmeijer, K., Castro, L., Cederberg, B.,
364 Dvorak, L., Fitzpatrick, U., Gonseth, Y., Haubruge, E., Mahe, G., Manino, A., Michez, D., Neumayer, J.,
365 Odegaard, F., Paukkunen, J., Pawlikowski, T., Potts, S., Reemer, M., Settele, J., Straka, J. and Schweiger,
366 O. (2015) Climatic risk and distribution atlas of European bumblebees. *BioRisk*, 10, 1–236.
- 367 Schuelke M. (2000) An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology*,
368 18: 233–234.
- 369 Stolle, E., Rohde, M., Vautrin, D., Solignac, M., Schmid-Hempel, P., Schmid-Hempel, R. and Moritz, R.F.A.
370 (2009) Novel microsatellite DNA loci for *Bombus terrestris* (Linnaeus, 1758). *Molecular Ecology*
371 *Resources*, 9, 1345–52.
- 372 Streinzer, M. and Spaethe, J. (2014) Functional morphology of the visual system and mating strategies in
373 bumblebees (Hymenoptera, Apidae, *Bombus*). *Zoological Journal of the Linnean Society*, 170, 735–747.
- 374 Takahashi, J., Ayabe, T., Mitsuhata, M., Shimizu, I. and Ono, M. (2008) Diploid male production in a rare and
375 locally distributed bumblebee, *Bombus florilegus* (Hymenoptera, Apidae). *Insectes Sociaux*, 55, 43–50.
- 376 Whidden, T.L. and Owen, R.E. (2011) Frequencies of Diploid Males in Natural Populations of Three North
377 American Bumble Bee (*Bombus*) Species (Hymenoptera: Apidae). *Annals of the Entomological Society of*
378 *America*, 104, 83–87.
- 379 Whitehorn, P.R., Tinsley, M.C., Brown, M.J.F., Darvill, B. and Goulson, D. (2009) Impacts of inbreeding on
380 bumblebee colony fitness under field conditions. *BMC Evolutionary Biology*, 9, 152.
- 381 Williams, P.H. (1985) A preliminary cladistic investigation of relationships among the bumble bees
382 (Hymenoptera, Apidae). *Systematic Entomology*, 10, 239–255.
- 383 Winfree, R. (2010) The conservation and restoration of wild bees. *Annals of the New York Academy of Sciences*,
384 1195, 169–197.
- 385 Woyke, J. (1963) Drone larvae from fertilized eggs of the honeybee. *Journal of Apicultural Research*, 1, 19–24.
- 386 Žáček, P., Kalinová, B., Šobotník, J., Hovorka, O., Ptáček, V., Coppée, A., Verheggen, F. and Valterová, I.
387 (2009) Comparison of age-dependent quantitative changes in the male labial gland secretion of *bombus*
388 *terrestris* and *Bombus lucorum*. *Journal of Chemical Ecology*, 35, 698–705.

- 389 Zayed, A. (2009) Bee genetics and conservation. *Apidologie*, 40, 237–262.
- 390 Zayed, A. and Packer, L. (2005) Complementary sex determination substantially increases extinction proneness
391 of haplodiploid populations. *Proceedings of the National Academy of Sciences of the United States of*
392 *America*, 102, 10742–6.
- 393


394 **Table**

395 **Table S1** Sampling table. The table gives the origin of samples and the ploidy determination by genotyping each
396 specimen with 13 microsatellite loci with indication of the fluorescent dye used in the PCR amplification.

397 **Table S2** Data matrix of cephalic labial gland secretions (relative amounts of each compound) and list of the
398 identified compounds in the sampling. MW is the molecular weight of the compound. IndVal results are
399 indicator value obtained in IndVal method.

400 **Figure captions**

401 **Fig. 1** Statistical analyses of cephalic labial gland secretion. Non-metric multidimensional scaling (nMDS)
402 ordination analysis of chemical reproductive trait of diploid and haploid *Bombus terrestris* males (n= 90). Three
403 first axes of the nMDS ordination plot based on Bray-Curtis distances calculated on a male cephalic labial gland
404 secretion matrix of *B terrestris*.



- Haploid males
- Diploid males (bred specimens)
- Diploid males (wild specimens)