First Chemical Analysis and Characterization of the Male Species-Specific Cephalic Labial-Gland Secretions of South American Bumblebees

Nicolas Braserio, Baptiste Martinet, Klara Urbanova, Irena Valterova, Alexandra Torres, Wolfgang Hoffmann, Pierre Rasmont, Thomas Lecocq

► To cite this version:

Nicolas Braserio, Baptiste Martinet, Klara Urbanova, Irena Valterova, Alexandra Torres, et al.. First Chemical Analysis and Characterization of the Male Species-Specific Cephalic Labial-Gland Secretions of South American Bumblebees. *Chemistry and Biodiversity*, 2015, 12 (10), pp.1535-1546. 10.1002/cbdv.201400375 . hal-01575790

HAL Id: hal-01575790

<https://hal.univ-lorraine.fr/hal-01575790>

Submitted on 11 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First chemical analysis and characterization of the male species-specific cephalic labial gland secretions of South American bumblebee

Nicolas Brasero^{a*)}, Baptiste Martinet^{a)}, Klára Urbanová^{b*)}, Irena Valterová^{b)}, Alexandra Torres^{c)}, Wolfgang Hoffmann^{c)}, Pierre Rasmont^{a*)}, Thomas Lecocq^{a)}

^{a)} University of Mons, Research institute of Biosciences, Laboratory of Zoology, Place du Parc 20, 7000 Mons, Belgium (phone: +3265373435; e-mail: nicolas.brasero@umons.ac.be)

^{b)} Academy of Sciences of the Czech Republic, Institute of Organic Chemistry and Biochemistry, Flamingovo nám 2, CZ-166 10 Prague, Czech Republic (e-mail: irena@uochb.cas.cz)

^{c)} Grupo de Biocalorimetría, Facultad de Ciencias Básicas, Universidad de Pamplona, Km 1 vía Bucaramanga, Ciudadela Universitaria, Pamplona, Colombia (e-mail: datorres@unipamplona.edu.co)

^{*)} Corresponding author

Abstract

The evolution of signals and reproductive traits involved in the pre-mating recognition has been in focus of abundant research in several model species group such as bumblebees (genus *Bombus*). However, the most studied bumblebee reproductive trait, the male cephalic labial gland secretions (CLGS), remains unknown among bumblebee species from South America. In this study we investigated the CLGS of five South American bumblebees of the subgenera *Thoracobombus* (*Bombus excellens*, *B. atratus*) and *Cullumanobombus* (*B. rubicundus*, *B. hortulanus* and *B. melaleucus*) comparing their secretions to those of European closely related species. Our results show an obvious interspecific differentiation in all subgenera. The

interspecific differentiation in *Thoracobombus* involves different compounds with a large abundance (main compounds) while all *Cullumanobombus* share the same main components. This suggests that in *Cullumanobombus* subgenus differentiation in minor components could lead to species discrimination.

Keywords

Cephalic labial gland, Chemical reproductive traits, South America, bumblebees

Introduction

The pre-mating communication is of particular interest for evolutionary biologists because of its key role in sexual selection [1,2]. Therefore, the evolution of signals and reproductive traits involved in the pre-mating recognition has been widely studied [3] in several model species groups such as birds (e.g., [4–6]), moths (e.g. [7–9]) flies (e.g., [10–12]), or more recently the bumblebees (e.g., [13–15]).

The pre-mating recognition of bumblebees (genus *Bombus*) includes both chemical and behavioral features [16]. Most bumblebee males patrol along paths (i.e. patrolling behavior) where they scent-mark objects with their cephalic labial gland secretions (CLGS) [17,18] that attract conspecific virgin females [13,19]. Other species exhibit less common behavior as perching behavior, in which males wait individually at prominent places and dart passing queens or other moving objects [20]. The CLGS is a complex mixture of (mainly aliphatic) compounds[18,21,22] produced *de novo* by the cephalic labial glands (acinar gland) [23] from saturated fatty-acids [24,25]. The resulting composition is species-specific [18]. The CLGS are thus an excellent tool for exploring interspecific and intraspecific variations of pre-mating recognition [13–15,26] as well as they are useful chemo-taxonomic characters [22,27–29].

Despite the usefulness of the CLGS as model in evolutionary biology and taxonomy, the male chemical secretions of a great number of species remain unknown [13]: the CLGS have been described for 50 species of a total of approximately 250 bumblebee species [30]. In several biogeographic regions such as South America, no studies have been published. Indeed, the CLGS chemical analyses require fresh samples [31] that are most of the time unavailable for uncommon taxa. In this paper, we investigated, the CLGS of males of five South American bumblebees: *Bombus melaleucus* Handlirsch, 1888 and *B. hortulanus* Friese, 1904 are near species belonging to the "*Robustus* group" (= *Robustobombus* sensu Richards, 1968 [32]) while *B. rubicundus* Smith, 1854 (Fig. 1 D) belongs to "*Rubicundus* group" (= *Rubicundobombus* sensu Richards, 1968 [32]), these species-groups being now included in *Cullumanobombus* subgenus [33]. *B. atratus* Franklin, 1913 (Fig. 1 B) and *B. excellens* Smith, 1879 (Fig. 1 A) both belong to the "*Fervidus*-species group" (= *Fervidobombus* sensu Richards, 1968 [32]) now all included in *Thoracobombus* subgenus [33]. We ultimately aim to compare their CLGS composition with those of related species from the same subgenus: *B. (Thoracobombus) muscorum liepetterseni* (L. 1758), *B. (Thoracobombus) pascuorum melleofacies* Vogt, 1911, *B. (Cullumanobombus) cullumanus apollineus* Skorikov, 1910 (Fig. 2 C) and *B. (Cullumanobombus) semenoviellus* Skorikov, 1910.

Results and discussion

Composition of cephalic labial gland secretions

Chemical analyses detected 146 compounds (33 from *B. excellens*, 35 from *B. atratus*, 53 from *B. pascuorum*, 19 from *B. muscorum*, 16 from *B. rubicundus*, 22 from *B. hortulanus*, 39 from *B. melaleucus*, 22 from *B. cullumanus* and 44 from *B. semenoviellus*) (see Table 1). The results showed that all *Thoracobombus* species were differentiated in their CLGS composition (seven compounds shared by all species and 17 compounds shared between the two South

American species), including changes in main compounds (octadec-11-enol (29% - 46%) in *B. excellens* and octadec-9-enol (54% - 91%) in *B. atratus* (Table 1). Inside the European taxa, hexadec-7-enol (10% - 31%) in *B. pascuorum* and octadec-9-enyl acetate in *B. muscorum* were the major compounds (Table 1). By main compounds of a taxon, we mean every compound that has the highest relative amount within the CLGS at least in one individual. Other differences between the South American *Thoracobombus* were mainly qualitative differences in minor compounds (Table 1). Similarly, the *Cullumanobombus* species displayed qualitative differences (only five compounds shared; six compounds shared between the three South American *Cullumanobombus*), but the main compound, geranylgeranyl acetate, is in common to all *Cullumanobombus* analyzed here: 81% - 94% in *B. rubicundus*, 76% - 83% in *B. hortulanus*, 18% - 59% in *B. melaleucus*, 51% - 92% in *B. cullumanus* and 37% - 60% in *B. semenoviellus*. Qualitative variations differentiated species in minor components (Table 1).

Statistical analysis

For both *Thoracobombus* and *Cullumanobombus* groups, the statistical analyses of the CLGS confirmed the interspecific differentiation (Fig. 2). This chemical differentiation is supported by high values of multiscale bootstrap resampling (> 80%, Fig. 2). The interspecific differentiations between all consubgeneric species observed in our results are in agreement with the species-specificity of CLGS observed in other bumblebee groups (e.g., [18,22,27,29]). Similarly to our results on *Thoracobombus* species, the comparison of CLGS between closely related bumblebee taxa with a commonly recognized species status allow suggesting that the interspecific differentiation involves the change in main compounds (e.g., [18,29,34]). These main compound differentiations can be (i) a significant shift (production of new types of compounds, e.g., *B. pascuorum* versus *B. muscorum*) consecutive to switching off and on the respective metabolic pathway (e.g., [7]), (ii) large increase of a minor

compound shared with closely related species (e.g., octadec-9-enyl acetate between *B. muscorum* versus *B. atratus*) [35], as well as (iii) difference in double bond position arisen by activity of different enzymes (e.g., $\Delta 11$ -desaturase and $\Delta 9$ -desaturase introducing a double bond in octadecenol in *B. excellens* and *B. atratus*, e.g., [8]) or by activation of a non-functional enzyme gene transcript present in a common ancestor, as observed in moths [9] and in other bumblebees [15]. In contrast, all *Cullumanobombus* species display the same main component (geranylgeranyl acetate) as observed in the North American *B. (Cullumanobombus) rufocinctus* Cresson, 1863 [20]. The geranylgeranyl acetate is also a main component of *Bombus (Alpigenobombus) wurflenii* [36], *B. (Kallobombus) soroensis* [37] and *B. (Separatobombus) griseocollis* [38] meaning that this component could not be considered as autapomorphic of *Cullumanobombus*. Assuming the key role of CLGS in pre-mating recognition, this suggests that differentiation in minor compounds could lead to species discrimination. Moreover, for each CLGS group, the IndVal method revealed several significantly indicator compounds (see supplementary file 1).

Conclusion

The cephalic labial gland secretions of males are widely acknowledge as a useful and practical diagnostic trait for bumblebee species. Among the European bumblebee fauna, several taxonomically doubtful taxa groups have been solved thank to this reproductive trait. The present study is the first study to examine this trait among South American bumblebee species. This paves the way to a massive taxonomic assessment of South American bumblebees.

122 *Acknowledgement*

123 The authors sincerely thank Peter Neumann (University of Bern, Switzerland) for his help in
124 providing contact for collecting the biological material. Financial support (IV and KU) was
125 provided by the Institute of Organic Chemistry and Biochemistry of the Academy of Sciences
126 of the Czech Republic (subvention for development of research organization RVO:
127 61388963). TL is post-doc in the context of the BELBEES project funded by the Belgian
128 Scientific Politic (BELSPO, BR/132/A1/BELBEES). The authors acknowledge the *Fonds*
129 *pour la recherche dans l'industrie et l'agriculture* (FRIA) for financial support (Project
130 2.4.564.06 F). Field trips of PR and TL in Sweden, Italy and in Turkey to collect material
131 have been funded by the Belgian FNRS. Thanks to B. Cederberg (Uppsala) and A.M. Aytekin
132 (Ankara) for their help. Special thanks to R. de Jonghe for the breeding of *B. muscorum*
133 *liepetterseni* males and to Nicolas R. Chimento (Laboratorio de Anatomía Comparada y
134 Evolución de los Vertebrados Museo Argentino de Ciencias Naturales "Bernardino
135 Rivadavia") for providing us pictures of *Bombus atratus*.

136

137

138

139

140

141

142

143

144

145

146

147

148 **Experimental part**

149 *Biological material*

150 Fifty males from two subgenera were used for this study (*Table 2*): *B. melaleucus* (n=2), *B.*
151 *hortulanus* (n=3), *B. rubicundus* (n=7), *B. atratus* (n=12), and *B. excellens* (n=2). As
152 comparison group of *Thoracobombus* we used specimens already described in Lecocq *et al.*
153 [28]: *B. muscorum liepetterseni* (n=5), *B. pascuorum melleofacies* (n=7). We used *B.*
154 *cullumanus apollineus* (n=9) and *B. semenoviellus* (n=3) already described in [39] as
155 comparison group of *Cullumanobombus*. We killed specimens by freezing at -20°C. The
156 CLGS were extracted in 400 ml of hexane [31]. Samples were stored at -40°C prior to the
157 analyses.

158 *Chemical analyses*

159 We identified the CLGS composition by gas chromatography-mass spectrometry (GC/MS) on
160 a quadrupol Finigan GCQ, with a DB-5ms non-polar capillary column (5% phenyl (methyl)
161 polysiloxane stationary phase; 30-m column length; 0.25-mm inner diameter; 0.25-µm film
162 thickness). One µl was injected in splitless mode at the injector temperature set to 220°C. The
163 temperature of the column was initially held to 70°C for two minutes and then increased at a
164 rate of 10°C/min to 320°C. The temperature was then held at 320°C for five minutes. The
165 carrier gas was helium at a constant velocity of 50 cm/s. Compounds were identified using the
166 retention times and mass spectra of each peak. Double bond positions were determined by
167 DMDS (dimethyl disulfide) derivatization [40].

168 We analyzed all samples by a gas chromatograph-flame ionization detector Shimadzu GC-
169 2010 with a SLB-5ms non-polar capillary column (5% phenyl (methyl) polysiloxane

stationary phase; 30-m column length; 0.25-mm inner diameter; 0.25- μ m film thickness) with the same chromatographic conditions as in GC/MS.

Peak areas of compounds were detected in GCsolution Postrun (Shimadzu Corporation) with automatic peak detection and noise measurement. We calculated relative amounts (RA in %) of compounds in each sample by dividing the peak areas of compounds by the total area of compounds in each sample. We did not use any correction factor to calculate the RA of individual compounds. We discarded all compounds for which RA were recorded as less than 0.1% for all specimens [31]. We elaborated the data matrix as the alignment of each compound between all samples performed with GCAAligner 1.0 [41,42](see supplementary file 1).

Statistical analyses

Statistical analyses were performed using R [43] to detected CLGS differentiations between specimen groups. Data consisting of the relative proportion of all compounds were transformed ($\log(x-1)$) to reduce the large difference of abundance between highly and slightly concentrated compounds, and we then standardized (mean= 0, standard deviation= 1) the data matrix to reduce the sample concentration effect [22]. Clustering method was used to detect the divergence between taxa. A Pearson Phi Correlation matrix based on CLGS data matrix (relative abundance of each compound) was computed. An unweighted pair group method with arithmetic mean (UPGMA) was used as clustering method (R-package ape,[44]). We assessed the uncertainty in hierarchical cluster analysis using p -values calculated via multiscale bootstrap resampling with 50.000 bootstrap replications (R-package pvclust,[44]). Only bootstrap values greater than 85% were considered significant. To determine compounds of each taxa (indicator compounds), we used the indicator value (IndVal) method [45,46]. The value given is the product of relative abundance and relative frequency of

occurrence of a compound within a group. We evaluated the statistical significance of a compound as an indicator at the 0.01 level with a randomization procedure.

- 220 [1] H. E. H. Paterson, *Evolution and the Recognition Concept of Species*, The Johns
221 Hopkins University Press, Baltimore, US, 1993.
- 222 [2] B. R. Grant, P. R. Grant, *Evolution* (N. Y). **1996**, 50, 2471–2487.
- 223 [3] T. D. Wyatt, *Pheromones and Animal Behaviour: Communication by Smell and Taste*,
224 Cambridge University Press, Cambridge, UK, 2003.
- 225 [4] M. I. Förschler, E. K. V Kalko, *J. Biogeogr.* **2007**, 34, 1591–1600.
- 226 [5] J. Martens, in: Kroodsma, D.E., Miller, E.H., Eds., *Ecology and Evolution of Acoustic*
227 *Communication in Birds*, Comstock Publishing, Ithaca, **1996**, 221–240.
- 228 [6] G. L. Patricelli, J. A. C. Uy, G. Borgia, *Proc. R. Soc. B Biol. Sci.* **2003**, 270, 2389–
229 2395.
- 230 [7] N. Glaser, B. Frérot, E. Leppik, C. Monsempes, C. Capdevielle-Dulac, B. Le Ru, T.
231 Lecocq, M. Harry, E. Jacquin-Joly, P.-A. Calatayud, *J. Chem. Ecol.* **2014**, 40, 923–927.
- 232 [8] C. Löfstedt, *Philos. Trans. R. Soc. B-Biological Sci.* **1993**, 340, 167–177.
- 233 [9] W. L. Roelofs, W. Liu, G. Hao, H. Jiao, A. P. Rooney, C. E. Linn Jr., *Proc. Natl. Acad.*
234 *Sci. U. S. A.* **2002**, 99, 13621–13626.
- 235 [10] M. R. E. Symonds, A. Moussalli, M. A. Elgar, *Biol. J. Linn. Soc.* 2009, 97, 594–603.
- 236 [11] C. R. B. Boake, *Genetica* **2002**, 116, 205–214.
- 237 [12] J. E. Blyth, D. Lachaise, M. G. Ritchie, *Ethology* **2008**, 114, 728–736.
- 238 [13] M. Ayasse, S. Jarau, *Annu. Rev. Entomol.* **2014**, 59, 299–319.
- 239 [14] T. Lecocq, N. J. Vereecken, D. Michez, S. Dellicour, P. Lhomme, I. Valterová, J.-Y.
240 Rasplus, P. Rasmont, *PLoS One* **2013**, 8, e65642.
- 241 [15] T. Lecocq, S. Dellicour, D. Michez, P. Lhomme, M. Vanderplanck, I. Valterová, J.-Y.
242 Rasplus, P. Rasmont, *BMC Evol. Biol.* **2013**, 13, 263.
- 243 [16] B. Baer, *Behav. Ecol. Sociobiol.* 2003, 54, 521–533.
- 244 [17] G. Bergström, *Entomol. Scand. Suppl.* **1981**, 15, 173–184.
- 245 [18] D. H. Calam, *Nature* **1969**, 221, 856–857.
- 246 [19] A. Coppée, *Bombus Terrestris* (L. 1758) : *A Complex Species or a Species Complex ? -*
247 *Intraspecific Pheromonal and Genetic Variations of Bombus Terrestris* (L.), Impacts
248 on the Speciation, Université de Mons Editor, 2010.

- 249 [20] A. Bertsch, H. Schweer, A. Titze, *J. Chem. Ecol.* **2008**, 34, 1268–1274.
- 250 [21] A. Coppée, M. Terzo, I. Valterová, P. Rasmont, *Chem. Biodivers.* **2008**, 5, 2654–2661.
- 251 [22] T. Lecocq, P. Lhomme, D. Michez, S. Dellicour, I. Valterová, P. Rasmont, *Syst.*
252 *Entomol.* **2011**, 36, 453–469.
- 253 [23] B. Kullenberg, G. Bergström, B. Bringer, B. Carlberg, B. Cederberg, *Zoon* **1973**,
254 Suppl. 1, 23–30.
- 255 [24] P. Žáček, D. Prchalová-Hornáková, R. Tykva, J. Kindl, H. Vogel, A. Svatoš, I. Pichová,
256 I. Valterová, *ChemBioChem* **2013**, 14, 361–371.
- 257 [25] A. Luxová, I. Valterová, K. Stránský, O. Hovorka, A. Svatoš, *Chemoecology* **2003**, 13,
258 81–87.
- 259 [26] P. Lhomme, M. Ayasse, I. Valterová, T. Lecocq, P. Rasmont, *PLoS One* **2012**, 7,
260 e43053.
- 261 [27] A. Bertsch, H. Schweer, A. Titze, H. Tanaka, *Insectes Soc.* **2005**, 52, 45–54.
- 262 [28] T. Lecocq, N. Brasero, T. De Meulemeester, D. Michez, S. Dellicour, P. Lhomme, R.
263 de Jonghe, I. Valterová, K. Urbanová, P. Rasmont, *Anim. Conserv.* **2014**,
264 DOI: 10.1111/acv.12164
- 265 [29] P. Rasmont, M. Terzo, A. M. Aytekin, H. Hines, K. Urbanova, L. Cahlikova, I.
266 Valterova, *Apidologie* **2005**, 36, 571–584.
- 267 [30] P. H. Williams, *Bull. Nat. Hist. Museum* **1998**, 67, 79–152.
- 268 [31] T. De Meulemeester, P. Gerbaux, M. Boulvin, A. Coppée, P. Rasmont, *Insectes Soc.*
269 2011, 58, 227–236.
- 270 [32] O. W. Richards, *Bull. Br. Museum Natural Hist. Entomol.* **1968**, 22, 211–276.
- 271 [33] P. H. Williams, S. A. Cameron, H. M. Hines, B. Cederberg, P. Rasmont, *Apidologie*
272 **2008**, 39, 46–74.
- 273 [34] A. Bertsch, H. Schweer, *Biochem. Syst. Ecol.* **2012**, 40, 103–111.
- 274 [35] M. Terzo, K. Urbanová, I. Valterová, P. Rasmont, *Apidologie* **2005**, 36, 85–96.
- 275 [36] B. G. Svensson, M. Appelgren, G. Bergstrom, *Nov. acta Regiae Soc. Sci.* **1984**, 3, 145–
276 147.
- 277 [37] B. Kullenberg, G. Bergström, S. Ställberg-Stenhagen, *Acta Chem. Scand.* **1970**,
278 24, 1481

- 280 [38] A. Bertsch, H. Schweer, A. Titze, *Zeitschrift fur Naturforsch. - Sect. C J. Biosci.* **2004**,
281 59, 701–707.
- 282 [39] O. Hovorka, I. Valterová, P. Rasmont, M. Terzo, *Chem. Biodivers.* **2006**, 3, 1015–22.
- 283 [40] J. Cvacka, E. Kofronová, S. Vasícková, K. Stránský, P. Jiros, O. Hovorka, J. Kindl, I.
284 Valterová, *Lipids* **2008**, 43, 441–50.
- 285 [41] S. Dellicour, T. Lecocq, *GALIGNER 1.0 and GCKOVATS 1.0 - Manual of a Software*
286 *Suite to Compute a Multiple Sample Comparison Data Matrix from Eco-Chemical*
287 *Datasets Obtained by Gas Chromatography*, University of Mons, Mons (Belgium),
288 2013.
- 289 [42] S. Dellicour, T. Lecocq, *J. Sep. Sci.* **2013**, 36, 3206–9.
- 290 [43] R Development Core Team (2013). *R: a language and environment for statistical*
291 *computing*. Vienna: R Foundation for Statistical Computing. ISBN 3-900051-07-0.
292 <http://www.R-project.org> (accessed 2 January 2014).
- 293 [44] R. Suzuki, H. Shimodaira, *Pvclust: Hierarchical Clustering with P-Values via*
294 *Multiscale Bootstrap Resampling* 2011, R package version 1.2-2.
- 295 [45] J. Claudet, D. Pelletier, J.-Y. Jouvenel, F. Bachet, R. Galzin, *Biol. Conserv.* **2006**, 130,
296 349–369.
- 297 [46] M. Dufrêne, P. Legendre, *Ecol. Monogr.* **1997**, 67, 345–366.

305 *Table*

306 Table 1. *List of the identified compounds in the Thoracobombus and Cullumanobombus subgenera.* Molecular Weight (MW), median (M [%]) of
 307 the 146 identified compounds. For each species, main compounds are shown on black background, characteristic compounds calculated with the
 308 Indicator value (IndVal) method (>0.70) are shown in grey background. The compounds we were not able to determine are indicated as *Ux*. The
 309 full table with first and fourth quartile (Q1 and Q4 [%]) are in the supplementary file 1

310

Compounds	MW	<i>B. atratus</i>	<i>B. excellens</i>	<i>B. pascuorum</i>	<i>B. muscorum</i>	<i>B. hortulanus</i>	<i>B. melaleucus</i>	<i>B. rubicundus</i>	<i>B. cullumanus</i>	<i>B. semenoviellus</i>
		M	M	M	M	M	M	M	M	M
Octanoic acid	144	-	-	-	-	0.40	0.10	-	-	-
Decanoic acid	172	-	-	-	-	-	0.30	-	-	-
Dodecanol	186	-	-	-	-	-	-	-	-	0.67
U1	?	-	-	-	-	0.16	-	-	-	-
Dodecanoic acid	200	-	-	-	-	-	-	-	-	0.07
Tetradecenol	212	-	-	0.05	-	-	-	-	-	-
Tetradecanol	214	-	-	0.07	-	-	-	-	-	-
Tetradecanoic acid	228	-	-	0.01	-	-	-	-	-	-

Dodecyl acetate	228	-	-	-	-	0.09	-	-	-	-
Hexadec-7-enal	238	-	-	22.49	-	-	-	-	-	-
Hexadec-7-enol	240	-	-	24.12	-	-	-	-	-	-
Heptadecane	240	-	-	-	-	-	0.07	-	-	-
Hexadecanol	242	-	-	0.31	-	-	-	-	-	-
U2	?	-	-	-	-	0.10	-	-	-	-
Tetradecyl acetate	256	-	-	-	-	1.52	0.13	-	-	-
Farnesyl acetate	264	-	-	-	-	1.49	0.10	-	0.11	0.31
Hexadecanol	242	-	-	-	-	-	-	-	-	3.32
Beta-Springene	272	-	-	-	-	-	1.42	0.08	-	0.48
Hexadecenoic acid	254	-	-	5.85	-	-	-	-	-	-
Hexadec-9-enoic acid	254	-	-	-	-	-	-	-	-	6.49
Alpha-Springene	?	-	-	-	-	-	-	-	-	0.19
Hexadecanoic acid	256	-	-	0.32	-	-	3.03	-	-	2.40
U3	?	-	-	-	-	-	-	0.48	-	-
U4	?	-	-	-	-	0.15	-	-	-	-
Octadecanal	268	-	-	-	-	-	-	0.12	-	-

Ethyl hexadec-9-enoate	282	-	-	-	-	0.09	-	-	-	-
Hexadecenyl acetate	282	-	-	1.07	-	-	-	-	-	-
Hexadec-9-enyl acetate	282	-	-	-	-	9.39	23.53	-	-	-
Hexadecyl acetate	284	-	-	0.03	-	1.76	1.28	-	-	0.12
U5	?	0.01	-	-	-	-	-	-	-	-
Octadec-9-enal	266	0.41	-	-	0.11	-	-	-	-	-
Octadec-11-enal	266	0.13	2.42	-	-	-	-	-	-	-
Octadec-13-enal	266	-	0.89	-	-	-	-	-	-	-
U6	?	-	0.99	-	-	-	-	-	-	-
U7	?	-	0.40	-	-	-	-	-	-	-
Octadecanal	268	-	-	-	-	0.02	-	-	-	-
Octadec-9-enol	268	74.04	-	1.25	4.14	-	-	-	-	-
Octadec-11-enol	268	4.72	37.59	-	-	-	-	-	-	-
Octadec-13-enol	268	-	0.07	-	-	-	-	-	-	-
Icosane	282	-	-	0.04	-	-	-	-	-	-
Heneicosene	294	-	-	-	-	-	0.11	-	-	-
Heneicosane	296	0.05	0.30	0.10	0.19	-	0.23	3.78	3.66	-

U8	?	-	-	0.08	-	-	-	-	-	-
U9	?	-	-	0.05	-	-	-	-	-	-
Ethyl octadecenoate	310	-	-	-	-	-	-	0.50	0.11	-
Geranyl geranial	288	-	-	-	-	-	0.19	0.29	-	-
Geranyl decanoate	308	-	-	-	-	-	-	-	-	0.31
Octadekadienoic acid	280	-	-	-	0.06	-	0.40	-	-	-
Octadecenoic acid	282	-	-	2.84	-	-	-	-	-	-
Octadec-9-enoic acid	282	-	-	-	-	-	-	-	-	1.73
Octadec-11-enoic acid	282	-	-	-	-	-	21.41	-	-	-
Docosane	310	-	0.13	0.24	-	-	0.19	0.08	0.18	-
U10	?	-	-	-	-	-	0.45	-	-	-
U11	?	-	-	-	-	-	0.07	-	-	-
Unidentified terpene 1	?	-	-	-	-	-	-	-	-	0.09
U12	?	-	0.07	-	-	-	-	-	-	-
Ethyl octadecatrienoate	306	-	3.92	-	-	-	0.23	-	0.07	-
U13	?	-	0.93	-	-	-	-	-	-	-
Ethyl octadec-9-enoate	310	0.00	-	-	-	-	-	-	-	-

Octadec-9-enyl acetate	310	1.16	-	-	88.19	-	2.12	-	-	-
Octadec-11-enyl acetate	310	0.16	5.39	-	-	-	-	-	-	-
Octadecadienyl acetate	308	0.02	-	-	-	-	-	-	-	-
Octadecenyl acetate	310	-	-	0.47	-	-	-	-	-	-
Icosenal	294	-	0.12	-	-	-	-	-	-	-
Octadecadienyl acetate1	308	-	0.29	-	-	-	-	-	-	-
Octadecadienyl acetate2	308	-	0.17	-	-	-	-	-	-	-
Dodecyl octanoate	312	-	-	-	-	-	-	-	-	0.14
Geranylgeranyl acetate	322	-	-	-	-	82.93	38.42	91.77	87.94	48.08
Tricos-9-ene	322	0.53	17.77	0.13	-	0.14	0.16	-	-	-
Tricos-7-ene	322	0.48	4.13	0.08	-	-	0.13	-	-	-
Tricos-5-ene	322	0.07	-	-	-	-	-	-	-	-
Tricosane	324	2.91	8.72	3.46	2.63	-	1.06	-	-	-
Unidentified terpene 2	?	-	-	-	-	-	-	-	-	0.00
Geranyl dodecanoate	336	-	-	-	-	-	-	-	-	0.99
Tetracos-9-ene	336	0.00	2.91	0.02	-	-	-	-	-	-
Tetracosene	336	-	-	0.07	-	-	-	-	-	-

[illegible]

Heptacos-9-ene	378	0.12	0.24	1.61	0.32	0.12	-	-	0.12	0.19
Heptacos-7-ene	378	0.08	0.87	-	0.13	0.28	0.22	0.16	1.92	0.67
Heptacosane	380	0.13	0.69	0.80	0.36	0.47	1.05	0.68	1.15	1.02
Geranyl hexadec-7-enoate	390	-	-	-	-	-	-	-	-	0.24
Octacosene	392	-	-	0.05	-	-	-	-	-	-
Octacosane	394	-	-	0.06	-	-	-	-	-	-
Unidentified terpene3	?	-	-	-	-	-	-	-	-	0.34
U14	?	-	-	-	-	-	-	-	-	0.05
Dodecyl tetradecanoate	396	-	-	-	-	-	-	-	-	1.19
Squalene	410	-	-	-	-	-	-	-	0.05	-
Nonacosadiene1	404	-	-	-	-	-	-	-	0.03	-
Nonacosadiene2	404	-	-	-	-	-	-	-	0.11	-
Nonacos-9-ene	406	0.06	0.15	0.94	0.12	0.07	-	0.10	0.50	0.21
Nonacos-7-ene	406	0.03	0.18	-	-	0.24	0.22	0.19	-	0.44
Nonacosane	408	0.06	-	0.57	0.08	0.24	0.40	0.26	0.23	0.21
Unidentified terpene 4	?	-	-	-	-	-	-	-	-	0.09
Triacontene1	422	-	-	0.02	-	-	-	-	-	-

[illegible]

Hexadecyl tetradecanoate	452	-	-	-	-	-	-	-	-	0.21
Octadecenyl tetradecenoate/tetradecanoate	476/ 478	-	0.12	-	-	-	-	-	-	-
Hexadecenyl hexadecenoate	476	-	-	16.19	-	-	0.06	-	-	-
Hexadecyl hexadec-9-enoate	478	-	-	-	-	-	-	-	-	2.90
Hexadecyl hexadecanoate	480	-	-	-	-	-	-	-	-	0.51
Hexadecenyl octadecatrienoate	500	-	-	0.33	-	-	0.11	-	-	-
Octadecadienyl hexadecenoate	502	-	1.66	-	-	-	-	-	-	-
Hexadecenyl octadecenoate	504	-	-	1.10	-	-	-	-	-	-
Octadecenyl hexadecenoate	504	-	3.92	-	-	-	-	-	-	-
Hexadecyl octadecenoate	506	-	-	0.04	-	-	0.06	-	-	0.09
Octadecenyl hexadecanoate1	506	0.23	-	-	0.05	-	-	-	-	-
Octadecenyl hexadecanoate2	506	0.13	-	-	-	-	-	-	-	-
U17	?	0.07	-	-	-	-	-	-	-	-

Geranylgeranyl hexadecanoate	528	-	-	-	-	-	0.41	-	-	-
U18	?	-	-	-	-	-	0.17	-	-	-
Octadecadienyl octadecenoate	530	-	-	2.02	-	-	-	-	-	-
Octadecenyl octadecenoate	532	-	-	3.37	0.57	-	-	-	-	-
Octadecenyl octadecanoate	534	2.18	-	-	0.20	-	-	-	-	-
Octadecenyl octadecatrienoate	528	1.04	-	-	-	-	-	-	-	-

311 Table 2. Data on Collection of Biological Material. Collecting sites and number (N) of
 312 samples collected for *Bombus atratus*, *B. excellens*, *B. pascuorum melleofacies*, *B. muscorum*
 313 *liepetterseni*, *B. hortulanus*, *B. melaleucus*, *B. rubicundus*, *B. cullumanus apollineus* and *B.*
 314 *semenoviellus*. Coordinate are given with the reference to the WGS84.

Taxa	Collecting sites	Coordinates	N
<i>Thoracobombus</i> subgenus			
<i>B. atratus</i>	Colombia, Pamplona	7°22'N 72°39'W	4
	Colombia, Tenjo	4°52'N 74°09'W	8
<i>B. excellens</i>	Colombia, Cundinamarca	4°34'N 74°20'W	1
	Colombia, Pamplona	7°22'N 72°39'W	1
<i>B. pascuorum melleofacies</i>	Italy, Torre d'Isola	45°13'N 9°02'E	4
	Italy, Cellara	39°13'N 16°20'E	3
<i>B. muscorum liepetterseni</i>	Norway, Flatanger	64°28'N 10°43'E	5
<i>Cullumanobombus</i> subgenus			
<i>B. hortulanus</i>	Colombia, Pamplona	7°21'N 72°41'W	2
	Colombia, Cundinamarca	4°34'N 74°01'W	1
<i>B. melaleucus</i>	Colombia, Pamplona	7°32'N 72°37'W	2
<i>B. rubicundus</i>	Colombia, Pamplona	7°20'N 72°37'W	7
<i>B. cullumanus apollineus</i>	Turkey, Kars, Çıldır gölü	40°55'N 43°16'E	2
	Turkey, Kars, Göldalı	40°58'N 43°18'E	5
	Turkey, Kars, Çanaksu	41°00'N 43°18'E	2
<i>B. semenoviellus</i>	Czech Republic, Krkonoše	50°41'N 15°51'E	3

315

316

317

318

319 Figure 1. A: *Bombus excellens* from Colombia (photo: W. Hoffmann). B: *Bombus atratus*
 320 from Argentina (photo: N. Chimento). C: *Bombus cullumanus apollineus* from Turkey
 321 (photo: P. Rasmont). D: *Bombus rubicundus* from Colombia (photo: W. Hoffmann)

322

Figure 2. Unweighted pair group method with arithmetic mean cluster based on a correlation matrix calculated from matrix of cephalic labial gland secretions of *Bombus atratus*, *B. excellens*, *B. pascuorum melleofacies*, *B. muscorum liepetterseni*, *B. hortulanus*, *B. melaleucus*, *B. rubicundus*, *B. cullumanus apollineus* and *B. semenoviellus*. Values above branch represent multiscale bootstrap resampling (only values >80% are given)