



**HAL**  
open science

## Identification of *Streptococcus thermophilus* CNRZ368 genes involved in

Annabelle Thibessard, Frédéric Borges, Annabelle Fernandez, Brigitte Gintz,  
Bernard Decaris, Nathalie N. Leblond-Bourget

► **To cite this version:**

Annabelle Thibessard, Frédéric Borges, Annabelle Fernandez, Brigitte Gintz, Bernard Decaris, et al..  
Identification of *Streptococcus thermophilus* CNRZ368 genes involved in. *Applied and Environmental  
Microbiology*, 2004, pp.2220-9. 10.1128/AEM.70.4.2220-2229.2004 . hal-01631257

**HAL Id: hal-01631257**

**<https://hal.univ-lorraine.fr/hal-01631257v1>**

Submitted on 8 Nov 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## ABSTRACT

19

20

21 To better understand the defense mechanism of *Streptococcus thermophilus* against  
22 superoxide stress, molecular analysis of 10 menadione-sensitive mutants, obtained by  
23 insertional mutagenesis, was undertaken. This analysis allowed the identification of 10 genes  
24 that, with respect to their putative function, were classified in 5 categories (i) those involved  
25 in cell wall metabolism, (ii) those involved in exopolysaccharide translocation, (iii) those  
26 involved in RNA modification (iv) those involved in iron homeostasis and (v) those whose  
27 function is still unknown.

28 The behavior of the 10 menadione sensitive mutants exposed to heat shock was investigated.  
29 Data from these experiments allowed us to distinguish genes whose action might be specific  
30 to oxidative stress defense (*tgt*, *ossF*, *ossG*) from those whose action may be generalized to  
31 other stressful conditions (*mreD*, *rodA*, *pbp2b*, *cpsX*, *iscU*).

32 Among the mutants, two harboured an independently inserted copy of pGh9:ISS1 in two loci  
33 close to each other. More precisely, these two loci are homologous to *sufD* and *iscU* genes,  
34 involved in the biosynthesis of iron-sulfur clusters. This region, called the *suf* region, was  
35 further characterized in *S. thermophilus* CNRZ368 by sequencing and by construction of  
36  $\Delta$ *sufD* and *iscU*<sub>97</sub> non polar mutants. Streptonigrin sensitivity level of both mutants suggested  
37 that these two genes were involved in iron metabolism.

## INTRODUCTION

38

39 Oxygen is an essential element for the viability of most living organisms through its involvement  
40 in oxidative phosphorylation that takes place during the aerobic respiration process. However,  
41 oxygen can also be harmful, due to the production, especially during respiration, of toxic  
42 derivatives namely ROS (reactive oxygen species) such as hydrogen peroxide, hydroxyl and  
43 superoxide radicals (for a review, (25)). Accumulation of these molecules, triggers lipid, protein  
44 and nucleic acid oxidation, causing lethal cellular damage (for reviews (7, 25, 53)). In response  
45 to such attack, aerobic micro-organisms have evolved anti-oxidant mechanisms (eg catalase,  
46 peroxidase and superoxide dismutase activities) to protect and adapt cells against ROS excess (6,  
47 18, 24, 65, 68).

48 Although they do not achieve aerobic respiration, anaerobic micro-organisms can be subject to  
49 ROS. The strict anaerobes *Bacteroides fragilis* and *Clostridium perfringens* are able to survive in  
50 the presence of a low amount of oxygen, due to the induction of an adaptative oxidative stress  
51 response that is activated under aerobic conditions (13, 46-49). The peroxide response of  
52 *B. fragilis* shares some characteristics with that of *E. coli* such as the involvement of the redox-  
53 sensitive transcriptional activator OxyR (46) and some antioxidant activities under its control  
54 (47, 49).

55 The lactic acid bacterium (LAB) *Streptococcus thermophilus* is a Gram-positive bacterium  
56 widely used in dairy factories, mainly as a starter of fermentation in yogurt and cheese  
57 production. During fermentation processes, *S. thermophilus* is exposed to oxygen and to its  
58 deleterious effects that can lead to a reduction of cell viability and consequently may have  
59 repercussions on texture, flavor and safety of the final product.

60 *S. thermophilus* is an anaerobic aerotolerant organism that can grow in presence of oxygen  
61 and survive to low concentrations of superoxide and hydroxyl radicals (59). This suggests that  
62 it displays a defense system that provides protection against the effects of ROS. In the  
63 literature, data about its oxidative stress response are scarce. The existence of at least one

64 ROS defense system and its inducibility by hydrogen peroxide are established in  
65 *S. thermophilus* CNRZ368 (58). This organism possesses a single manganese-containing  
66 superoxide dismutase (19) and a glutathione reductase (43). However, no catalase activity was  
67 found. In other streptococci, antioxidant enzymes were reported to play a role in the defense  
68 against oxidative stress (27, 33, 44, 66) and in pathogenesis (45, 67, 69).  
69 The aim of this work was to identify genes involved in the oxidative stress defense of  
70 *S. thermophilus* CNRZ368 without presuming their function. For this purpose a collection of  
71 mutants was created by random insertional mutagenesis and the mutants were screened for  
72 their sensitivity to menadione (a superoxide generating agent). The locus disrupted in 10  
73 *S. thermophilus* menadione sensitive clones was identified and the potential physiological role  
74 of the mutated genes in oxidative stress defense is discussed. The implication of these 10 loci  
75 during heat shock was also evaluated to assess their specificity to oxidative stress defense.  
76 Moreover, the characterization of two of these mutants, disrupted in *sufD* and *iscU* genes  
77 which may participate in Fe-S clusters assembly or repair, is more detailed. The results  
78 presented in this paper indicate that the inactivation of *sufD* and *iscU* genes caused a higher  
79 sensitivity to superoxide stress and suggest that Fe-S cluster repair or assembly could play an  
80 important role in the oxidative stress defense of anaerobic bacteria.

81

82

## MATERIALS AND METHODS

83 **Bacterial strains, growth conditions.** *S. thermophilus* CNRZ368 was purchased from the  
84 INRA-CNRZ strain collection (Jouy en Josas, France). Depending on the experiments,  
85 *S. thermophilus* CNRZ368 and its derivatives were cultivated at 30°C or 42°C in milk  
86 medium (for pre-cultures), TPPY (12) or M17 (55) media, without shaking. Erythromycin was  
87 added at 2 µg.ml<sup>-1</sup>, when required. Streptonigrin sensitivity was monitored using a TPPYC  
88 medium which contained a reduced concentration of metallic dications (including Fe<sup>2+</sup>).  
89 TPPYC was performed according to the protocol described by Visca *et al.* (62) and modified as

90 follow: Chelex beads (20 g/l, sigma) were added to TPPY medium and stirred at 4°C for 48 h  
91 and the medium was then supplemented with 1 mM MgCl<sub>2</sub> and 1 mM CaCl<sub>2</sub>.  
92 Recombinant plasmids were transformed into *E. coli* EC101, a TG1 strain [*supE hsd-5 thi Δ(lac-*  
93 *proAB)* F' (*traD6 proAB<sup>+</sup> lacI<sup>f</sup> lacZΔM15*)] containing a chromosomal copy of the pWV01 *repA*  
94 gene (34) and selected at 37°C on Luria-Bertani (LB) (50) agar plates containing 150 µg/ml of  
95 erythromycin.

96

97 **Insertional mutagenesis.** The collection of mutants was made by insertional mutagenesis using the  
98 plasmid pGh9:ISSI that carries Em<sup>R</sup> gene (35). The mutagenesis procedure and the cloning of the  
99 fragments flanking the pGh9:ISSI insertion locus were carried out on *S. thermophilus* CNRZ368  
100 genome, as previously described (56).

101

102 **Selection of mutants impaired in their ability to tolerate menadione.** For selection of  
103 menadione sensitive clones, two different procedures, differing mainly by the length of the  
104 menadione-exposure, were assayed. Both assays were preceded by a pre-culture in milk medium  
105 containing 2 µg/ml erythromycin: pre-cultures of the 2112 mutants were performed in 96-well  
106 microtitre plates.

107 For “long” time menadione exposure, each pre-culture was transferred, using a hedgehog, onto  
108 M17 agar plates containing 2 µg/ml erythromycin and increasing concentrations of menadione  
109 (from 0 µg/ml to 120 µg/ml). After 20h incubation at 42°C, the presence (or the absence) of  
110 colonies was observed.

111 For “short” time menadione exposure, each pre-culture was diluted (with a 1/100 ratio) in 5  
112 different 96-well microtitre plates containing liquid TPPY medium supplemented with 2 µg/ml  
113 erythromycin (in final volume of 100µl, resulting in an approximately 1/1 liquid to air space  
114 ratio). After a 3h30 growth period at 42°C, 5 different concentrations of menadione (from  
115 0 mg/ml to 15 mg/ml) were added to each of the 5 microtitre plates. Microtitre plates were

116 incubated at 42°C for 3h. The cultures were then transferred on TPPY agar plates containing  
117 erythromycin (2 µg/ml) using a hedgehog. The ability of the mutants to survive to the menadione  
118 stress was assessed, after 20 h incubation at 42°C, by observation of the presence or the absence  
119 of colonies.

120 For each experiment, clones were considered as sensitive when they did not grow on a plate  
121 where more than 90% of the clones were growing. Both procedures were applied twice and  
122 the results had been compiled to reveal the mutants raising the more severe phenotype.

123

124 **DNA manipulations.** Preparation of chromosomal, plasmid and recombinant DNA and  
125 Southern analysis were performed according to standard protocols (50). All cloning junctions  
126 and cloned PCR products were checked by sequencing using Dye-Terminator chemistry on an  
127 ABI Prism 377 Genetic Analyser (PE Biosystems). Sequence data were analysed using  
128 Genbank, BLAST (2, 3), TMHMM (51) and HMMTOP (61) software. For all analysed  
129 sequences, The DNA sequences reported in this paper have been deposited in GenBank under  
130 accession numbers AY386239, AY386240, AY386241, AY386242, AY386243, AY386244,  
131 AY386245.

132

133 **Construction of the *ΔsufD* and *iscU<sub>97</sub>* mutants.** Oligonucleotides used to construct the  
134 *ΔsufD* and *iscU<sub>97</sub>* mutants were purchased from MWG Biotech AG (Ebersberg, Germany). To  
135 construct non-polar truncated *ΔsufD* mutant, the 5' and 3' ends of the *sufD* gene were  
136 independently amplified by PCR. The 5' end was amplified by using the primers  
137 GAGCGTGTT**GAATTC**ACCG containing an *EcoRI* site, symbolized in bold characters  
138 and AACTTAACT**GCAGC**ACCGTCT containing a *PstI* site. The 3' end was amplified by  
139 using the primers GGTGTTCCAGAATTC**AAATAACC** containing an *EcoRI* site and  
140 TGGTCTGTCATCAGGTCG. This last primer does not contain any *PstI* site, but the PCR  
141 product generated does contain a *PstI* restriction site. These two PCR products were digested

142 by *Pst*I and ligated with each other. The resulting fragment was amplified by PCR, then  
143 digested by *Eco*RI and cloned into pGh9. The final construct (pGh9 $\Delta$ *su*fD) was used to  
144 transform *S. thermophilus* CNRZ368. For IscU protein inactivation, the strategy consisted in  
145 replacing the TGT codon encoding the conserved Cys<sub>97</sub> by a TAT codon encoding a Tyr. This  
146 was achieved by replacing, in oligonucleotides, the 3 underlined nucleotides of the sequence  
147 (AGG TGT ACA) generating an *Eco*RV restriction site (AGA TAT CCA). Thus, the three  
148 amino-acids Gly<sub>96</sub> Cys<sub>97</sub> Tyr<sub>98</sub> were replaced by Arg<sub>96</sub> Tyr<sub>97</sub> Thr<sub>98</sub> in the mutated protein. Two  
149 internal and complementary primers containing the mutation  
150 (GCTGGTGATGATATCACCATTTCACA and  
151 GAAATGGTGATATCATCACCAGCAAAGG) were used with two additional external  
152 primers (AAGCGGAATTCAAGCATCCACTCAGGC containing an *Eco*RI site and  
153 TGTGACAAGCTTGTAGAGGCAATTCTT containing an *Hind*III site), respectively, to  
154 amplify 5' and 3' end of *iscU*. The two PCR products of 515 bp and 262 bp were digested by  
155 *Eco*RI and *Eco*RV and by *Hind*III and *Eco*RV respectively before being ligated to each other.  
156 The resulting fragment was cloned into pGh9 digested by *Eco*RI and *Hind*III. The final  
157 construct, (pGh9*iscU*<sub>97</sub>) was used to transform *S. thermophilus* CNRZ368. Preparation of  
158 *S. thermophilus* competent cells was performed as previously described by O'sullivan and  
159 Fitzgerald (41). The following gene replacements were done as previously described (8) but  
160 in TPPY medium and at 30°C/42°C as permissive and restrictive temperature, respectively.

161

## 162 **Physiological characterization of *S. thermophilus* mutant strains.**

163 **Exposure to methyl viologen.** For “short” time exposure, cultures of CNRZ368 and its  
164 derivative strains were grown in TPPY medium. At an OD<sub>600</sub> of 0.6, cultures were divided into  
165 two subcultures (in a ratio of liquid to air space of approximately 1/1). Methyl viologen  
166 (paraquat; Sigma-Aldrich) was added to one of them with a final concentration of 50 mM. After  
167 3 h incubation at 42°C, cultures were diluted to appropriate concentrations and spread onto



168 TPPY agar plates. For “long” time methyl viologen exposure, strains were grown in TPPY  
169 medium until OD<sub>600</sub> of 0.6. Cultures were then diluted to a final theoretical OD<sub>600</sub> of 0.001 in  
170 TPPY medium (in a ratio of liquid to air space of approximately 5/6) supplemented with 1.7 mM  
171 methyl viologen, or not for control experiment. After 18h incubation at 42°C, cultures were  
172 diluted in TPPY to appropriate concentrations and spread onto TPPY agar plates. In both  
173 procedures, methyl viologen sensitivity was estimated by comparing CFU/ml obtained with  
174 treated cultures, with respect to non treated cultures.

175

176 **Heat shock.** The ability of *S. thermophilus* and its derivatives to respond to a heat stress was  
177 assessed as follow. Each strain is grown at 42°C in TPPY medium until early log phase  
178 (OD<sub>600</sub> = 0.3). Then the culture was divided into two. One part was kept at 42°C (control  
179 experiment); the other one was immediately transferred to 60°C for 15 min. Appropriate  
180 dilutions were spread on TPPY agar plates. The fraction of surviving cells was calculated by  
181 dividing the surviving CFU/ml by the count for the control experiment.

182

183 **Streptonigrin sensitivity.** Streptonigrin sensitivity was monitored on cells grown overnight  
184 in TPPYC medium at 42°C. Cells were harvested by centrifugation, washed twice in TPPYC  
185 medium, resuspended in TPPYC, diluted to OD<sub>600</sub> = 0.01 and cultivated in TPPYC for 9 h at  
186 42°C. Then, the cultures were diluted again to OD<sub>600</sub> = 0.005 and divided into 4 parts which  
187 were all incubated for 21 h at 42°C. One part was supplemented with 0.4% (v/v) dimethyl  
188 formamide (control experiment), the 3 other parts were supplemented with different  
189 concentrations of streptonigrin (1 µg/ml, 2 µg/ml and 4 µg/ml dissolved in dimethyl  
190 formamide). As a control experiment, cultures were also carried out in TPPYC supplemented  
191 with 1 mM of 2'2-dipyridyl (an Fe chelator). In each experiment, OD<sub>600</sub> value was used to  
192 estimate the cellular density after 21h incubation. Relative optic density was determined by

193 comparing cellular density obtained with and without streptonigrin. All experiments were  
194 performed in triplicate.

195

196

## RESULTS

### 197 **1. Selection of *S. thermophilus* CNRZ368 mutants with altered oxidative stress response.**

198 A collection of 2112 mutants of *S. thermophilus* CNRZ368 was generated by insertional  
199 mutagenesis using the plasmid pGh9:ISS1 (35, 56). This plasmid carries an erythromycin  
200 resistance gene for its selection and the insertional element ISS1 that allows random  
201 integration of the plasmid into *S. thermophilus* genome. The 2112 mutants obtained were  
202 screened to select clones sensitive to menadione (a superoxide-generating compound). Two  
203 distinct selection procedures were used to identify mutants susceptible to menadione (see  
204 materials and methods). The characterization of 10 mutants, disrupted in loci originally named  
205 *ossA* to *ossJ* (for oxidative stress sensitive locus) is described in this paper.

206

### 207 **2. Sensitivity of *S. thermophilus* CNRZ368 mutants to methyl viologen.** Previously, we

208 demonstrated that *ossB* and *ossC* mutants (previously named 6G4 and 18C3, respectively) are  
209 compromised in their ability to survive superoxide stress (57). To confirm that the other 8  
210 mutants selected were sensitive to superoxide stress, their sensitivity to methyl viologen was  
211 examined (Fig. 1). In this experiment, we used the non-polar mutants *ΔsufD* and *iscU<sub>97</sub>*  
212 constructed in this work (see further) rather than the potential polar clones *ossI* and *ossJ*.  
213 Short-term methyl viologen exposure revealed the extreme sensitivity of *ossG*, as well as the  
214 4 to 5-fold increased sensitivity of *ossD*, *ossH* and *iscU<sub>97</sub>* mutants (Fig. 1A). Although less  
215 severe the *ΔsufD* mutant exhibited about 2-fold less survival following such stress. Under  
216 these conditions, the other 3 mutants (*ossA*, *ossE* and *ossF*) did not exhibit any significant  
217 sensitivity, however long term methyl viologen exposure confirmed their sensitivity. Indeed,  
218 long term methyl viologen exposure (Fig. 1B) showed that the *ossA* mutant strain displayed a

219 relative survival more than 10-fold lower than that of the wild type strain. The *ossE* and *ossF*  
220 mutants also showed a 2-fold lower growth compared to the wild type strain.

221

222 **3. Survival of the mutants exposed to heat stress.** To assess the specificity of these 10 loci  
223 to oxidative stress defense, the mutants ability to survive to heat shock was estimated. Five  
224 out of the 10 mutants (*ossA*, *ossB*, *ossC*, *ossD*, and *ossJ*) isolated as oxidative stress sensitive  
225 clones also showed a reduced survival to heat shock compared with WT (Fig. 2). These data  
226 suggest that the genes impaired in these mutants are not specific to oxidative stress defense  
227 process but are probably involved in a more global response to stresses.

228 In contrast, the *ossE*, *ossF* and *ossG* mutants did not differ significantly from the wild type,  
229 indicating that the genes disrupted in these strains did not influence cells ability to respond to  
230 heat shock. Interestingly, the *ossG* mutant, that was highly sensitive to methyl viologen,  
231 showed no significant decrease of survival to heat shock compared with WT, suggesting that  
232 *ossG* gene, essential for oxidative stress defense, could be specific to such a stress.

233

234 **4. Identification of genes implicated in oxidative stress tolerance.** The disrupted locus was  
235 identified for each of the 10 mutants by cloning and sequencing the junctions as previously  
236 described (56). All of the insertions were mapped within coding sequences. Sequence  
237 similarity searches and analysis revealed similar sequences in other bacteria (see Table 1)  
238 with an E-value lower than  $10^{-10}$  for BLASTX searches. In 9/10 cases, data about the function  
239 of these genes were available. To facilitate their description, these genes were grouped into 5  
240 classes (from I to V) according to the putative function and renamed when relevant (Table 1).

241

242 **Class I: Genes potentially involved in cell wall structure and cell shape determination.**

243 The *ossA*, *ossB* and *ossC* putative gene products shared sequence identity with MreD, RodA  
244 and Pbp2b, respectively, which putatively direct the elongation of the lateral cell wall in

245 *E. coli*, conferring the cell its characteristic rod shape (5, 28, 29). The *ossA* mutant was  
246 defective in a gene encoding a protein, with 30% identity with MreD of *Streptococcus*  
247 *mutans*. Sequence analysis of the surrounding region indicated that *S. thermophilus mreD*  
248 gene (*ossA*) was clustered with a gene encoding a putative protein similar to MreC. Moreover,  
249 hydrophopathy profiles of MreD (OssA) determined by using TMHMM and HMMTOP  
250 showed 5 regions of sufficient length and hydrophobicity to be membrane-spanning segments  
251 (data not shown), suggesting that this protein is the counterpart of MreD. The relatively low  
252 identity (30%) observed between *S. thermophilus* and *S. mutans* MreD was in agreement with  
253 the general idea that the structure of this protein is more highly conserved than its primary  
254 sequence (17). To study the effect of *mreD* disruption in *S. thermophilus*, TPPY cultures of  
255 the wild type and *mreD* mutant strains were observed under light microscopy. In contrast to  
256 wild type cells, *mreD* mutant cells formed very long chains, up to 180 cells per chain (Fig. 3).  
257 Moreover, a reduction of about 25% of the cell size was observed in the mutant. Growth of  
258 the *mreD* mutant and of the wild type in TPPY medium were monitored by measuring the  
259 OD<sub>600</sub>. Both strains displayed a similar doubling time of 30 min, but the maximal OD<sub>600</sub> of the  
260 *mreD* mutant (OD<sub>max</sub> = 1.7 ± 0.1) was lower than that of the wild type (OD<sub>max</sub> = 2.4 ± 0.2)  
261 showing that the *mreD* mutant entered stationary phase earlier than wild type cells. The  
262 characterization of *rodA* and *pbp2b* mutants of *S. thermophilus* CNRZ368 is described in a  
263 previous paper (57).

264

265 **Class II: Gene potentially involved in exopolysaccharide translocation.** The sequence  
266 disrupted in the mutant *ossD* was partially determined. Its analysis indicated that the  
267 C-terminal portion of the predicted encoded protein showed 98% identity to CpsX, a putative  
268 membrane protein from *S. thermophilus* MR-1C. The role of CpsX is still unknown but the  
269 conserved location of *cpsX* near *eps* genes of *S. thermophilus* (14) and the presence of a  
270 putative gene related to an ABC transport system in its vicinity argue in favor of CpsX being

271 involved in exopolysaccharide translocation (15). Moreover, the TMHMM and HMMTOP  
272 software predictions indicated that CpsX (OssD) from *S. thermophilus* CNRZ368 would  
273 contain highly hydrophobic domains compatible with a location in the cytoplasmic  
274 membrane. Although the *cpsX* function is not known yet, our work indicate that its disruption  
275 is associated with sensitivity not only to superoxide stress but also to heat shock.

276

277 **Class III: Genes involved in RNA maturation.** In the *ossE* and *ossF* mutants, the plasmid  
278 inserted into genes whose predicted products share sequence identity with enzymes involved  
279 in RNA maturation. The *ossE* gene encodes a putative tRNA guanine transglycosylase (TGT)  
280 which shares 92% identity with the putative *Streptococcus pneumoniae* TGT. TGT enzyme  
281 takes part in the process which removes the guanine residue at position 34 (wobble position of  
282 the anticodon loop) of tRNA<sup>His</sup>, tRNA<sup>Asp</sup>, tRNA<sup>Asn</sup> and tRNA<sup>Tyr</sup> and replaces it by the  
283 modified residue queuosine (39, 40).

284 *ossF* potentially encodes a protein that shows 77 % identity with a putative RNA  
285 methyltransferase of *Streptococcus agalactiae* (Gbs0448) belonging to the TrmA family.  
286 Enzymes of the TrmA family catalyze the formation of the 5-methyluridine in position 54  
287 (localized in the TΨC loop) of tRNA (9). The presence of a TRAM-like domain (RNA  
288 binding domain) in the N-terminal part of OssF supports the idea of an interaction between  
289 this protein and RNA.

290

291 **Class IV: Gene of unknown function.** The *ossG* product shows 67% identity with the  
292 product of a gene of unknown function (*smu.1428c*) of *S. mutans* (Table 1). Although *ossG*  
293 function is not known yet, the mutation of this gene is now associated with a sensitivity  
294 phenotype to superoxide stress. The TMHMM and HMMTOP software predictions suggest  
295 the existence of two transmembrane domains in the N-terminal region of OssG, making of  
296 OssG a potential membrane protein.

297

298 **Class V: Genes involved in iron metabolism.**

299 **(i) Ferric iron transport.** The putative product of *ossH* shows 55% identity with FatD from  
300 *S. pneumoniae*, a potential membrane spanning permease of an Fe<sup>3+</sup> ABC-transporter. Analysis  
301 of the adjacent sequence reveals the presence of a gene encoding a putative FatC similar protein  
302 downstream of *fatD* (*ossH*), in the same orientation. The *fatC* gene is also clustered with *fatD* in  
303 *S. pneumoniae* genome. Hydropathy profile of FatD (OssH) from *S. thermophilus*, assessed with  
304 TMHMM and HMMTOP, reveals that this protein displays ten putative membrane spanning  
305 segments compatible with its expected cytoplasmic membrane location.

306

307 **(ii) [Fe-S] cluster biosynthesis.** The sequence affected in the *ossI* mutant encodes a product  
308 that shows 75% identity with Sag0142, a putative protein deduced from the complete genome  
309 sequence of *S. agalactiae* and similar to SufD-like proteins. Analysis of totally sequenced  
310 microbial genomes reveals that *sufD*-like genes are conserved in bacteria and also in some  
311 archaea such as *Halobacterium* sp. NRC-1, *Thermoplasma acidophilum* and *Aeropyrum*  
312 *pernix* (23). In *E. coli* and *Erwinia chrysanthemi*, the *sufD* gene belongs to the *sufABCDSE*  
313 cluster identified as being involved in iron metabolism and more precisely in [Fe-S] cluster  
314 biogenesis and/or repair (37, 42). The *ossJ* gene product shows sequence identity to the  
315 *gbs0140* product from *S. agalactiae* (Table 1), probably encoding a IscU protein, a NifU  
316 family protein. NifU proteins have been proposed to serve as scaffold proteins in [Fe-S]  
317 cluster assembly in *Synechocystis* PCC6803 and would function as a mediator for iron-sulfur  
318 cluster delivery (38).

319 In some genomes including that of *S. pyogenes*, *nifU*-like genes are clustered with the *suf*  
320 genes (23). To determine if *sufD* (*ossI*) and *iscU* (*ossJ*) from *S. thermophilus* were genetically  
321 linked together, the chromosomal DNA regions flanking the pGhost9:ISSI insertion site of  
322 both mutants were sequenced until overlapping. Examination of the genes in the

323 neighborhood of *sufD* and *iscU* revealed that they both belong to a cluster of 5 genes all  
324 orientated in the same direction. This cluster, named *sufCDS iscU sufB* cluster, is shown in  
325 Fig. 4. The highest sequence identities of these gene products are found with putative proteins  
326 from *S. agalactiae* (Table 2). They also share sequence similarity with proteins encoded by  
327 *E. coli* and *Er. chrysanthemi* *suf* gene clusters. The putative SufC, SufD, SufS and SufB  
328 proteins encoded by the *S. thermophilus* cluster present respectively 72%, 48%, 65%, and  
329 57% similarity to their *E. coli* counterparts. Moreover, IscU presents 47% similarity with IscU  
330 from *E. coli*.

331 Sequences corresponding to promoter regions were searched by inspection of the total  
332 sequence of the *sufCDS iscU sufB* cluster. A unique extended -10 sequence  
333 (5'-TTTGGTATAAT-3') is found upstream *sufC*, the first ORF of the cluster. A potential  
334 rho-independent transcriptional terminator is identified 63 bp downstream *sufB*, the last ORF  
335 of the cluster ( $\Delta G = -10$  kcal/mol). These features, together with the short or missing  
336 intergenic regions and the same orientation of the genes, suggest that these 5 ORFs form an  
337 operon, as it is the case for the *Er. chrysanthemi* *suf* locus.

338

## 339 **5. Molecular characterization of the *sufCDS iscU sufB* cluster.**

340 **5.1. Construction of  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutants.** Since *S. thermophilus* *suf* genes are  
341 potentially clustered in an operon, non polar  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutants were constructed and  
342 analysed. The *sufD* mutation consists of an internally in frame deletion leading to the loss of  
343 43% of the whole protein (see materials and methods and Fig. 4). Sequence comparison of  
344 NifU family proteins from various organisms indicates that *S. thermophilus* IscU protein  
345 contains the three highly conserved cystein residues that have been shown to be the assembly  
346 site of a "transitory" [2Fe-2S] center within NifU (70). Thus, the construction of a null *iscU*  
347 mutant was achieved by replacing one of these conserved residues, the Cys<sub>97</sub>, by a Tyr (see

348 materials and methods and Fig. 4). Sequencing of these two constructs confirmed that they  
349 did not contain any secondary mutation.

350

## 351 **5.2. Role of *sufD* and *iscU* in iron metabolism.**

352 To investigate the possibility that  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutations impair growth due to the lack  
353 of functional [Fe-S] cluster biosynthesis, the mutants were grown in TPPY medium and  
354 compared to wild type. Growth of the  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutants showed no significant  
355 difference compared to the WT strain. Assuming that the high iron concentration of the TPPY  
356 medium could allow self assembling of the [Fe-S] clusters (71), and consequently could be  
357 responsible for the absence of difference in growth between strains, the same experiment was  
358 conducted in TPPYC, a divalent cation depleted medium. Under these conditions, the mutants  
359 displayed a growth rate similar of that of the WT strain but their maximal OD<sub>600</sub> was lower  
360 than that of the wild type, showing their premature entry in stationary phase (Table 3).  
361 Moreover, this difference disappeared when the cultures were grown on TPPYC  
362 supplemented by 100  $\mu$ M of FeSO<sub>4</sub>. Taken together, these results suggest that (i) *iscU* and  
363 *sufD* genes of *S. thermophilus* are involved in iron metabolism and (ii) their presence would  
364 be of particular importance under iron-limiting conditions.

365 To confirm the involvement of *sufD* and *iscU* in iron metabolism, the iron intracellular  
366 concentration of  $\Delta$ *sufD* and *iscU*<sub>97</sub> was checked by measuring the sensitivity of these strains  
367 to streptonigrin (an iron-activated antibiotic). As shown in Fig. 5 A, the *iscU* mutant was more  
368 sensitive to streptonigrin than the wild type strain whereas the  $\Delta$ *sufD* mutant was more  
369 resistant. Moreover, addition of 2,2'-dipyridyl (a ferrous iron chelator) abolished the  
370 difference of growth between the strains (Fig. 5 B) indicating that iron was responsible for the  
371 streptonigrin sensitivity observed without 2,2'-dipyridyl. Taken altogether these results  
372 suggest that both *iscU* and *sufD* of *S. thermophilus* play a role in iron metabolism.

373



## Discussion

374

375 To investigate the oxidative stress defense system of *S. thermophilus*, a collection of random  
376 insertion mutants was screened to select clones with high sensitivity to menadione. Using this  
377 approach, we aimed to identify genes involved in superoxide stress defense without  
378 presuming their function. It can be noted that no gene encoding a protein considered, in other  
379 organisms, as major antioxidant enzyme were found using this approach. This can be due to  
380 the fact that (i) the collection of mutants was non-exhaustive (ii) the corresponding mutant  
381 could have a weak phenotype (iii) such mutations could be lethal (iv) except from SOD, most  
382 anti-oxidant enzymes are protective against H<sub>2</sub>O<sub>2</sub> and not O<sub>2</sub><sup>-</sup>. The genes identified in the  
383 present work were classified into 5 groups according to their function: genes potentially  
384 involved in cell wall metabolism, exopolysaccharide translocation, tRNA modification, and  
385 iron metabolism and gene of unknown function.

386 Perhaps the most striking observation of this study is that 5 out of the 10 sensitive mutants  
387 selected are disrupted in a gene encoding an membrane protein (OssH, MreD, RodA, PBP2b  
388 and CpsX). This could reflect that the cell envelope is the main target of the stress applied.  
389 Among these mutants, three (*mreD*, *rodA* and *pbp2b*) are impaired in genes potentially  
390 encoding proteins described in *E. coli* as playing a role in peptidoglycan biosynthesis during  
391 cell elongation (28, 52, 63). Phenotypic analysis of these three *S. thermophilus* mutants  
392 showed that they all present a reduced cell size associated with a modification of the cell  
393 shape, as observed in *rodA*, *pbp2* and *mre* mutants of *E. coli* or *Salmonella typhimurium* (5,  
394 20, 28, 29, 63, 64). Moreover, the 3 *S. thermophilus* mutants also presented an increased  
395 sensitivity to heat shock compared to the wild type strain suggesting that the protective role of  
396 MreD, RodA and PBP2b is not specific to oxidative stress but could be enlarged to a more  
397 global stress response. In support of this hypothesis is the existence in *Lactococcus lactis* of  
398 mutations in *pbp* genes that negatively affect acid tolerance of a *guaA* mutant (16).

399 The major function assigned to cell wall is the maintenance of cell integrity by protecting  
400 cells not only against osmotic stress but also against other stress conditions. For instance, in  
401 LAB, it has already been shown that compromised biosynthesis of peptidoglycan results in an  
402 increased sensitivity to either UV (22) or acid (11) stress. Additionally, disruption of *mreD*,  
403 *rodA* or *pbp2b* may cause a modification of the peptidoglycan architecture in *S. thermophilus*,  
404 reducing the cell protection against aggressions such as oxidative and heat stress.

405

406 Transfer RNA modification is a universal feature of all living organisms, although some  
407 described modifications have no assigned function yet. Modifying enzymes affect the tRNA  
408 precursor either by chemical modification of an existing residue (in the base or in the sugar)  
409 or by nucleotide substitution. These modifications, in particular those present in the anticodon  
410 region can affect translation efficiency and fidelity (for a review (10)). In this study, two  
411 clones disrupted in the loci *tgt* and *ossF*, encoding proteins showing high identity with tRNA  
412 modifying enzymes were selected for their high sensitivity to oxidative stress. This sensitivity  
413 could suggest that the expression of genes involved in oxidative stress defense may require  
414 tRNA maturation or, that such modifications could help tRNA stabilization under stress  
415 conditions. In agreement with the former hypothesis *Shigella flexneri* pathogenicity is  
416 regulated by *virF* mRNA translation efficiency under TGT activity control (21). Additionally,  
417 several environmental conditions are suggested to influence tRNA modification: in the case of  
418 queuosine (whose synthesis and substitution is carried out by several enzymes among which  
419 is TGT), synthesis is sensitive to the presence of iron (31) and TGT requires oxygen for its  
420 full activity (54). Therefore, further investigations on this topic should provide an interesting  
421 new regulatory model and could contribute towards the emerging observation that  
422 translational regulation plays an important role in oxidative stress response, a role mainly  
423 ascribed to transcriptional regulation thus far.

424

425 Iron is essential for cellular metabolism, due to its requirement as a cofactor for a large number  
426 of enzymes. However, an excess of iron is toxic because of its involvement in the Fenton  
427 reaction that enhances the formation of ROS. In this work, a gene homologous to *fatD*, encoding  
428 a putative Fe<sup>3+</sup> ABC-transporter was found to be involved in *S. thermophilus* oxidative stress  
429 response. To test whether or not *S. thermophilus* FatD is indeed an iron transporter, the  
430 sensitivity of the *fatD* mutant to streptonigrin (an iron activated-antibiotic) was assessed. No  
431 significant difference between the mutant and the wild type strains was detected (data not  
432 shown) suggesting that disruption of *fatD* would have no effect on intracellular iron  
433 concentration. Thus, either FatD is not a major iron transporter under the experimental procedure  
434 we used or *S. thermophilus fatD* encodes an ABC transporter for another metallic cation. A  
435 possible cation could be Mn<sup>2+</sup> known to have a protective function against oxidative damage  
436 either via direct dismutase activity, such as reported in other organisms (4, 26, 60), or via its role  
437 as a cofactor of the only SOD reported into *S. thermophilus* (19). Thus, inactivation of a Mn<sup>2+</sup>  
438 ABC transporter could trigger sensitivity to oxidative stress.

439

440 [Fe-S] clusters are oxygen sensitive prosthetic groups, incorporated into proteins via covalent  
441 bonds to cysteines. Iron-sulfur proteins, which are found in more than 120 classes of enzymes  
442 and proteins (30), are known to play important physiological roles in electron transfer,  
443 metabolic reactions and in transcriptional regulation (30, 32). Indeed, because of their  
444 intrinsic redox properties, they are likely to sense oxygen or its derivatives as it is the case in  
445 SoxR and FNR transcriptional regulators, responding to oxidative stress and oxygen  
446 availability, (for a review (32)). IscR is also a iron-sulfur transcriptional regulator that is  
447 suggested to respond to Fe-S cluster limiting conditions and potentially to hydrogen peroxide  
448 in *E. coli* (32). Although it has been shown that [Fe-S] cluster assembly can occur  
449 spontaneously *in vitro* under anaerobic conditions (36), this process can also be catalyzed  
450 enzymatically *in vivo*. Genes of the *suf* operon and of the *isc* cluster (iron sulfur cluster) are

451 conserved and encode [Fe-S] cluster biosynthesis and/or repair pathways (23). More  
452 precisely, it has been recently proposed that Isc proteins are necessary for biosynthesis, while  
453 Suf proteins are essentially involved in repair of oxidatively damaged [Fe-S] clusters (37).  
454 Study of  $\Delta$ *sufD* and *iscU*<sub>97</sub> non-polar mutants indicates that SufD and IscU proteins are  
455 required for growth in an iron-deficient medium. SufD and IscU inactivation conferred an  
456 increased resistance and sensitivity respectively to streptonigrin. These results indicate that  
457 *S. thermophilus* SufD and IscU are involved in intracellular iron balance. Additionally, SufD  
458 and IscU of *S. thermophilus*, similar to SufD and NifU family proteins from other  
459 microorganisms, are encoded by genes clustered with *sufB*, *sufC* and *sufS* homologues, also  
460 potentially involved in [Fe-S] cluster repair. Hence the participation of *S. thermophilus* SufD  
461 and IscU proteins in iron intracellular balance is probably the result of their involvement in  
462 [Fe-S] cluster repair, as in *Er. chrysanthemi* (37). The NifU family protein function is to  
463 provide a scaffold for assembly of clusters mediated by NifS activity, and such clusters can  
464 then be transferred into apo forms of [Fe-S] cluster containing proteins (1, 70). The role of the  
465 SufD in [Fe-S] cluster biosynthesis is not well established. However, in *S. thermophilus*, the  
466 increased resistance to streptonigrin of the  $\Delta$ *sufD* mutant suggests that its intracellular iron  
467 concentration is lower than in the wild type. At least two alternative hypotheses can be drawn:  
468 (i) SufD could be directly involved in the uptake or assimilation of extracellular iron, (ii)  
469 SufD could inhibit the synthesis and/or repair of the [Fe-S] clusters. In this latter case, SufD  
470 inactivation would result in an increase synthesis of [Fe-S] clusters, and consequently in a  
471 reduced intracellular free iron concentration.  
472 Additionally, *S. thermophilus* *sufD* and *iscU* mutants displayed an increased sensitivity to  
473 superoxide radicals compared to wild type. Thus, SufD and IscU are likely to play a major  
474 role in oxidative stress defense, probably resulting from their predicted role in [Fe-S] cluster  
475 assembly and/or repair. The involvement of the *suf* operon in oxidative stress resistance  
476 during the early steps of infection by *Er. chrysanthemi* has also been reported (37), as well as

477 the induction of the *suf* gene cluster of *E. coli* by the hydrogen peroxide response regulator  
478 OxyR (72). In spite of a possible spontaneous formation, our data suggest that under oxidative  
479 stress, [Fe-S] cluster assembly and/or repair becomes a limiting step in the formation of iron-  
480 sulfur proteins, and that the cell requires an additional, dedicated system, encoded by the *suf*  
481 gene cluster, to optimize [Fe-S] cluster formation or repair.

482

483 Conclusion: The strategy used in this work allowed the identification of genes involved in  
484 *S. thermophilus* oxidative stress defense system. Data concerning the function of some loci is  
485 already available in the literature, whereas the role of others is not known. In both cases, the  
486 exact roles of these genes in the defense of *S. thermophilus* against oxidative stress remains to  
487 be elucidated.

## ACKNOWLEDGEMENTS

488

489

490 A. T and A. F. were supported by grants from the Ministère de la recherche. F. B. was  
491 supported by a grant from the Institut National de la Recherche Agronomique.

492 We are grateful to Paul Hoskinson for his advice regarding the English formulation of the  
493 manuscript.

494

## References

495

496

- 497 1. **Agar, J. N., C. Krebs, J. Frazzon, B. H. Huynh, D. R. Dean, and M. K. Johnson.**  
498 2000. IscU as a scaffold for iron-sulfur cluster biosynthesis: sequential assembly of  
499 [2Fe-2S] and [4Fe-4S] clusters in IscU. *Biochemistry* **39**:7856-7862.
- 500 2. **Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman.** 1990. Basic  
501 local alignment search tool. *J Mol Biol* **215**:403-410.
- 502 3. **Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and**  
503 **D. J. Lipman.** 1997. Gapped BLAST and PSI-BLAST: a new generation of protein  
504 database search programs. *Nucleic Acids Res* **25**:3389-3402.
- 505 4. **Archibald, F. S., and I. Fridovich.** 1981. Manganese, superoxide dismutase, and  
506 oxygen tolerance in some lactic acid bacteria. *J Bacteriol* **146**:928-36.
- 507 5. **Begg, K. J., and W. D. Donachie.** 1985. Cell shape and division in *Escherichia coli*:  
508 experiments with shape and division mutants. *J Bacteriol* **163**:615-622.
- 509 6. **Benov, L. T., and I. Fridovich.** 1994. *Escherichia coli* expresses a copper- and zinc-  
510 containing superoxide dismutase. *J Biol Chem* **269**:25310-25314.
- 511 7. **Berlett, B. S., and E. R. Stadtman.** 1997. Protein oxidation in aging, disease, and  
512 oxidative stress. *J Biol Chem* **272**:20313-20316.

- 513 8. **Biswas, I., A. Gruss, S. D. Ehrlich, and E. Maguin.** 1993. High-efficiency gene  
514 inactivation and replacement system for gram-positive bacteria. *J Bacteriol* **175**:3628-  
515 3635.
- 516 9. **Bjork, G. R.** 1975. Transductional mapping of gene *trmA* responsible for the  
517 production of 5-methyluridine in transfer ribonucleic acid of *Escherichia coli*. *J*  
518 *Bacteriol* **124**:92-98.
- 519 10. **Bjork, G. R., J. M. Durand, T. G. Hagervall, R. Leipuviene, H. K. Lundgren, K.**  
520 **Nilsson, P. Chen, Q. Qian, and J. Urbonavicius.** 1999. Transfer RNA modification:  
521 influence on translational frameshifting and metabolism. *FEBS Lett* **452**:47-51.
- 522 11. **Boyd, D. A., D. G. Cvitkovitch, A. S. Bleiweis, M. Y. Kiriukhin, D. V. Debabov, F.**  
523 **C. Neuhaus, and I. R. Hamilton.** 2000. Defects in D-alanyl-lipoteichoic acid  
524 synthesis in *Streptococcus mutans* results in acid sensitivity. *J Bacteriol* **182**:6055-  
525 6065.
- 526 12. **Bracquart, P.** 1981. An agar medium for the differential enumeration of  
527 *Streptococcus thermophilus* and *Lactobacillus bulgaricus* in yogurth. *J appl bacteriol*  
528 **51**:303-305.
- 529 13. **Briolat, V., and G. Reyssset.** 2002. Identification of the *Clostridium perfringens* genes  
530 involved in the adaptive response to oxidative stress. *J Bacteriol* **184**:2333-2343.
- 531 14. **Broadbent, J., D. McMahon, C. Oberg, and D. Welker.** 2001. Use of  
532 exopolysaccharide-producing cultures to improve the functionality of low fat cheese.  
533 *Int Dairy J* **11**:433-439.
- 534 15. **Broadbent, J., D. McMahon, D. Welker, C. Oberg, and S. Moineau.** 2003.  
535 Biochemistry, genetics, and applications of exopolysaccharide production in  
536 *Streptococcus thermophilus*: a review. *J Dairy Sci* **86**:407-423.

- 537 16. **Budin, A., V. Pichereau, E. Domakova, J. Tremblay, R. Dervyn, E. Auffray, S. D.**  
538 **Ehrlich, S. Kulakauskas, and E. Maguin.** 2002. Effect of a *guaA* mutation on the  
539 acid tolerance of *L. lactis*. *Sciences des aliments* **22**:67-74.
- 540 17. **Burger, A., K. Sichler, G. Kelemen, M. Buttner, and W. Wohlleben.** 2000.  
541 Identification and characterization of the *mre* gene region of *Streptomyces coelicolor*  
542 A3(2). *Mol Gen Genet* **263**:1053-1060.
- 543 18. **Chae, H. Z., K. Robison, L. B. Poole, G. Church, G. Storz, and S. G. Rhee.** 1994.  
544 Cloning and sequencing of thiol-specific antioxidant from mammalian brain: alkyl  
545 hydroperoxide reductase and thiol-specific antioxidant define a large family of  
546 antioxidant enzymes. *Proc Natl Acad Sci U S A* **91**:7017-7021.
- 547 19. **Chang, S. K., and H. M. Hassan.** 1997. Characterization of superoxide dismutase in  
548 *Streptococcus thermophilus*. *Appl Environ Microbiol* **63**:3732-3735.
- 549 20. **Costa, C. S., and D. N. Anton.** 1993. Round-cell mutants of *Salmonella typhimurium*  
550 produced by transposition mutagenesis: lethality of *rodA* and *mre* mutations. *Mol Gen*  
551 *Genet* **236**:387-394.
- 552 21. **Durand, J. M., B. Dagberg, B. E. Uhlin, and G. R. Bjork.** 2000. Transfer RNA  
553 modification, temperature and DNA superhelicity have a common target in the  
554 regulatory network of the virulence of *Shigella flexneri*: the expression of the *virF*  
555 gene. *Mol Microbiol* **35**:924-35.
- 556 22. **Duwat, P., A. Cochu, S. D. Ehrlich, and A. Gruss.** 1997. Characterization of  
557 *Lactococcus lactis* UV-sensitive mutants obtained by *ISS1* transposition. *J Bacteriol*  
558 **179**:4473-9.
- 559 23. **Ellis, K. E., B. Clough, J. W. Saldanha, and R. J. Wilson.** 2001. *Nifs* and *Sufs* in  
560 malaria. *Mol Microbiol* **41**:973-981.
- 561 24. **Faguy, D. M., and W. F. Doolittle.** 2000. Horizontal transfer of catalase-peroxidase  
562 genes between archaea and pathogenic bacteria. *Trends Genet* **16**:196-197.



- 563 25. **Farr, S. B., and T. Kogoma.** 1991. Oxidative stress responses in *Escherichia coli* and  
564 *Salmonella typhimurium*. Microbiol Rev **55**:561-585.
- 565 26. **Fridovich, I.** 1983. Superoxide radical: an endogenous toxicant. Annu Rev Pharmacol  
566 Toxicol **23**:239-57.
- 567 27. **Gibson, C. M., T. C. Mallett, A. Claiborne, and M. G. Caparon.** 2000.  
568 Contribution of NADH oxidase to aerobic metabolism of *Streptococcus pyogenes*. J  
569 Bacteriol **182**:448-55.
- 570 28. **Ishino, F., W. Park, S. Tomioka, S. Tamaki, I. Takase, K. Kunugita, H.**  
571 **Matsuzawa, S. Asoh, T. Ohta, B. G. Spratt, and et al.** 1986. Peptidoglycan  
572 synthetic activities in membranes of *Escherichia coli* caused by overproduction of  
573 penicillin-binding protein 2 and RodA protein. J Biol Chem **261**:7024-7031.
- 574 29. **Iwaya, M., R. Goldman, D. J. Tipper, B. Feingold, and J. L. Strominger.** 1978.  
575 Morphology of an *Escherichia coli* mutant with a temperature-dependent round cell  
576 shape. J Bacteriol **136**:1143-1158.
- 577 30. **Johnson, M. K.** 1998. Iron-sulfur proteins: new roles for old clusters. Curr Opin  
578 Chem Biol **2**:173-81.
- 579 31. **Kersten, H., and W. Kersten.** 1990. Biosynthesis and function of queuine and  
580 queuosine tRNAs, p. B69-B108. In C. W. G. a. K. C. T. K. (ed.) (ed.),  
581 Chromatography and modification of nucleosides. Elsevier, Amsterdam.
- 582 32. **Kiley, P. J., and H. Beinert.** 2003. The role of Fe-S proteins in sensing and regulation  
583 in bacteria. Curr Opin Microbiol **6**:181-5.
- 584 33. **King, K. Y., J. A. Horenstein, and M. G. Caparon.** 2000. Aerotolerance and  
585 peroxide resistance in peroxidase and PerR mutants of *Streptococcus pyogenes*. J  
586 Bacteriol **182**:5290-5299.
- 587 34. **Leenhouts, K.** 1995. Integration strategies and vectors. Dev Biol Stand **85**:523-530.

- 588 35. **Maguin, E., H. Prevost, S. D. Ehrlich, and A. Gruss.** 1996. Efficient insertional  
589 mutagenesis in lactococci and other gram-positive bacteria. *J Bacteriol* **178**:931-935.
- 590 36. **Merchant, S., and B. W. Dreyfuss.** 1998. Metalloprotein and Cytochrome Assembly  
591 in Chloroplasts. *Annu Rev Plant Physiol Plant Mol Biol* **49**:25-51.
- 592 37. **Nachin, L., M. El Hassouni, L. Loiseau, D. Expert, and F. Barras.** 2001. SoxR-  
593 dependent response to oxidative stress and virulence of *Erwinia chrysanthemi*: the key  
594 role of SufC, an orphan ABC ATPase. *Mol Microbiol* **39**:960-972.
- 595 38. **Nishio, K., and M. Nakai.** 2000. Transfer of iron-sulfur cluster from NifU to  
596 apoferredoxin. *J Biol Chem* **275**:22615-22618.
- 597 39. **Noguchi, S., Y. Nishimura, Y. Hirota, and S. Nishimura.** 1982. Isolation and  
598 characterization of an *Escherichia coli* mutant lacking tRNA-guanine  
599 transglycosylase. Function and biosynthesis of queuosine in tRNA. *J Biol Chem*  
600 **257**:6544-6550.
- 601 40. **Okada, N., S. Noguchi, H. Kasai, N. Shindo-Okada, T. Ohgi, T. Goto, and S.**  
602 **Nishimura.** 1979. Novel mechanism of post-transcriptional modification of tRNA.  
603 Insertion of bases of Q precursors into tRNA by a specific tRNA transglycosylase  
604 reaction. *J Biol Chem* **254**:3067-3073.
- 605 41. **O'Sullivan, T., and G. Fitzgerald.** 1999. Electrotransformation of industrial strains  
606 of *Streptococcus thermophilus*. *J Appl Microbiol* **86**:275-283.
- 607 42. **Patzer, S. I., and K. Hantke.** 1999. SufS is a NifS-like protein, and SufD is necessary  
608 for stability of the [2Fe-2S] FhuF protein in *Escherichia coli*. *J Bacteriol* **181**:3307-  
609 3309.
- 610 43. **Pébay, M., A. C. Holl, J. M. Simonet, and B. Decaris.** 1995. Characterization of the  
611 *gor* gene of the lactic acid bacterium *Streptococcus thermophilus* CNRZ368. *Res*  
612 *Microbiol* **146**:371-383.

- 613 44. **Poole, L. B., M. Higuchi, M. Shimada, M. L. Calzi, and Y. Kamio.** 2000.  
614 *Streptococcus mutans* H<sub>2</sub>O<sub>2</sub>-forming NADH oxidase is an alkyl hydroperoxide  
615 reductase protein. *Free Radic Biol Med* **28**:108-120.
- 616 45. **Poyart, C., E. Pellegrini, O. Gaillot, C. Boumaila, M. Baptista, and P. Trieu-Cuot.**  
617 2001. Contribution of Mn-cofactored superoxide dismutase (SodA) to the virulence of  
618 *Streptococcus agalactiae*. *Infect Immun* **69**:5098-5106.
- 619 46. **Rocha, E. R., G. Owens, Jr., and C. J. Smith.** 2000. The redox-sensitive  
620 transcriptional activator OxyR regulates the peroxide response regulon in the obligate  
621 anaerobe *Bacteroides fragilis*. *J Bacteriol* **182**:5059-5069.
- 622 47. **Rocha, E. R., T. Selby, J. P. Coleman, and C. J. Smith.** 1996. Oxidative stress  
623 response in an anaerobe, *Bacteroides fragilis*: a role for catalase in protection against  
624 hydrogen peroxide. *J Bacteriol* **178**:6895-6903.
- 625 48. **Rocha, E. R., and C. J. Smith.** 1998. Characterization of a peroxide-resistant mutant  
626 of the anaerobic bacterium *Bacteroides fragilis*. *J Bacteriol* **180**:5906-5912.
- 627 49. **Rocha, E. R., and C. J. Smith.** 1999. Role of the alkyl hydroperoxide reductase  
628 (*ahpCF*) gene in oxidative stress defense of the obligate anaerobe *Bacteroides fragilis*.  
629 *J Bacteriol* **181**:5701-10.
- 630 50. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. Molecular cloning : a laboratory  
631 manual. Cold Spring Harbor Laboratory, New York.
- 632 51. **Sonnhammer, E. L. L., G. Von Heijne, and A. Krogh.** 1998. A hidden Markov  
633 model for predicting transmembrane helices in protein sequences, p. 175-182. *In* J.  
634 Glasgow, T. Littlejohn, F. Major, L. R., D. Sankoff, and C. Sensen (ed.), Proceedings  
635 of the sixth international conference on intelligent systems for molecular  
636 microbiology. AAAI Press, Menlo park, CA.

- 637 52. **Spratt, B. G.** 1975. Distinct penicillin binding proteins involved in the division,  
638 elongation, and shape of *Escherichia coli* K12. *Proc Natl Acad Sci U S A* **72**:2999-  
639 3003.
- 640 53. **Stadtman, E. R.** 1993. Oxidation of free amino acids and amino acid residues in  
641 proteins by radiolysis and by metal-catalyzed reactions. *Annu Rev Biochem* **62**:797-  
642 821.
- 643 54. **Szabo, L., S. Nishimura, and W. R. Farkas.** 1988. Possible involvement of queuine  
644 in oxidative metabolism. *Biofactors* **1**:241-4.
- 645 55. **Terzaghi, B., and W. Sandine.** 1975. Improved medium for lactic *streptococci* and  
646 their bacteriophages. *Appl Environ Microbiol* **29**:807-813.
- 647 56. **Thibessard, A., A. Fernandez, B. Gintz, B. Decaris, and N. Leblond-Bourget.**  
648 2002. Transposition of pGh9:ISS1 is random and efficient in *Streptococcus*  
649 *thermophilus* CNRZ368. *Can J Microbiol* **48**:473-478.
- 650 57. **Thibessard, A., A. Fernandez, B. Gintz, N. Leblond-Bourget, and B. Decaris.**  
651 2002. Effects of *rodA* and *pbp2b* disruption on cell morphology and oxidative stress  
652 response of *Streptococcus thermophilus* CNRZ368. *J Bacteriol* **184**:2821-2826.
- 653 58. **Thibessard, A., A. Fernandez, B. Gintz, N. Leblond-Bourget, and B. Decaris.**  
654 2001. Hydrogen peroxide effects on *Streptococcus thermophilus* CNRZ368 cell  
655 viability. *Res Microbiol* **152**:593-596.
- 656 59. **Thibessard, A., N. Leblond-Bourget, A. Fernandez, B. Gintz, and B. Decaris.**  
657 2001. Response of *Streptococcus thermophilus* CNRZ368 and its colonial variants to  
658 oxidative stress : evidence for an inducible defence system. *Lait* **81**:311-316.
- 659 60. **Tseng, H., Y. Srikhanta, A. McEwan, and M. Jennings.** 2001. Accumulation of  
660 manganese in *Neisseria gonorrhoeae* correlates with resistance to oxidative killing by  
661 superoxide anion and is independent of superoxide dismutase activity. *Mol Microbiol*  
662 **40**:1175-1186.

- 663 61. **Tusnady, G. E., and I. Simon.** 1998. Principles governing amino acid composition of  
664 integral membrane proteins: application to topology prediction. *J Mol Biol* **283**:489-  
665 506.
- 666 62. **Visca, P., A. Ciervo, V. Sanfilippo, and N. Orsi.** 1993. Iron-regulated salicylate  
667 synthesis by *Pseudomonas spp.* *J Gen Microbiol* **139**:1995-2001.
- 668 63. **Wachi, M., M. Doi, Y. Okada, and M. Matsuhashi.** 1989. New mre genes *mreC* and  
669 *mreD*, responsible for formation of the rod shape of *Escherichia coli* cells. *J Bacteriol*  
670 **171**:6511-6516.
- 671 64. **Wachi, M., M. Doi, S. Tamaki, W. Park, S. Nakajima-Iijima, and M. Matsuhashi.**  
672 1987. Mutant isolation and molecular cloning of *mre* genes, which determine cell  
673 shape, sensitivity to mecillinam, and amount of penicillin-binding proteins in  
674 *Escherichia coli*. *J Bacteriol* **169**:4935-4940.
- 675 65. **Wu, J., and B. Weiss.** 1992. Two-stage induction of the *soxRS* (superoxide response)  
676 regulon of *Escherichia coli*. *J Bacteriol* **174**:3915-3920.
- 677 66. **Yamamoto, Y., M. Higuchi, L. B. Poole, and Y. Kamio.** 2000. Role of the *dpr*  
678 product in oxygen tolerance in *Streptococcus mutans*. *J Bacteriol* **182**:3740-3747.
- 679 67. **Yesilkaya, H., A. Kadioglu, N. Gingles, J. E. Alexander, T. J. Mitchell, and P. W.**  
680 **Andrew.** 2000. Role of manganese-containing superoxide dismutase in oxidative  
681 stress and virulence of *Streptococcus pneumoniae*. *Infect Immun* **68**:2819-2826.
- 682 68. **Youn, H. D., H. Youn, J. W. Lee, Y. I. Yim, J. K. Lee, Y. C. Hah, and S. O. Kang.**  
683 1996. Unique isozymes of superoxide dismutase in *Streptomyces griseus*. *Arch*  
684 *Biochem Biophys* **334**:341-348.
- 685 69. **Yu, J., A. P. Bryant, A. Marra, M. A. Lonetto, K. A. Ingraham, A. F. Chalker, D.**  
686 **J. Holmes, D. Holden, M. Rosenberg, and D. McDevitt.** 2001. Characterization of  
687 the *Streptococcus pneumoniae* NADH oxidase that is required for infection.  
688 *Microbiology* **147**:431-438.

- 689 70. **Yuvaniyama, P., J. N. Agar, V. L. Cash, M. K. Johnson, and D. R. Dean.** 2000.  
690 NifS-directed assembly of a transient [2Fe-2S] cluster within the NifU protein. Proc  
691 Natl Acad Sci U S A **97**:599-604.
- 692 71. **Zheng, L., R. H. White, V. L. Cash, R. F. Jack, and D. R. Dean.** 1993. Cysteine  
693 desulfurase activity indicates a role for NIFS in metallocluster biosynthesis. Proc Natl  
694 Acad Sci U S A **90**:2754-2758.
- 695 72. **Zheng, M., X. Wang, L. J. Templeton, D. R. Smulski, R. A. LaRossa, and G.**  
696 **Storz.** 2001. DNA microarray-mediated transcriptional profiling of the *Escherichia*  
697 *coli* response to hydrogen peroxide. J Bacteriol **183**:4562-70.

698

699

#### LEGENDS FIGURES

700

701 **FIG. 1. Relative methyl viologen sensitivity.** (A). Results of short term methyl viologen  
702 exposure. (B) Results of long term methyl viologen exposure (see materials and methods).

703 The relative methyl viologen sensitivity was determined by comparing viability with and  
704 without treatment. Levels are shown as values relative to the survival level of the wild type  
705 strain. All experiments were performed at least in duplicate. Error bars show two standard  
706 deviations of the data.

707

708 **FIG. 2. Survival to heat shock.** The heat shock consisted of 15 min treatment at 60°C. The  
709 percentage of surviving cells corresponds to the number of colony forming units (CFUs)  
710 counted on plates after the heat-shock exposure divided by the number of CFU before  
711 treatment. Error bars show two standard deviations of the data. The dotted line represents the  
712 surviving percentage observed for the wild type strain and the grey band comprises 2 standard  
713 deviations of the data.

714

715 **FIG. 3. Influence of *mreD* disruption on chain length.** Cell number per chains was  
716 determined for 70 chains of the wild type strain (■) and of the *mreD* mutant (■) grown in  
717 liquid TPPY medium.

718

719 **FIG. 4. Schematic map of the *S. thermophilus* *sufCDS iscU sufB* cluster.** Genes are  
720 indicated by grey arrows, and the positions of putative start and stop codons are given (in bp)  
721 below the map. The site of insertion of the pGh9:ISS1 plasmid in *sufD* (*ossI*) and *iscU* (*ossJ*)  
722 disrupted mutants is indicated by the solid black triangles. The broken arrow indicates the  
723 position of the *suf* cluster putative promoter. Putative rho-independent terminator is  
724 represented as hairpin loop. Fragments used for generation of the non-polar mutants are  
725 indicated as open boxes. The star indicates the position of the *EcoRV* restriction site causing  
726 the Cys<sub>97</sub> replacement by a Tyr in the *iscU*<sub>97</sub> mutant.

727

728 **FIG. 5. Streptonigrin tolerance.** *S. thermophilus* strain tolerance to streptonigrin was  
729 assessed by measuring OD<sub>600</sub> after growth in TPPYC medium containing various  
730 concentrations of streptonigrin without (A) or supplemented with 1 mM of 2,2'-dipyridyl (B)  
731 (see material and methods for details). Relative OD<sub>600nm</sub> was determined by comparing the  
732 OD<sub>600</sub> obtained with or without streptonigrin. Wild type strain (○ and ●); *iscU*<sub>97</sub> (□ and ■);  
733  $\Delta$ *sufD* (Δ and ▲).

734 TABLE 1. Products of *oss* genes: database similarities and putative fonctions.

735

Gene name <sup>a</sup>	Accession number	Gene encoding the most similar product <sup>c</sup>	amino acid identity <sup>d</sup>	Insertion site <sup>e</sup>	Putative function	Class
<i>ossA</i> ( <i>mreD</i> )	AY386242	<i>S. mutans mreD</i> (168)	30% (1-168)	127	Cell shape maintenance	I
<i>ossB</i> ( <i>rodA</i> )	AF399832	<i>S. mutans rodA</i> (408)	65% (1-391)	304	Peptidoglycan elongation	
<i>ossC</i> ( <i>pbp2b</i> ) <sup>b</sup>	AF399833	<i>S. thermophilus Sfi6 pbp2b</i> (704)	98% (641-704)	640	Peptidoglycan elongation	
<i>ossD</i> ( <i>cpsX</i> ) <sup>b</sup>	AY386241	<i>S. thermophilus cpsX</i> (378)	98% (212-378)	264	Membrane translocation of exopolysaccharide	II
<i>ossE</i> ( <i>tgt</i> ) <sup>b</sup>	AY386243	<i>S. pneumoniae tgt</i> (380)	92% (1-333)	99	tRNA-guanine transglycosylase	III
<i>ossF</i> <sup>b</sup>	AY386244	<i>S. agalactiae gbs0448</i> (451)	77% (1-204)	51	RNA methyltransferase	
<i>ossG</i>	AY386239	<i>S. mutans smu.1428c</i> (284)	67% (1-280)	170	Unknown conserved function	IV
<i>ossH</i> ( <i>fatD</i> )	AY386240	<i>S. pneumoniae fatD</i> (324)	55% (22-323)	119	Iron ABC transporter	V
<i>ossI</i> ( <i>sufD</i> )	AY386245	<i>S. agalactiae sag0142</i> (420)	76% (1-420)	144	Assembly of Fe/S clusters	
<i>ossJ</i> ( <i>iscU</i> )	AY386245	<i>S. agalactiae gbs0140</i> (147)	78% (1-141)	57	Assembly of Fe/S clusters	

736

737 a. Initial gene name with, in brackets and when applicable, the definitive gene name.

738 b. Disrupted ORF of *S. thermophilus* CNRZ368 that have been partially sequenced.739 c. Organism, gene name and, in brackets, length of the protein sequence giving the best alignment  
740 score.741 d. Percentage of sequence identity with the protein showing the best alignment score to each  
742 *S. thermophilus* Oss protein. In brackets is indicated the portion of the protein sharing this identity  
743 (numbers correspond to the amino acid positions).744 e. The numbers correspond to the deduced amino acid position of the pGh9:ISS1-generated  
745 disruption in the full length protein showing the best alignment score to the Oss proteins.



746 TABLE 2. *suf* cluster gene products of *S. thermophilus* CNRZ368

747

Gene product	Closest similar protein (organism, accession number) <sup>a</sup>	Identity/similarity(%)	Function
SufC	CAD45782 ( <i>S. agalactiae</i> NEM316)	88/94	Putative ABC transporter (ATP-binding protein)
SufD	AAM99050 ( <i>S. agalactiae</i> 2603V/R)	76/87	Unknown, conserved protein
SufS	CAD45784 ( <i>S. agalactiae</i> NEM316)	79/89	Similar to SufS
IscU	CAD45785 ( <i>S. agalactiae</i> NEM316)	78/90	Similar to NifU family proteins
SufB	CAD45786 ( <i>S. agalactiae</i> NEM316)	89/94	Unknown, conserved protein

748

749 a. Accession number of the protein giving the best alignment score. In brackets, the name of the

750 organism is indicated.

751 TABLE 3. Growth of wild type and mutant strains in TPPYC and in TPPYC supplemented with  
752 iron.


753


Strain	Final OD <sub>600</sub> resulting from growth <sup>a</sup> in:	
	TPPYC	TPPYC + FeSO <sub>4</sub>
WT	0.96	1.32
<i>ΔsufD</i>	0.58	1.48
<i>iscU<sub>97</sub></i>	0.58	1.15


754


755 <sup>a</sup> Cells were grown in milk medium for 15<sup>o</sup>h at 42°C and diluted in FeSO<sub>4</sub>-deficient medium  
756 (TPPYC medium supplemented with MgCl<sub>2</sub> (1 mM) and CaCl<sub>2</sub> (1 mM)). After one night  
757 growth, cells were washed twice and resuspend either in FeSO<sub>4</sub>-deficient medium or  
758 supplemented with 100 μM FeSO<sub>4</sub>. The growth of the cells was then followed by measuring  
759 the OD<sub>600</sub>. The OD<sub>600</sub> were measured after 24h incubation at 42°C. This experiment was done  
760 in triplicate.



761















## ABSTRACT

19

20

21 To better understand the defense mechanism of *Streptococcus thermophilus* against  
22 superoxide stress, molecular analysis of 10 menadione-sensitive mutants, obtained by  
23 insertional mutagenesis, was undertaken. This analysis allowed the identification of 10 genes  
24 that, with respect to their putative function, were classified in 5 categories (i) those involved  
25 in cell wall metabolism, (ii) those involved in exopolysaccharide translocation, (iii) those  
26 involved in RNA modification (iv) those involved in iron homeostasis and (v) those whose  
27 function is still unknown.

28 The behavior of the 10 menadione sensitive mutants exposed to heat shock was investigated.  
29 Data from these experiments allowed us to distinguish genes whose action might be specific  
30 to oxidative stress defense (*tgt*, *ossF*, *ossG*) from those whose action may be generalized to  
31 other stressful conditions (*mreD*, *rodA*, *pbp2b*, *cpsX*, *iscU*).

32 Among the mutants, two harboured an independently inserted copy of pGh9:ISS1 in two loci  
33 close to each other. More precisely, these two loci are homologous to *sufD* and *iscU* genes,  
34 involved in the biosynthesis of iron-sulfur clusters. This region, called the *suf* region, was  
35 further characterized in *S. thermophilus* CNRZ368 by sequencing and by construction of  
36  $\Delta$ *sufD* and *iscU*<sub>97</sub> non polar mutants. Streptonigrin sensitivity level of both mutants suggested  
37 that these two genes were involved in iron metabolism.

## INTRODUCTION

39 Oxygen is an essential element for the viability of most living organisms through its involvement  
40 in oxidative phosphorylation that takes place during the **aerobic** respiration process. However,  
41 oxygen can also be harmful, due to the production, especially during respiration, of toxic  
42 derivatives namely ROS (reactive oxygen species) such as hydrogen peroxide, hydroxyl and  
43 superoxide radicals (for a review, (25)). Accumulation of these molecules, triggers lipid, protein  
44 and nucleic acid oxidation, causing lethal cellular damage (for reviews (7, 25, 53)). In response  
45 to such attack, aerobic micro-organisms have evolved anti-oxidant mechanisms (eg catalase,  
46 peroxidase and superoxide dismutase activities) to protect and adapt cells against ROS excess (6,  
47 18, 24, 65, 68).

48 Although **they do not achieve aerobic respiration**, anaerobic micro-organisms can be subject to  
49 ROS. The strict anaerobes *Bacteroides fragilis* and *Clostridium perfringens* are able to survive in  
50 the presence of a low amount of oxygen, due to the induction of an adaptative oxidative stress  
51 response that is activated under aerobic conditions (13, 46-49). The peroxide response of  
52 *B. fragilis* shares some characteristics with that of *E. coli* such as the involvement of the redox-  
53 sensitive transcriptional activator OxyR (46) and some antioxidant activities under its control  
54 (47, 49).

55 The lactic acid bacterium (LAB) *Streptococcus thermophilus* is a **Gram-positive bacterium**  
56 widely used in dairy factories, mainly as a starter of fermentation in yogurt and cheese  
57 production. During fermentation processes, *S. thermophilus* is exposed to oxygen and to its  
58 deleterious effects that can lead to a reduction of cell viability and consequently may have  
59 repercussions on texture, flavor and safety of the final product.

60 *S. thermophilus* is an anaerobic aerotolerant organism that can grow in presence of oxygen  
61 and survive to low concentrations of superoxide and hydroxyl radicals (59). This suggests that  
62 it displays a defense system that provides protection against the effects of ROS. In the  
63 literature, data about its oxidative stress response are scarce. The existence of at least one

64 ROS defense system and its inducibility by hydrogen peroxide are established in  
65 *S. thermophilus* CNRZ368 (58). This organism possesses a single manganese-containing  
66 superoxide dismutase (19) and a glutathione reductase (43). However, no catalase activity was  
67 found. In other streptococci, antioxidant enzymes were reported to play a role in the defense  
68 against oxidative stress (27, 33, 44, 66) and in pathogenesis (45, 67, 69).  
69 The aim of this work was to identify genes involved in the oxidative stress defense of  
70 *S. thermophilus* CNRZ368 without presuming their function. For this purpose a collection of  
71 mutants was created by random insertional mutagenesis and the mutants were screened for  
72 their sensitivity to menadione (a superoxide generating agent). The locus disrupted in 10  
73 *S. thermophilus* menadione sensitive clones was identified and the potential physiological role  
74 of the mutated genes in oxidative stress defense is discussed. The implication of these 10 loci  
75 during heat shock was also evaluated to assess their specificity to oxidative stress defense.  
76 Moreover, the characterization of two of these mutants, disrupted in *sufD* and *iscU* genes  
77 which may participate in Fe-S clusters assembly or repair, is more detailed. The results  
78 presented in this paper indicate that the inactivation of *sufD* and *iscU* genes caused a higher  
79 sensitivity to superoxide stress and suggest that Fe-S cluster repair or assembly could play an  
80 important role in the oxidative stress defense of anaerobic bacteria.

81

82

## MATERIALS AND METHODS

83 **Bacterial strains, growth conditions.** *S. thermophilus* CNRZ368 was purchased from the  
84 INRA-CNRZ strain collection (Jouy en Josas, France). Depending on the experiments,  
85 *S. thermophilus* CNRZ368 and its derivatives were cultivated at 30°C or 42°C in milk  
86 medium (for pre-cultures), TPPY (12) or M17 (55) media, **without shaking**. Erythromycin was  
87 added at 2 µg.ml<sup>-1</sup>, when required. Streptonigrin sensitivity was monitored using a TPPYC  
88 medium which contained a reduced concentration of metallic dications (including Fe<sup>2+</sup>).  
89 TPPYC was performed according to the protocol described by Visca *et al.* (62) and modified as

90 follow: Chelex beads (20 g/l, sigma) were added to TPPY medium and stirred at 4°C for 48 h  
91 and the medium was then supplemented with 1 mM MgCl<sub>2</sub> and 1 mM CaCl<sub>2</sub>.

92 Recombinant plasmids were transformed into *E. coli* EC101, a TG1 strain [*supE hsd-5 thi Δ(lac-*  
93 *proAB)* F' (*traD6 proAB<sup>+</sup> lacI<sup>f</sup> lacZΔM15*)] containing a chromosomal copy of the pWV01 *repA*  
94 gene (34) and selected at 37°C on Luria-Bertani (LB) (50) agar plates containing 150 µg/ml of  
95 erythromycin.

96

97 **Insertional mutagenesis.** The collection of mutants was made by insertional mutagenesis using the  
98 plasmid pGh9:ISSI that carries Em<sup>R</sup> gene (35). The mutagenesis procedure and the cloning of the  
99 fragments flanking the pGh9:ISSI insertion locus were carried out on *S. thermophilus* CNRZ368  
100 genome, as previously described (56).

101

102 **Selection of mutants impaired in their ability to tolerate menadione.** For selection of  
103 menadione sensitive clones, two different procedures, differing mainly by the length of the  
104 menadione-exposure, were assayed. Both assays were preceded by a pre-culture in milk medium  
105 containing 2 µg/ml erythromycin: pre-cultures of the 2112 mutants were performed in 96-well  
106 microtitre plates.

107 For “long” time menadione exposure, each pre-culture was transferred, using a hedgehog, onto  
108 M17 agar plates containing 2 µg/ml erythromycin and increasing concentrations of menadione  
109 (from 0 µg/ml to 120 µg/ml). After 20h incubation at 42°C, the presence (or the absence) of  
110 colonies was observed.

111 For “short” time menadione exposure, each pre-culture was diluted (with a 1/100 ratio) in 5  
112 different 96-well microtitre plates containing liquid TPPY medium supplemented with 2 µg/ml  
113 erythromycin (in final volume of 100µl, resulting in an approximately 1/1 liquid to air space  
114 ratio). After a 3h30 growth period at 42°C, 5 different concentrations of menadione (from  
115 0 mg/ml to 15 mg/ml) were added to each of the 5 microtitre plates. Microtitre plates were

116 incubated at 42°C for 3h. The cultures were then transferred on TPPY agar plates containing  
117 erythromycin (2 µg/ml) using a hedgehog. The ability of the mutants to survive to the menadione  
118 stress was assessed, after 20 h incubation at 42°C, by observation of the presence or the absence  
119 of colonies.

120 For each experiment, clones were considered as sensitive when they did not grow on a plate  
121 where more than 90% of the clones were growing. Both procedures were applied twice and  
122 the results had been compiled to reveal the mutants raising the more severe phenotype.

123

124 **DNA manipulations.** Preparation of chromosomal, plasmid and recombinant DNA and  
125 Southern analysis were performed according to standard protocols (50). All cloning junctions  
126 and cloned PCR products were checked by sequencing using Dye-Terminator chemistry on an  
127 ABI Prism 377 Genetic Analyser (PE Biosystems). Sequence data were analysed using  
128 Genbank, BLAST (2, 3), TMHMM (51) and HMMTOP (61) software. For all analysed  
129 sequences, The DNA sequences reported in this paper have been deposited in GenBank under  
130 accession numbers AY386239, AY386240, AY386241, AY386242, AY386243, AY386244,  
131 AY386245.

132

133 **Construction of the *ΔsufD* and *iscU<sub>97</sub>* mutants.** Oligonucleotides used to construct the  
134 *ΔsufD* and *iscU<sub>97</sub>* mutants were purchased from MWG Biotech AG (Ebersberg, Germany). To  
135 construct non-polar truncated *ΔsufD* mutant, the 5' and 3' ends of the *sufD* gene were  
136 independently amplified by PCR. The 5' end was amplified by using the primers  
137 GAGCGTGTTGAATTCCACCG containing an *EcoRI* site, symbolized in bold characters  
138 and AACTTAACTGCAGCACCGTCT containing a *PstI* site. The 3' end was amplified by  
139 using the primers GGTGTTCCAGAATTCAAATAACC containing an *EcoRI* site and  
140 TGGTCTGTCATCAGGTCG. This last primer does not contain any *PstI* site, but the PCR  
141 product generated does contain a *PstI* restriction site. These two PCR products were digested

142 by *Pst*I and ligated with each other. The resulting fragment was amplified by PCR, then  
143 digested by *Eco*RI and cloned into pGh9. The final construct (pGh9 $\Delta$ *scfD*) was used to  
144 transform *S. thermophilus* CNRZ368. For IscU protein inactivation, the strategy consisted in  
145 replacing the TGT codon encoding the conserved Cys<sub>97</sub> by a TAT codon encoding a Tyr. This  
146 was achieved by replacing, in oligonucleotides, the 3 **underlined** nucleotides of the sequence  
147 (**AGG TGT ACA**) generating an *Eco*RV restriction site (**AGA TAT CCA**). Thus, the three  
148 amino-acids Gly<sub>96</sub> Cys<sub>97</sub> Tyr<sub>98</sub> were replaced by Arg<sub>96</sub> Tyr<sub>97</sub> Thr<sub>98</sub> in the mutated protein. Two  
149 internal and complementary primers containing the mutation  
150 (**GCTGGTGATGATATCACCATTTCACA** and  
151 **GAAATGGTGATATCATCACCAGCAAAGG**) were used with two additional external  
152 primers (**AAGCGGAATTCAAGCATCCACTCAGGC** containing an *Eco*RI site and  
153 **TGTGACAAGCTTGTAGAGGCAATTCTT** containing an *Hind*III site), respectively, to  
154 amplify 5' and 3' end of *iscU*. The two PCR products of 515 bp and 262 bp were digested by  
155 *Eco*RI and *Eco*RV and by *Hind*III and *Eco*RV respectively before being ligated to each other.  
156 The resulting fragment was cloned into pGh9 digested by *Eco*RI and *Hind*III. The final  
157 construct, (pGh9 $\Delta$ *iscU*<sub>97</sub>) was used to transform *S. thermophilus* CNRZ368. Preparation of  
158 *S. thermophilus* competent cells was performed as previously described by O'sullivan and  
159 Fitzgerald (41). The following gene replacements were done as previously described (8) but  
160 in TPPY medium and at 30°C/42°C as permissive and restrictive temperature, respectively.

161

### 162 **Physiological characterization of *S. thermophilus* mutant strains.**

163 **Exposure to methyl viologen.** For “short” time exposure, cultures of CNRZ368 and its  
164 derivative strains were grown in TPPY medium. At an OD<sub>600</sub> of 0.6, cultures were divided into  
165 two subcultures (in a ratio of liquid to air space of approximately 1/1). Methyl viologen  
166 (paraquat; Sigma-Aldrich) was added to one of them with a final concentration of 50 mM. After  
167 3 h incubation at 42°C, cultures were diluted to appropriate concentrations and spread onto

168 TPPY agar plates. For “long” time methyl viologen exposure, strains were **grown** in TPPY  
169 medium until OD<sub>600</sub> of 0.6. Cultures were then diluted to a final theoretical OD<sub>600</sub> of 0.001 in  
170 TPPY medium **(in a ratio of liquid to air space of approximately 5/6)** supplemented with 1.7 mM  
171 methyl viologen, or not for control experiment. After 18h incubation at 42°C, cultures were  
172 diluted in TPPY to appropriate concentrations and spread onto TPPY agar plates. In both  
173 procedures, methyl viologen sensitivity was estimated by comparing CFU/ml obtained with  
174 treated cultures, with respect to non treated cultures.

175

176 **Heat shock.** The ability of *S. thermophilus* and its derivatives to respond to a heat stress was  
177 assessed as follow. Each strain is grown at 42°C in TPPY medium until early log phase  
178 (OD<sub>600</sub> = 0.3). Then the culture was divided into two. One part was kept at 42°C (control  
179 experiment); the other one was immediately transferred to 60°C for 15 min. Appropriate  
180 dilutions were spread on TPPY agar plates. The fraction of surviving cells was calculated by  
181 dividing the surviving CFU/ml by the count for the control experiment.

182

183 **Streptonigrin sensitivity.** Streptonigrin sensitivity was monitored on cells grown overnight  
184 in TPPYC medium at 42°C. Cells were harvested by centrifugation, washed twice in TPPYC  
185 medium, resuspended in TPPYC, diluted to OD<sub>600</sub> = 0.01 and cultivated in TPPYC for 9 h at  
186 42°C. Then, the cultures were diluted again to OD<sub>600</sub> = 0.005 and divided into 4 parts which  
187 were all incubated for 21 h at 42°C. One part was supplemented with 0.4% (v/v) dimethyl  
188 formamide (control experiment), the 3 other parts were supplemented with different  
189 concentrations of streptonigrin (1 µg/ml, 2 µg/ml and 4 µg/ml dissolved in dimethyl  
190 formamide). As a control experiment, cultures were also carried out in TPPYC supplemented  
191 with 1 mM of 2'2-dipyridyl (an Fe chelator). In each experiment, OD<sub>600</sub> value was used to  
192 estimate the cellular density after 21h incubation. Relative optic density was determined by

193 comparing cellular density obtained with and without streptonigrin. All experiments were  
194 performed in triplicate.

195

196

## RESULTS

### 197 **1. Selection of *S. thermophilus* CNRZ368 mutants with altered oxidative stress response.**

198 A collection of 2112 mutants of *S. thermophilus* CNRZ368 was generated by insertional  
199 mutagenesis using the plasmid pGh9:ISS1 (35, 56). This plasmid carries an erythromycin  
200 resistance gene for its selection and the insertional element ISS1 that allows random  
201 integration of the plasmid into *S. thermophilus* genome. The 2112 mutants obtained were  
202 screened to select clones sensitive to menadione (a superoxide-generating compound). Two  
203 distinct selection procedures were used to identify mutants susceptible to menadione (see  
204 materials and methods). The characterization of 10 mutants, disrupted in loci originally named  
205 *ossA* to *ossJ* (for oxidative stress sensitive locus) is described in this paper.

206

### 207 **2. Sensitivity of *S. thermophilus* CNRZ368 mutants to methyl viologen.** Previously, we

208 demonstrated that *ossB* and *ossC* mutants (previously named 6G4 and 18C3, respectively) are  
209 compromised in their ability to survive superoxide stress (57). To confirm that the other 8  
210 mutants selected were sensitive to superoxide stress, their sensitivity to methyl viologen was  
211 examined (Fig. 1). In this experiment, we used the non-polar mutants *ΔsufD* and *iscU<sub>97</sub>*  
212 constructed in this work (see further) rather than the potential polar clones *ossI* and *ossJ*.  
213 Short-term methyl viologen exposure revealed the extreme sensitivity of *ossG*, as well as the  
214 4 to 5-fold increased sensitivity of *ossD*, *ossH* and *iscU<sub>97</sub>* mutants (Fig. 1A). Although less  
215 severe the *ΔsufD* mutant exhibited about 2-fold less survival following such stress. Under  
216 these conditions, the other 3 mutants (*ossA*, *ossE* and *ossF*) did not exhibit any significant  
217 sensitivity, however long term methyl viologen exposure confirmed their sensitivity. Indeed,  
218 long term methyl viologen exposure (Fig. 1B) showed that the *ossA* mutant strain displayed a



219 relative survival more than 10-fold lower than that of the wild type strain. The *ossE* and *ossF*  
220 mutants also showed a 2-fold lower growth compared to the wild type strain.

221

222 **3. Survival of the mutants exposed to heat stress.** To assess the specificity of these 10 loci  
223 to oxidative stress defense, the mutants ability to survive to heat shock was estimated. Five  
224 out of the 10 mutants (*ossA*, *ossB*, *ossC*, *ossD*, and *ossJ*) isolated as oxidative stress sensitive  
225 clones also showed a reduced survival to heat shock compared with WT (Fig. 2). These data  
226 suggest that the genes impaired in these mutants are not specific to oxidative stress defense  
227 process but are probably involved in a more global response to stresses.

228 In contrast, the *ossE*, *ossF* and *ossG* mutants did not differ significantly from the wild type,  
229 indicating that the genes disrupted in these strains did not influence cells ability to respond to  
230 heat shock. Interestingly, the *ossG* mutant, that was highly sensitive to methyl viologen,  
231 showed no significant decrease of survival to heat shock compared with WT, suggesting that  
232 *ossG* gene, essential for oxidative stress defense, could be specific to such a stress.

233

234 **4. Identification of genes implicated in oxidative stress tolerance.** The disrupted locus was  
235 identified for each of the 10 mutants by cloning and sequencing the junctions as previously  
236 described (56). All of the insertions were mapped within coding sequences. Sequence  
237 similarity searches and analysis revealed similar sequences in other bacteria (see Table 1)  
238 with an E-value lower than  $10^{-10}$  for BLASTX searches. In 9/10 cases, data about the function  
239 of these genes were available. To facilitate their description, these genes were grouped into 5  
240 classes (from I to V) according to the putative function and renamed when relevant (Table 1).

241

242 **Class I: Genes potentially involved in cell wall structure and cell shape determination.**

243 The *ossA*, *ossB* and *ossC* putative gene products shared sequence identity with MreD, RodA  
244 and Pbp2b, respectively, which putatively direct the elongation of the lateral cell wall in

245 *E. coli*, conferring the cell its characteristic rod shape (5, 28, 29). The *ossA* mutant was  
246 defective in a gene encoding a protein, with 30% identity with MreD of *Streptococcus*  
247 *mutans*. Sequence analysis of the surrounding region indicated that *S. thermophilus mreD*  
248 gene (*ossA*) was clustered with a gene encoding a putative protein similar to MreC. Moreover,  
249 hydrophopathy profiles of MreD (OssA) determined by using TMHMM and HMMTOP  
250 showed 5 regions of sufficient length and hydrophobicity to be membrane-spanning segments  
251 (data not shown), suggesting that this protein is the counterpart of MreD. The relatively low  
252 identity (30%) observed between *S. thermophilus* and *S. mutans* MreD was in agreement with  
253 the general idea that the structure of this protein is more highly conserved than its primary  
254 sequence (17). To study the effect of *mreD* disruption in *S. thermophilus*, TPPY cultures of  
255 the wild type and *mreD* mutant strains were observed under light microscopy. In contrast to  
256 wild type cells, *mreD* mutant cells formed very long chains, up to 180 cells per chain (Fig. 3).  
257 Moreover, a reduction of about 25% of the cell size was observed in the mutant. Growth of  
258 the *mreD* mutant and of the wild type in TPPY medium were monitored by measuring the  
259 OD<sub>600</sub>. Both strains displayed a similar doubling time of 30 min, but the maximal OD<sub>600</sub> of the  
260 *mreD* mutant (OD<sub>max</sub> = 1.7 ± 0.1) was lower than that of the wild type (OD<sub>max</sub> = 2.4 ± 0.2)  
261 showing that the *mreD* mutant entered stationary phase earlier than wild type cells. The  
262 characterization of *rodA* and *pbp2b* mutants of *S. thermophilus* CNRZ368 is described in a  
263 previous paper (57).

264

265 **Class II: Gene potentially involved in exopolysaccharide translocation.** The sequence  
266 disrupted in the mutant *ossD* was partially determined. Its analysis indicated that the  
267 C-terminal portion of the predicted encoded protein showed 98% identity to CpsX, a putative  
268 membrane protein from *S. thermophilus* MR-1C. The role of CpsX is still unknown but the  
269 conserved location of *cpsX* near *eps* genes of *S. thermophilus* (14) and the presence of a  
270 putative gene related to an ABC transport system in its vicinity argue in favor of CpsX being

271 involved in exopolysaccharide translocation (15). Moreover, the TMHMM and HMMTOP  
272 software predictions indicated that CpsX (OssD) from *S. thermophilus* CNRZ368 would  
273 contain highly hydrophobic domains compatible with a location in the cytoplasmic  
274 membrane. Although the *cpsX* function is not known yet, our work indicate that its disruption  
275 is associated with sensitivity not only to superoxide stress but also to heat shock.

276

277 **Class III: Genes involved in RNA maturation.** In the *ossE* and *ossF* mutants, the plasmid  
278 inserted into genes whose predicted products **share sequence identity with** enzymes involved  
279 in RNA maturation. The *ossE* gene encodes a putative tRNA guanine transglycosylase (TGT)  
280 which shares 92% identity with the putative *Streptococcus pneumoniae* TGT. TGT enzyme  
281 takes part in the process which removes the guanine residue at position 34 (wobble position of  
282 the anticodon loop) of tRNA<sup>His</sup>, tRNA<sup>Asp</sup>, tRNA<sup>Asn</sup> and tRNA<sup>Tyr</sup> and replaces it by the  
283 modified residue queuosine (39, 40).

284 *ossF* potentially encodes a protein that shows 77 % identity with a putative RNA  
285 methyltransferase of *Streptococcus agalactiae* (Gbs0448) belonging to the TrmA family.  
286 Enzymes of the TrmA family catalyze the formation of the 5-methyluridine in position 54  
287 (localized in the TΨC loop) of tRNA (9). The presence of a TRAM-like domain (RNA  
288 binding domain) in the N-terminal part of OssF supports the idea of an interaction between  
289 this protein and RNA.

290

291 **Class IV: Gene of unknown function.** The *ossG* **product shows 67% identity with the**  
292 **product of** a gene of unknown function (*smu.1428c*) of *S. mutans* (Table 1). Although *ossG*  
293 function is not known yet, the mutation of this gene is now associated with a sensitivity  
294 phenotype to superoxide stress. The TMHMM and HMMTOP software predictions suggest  
295 the existence of two transmembrane domains in the N-terminal region of OssG, making of  
296 OssG a potential membrane protein.

297

298 **Class V: Genes involved in iron metabolism.**

299 **(i) Ferric iron transport.** The putative product of *ossH* shows 55% identity with FatD from  
300 *S. pneumoniae*, a potential membrane spanning permease of an Fe<sup>3+</sup> ABC-transporter. Analysis  
301 of the adjacent sequence reveals the presence of a **gene encoding a putative FatC similar protein**  
302 downstream of *fatD* (*ossH*), in the same orientation. The *fatC* gene is also clustered with *fatD* in  
303 *S. pneumoniae* genome. Hydropathy profile of FatD (OssH) from *S. thermophilus*, assessed with  
304 TMHMM and HMMTOP, reveals that this protein displays ten putative membrane spanning  
305 segments compatible with its expected cytoplasmic membrane location.

306

307 **(ii) [Fe-S] cluster biosynthesis.** The sequence affected in the *ossI* mutant encodes a product  
308 that shows 75% identity with Sag0142, a putative protein deduced from the complete genome  
309 sequence of *S. agalactiae* and **similar** to SufD-like proteins. Analysis of totally sequenced  
310 microbial genomes reveals that *sufD*-like genes are conserved in bacteria and also in some  
311 archaea such as *Halobacterium* sp. NRC-1, *Thermoplasma acidophilum* and *Aeropyrum*  
312 *pernix* (23). In *E. coli* and *Erwinia chrysanthemi*, the *sufD* gene belongs to the *sufABCDSE*  
313 cluster identified as being involved in iron metabolism and more precisely in [Fe-S] cluster  
314 biogenesis and/or repair (37, 42). The *ossJ* **gene product shows sequence identity to the**  
315 ***gbs0140* product** from *S. agalactiae* (Table 1), probably encoding a IscU protein, a NifU  
316 family protein. NifU proteins have been proposed to serve as scaffold proteins in [Fe-S]  
317 cluster assembly in *Synechocystis* PCC6803 and would function as a mediator for iron-sulfur  
318 cluster delivery (38).

319 In some genomes including that of *S. pyogenes*, *nifU*-like genes are clustered with the *suf*  
320 genes (23). To determine if *sufD* (*ossI*) and *iscU* (*ossJ*) from *S. thermophilus* were genetically  
321 linked together, the chromosomal DNA regions flanking the pGhost9:ISSI insertion site of  
322 both mutants were sequenced until overlapping. Examination of the genes in the

323 neighborhood of *sufD* and *iscU* revealed that they both belong to a cluster of 5 genes all  
324 orientated in the same direction. This cluster, named *sufCDS iscU sufB* cluster, is shown in  
325 Fig. 4. The highest **sequence identities** of these gene products are found with putative proteins  
326 from *S. agalactiae* (Table 2). They also share sequence similarity with proteins encoded by  
327 *E. coli* and *Er. chrysanthemi suf* gene clusters. The putative SufC, SufD, SufS and SufB  
328 proteins encoded by the *S. thermophilus* cluster present respectively 72%, 48%, 65%, and  
329 57% similarity to their *E. coli* **counterparts**. Moreover, IscU presents 47% similarity with IscU  
330 from *E. coli*.

331 Sequences corresponding to promoter regions were searched by inspection of the total  
332 sequence of the *sufCDS iscU sufB* cluster. A unique extended -10 sequence  
333 (5'-TTTGGTATAAT-3') is found upstream *sufC*, the first ORF of the cluster. A potential  
334 rho-independent transcriptional terminator is identified 63 bp downstream *sufB*, the last ORF  
335 of the cluster ( $\Delta G = -10$  kcal/mol). These features, together with the short or missing  
336 intergenic regions and the same orientation of the genes, suggest that these 5 ORFs form an  
337 operon, as it is the case for the *Er. chrysanthemi suf* locus.

338

## 339 **5. Molecular characterization of the *sufCDS iscU sufB* cluster.**

340 **5.1. Construction of  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutants.** Since *S. thermophilus suf* genes are  
341 potentially clustered in an operon, non polar  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutants were constructed and  
342 analysed. The *sufD* mutation consists of an internally in frame deletion leading to the loss of  
343 43% of the whole protein (see materials and methods and Fig. 4). Sequence comparison of  
344 NifU family proteins from various organisms indicates that *S. thermophilus* IscU protein  
345 contains the three highly conserved cystein residues that have been shown to be the assembly  
346 site of a "transitory" [2Fe-2S] center within NifU (70). Thus, the construction of a null *iscU*  
347 mutant was achieved by replacing one of these conserved residues, the Cys<sub>97</sub>, by a Tyr (see

348 materials and methods and Fig. 4). Sequencing of these two constructs confirmed that they  
349 did not contain any secondary mutation.

350

## 351 **5.2. Role of *sufD* and *iscU* in iron metabolism.**

352 To investigate the possibility that  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutations impair growth due to the lack  
353 of functional [Fe-S] cluster biosynthesis, the mutants were grown in TPPY medium and  
354 compared to wild type. Growth of the  $\Delta$ *sufD* and *iscU*<sub>97</sub> mutants showed no significant  
355 difference compared to the WT strain. Assuming that the high iron concentration of the TPPY  
356 medium could allow self assembling of the [Fe-S] clusters (71), and consequently could be  
357 responsible for the absence of difference in growth between strains, the same experiment was  
358 conducted in TPPYC, a divalent cation depleted medium. Under these conditions, the mutants  
359 displayed a growth rate similar of that of the WT strain but their maximal OD<sub>600</sub> was lower  
360 than that of the wild type, showing their premature entry in stationary phase (Table 3).  
361 Moreover, this difference disappeared when the cultures were grown on TPPYC  
362 supplemented by 100  $\mu$ M of FeSO<sub>4</sub>. Taken together, these results suggest that (i) *iscU* and  
363 *sufD* genes of *S. thermophilus* are involved in iron metabolism and (ii) their presence would  
364 be of particular importance under iron-limiting conditions.

365 To confirm the involvement of *sufD* and *iscU* in iron metabolism, the iron intracellular  
366 concentration of  $\Delta$ *sufD* and *iscU*<sub>97</sub> was checked by measuring the sensitivity of these strains  
367 to streptonigrin (an iron-activated antibiotic). As shown in Fig. 5 A, the *iscU* mutant was more  
368 sensitive to streptonigrin than the wild type strain whereas the  $\Delta$ *sufD* mutant was more  
369 resistant. Moreover, addition of 2,2'-dipyridyl (a ferrous iron chelator) abolished the  
370 difference of growth between the strains (Fig. 5 B) indicating that iron was responsible for the  
371 streptonigrin sensitivity observed without 2,2'-dipyridyl. Taken altogether these results  
372 suggest that both *iscU* and *sufD* of *S. thermophilus* play a role in iron metabolism.

373

## Discussion

374

375 To investigate the oxidative stress defense system of *S. thermophilus*, a collection of random  
376 insertion mutants was screened to select clones with high sensitivity to menadione. Using this  
377 approach, we aimed to identify genes involved in superoxide stress defense without  
378 presuming their function. It can be noted that no gene encoding a protein considered, in other  
379 organisms, as major antioxidant enzyme were found using this approach. This can be due to  
380 the fact that (i) the collection of mutants was non-exhaustive (ii) the corresponding mutant  
381 could have a weak phenotype (iii) such mutations could be lethal (iv) except from SOD, most  
382 anti-oxidant enzymes are protective against H<sub>2</sub>O<sub>2</sub> and not O<sub>2</sub><sup>-</sup>. The genes identified in the  
383 present work were classified into 5 groups according to their function: genes potentially  
384 involved in cell wall metabolism, exopolysaccharide translocation, tRNA modification, and  
385 iron metabolism and gene of unknown function.

386 Perhaps the most striking observation of this study is that 5 out of the 10 sensitive mutants  
387 selected are disrupted in a gene encoding an membrane protein (OssH, MreD, RodA, PBP2b  
388 and CpsX). This could reflect that the cell envelope is the main target of the stress applied.  
389 Among these mutants, three (*mreD*, *rodA* and *pbp2b*) are impaired in genes potentially  
390 encoding proteins described in *E. coli* as playing a role in peptidoglycan biosynthesis during  
391 cell elongation (28, 52, 63). Phenotypic analysis of these three *S. thermophilus* mutants  
392 showed that they all present a reduced cell size associated with a modification of the cell  
393 shape, as observed in *rodA*, *pbp2* and *mre* mutants of *E. coli* or *Salmonella typhimurium* (5,  
394 20, 28, 29, 63, 64). Moreover, the 3 *S. thermophilus* mutants also presented an increased  
395 sensitivity to heat shock compared to the wild type strain suggesting that the protective role of  
396 MreD, RodA and PBP2b is not specific to oxidative stress but could be enlarged to a more  
397 global stress response. In support of this hypothesis is the existence in *Lactococcus lactis* of  
398 mutations in *pbp* genes that negatively affect acid tolerance of a *guaA* mutant (16).

399 The major function assigned to cell wall is the maintenance of cell integrity by protecting  
400 cells not only against osmotic stress but also against other stress conditions. For instance, in  
401 LAB, it has already been shown that compromised biosynthesis of peptidoglycan results in an  
402 increased sensitivity to either UV (22) or acid (11) stress. Additionally, disruption of *mreD*,  
403 *rodA* or *pbp2b* may cause a modification of the peptidoglycan architecture in *S. thermophilus*,  
404 reducing the cell protection against aggressions such as oxidative and heat stress.

405

406 Transfer RNA modification is a universal feature of all living organisms, although some  
407 described modifications have no assigned function yet. Modifying enzymes affect the tRNA  
408 precursor either by chemical modification of an existing residue (in the base or in the sugar)  
409 or by nucleotide substitution. These modifications, in particular those present in the anticodon  
410 region can affect translation efficiency and fidelity (for a review (10)). In this study, two  
411 clones disrupted in the loci *tgt* and *ossF*, encoding proteins showing high identity with tRNA  
412 modifying enzymes were selected for their high sensitivity to oxidative stress. This sensitivity  
413 could suggest that the expression of genes involved in oxidative stress defense may require  
414 tRNA maturation or, that such modifications could help tRNA stabilization under stress  
415 conditions. In agreement with the former hypothesis *Shigella flexneri* pathogenicity is  
416 regulated by *virF* mRNA translation efficiency under TGT activity control (21). Additionally,  
417 several environmental conditions are suggested to influence tRNA modification: in the case of  
418 queuosine (whose synthesis and substitution is carried out by several enzymes among which  
419 is TGT), synthesis is sensitive to the presence of iron (31) and TGT requires oxygen for its  
420 full activity (54). Therefore, further investigations on this topic should provide an interesting  
421 new regulatory model and could contribute towards the emerging observation that  
422 translational regulation plays an important role in oxidative stress response, a role mainly  
423 ascribed to transcriptional regulation thus far.

424



425 Iron is essential for cellular metabolism, due to its requirement as a cofactor for a large number  
426 of enzymes. However, an excess of iron is toxic because of its involvement in the Fenton  
427 reaction that enhances the formation of ROS. In this work, a gene homologous to *fatD*, encoding  
428 a putative Fe<sup>3+</sup> ABC-transporter was found to be involved in *S. thermophilus* oxidative stress  
429 response. To test whether or not *S. thermophilus* FatD is indeed an iron transporter, the  
430 sensitivity of the *fatD* mutant to streptonigrin (an iron activated-antibiotic) was assessed. No  
431 significant difference between the mutant and the wild type strains was detected (data not  
432 shown) suggesting that disruption of *fatD* would have no effect on intracellular iron  
433 concentration. Thus, either FatD is not a major iron transporter under the experimental procedure  
434 we used or *S. thermophilus fatD* encodes an ABC transporter for another metallic cation. A  
435 possible cation could be Mn<sup>2+</sup> known to have a protective function against oxidative damage  
436 either via direct dismutase activity, such as reported in other organisms (4, 26, 60), or via its role  
437 as a cofactor of the only SOD reported into *S. thermophilus* (19). Thus, inactivation of a Mn<sup>2+</sup>  
438 ABC transporter could trigger sensitivity to oxidative stress.

439

440 [Fe-S] clusters are oxygen sensitive prosthetic groups, incorporated into proteins via covalent  
441 bonds to cysteines. Iron-sulfur proteins, which are found in more than 120 classes of enzymes  
442 and proteins (30), are known to play important physiological roles in electron transfer,  
443 metabolic reactions and in transcriptional regulation (30, 32). Indeed, because of their  
444 intrinsic redox properties, they are likely to sense oxygen or its derivatives as it is the case in  
445 SoxR and FNR transcriptional regulators, responding to oxidative stress and oxygen  
446 availability, (for a review (32)). IscR is also a iron-sulfur transcriptional regulator that is  
447 suggested to respond to Fe-S cluster limiting conditions and potentially to hydrogen peroxide  
448 in *E. coli* (32). Although it has been shown that [Fe-S] cluster assembly can occur  
449 spontaneously *in vitro* under anaerobic conditions (36), this process can also be catalyzed  
450 enzymatically *in vivo*. Genes of the *suf* operon and of the *isc* cluster (iron sulfur cluster) are

451 conserved and encode [Fe-S] cluster biosynthesis and/or repair pathways (23). More  
452 precisely, it has been recently proposed that Isc proteins are necessary for biosynthesis, while  
453 Suf proteins are essentially involved in repair of oxidatively damaged [Fe-S] clusters (37).  
454 Study of  $\Delta$ *sufD* and *iscU*<sub>97</sub> non-polar mutants indicates that SufD and IscU proteins are  
455 required for growth in an iron-deficient medium. SufD and IscU inactivation conferred an  
456 increased resistance and sensitivity respectively to streptonigrin. These results indicate that  
457 *S. thermophilus* SufD and IscU are involved in intracellular iron balance. Additionally, SufD  
458 and IscU of *S. thermophilus*, similar to SufD and NifU family proteins from other  
459 microorganisms, are encoded by genes clustered with *sufB*, *sufC* and *sufS* homologues, also  
460 potentially involved in [Fe-S] cluster repair. Hence the participation of *S. thermophilus* SufD  
461 and IscU proteins in iron intracellular balance is probably the result of their involvement in  
462 [Fe-S] cluster repair, as in *Er. chrysanthemi* (37). The NifU family protein function is to  
463 provide a scaffold for assembly of clusters mediated by NifS activity, and such clusters can  
464 then be transferred into apo forms of [Fe-S] cluster containing proteins (1, 70). The role of the  
465 SufD in [Fe-S] cluster biosynthesis is not well established. However, in *S. thermophilus*, the  
466 increased resistance to streptonigrin of the  $\Delta$ *sufD* mutant suggests that its intracellular iron  
467 concentration is lower than in the wild type. At least two alternative hypotheses can be drawn:  
468 (i) SufD could be directly involved in the uptake or assimilation of extracellular iron, (ii)  
469 SufD could inhibit the synthesis and/or repair of the [Fe-S] clusters. In this latter case, SufD  
470 inactivation would result in an increase synthesis of [Fe-S] clusters, and consequently in a  
471 reduced intracellular free iron concentration.  
472 Additionally, *S. thermophilus* *sufD* and *iscU* mutants displayed an increased sensitivity to  
473 superoxide radicals compared to wild type. Thus, SufD and IscU are likely to play a major  
474 role in oxidative stress defense, probably resulting from their predicted role in [Fe-S] cluster  
475 assembly and/or repair. The involvement of the *suf* operon in oxidative stress resistance  
476 during the early steps of infection by *Er. chrysanthemi* has also been reported (37), as well as

477 the induction of the *suf* gene cluster of *E. coli* by the hydrogen peroxide response regulator  
478 OxyR (72). In spite of a possible spontaneous formation, our data suggest that under oxidative  
479 stress, [Fe-S] cluster assembly and/or repair becomes a limiting step in the formation of iron-  
480 sulfur proteins, and that the cell requires an additional, dedicated system, encoded by the *suf*  
481 gene cluster, to optimize [Fe-S] cluster formation or repair.

482

483 Conclusion: The strategy used in this work allowed the identification of genes involved in  
484 *S. thermophilus* oxidative stress defense system. Data concerning the function of some loci is  
485 already available in the literature, whereas the role of others is not known. In both cases, the  
486 exact roles of these genes in the defense of *S. thermophilus* against oxidative stress remains to  
487 be elucidated.

## ACKNOWLEDGEMENTS

488

489

490 A. T and A. F. were supported by grants from the Ministère de la recherche. F. B. was  
491 supported by a grant from the Institut National de la Recherche Agronomique.

492 We are grateful to Paul Hoskinson for his advice regarding the English formulation of the  
493 manuscript.

494

## References

495

496

- 497 1. **Agar, J. N., C. Krebs, J. Frazzon, B. H. Huynh, D. R. Dean, and M. K. Johnson.**  
498 2000. IscU as a scaffold for iron-sulfur cluster biosynthesis: sequential assembly of  
499 [2Fe-2S] and [4Fe-4S] clusters in IscU. *Biochemistry* **39**:7856-7862.
- 500 2. **Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman.** 1990. Basic  
501 local alignment search tool. *J Mol Biol* **215**:403-410.
- 502 3. **Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and**  
503 **D. J. Lipman.** 1997. Gapped BLAST and PSI-BLAST: a new generation of protein  
504 database search programs. *Nucleic Acids Res* **25**:3389-3402.
- 505 4. **Archibald, F. S., and I. Fridovich.** 1981. Manganese, superoxide dismutase, and  
506 oxygen tolerance in some lactic acid bacteria. *J Bacteriol* **146**:928-36.
- 507 5. **Begg, K. J., and W. D. Donachie.** 1985. Cell shape and division in *Escherichia coli*:  
508 experiments with shape and division mutants. *J Bacteriol* **163**:615-622.
- 509 6. **Benov, L. T., and I. Fridovich.** 1994. *Escherichia coli* expresses a copper- and zinc-  
510 containing superoxide dismutase. *J Biol Chem* **269**:25310-25314.
- 511 7. **Berlett, B. S., and E. R. Stadtman.** 1997. Protein oxidation in aging, disease, and  
512 oxidative stress. *J Biol Chem* **272**:20313-20316.

- 513 8. **Biswas, I., A. Gruss, S. D. Ehrlich, and E. Maguin.** 1993. High-efficiency gene  
514 inactivation and replacement system for gram-positive bacteria. *J Bacteriol* **175**:3628-  
515 3635.
- 516 9. **Bjork, G. R.** 1975. Transductional mapping of gene *trmA* responsible for the  
517 production of 5-methyluridine in transfer ribonucleic acid of *Escherichia coli*. *J*  
518 *Bacteriol* **124**:92-98.
- 519 10. **Bjork, G. R., J. M. Durand, T. G. Hagervall, R. Leipuviene, H. K. Lundgren, K.**  
520 **Nilsson, P. Chen, Q. Qian, and J. Urbonavicius.** 1999. Transfer RNA modification:  
521 influence on translational frameshifting and metabolism. *FEBS Lett* **452**:47-51.
- 522 11. **Boyd, D. A., D. G. Cvitkovitch, A. S. Bleiweis, M. Y. Kiriukhin, D. V. Debabov, F.**  
523 **C. Neuhaus, and I. R. Hamilton.** 2000. Defects in D-alanyl-lipoteichoic acid  
524 synthesis in *Streptococcus mutans* results in acid sensitivity. *J Bacteriol* **182**:6055-  
525 6065.
- 526 12. **Bracquart, P.** 1981. An agar medium for the differential enumeration of  
527 *Streptococcus thermophilus* and *Lactobacillus bulgaricus* in yogurth. *J appl bacteriol*  
528 **51**:303-305.
- 529 13. **Briolat, V., and G. Reyssset.** 2002. Identification of the *Clostridium perfringens* genes  
530 involved in the adaptive response to oxidative stress. *J Bacteriol* **184**:2333-2343.
- 531 14. **Broadbent, J., D. McMahon, C. Oberg, and D. Welker.** 2001. Use of  
532 exopolysaccharide-producing cultures to improve the functionality of low fat cheese.  
533 *Int Dairy J* **11**:433-439.
- 534 15. **Broadbent, J., D. McMahon, D. Welker, C. Oberg, and S. Moineau.** 2003.  
535 Biochemistry, genetics, and applications of exopolysaccharide production in  
536 *Streptococcus thermophilus*: a review. *J Dairy Sci* **86**:407-423.

- 537 16. **Budin, A., V. Pichereau, E. Domakova, J. Tremblay, R. Dervyn, E. Auffray, S. D.**  
538 **Ehrlich, S. Kulakauskas, and E. Maguin.** 2002. Effect of a *guaA* mutation on the  
539 acid tolerance of *L. lactis*. *Sciences des aliments* **22**:67-74.
- 540 17. **Burger, A., K. Sichler, G. Kelemen, M. Buttner, and W. Wohlleben.** 2000.  
541 Identification and characterization of the *mre* gene region of *Streptomyces coelicolor*  
542 A3(2). *Mol Gen Genet* **263**:1053-1060.
- 543 18. **Chae, H. Z., K. Robison, L. B. Poole, G. Church, G. Storz, and S. G. Rhee.** 1994.  
544 Cloning and sequencing of thiol-specific antioxidant from mammalian brain: alkyl  
545 hydroperoxide reductase and thiol-specific antioxidant define a large family of  
546 antioxidant enzymes. *Proc Natl Acad Sci U S A* **91**:7017-7021.
- 547 19. **Chang, S. K., and H. M. Hassan.** 1997. Characterization of superoxide dismutase in  
548 *Streptococcus thermophilus*. *Appl Environ Microbiol* **63**:3732-3735.
- 549 20. **Costa, C. S., and D. N. Anton.** 1993. Round-cell mutants of *Salmonella typhimurium*  
550 produced by transposition mutagenesis: lethality of *rodA* and *mre* mutations. *Mol Gen*  
551 *Genet* **236**:387-394.
- 552 21. **Durand, J. M., B. Dagberg, B. E. Uhlin, and G. R. Bjork.** 2000. Transfer RNA  
553 modification, temperature and DNA superhelicity have a common target in the  
554 regulatory network of the virulence of *Shigella flexneri*: the expression of the *virF*  
555 gene. *Mol Microbiol* **35**:924-35.
- 556 22. **Duwat, P., A. Cochu, S. D. Ehrlich, and A. Gruss.** 1997. Characterization of  
557 *Lactococcus lactis* UV-sensitive mutants obtained by *ISS1* transposition. *J Bacteriol*  
558 **179**:4473-9.
- 559 23. **Ellis, K. E., B. Clough, J. W. Saldanha, and R. J. Wilson.** 2001. *Nifs* and *Sufs* in  
560 malaria. *Mol Microbiol* **41**:973-981.
- 561 24. **Faguy, D. M., and W. F. Doolittle.** 2000. Horizontal transfer of catalase-peroxidase  
562 genes between archaea and pathogenic bacteria. *Trends Genet* **16**:196-197.

- 563 25. **Farr, S. B., and T. Kogoma.** 1991. Oxidative stress responses in *Escherichia coli* and  
564 *Salmonella typhimurium*. Microbiol Rev **55**:561-585.
- 565 26. **Fridovich, I.** 1983. Superoxide radical: an endogenous toxicant. Annu Rev Pharmacol  
566 Toxicol **23**:239-57.
- 567 27. **Gibson, C. M., T. C. Mallett, A. Claiborne, and M. G. Caparon.** 2000.  
568 Contribution of NADH oxidase to aerobic metabolism of *Streptococcus pyogenes*. J  
569 Bacteriol **182**:448-55.
- 570 28. **Ishino, F., W. Park, S. Tomioka, S. Tamaki, I. Takase, K. Kunugita, H.**  
571 **Matsuzawa, S. Asoh, T. Ohta, B. G. Spratt, and et al.** 1986. Peptidoglycan  
572 synthetic activities in membranes of *Escherichia coli* caused by overproduction of  
573 penicillin-binding protein 2 and RodA protein. J Biol Chem **261**:7024-7031.
- 574 29. **Iwaya, M., R. Goldman, D. J. Tipper, B. Feingold, and J. L. Strominger.** 1978.  
575 Morphology of an *Escherichia coli* mutant with a temperature-dependent round cell  
576 shape. J Bacteriol **136**:1143-1158.
- 577 30. **Johnson, M. K.** 1998. Iron-sulfur proteins: new roles for old clusters. Curr Opin  
578 Chem Biol **2**:173-81.
- 579 31. **Kersten, H., and W. Kersten.** 1990. Biosynthesis and function of queuine and  
580 queuosine tRNAs, p. B69-B108. In C. W. G. a. K. C. T. K. (ed.) (ed.),  
581 Chromatography and modification of nucleosides. Elsevier, Amsterdam.
- 582 32. **Kiley, P. J., and H. Beinert.** 2003. The role of Fe-S proteins in sensing and regulation  
583 in bacteria. Curr Opin Microbiol **6**:181-5.
- 584 33. **King, K. Y., J. A. Horenstein, and M. G. Caparon.** 2000. Aerotolerance and  
585 peroxide resistance in peroxidase and PerR mutants of *Streptococcus pyogenes*. J  
586 Bacteriol **182**:5290-5299.
- 587 34. **Leenhouts, K.** 1995. Integration strategies and vectors. Dev Biol Stand **85**:523-530.

- 588 35. **Maguin, E., H. Prevost, S. D. Ehrlich, and A. Gruss.** 1996. Efficient insertional  
589 mutagenesis in lactococci and other gram-positive bacteria. *J Bacteriol* **178**:931-935.
- 590 36. **Merchant, S., and B. W. Dreyfuss.** 1998. Metalloprotein and Cytochrome Assembly  
591 in Chloroplasts. *Annu Rev Plant Physiol Plant Mol Biol* **49**:25-51.
- 592 37. **Nachin, L., M. El Hassouni, L. Loiseau, D. Expert, and F. Barras.** 2001. SoxR-  
593 dependent response to oxidative stress and virulence of *Erwinia chrysanthemi*: the key  
594 role of SufC, an orphan ABC ATPase. *Mol Microbiol* **39**:960-972.
- 595 38. **Nishio, K., and M. Nakai.** 2000. Transfer of iron-sulfur cluster from NifU to  
596 apoferredoxin. *J Biol Chem* **275**:22615-22618.
- 597 39. **Noguchi, S., Y. Nishimura, Y. Hirota, and S. Nishimura.** 1982. Isolation and  
598 characterization of an *Escherichia coli* mutant lacking tRNA-guanine  
599 transglycosylase. Function and biosynthesis of queuosine in tRNA. *J Biol Chem*  
600 **257**:6544-6550.
- 601 40. **Okada, N., S. Noguchi, H. Kasai, N. Shindo-Okada, T. Ohgi, T. Goto, and S.**  
602 **Nishimura.** 1979. Novel mechanism of post-transcriptional modification of tRNA.  
603 Insertion of bases of Q precursors into tRNA by a specific tRNA transglycosylase  
604 reaction. *J Biol Chem* **254**:3067-3073.
- 605 41. **O'Sullivan, T., and G. Fitzgerald.** 1999. Electrotransformation of industrial strains  
606 of *Streptococcus thermophilus*. *J Appl Microbiol* **86**:275-283.
- 607 42. **Patzer, S. I., and K. Hantke.** 1999. SufS is a NifS-like protein, and SufD is necessary  
608 for stability of the [2Fe-2S] FhuF protein in *Escherichia coli*. *J Bacteriol* **181**:3307-  
609 3309.
- 610 43. **Pébay, M., A. C. Holl, J. M. Simonet, and B. Decaris.** 1995. Characterization of the  
611 *gor* gene of the lactic acid bacterium *Streptococcus thermophilus* CNRZ368. *Res*  
612 *Microbiol* **146**:371-383.



- 613 44. **Poole, L. B., M. Higuchi, M. Shimada, M. L. Calzi, and Y. Kamio.** 2000.  
614 *Streptococcus mutans* H<sub>2</sub>O<sub>2</sub>-forming NADH oxidase is an alkyl hydroperoxide  
615 reductase protein. *Free Radic Biol Med* **28**:108-120.
- 616 45. **Poyart, C., E. Pellegrini, O. Gaillot, C. Boumaila, M. Baptista, and P. Trieu-Cuot.**  
617 2001. Contribution of Mn-cofactored superoxide dismutase (SodA) to the virulence of  
618 *Streptococcus agalactiae*. *Infect Immun* **69**:5098-5106.
- 619 46. **Rocha, E. R., G. Owens, Jr., and C. J. Smith.** 2000. The redox-sensitive  
620 transcriptional activator OxyR regulates the peroxide response regulon in the obligate  
621 anaerobe *Bacteroides fragilis*. *J Bacteriol* **182**:5059-5069.
- 622 47. **Rocha, E. R., T. Selby, J. P. Coleman, and C. J. Smith.** 1996. Oxidative stress  
623 response in an anaerobe, *Bacteroides fragilis*: a role for catalase in protection against  
624 hydrogen peroxide. *J Bacteriol* **178**:6895-6903.
- 625 48. **Rocha, E. R., and C. J. Smith.** 1998. Characterization of a peroxide-resistant mutant  
626 of the anaerobic bacterium *Bacteroides fragilis*. *J Bacteriol* **180**:5906-5912.
- 627 49. **Rocha, E. R., and C. J. Smith.** 1999. Role of the alkyl hydroperoxide reductase  
628 (*ahpCF*) gene in oxidative stress defense of the obligate anaerobe *Bacteroides fragilis*.  
629 *J Bacteriol* **181**:5701-10.
- 630 50. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. Molecular cloning : a laboratory  
631 manual. Cold Spring Harbor Laboratory, New York.
- 632 51. **Sonnhammer, E. L. L., G. Von Heijne, and A. Krogh.** 1998. A hidden Markov  
633 model for predicting transmembrane helices in protein sequences, p. 175-182. *In* J.  
634 Glasgow, T. Littlejohn, F. Major, L. R., D. Sankoff, and C. Sensen (ed.), Proceedings  
635 of the sixth international conference on intelligent systems for molecular  
636 microbiology. AAAI Press, Menlo park, CA.

- 637 52. **Spratt, B. G.** 1975. Distinct penicillin binding proteins involved in the division,  
638 elongation, and shape of *Escherichia coli* K12. *Proc Natl Acad Sci U S A* **72**:2999-  
639 3003.
- 640 53. **Stadtman, E. R.** 1993. Oxidation of free amino acids and amino acid residues in  
641 proteins by radiolysis and by metal-catalyzed reactions. *Annu Rev Biochem* **62**:797-  
642 821.
- 643 54. **Szabo, L., S. Nishimura, and W. R. Farkas.** 1988. Possible involvement of queuine  
644 in oxidative metabolism. *Biofactors* **1**:241-4.
- 645 55. **Terzaghi, B., and W. Sandine.** 1975. Improved medium for lactic *streptococci* and  
646 their bacteriophages. *Appl Environ Microbiol* **29**:807-813.
- 647 56. **Thibessard, A., A. Fernandez, B. Gintz, B. Decaris, and N. Leblond-Bourget.**  
648 2002. Transposition of pGh9:ISS1 is random and efficient in *Streptococcus*  
649 *thermophilus* CNRZ368. *Can J Microbiol* **48**:473-478.
- 650 57. **Thibessard, A., A. Fernandez, B. Gintz, N. Leblond-Bourget, and B. Decaris.**  
651 2002. Effects of *rodA* and *pbp2b* disruption on cell morphology and oxidative stress  
652 response of *Streptococcus thermophilus* CNRZ368. *J Bacteriol* **184**:2821-2826.
- 653 58. **Thibessard, A., A. Fernandez, B. Gintz, N. Leblond-Bourget, and B. Decaris.**  
654 2001. Hydrogen peroxide effects on *Streptococcus thermophilus* CNRZ368 cell  
655 viability. *Res Microbiol* **152**:593-596.
- 656 59. **Thibessard, A., N. Leblond-Bourget, A. Fernandez, B. Gintz, and B. Decaris.**  
657 2001. Response of *Streptococcus thermophilus* CNRZ368 and its colonial variants to  
658 oxidative stress : evidence for an inducible defence system. *Lait* **81**:311-316.
- 659 60. **Tseng, H., Y. Srikhanta, A. McEwan, and M. Jennings.** 2001. Accumulation of  
660 manganese in *Neisseria gonorrhoeae* correlates with resistance to oxidative killing by  
661 superoxide anion and is independent of superoxide dismutase activity. *Mol Microbiol*  
662 **40**:1175-1186.

- 663 61. **Tusnady, G. E., and I. Simon.** 1998. Principles governing amino acid composition of  
664 integral membrane proteins: application to topology prediction. *J Mol Biol* **283**:489-  
665 506.
- 666 62. **Visca, P., A. Ciervo, V. Sanfilippo, and N. Orsi.** 1993. Iron-regulated salicylate  
667 synthesis by *Pseudomonas spp.* *J Gen Microbiol* **139**:1995-2001.
- 668 63. **Wachi, M., M. Doi, Y. Okada, and M. Matsuhashi.** 1989. New mre genes *mreC* and  
669 *mreD*, responsible for formation of the rod shape of *Escherichia coli* cells. *J Bacteriol*  
670 **171**:6511-6516.
- 671 64. **Wachi, M., M. Doi, S. Tamaki, W. Park, S. Nakajima-Iijima, and M. Matsuhashi.**  
672 1987. Mutant isolation and molecular cloning of *mre* genes, which determine cell  
673 shape, sensitivity to mecillinam, and amount of penicillin-binding proteins in  
674 *Escherichia coli*. *J Bacteriol* **169**:4935-4940.
- 675 65. **Wu, J., and B. Weiss.** 1992. Two-stage induction of the *soxRS* (superoxide response)  
676 regulon of *Escherichia coli*. *J Bacteriol* **174**:3915-3920.
- 677 66. **Yamamoto, Y., M. Higuchi, L. B. Poole, and Y. Kamio.** 2000. Role of the *dpr*  
678 product in oxygen tolerance in *Streptococcus mutans*. *J Bacteriol* **182**:3740-3747.
- 679 67. **Yesilkaya, H., A. Kadioglu, N. Gingles, J. E. Alexander, T. J. Mitchell, and P. W.**  
680 **Andrew.** 2000. Role of manganese-containing superoxide dismutase in oxidative  
681 stress and virulence of *Streptococcus pneumoniae*. *Infect Immun* **68**:2819-2826.
- 682 68. **Youn, H. D., H. Youn, J. W. Lee, Y. I. Yim, J. K. Lee, Y. C. Hah, and S. O. Kang.**  
683 1996. Unique isozymes of superoxide dismutase in *Streptomyces griseus*. *Arch*  
684 *Biochem Biophys* **334**:341-348.
- 685 69. **Yu, J., A. P. Bryant, A. Marra, M. A. Lonetto, K. A. Ingraham, A. F. Chalker, D.**  
686 **J. Holmes, D. Holden, M. Rosenberg, and D. McDevitt.** 2001. Characterization of  
687 the *Streptococcus pneumoniae* NADH oxidase that is required for infection.  
688 *Microbiology* **147**:431-438.

- 689 70. **Yuvaniyama, P., J. N. Agar, V. L. Cash, M. K. Johnson, and D. R. Dean.** 2000.  
690 NifS-directed assembly of a transient [2Fe-2S] cluster within the NifU protein. Proc  
691 Natl Acad Sci U S A **97**:599-604.
- 692 71. **Zheng, L., R. H. White, V. L. Cash, R. F. Jack, and D. R. Dean.** 1993. Cysteine  
693 desulfurase activity indicates a role for NIFS in metallocluster biosynthesis. Proc Natl  
694 Acad Sci U S A **90**:2754-2758.
- 695 72. **Zheng, M., X. Wang, L. J. Templeton, D. R. Smulski, R. A. LaRossa, and G.**  
696 **Storz.** 2001. DNA microarray-mediated transcriptional profiling of the *Escherichia*  
697 *coli* response to hydrogen peroxide. J Bacteriol **183**:4562-70.

698

699

#### LEGENDS FIGURES

700

701 **FIG. 1. Relative methyl viologen sensitivity.** (A). Results of short term methyl viologen  
702 exposure. (B) Results of long term methyl viologen exposure (see materials and methods).

703 The relative methyl viologen sensitivity was determined by comparing viability with and  
704 without treatment. Levels are shown as values relative to the survival level of the wild type  
705 strain. All experiments were performed at least in duplicate. Error bars show two standard  
706 deviations of the data.

707

708 **FIG. 2. Survival to heat shock.** The heat shock consisted of 15 min treatment at 60°C. The  
709 percentage of surviving cells corresponds to the number of colony forming units (CFUs)  
710 counted on plates after the heat-shock exposure divided by the number of CFU before  
711 treatment. Error bars show two standard deviations of the data. The dotted line represents the  
712 surviving percentage observed for the wild type strain and the grey band comprises 2 standard  
713 deviations of the data.

714

715 **FIG. 3. Influence of *mreD* disruption on chain length.** Cell number per chains was  
716 determined for 70 chains of the wild type strain (■) and of the *mreD* mutant (■) grown in  
717 liquid TPPY medium.

718

719 **FIG. 4. Schematic map of the *S. thermophilus* *sufCDS iscU sufB* cluster.** Genes are  
720 indicated by grey arrows, and the positions of putative start and stop codons are given (in bp)  
721 below the map. The site of insertion of the pGh9:ISSI plasmid in *sufD* (*ossI*) and *iscU* (*ossJ*)  
722 disrupted mutants is indicated by the solid black triangles. The broken arrow indicates the  
723 position of the *suf* cluster putative promoter. Putative rho-independent terminator is  
724 represented as hairpin loop. Fragments used for generation of the non-polar mutants are  
725 indicated as open boxes. The star indicates the position of the *EcoRV* restriction site causing  
726 the Cys<sub>97</sub> replacement by a Tyr in the *iscU*<sub>97</sub> mutant.

727

728 **FIG. 5. Streptonigrin tolerance.** *S. thermophilus* strain tolerance to streptonigrin was  
729 assessed by measuring OD<sub>600</sub> after growth in TPPYC medium containing various  
730 concentrations of streptonigrin without (A) or supplemented with 1 mM of 2,2'-dipyridyl (B)  
731 (see material and methods for details). Relative OD<sub>600nm</sub> was determined by comparing the  
732 OD<sub>600</sub> obtained with or without streptonigrin. Wild type strain (○ and ●); *iscU*<sub>97</sub> (□ and ■);  
733  $\Delta$ *sufD* (Δ and ▲).

734 TABLE 1. Products of *oss* genes: database similarities and putative functions.

735

Gene name <sup>a</sup>	Accession number	Gene encoding the most similar product <sup>c</sup>	amino acid identity <sup>d</sup>	Insertion site <sup>e</sup>	Putative function	Class
<i>ossA</i> ( <i>mreD</i> )	AY386242	<i>S. mutans mreD</i> (168)	30% (1-168)	127	Cell shape maintenance	I
<i>ossB</i> ( <i>rodA</i> )	AF399832	<i>S. mutans rodA</i> (408)	65% (1-391)	304	Peptidoglycan elongation	
<i>ossC</i> ( <i>pbp2b</i> ) <sup>b</sup>	AF399833	<i>S. thermophilus Sfi6 pbp2b</i> (704)	98% (641-704)	640	Peptidoglycan elongation	
<i>ossD</i> ( <i>cpsX</i> ) <sup>b</sup>	AY386241	<i>S. thermophilus cpsX</i> (378)	98% (212-378)	264	Membrane translocation of exopolysaccharide	II
<i>ossE</i> ( <i>tgt</i> ) <sup>b</sup>	AY386243	<i>S. pneumoniae tgt</i> (380)	92% (1-333)	99	tRNA-guanine transglycosylase	III
<i>ossF</i> <sup>b</sup>	AY386244	<i>S. agalactiae gbs0448</i> (451)	77% (1-204)	51	RNA methyltransferase	
<i>ossG</i>	AY386239	<i>S. mutans smu.1428c</i> (284)	67% (1-280)	170	Unknown conserved function	IV
<i>ossH</i> ( <i>fatD</i> )	AY386240	<i>S. pneumoniae fatD</i> (324)	55% (22-323)	119	Iron ABC transporter	V
<i>ossI</i> ( <i>sufD</i> )	AY386245	<i>S. agalactiae sag0142</i> (420)	76% (1-420)	144	Assembly of Fe/S clusters	
<i>ossJ</i> ( <i>iscU</i> )	AY386245	<i>S. agalactiae gbs0140</i> (147)	78% (1-141)	57	Assembly of Fe/S clusters	

736

737 a. Initial gene name with, in brackets and when applicable, the definitive gene name.

738 b. Disrupted ORF of *S. thermophilus* CNRZ368 that have been partially sequenced.

739 c. Organism, gene name and, in brackets, length of the protein sequence giving the best alignment  
740 score.

741 d. Percentage of sequence identity with the protein showing the best alignment score to each  
742 *S. thermophilus* Oss protein. In brackets is indicated the portion of the protein sharing this identity  
743 (numbers correspond to the amino acid positions).

744 e. The numbers correspond to the deduced amino acid position of the pGh9:ISS1-generated  
745 disruption in the full length protein showing the best alignment score to the Oss proteins.

746 TABLE 2. *suf* cluster gene products of *S. thermophilus* CNRZ368

747

Gene product	Closest similar protein (organism, accession number) <sup>a</sup>	Identity/similarity(%)	Function
SufC	CAD45782 ( <i>S. agalactiae</i> NEM316)	88/94	Putative ABC transporter (ATP-binding protein)
SufD	AAM99050 ( <i>S. agalactiae</i> 2603V/R)	76/87	Unknown, conserved protein
SufS	CAD45784 ( <i>S. agalactiae</i> NEM316)	79/89	Similar to SufS
IscU	CAD45785 ( <i>S. agalactiae</i> NEM316)	78/90	Similar to NifU family proteins
SufB	CAD45786 ( <i>S. agalactiae</i> NEM316)	89/94	Unknown, conserved protein

748

749 a. Accession number of the protein giving the best alignment score. In brackets, the name of the

750 organism is indicated.

751 TABLE 3. Growth of wild type and mutant strains in TPPYC and in TPPYC supplemented with  
752 iron.

753

Strain	Final OD <sub>600</sub> resulting from growth <sup>a</sup> in:	
	TPPYC	TPPYC + FeSO <sub>4</sub>
WT	0.96	1.32
<i>ΔsufD</i>	0.58	1.48
<i>iscU<sub>97</sub></i>	0.58	1.15

754

755 <sup>a</sup> Cells were grown in milk medium for 15<sup>o</sup>h at 42°C and diluted in FeSO<sub>4</sub>-deficient medium  
756 (TPPYC medium supplemented with MgCl<sub>2</sub> (1 mM) and CaCl<sub>2</sub> (1 mM)). After one night  
757 growth, cells were washed twice and resuspend either in FeSO<sub>4</sub>-deficient medium or  
758 supplemented with 100 μM FeSO<sub>4</sub>. The growth of the cells was then followed by measuring  
759 the OD<sub>600</sub>. The OD<sub>600</sub> were measured after 24h incubation at 42°C. This experiment was done  
760 in triplicate.

761