

The N-terminal Domain of PILB from *Neisseria meningitidis* Is a Disulfide Reductase That Can Recycle Methionine Sulfoxide Reductases

Junzhu Wu, Fabrice Neiers, Sandrine Boschi-Muller, Guy Branlant

► **To cite this version:**

Junzhu Wu, Fabrice Neiers, Sandrine Boschi-Muller, Guy Branlant. The N-terminal Domain of PILB from *Neisseria meningitidis* Is a Disulfide Reductase That Can Recycle Methionine Sulfoxide Reductases. *Journal of Biological Chemistry*, American Society for Biochemistry and Molecular Biology, 2005, 280 (13), pp.12344-12350. 10.1074/jbc.M500385200 . hal-01690798

HAL Id: hal-01690798

<https://hal.univ-lorraine.fr/hal-01690798>

Submitted on 23 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The N-terminal Domain of PilB from *Neisseria meningitidis* Is a Disulfide Reductase That Can Recycle Methionine Sulfoxide Reductases*

Received for publication, January 11, 2005, and in revised form, January 21, 2005
Published, JBC Papers in Press, January 24, 2005, DOI 10.1074/jbc.M500385200

Junzhu Wu‡§, Fabrice Neiers‡, Sandrine Boschi-Muller, and Guy Branlant¶

From the UMR CNRS-UHP 7567, Maturation des ARN et Enzymologie Moléculaire, Faculté des Sciences, Bld des Aiguillettes, BP 239, 54506 Vandoeuvre-les-Nancy, France

The PilB protein of the *Neisseria* genus comprises three domains. Two forms have been recently reported to be produced *in vivo*. One form, containing the three domains, is secreted from the bacterial cytoplasm to the outer membrane, whereas the second form, which is cytoplasmic, only contains the central and the C-terminal domains. The secreted form was shown to be involved in survival under oxidative conditions. Although previous studies indicated that the central and the C-terminal domains display methionine sulfoxide reductase A and B activities, respectively, no function was described so far for the N-terminal domain. In the present study, the N-terminal domain of the PilB of *Neisseria meningitidis* was produced as a folded entity, and its biochemical and enzymatic properties have been determined. The data show that the N-terminal domain possesses a disulfide redox-active site with a redox potential in the range of that of thioredoxin. Moreover, the N-terminal domain, either as an isolated form or included in PilB, recycles the oxidized forms of the methionine sulfoxide reductases like thioredoxin. These results, which show that the N-terminal domain exhibits a disulfide reductase activity and probably has a thioredoxin-fold, are discussed in relation to its possible functional role in *Neisseria*.

The obligate human pathogens, *Neisseria gonorrhoeae* and *Neisseria meningitidis* are the only two pathogenic members of the *Neisseria* family of Gram-negative bacteria. *N. gonorrhoeae*, which colonizes mucosal epithelia of the genitourinary tract, is the causative agent of the disease gonorrhea, whereas *N. meningitidis*, which colonizes the nasopharynx, is the cause of two serious human diseases, pyogenic meningitis and meningococcal septicemia. Like many bacterial pathogens, the ability of these *Neisseriae* to infect their host is conditioned by successive steps of interactions with host cells and by the

survival of the bacteria in the host environment (for review see Refs. 1 and 2). In particular, the bacteria have to resist the oxidative burst of the host, which generates a variety of reactive oxygen species. *N. gonorrhoeae* and *N. meningitidis* possess several antioxidant defense mechanisms among which the PilB protein was recently shown to play an essential role (3).

PilB is composed of three domains. The central and the C-terminal domains were shown to display methionine sulfoxide reductase (Msr) A and B activities, specific for the S and the R-isomers of the sulfur of MetSO, respectively (4–8). Although MsrA and MsrB belong to two structurally unrelated classes of enzymes (9–13), they share a similar catalytic mechanism consisting of three steps, the third of which permits oxidized Msrs under disulfide state to return back to reduced forms *via* reduction by thioredoxin (Trx) (8, 14, 15). The enzymology of the methionine sulfoxide reductase step and of the Trx-recycling process was recently well characterized (16, 17).

Different roles have been assigned to Msrs. One of the most important roles is to restore the function of proteins oxidized on their methionine residues (18, 19). In contrast, no role has been assigned so far to the N-terminal domain of PilB. Recently, two forms of PilB from *N. gonorrhoeae* have been characterized (3). One is secreted from the cytoplasm to the outer membrane as an entire polypeptide composed of the three domains. The second one is a truncated cytoplasmic form corresponding to amino acids 196–521 and therefore lacks the N-terminal domain. This form is produced from an internal AUG initiation codon corresponding to Met-195 (Ref. 3, see also “Results”). The fact that the extracytoplasmic localization of PilB was shown to be required for survival in the presence of oxidative damage raises the question of the function of the N-terminal domain in the periplasm and of its relationship with the MsrA and MsrB activities.

Comparison of the primary structure of the N-terminal domain with those of the known proteins in the public data bases shows no significant identity with any protein excepted one from *Fusobacterium nucleatum* (see Fig. 1). The only peculiar feature is the presence of a CXXC signature.

In an attempt to identify the role of the N-terminal domain of PilB, a soluble form of the *N. meningitidis* domain of 143 amino acids, which only differs from its *N. gonorrhoeae* counterpart by four amino acids (see Fig. 1), has been overproduced and purified. Its biochemical and enzymatic properties have been determined. The results demonstrate that the N-terminal domain displays a disulfide reductase activity that is able to recycle

* This research was supported by the Centre National de la Recherche Scientifique, the Ministère délégué à la Recherche (ACI BCMS047), the University Henry Poincaré Nancy I, the Association pour la Recherche sur le Cancer (Association pour la Recherche sur le Cancer Grant 5436), and the Institut Fédératif de Recherches 111 Bioingénierie. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

‡ Both authors contributed equally to this work.

§ Present address: Dept. of Biochemistry, College of Medicine, Wuhan University, Wuhan, 430071, China.

¶ To whom correspondence should be addressed: Laboratoire MAEM, UMR CNRS-UHP 7567, Bld des Aiguillettes, BP 239, 54506 Vandoeuvre-les-Nancy, France. Tel.: 33-3-83-68-43-05; Fax: 33-3-83-68-43-07; E-mail: guy.branlant@maem.uhp-nancy.fr.

¹ The abbreviations used are: Msr, methionine sulfoxide reductase; DTNB, 5,5'-dithiobis(2-nitrobenzoic acid); DTT, dithiothreitol; 2PDS, 2,2'-dipyridyl disulfide; Trx, thioredoxin; GSH, glutathione; GSSG, oxidized glutathione.

Msr activities. In particular, the *N. meningitidis* MsrB activity is shown to be recycled with a catalytic efficiency similar to that observed with Trx1 from *Escherichia coli* (8).

MATERIALS AND METHODS

Plasmid Constructions, Site-directed Mutageneses, Productions, and Purifications of the N-terminal Domain and of the Entire Form of PilB from *N. meningitidis*—Plasmid pETPilB was obtained by cloning the internal fragment of the *pilB* open reading frame synthesized by PCR (sequences of oligonucleotides not shown) using *N. meningitidis* Z2491 genomic DNA, kindly provided by Dr. M. K. Taha, between the NdeI and the SacI sites of the pET24c plasmid. To produce the N-terminal domain truncated after position 32, a deletion was done by NdeI digestion of the plasmid in which a second NdeI site has been introduced between the 100th and the 105th bp. Truncation at the C terminus after position 175 was obtained by site-directed mutagenesis of the Gly-176 codon into a stop codon TTA. The resulting plasmid was named pET-NterPilB. To produce a soluble form of PilB, a similar plasmid was used except that instead of introducing a stop codon at codon 176, the Met-195 codon was changed into an Ala. The resulting plasmid was named pET195PilB. Site-directed mutageneses were performed using the QuikChange site-directed mutagenesis kit (Stratagene).

The *E. coli* strain used for N-terminal domain and entire PilB productions was BL21 (DE3) pLysS transformed with the pETNterPilB and pET195PilB plasmids, respectively. The overexpression of the N-terminal domain and of entire PilB was performed by addition of 1 mM isopropyl 1-thio- β -D-galactopyranoside in the culture medium at 0.6 A₆₀₀. After 3 h of induction, cells were harvested by centrifugation, resuspended in minimal volume of buffer A (50 mM Tris-HCl, 2 mM EDTA, pH 8) containing 20 mM dithiothreitol (DTT), and sonicated. For the N-terminal domain and entire PilB, the supernatant was then precipitated at 70% ammonium sulfate saturation. The contaminating proteins were removed by exclusion size chromatography on ACA 54 gel (IBF) in buffer A. Purified fractions were then pooled and applied to a phenyl-Sepharose column equilibrated with buffer A, followed by a linear gradient of ammonium sulfate (0–1.0 M) connected to a fast protein liquid chromatography system (Amersham Biosciences). The two proteins were pure as checked by SDS-PAGE electrophoresis. The protein mass determined by electrospray mass spectrometry corresponded to that expected with no Met at the N terminus. The molecular concentrations were determined spectrophotometrically, using the extinction coefficient at 280 nm of 33,690 M⁻¹·cm⁻¹ for the N-terminal domain and 77,130 M⁻¹·cm⁻¹ for the entire PilB as deduced from the method of Scopes (20).

Trx1 from *E. coli* and MsrA and MsrB domains from *N. meningitidis* were prepared following experimental procedures already published (8, 21). The cysteine contents of the N-terminal domain and of *E. coli* Trx1 were determined using 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB) under non-denaturing conditions in buffer A as described previously (14).

Preparation of the Oxidized Forms of the N-terminal Domain and of *E. coli* Trx1—The oxidation of the N-terminal domain and Trx1 was achieved by incubating a reduced N-terminal domain (300 μ M) or reduced Trx1 (500 μ M) with DTNB (1 mM) in degassed and He₂-purged solution of buffer B (100 mM phosphate, pH 7) at room temperature for 10 min. Both oxidized forms were then isolated by gel filtration on Econo-Pac 10 DG Column (Bio-Rad Laboratories) equilibrated with buffer B. The oxidation states of the N-terminal domain and of Trx1 were checked by titration with DTNB.

Fluorescence Properties of the N-terminal Domain under Oxidized and Reduced Forms—The fluorescence spectra of the reduced and oxidized N-terminal domains (10 μ M) in buffer A were recorded on a spectrofluorometer (flx SAFAS) thermostated at 25 °C. Excitation spectra were recorded at maximum emission (347 nm), and emission spectra were recorded at maximum excitation (295 nm).

Determination of the Rate of Reduction of Oxidized N-terminal Domain by DTT—The rate was determined by recording the change in fluorescence intensity at 347 nm at an excitation wavelength of 295 nm. The change in fluorescence intensity was recorded for 3 min after the addition of DTT at various concentrations (5–40 μ M) to oxidized N-terminal domain (2 μ M) in degassed buffer B. The apparent second-order rate constant was calculated from the measured pseudo-first-order rate constants by dividing k_{obs} value by the concentration of DTT.

Determination of the Redox Potential of the N-terminal Domain, the GSH/GSSG System—The change in fluorescence intensity (excitation wavelength 295 nm) was measured at the wavelength of maximum emission (347 nm). Experiments were carried out in degassed and He₂-purged buffer B, with 2 mM EDTA. Oxidized N-terminal domain

(1 μ M) was incubated at 30 °C in the presence of GSSG (0.1 mM) and varying concentrations of reduced glutathione (GSH) (0–180 mM) for 20 h before recording the fluorescence emission spectra. The equilibrium concentrations of GSH and GSSG were calculated according to Equations 1–3,

$$[\text{GSH}] = [\text{GSH}]_0 - 2R [\text{N-terminal domain}]_0 \quad (\text{Eq. 1})$$

$$[\text{GSSG}] = [\text{GSSG}]_0 + R [\text{N-terminal domain}]_0 \quad (\text{Eq. 2})$$

$$R = (F - F_{\text{ox}})/(F_{\text{red}} - F_{\text{ox}}) \quad (\text{Eq. 3})$$

where $[\text{GSH}]_0$ and $[\text{GSSG}]_0$ represent the initial concentrations of GSH and GSSG, respectively, R is the relative amount of reduced protein at equilibrium, $[\text{N-terminal domain}]_0$ is the initial concentration of the N-terminal domain under the oxidized state, F is the fluorescence intensity, and F_{ox} and F_{red} are the fluorescence intensities of the completely oxidized and reduced protein. The equilibrium constant K_{eq} was estimated according to Equation 4,

$$R = ([\text{GSH}]^2/[\text{GSSG}])/([K_{\text{eq}} + [\text{GSH}]/[\text{GSSG}]) \quad (\text{Eq. 4})$$

from a non-linear regression analysis of the data.

From the equilibrium constant and using the glutathione standard potential ($E'_{\text{GSH/GSSG}} = -0.240$ V) (22), the standard redox potential (E'_{o}) was calculated with the Nernst equation (Equation 5),

$$E'_{\text{o}} = E'_{\text{GSH/GSSG}} - (RT/nF) \ln K_{\text{eq}} \quad (\text{Eq. 5})$$

in which F represents Faraday's constant (23,040.612 cal·mol⁻¹·V⁻¹), n is the number of electrons transferred (here $n = 2$), and RT is the product of the gas constant (1.987 cal·K⁻¹·mol⁻¹) and the absolute temperature.

Redox Equilibrium between the N-terminal Domain and Trx1—Equilibrium reactions (250 μ l) typically contained 15 μ M each redox-active protein in degassed and He₂-purged solution of buffer B with 2 mM EDTA. The reduced form of each protein (1 mM) was prepared immediately before use by the incubation of protein for 1 h at room temperature in the presence of 50 mM DTT, followed by desalting on an Econo-Pac 10 DG column. Redox reactions between both proteins were initiated by mixing one protein in the reduced state and the other in the oxidized state. After 15 h of equilibration at 25 °C, each sample was quenched by the addition of 0.1% trifluoroacetic acid final. The oxidized and reduced forms of the proteins present in the samples were separated by reverse-phase high pressure liquid chromatography on a C8 column (4.6 mm \times 100 mm) using a gradient from 30 to 80% (v/v) acetonitrile, 0.1% (v/v) trifluoroacetic acid in 25 min at a flow rate of 0.8 ml/min at room temperature. Column effluent was monitored at 215 and 280 nm. The amounts of oxidized and reduced forms of each protein in the quenched equilibrium mixture were obtained from peak areas after integration of the chromatograms. Essentially identical results were obtained from analysis of chromatograms recorded at 215 and 280 nm, and the former was used in a subsequent analysis. The mechanism of thiol-disulfide exchange is described as a two-step reaction proceeding through a mixed disulfide intermediate (Equation 6). It is assumed that the mixed disulfide intermediate is not populated under standard conditions.

The apparent concentration equilibrium constant, K_{eq} , was calculated according to Equation 7,



$$K_{\text{eq}} = ([\text{Trx}_{\text{S}}^{\text{S}}][\text{Nter}_{\text{SH}}^{\text{SH}}])/([\text{Trx}_{\text{SH}}^{\text{SH}}][\text{Nter}_{\text{S}}^{\text{S}}]) \quad (\text{Eq. 7})$$

$$E'_{\text{o}} = E'_{\text{Trx1}} - (RT/nF) \ln K_{\text{eq}} \quad (\text{Eq. 8})$$

and the redox potential of the N-terminal domain (E'_{o}) was then calculated using the Nernst equation (Equation 8), using the Trx1 standard potential E'_{Trx1} of -0.270 V at pH 7 and 25 °C (23), in which n is the number of electrons transferred in the reaction (here $n = 2$), F is Faraday's constant (23,040.612 cal·mol⁻¹·V⁻¹), and R is the gas constant (1.987 cal·K⁻¹·mol⁻¹).

Stoichiometry of Met Formation—In the case of MsrA/B domains alone, the reaction mixture containing DL-Met-(R,S)-SO (100 mM) and MsrA/B (at two concentrations of 50 and 100 μ M), in the absence or presence of stoichiometric concentration of N-terminal domain (*i.e.* 50 and 100 μ M), was incubated at 25 °C for 60 min in buffer A. In the case of entire PilB, the reaction mixture containing L-Met-(R)-SO or L-Met-(S)-SO (200 μ M) and PilB (at two concentrations of 50 and 100 μ M), was incubated at 25 °C for 120 min in buffer A. Then, 250 μ l of each reaction



FIG. 1. Sequence alignment of the N-terminal domain of PilB from *N. meningitidis*, *N. gonorrhoeae*, and *F. nucleatum*. Sequences are from the PEDANT data base. The numbering of amino acid residues is based on the PilB from *N. meningitidis* sequence. The cysteine residues from catalytic motif are shown in white on black boxes. Conserved residues in the three PilB sequences are shown in black on gray boxes. The diagram down the alignment depicts the prediction of *N. meningitidis* PilB N-terminal domain secondary structure elements. The prediction was obtained with Network Protein Sequence Analysis at the IBCP web interface. The prediction for *N. meningitidis* PilB signal peptide region was obtained with SignalP V1.1 software and is indicated by a gray box. The hatched gray box indicates different possible ends of signal peptide. The vertical black arrows indicate the site of cleavage used for production of the soluble form of the N-terminal domain. *N. menin.*, *N. meningitidis*; *N. gono.*, *N. gonorrhoeae*; *F. nucl.*, *F. nucleatum*.

mixture was injected onto a Sephasil C18 column for Met quantification as already described (14).

Determination of the Kinetic Parameters—Msr activities were determined with Ac-L-Met-(R,S)-SO-NHMe, which is a better substrate than Met-(R,S)-SO (8), at a saturating concentration of 200 mM. The reaction mixture also contained MsrA (2 μ M) or MsrB (1 μ M), and N-terminal domains at various concentrations (10–800 μ M). Initial rate measurements were carried out at 25 °C by following the appearance of Ac-L-Met-NHMe. To do so, aliquots of 250 μ l of the reaction mixture were removed at different times of incubation up to 2 min for MsrB and 3 h for MsrA, and the reaction was stopped by the addition of trifluoroacetic acid to a final concentration of 1% (v/v). In each aliquot, Ac-L-Met-NHMe quantification was carried out by reverse phase chromatography as described previously (17). The concentration of Ac-L-Met-NHMe formed was plotted against the time, and the data were fit to a linear model to attain initial rates. The initial rate data were fit to the Michaelis-Menten relationship using least squares analysis to determine k_{cat} and K_m .

Determination of pK_a of Cysteine Residues with 2,2'-Dipyridyl Disulfide (2PDS)—Fast kinetic measurements were carried out on a SX18MV-R stopped-flow apparatus (Applied PhotoPhysics). Kinetic reactions were performed at 25 °C, under pseudo-first-order conditions at a constant ionic strength of 0.15 M over the pH range 6–10.5 (Tris 120 mM, imidazole 30 mM, acetic acid 30 mM). One syringe was filled with wild type or Cys to Ser mutated N-terminal domain, and the other was filled with 2PDS. The reaction was monitored at 343 nm with 6.2 μ M N-terminal domain and 310 μ M 2PDS. The release of pyridine-2-thione was quantified using extinction coefficient of 8080 $M^{-1}cm^{-1}$ at 343 nm.

The pseudo-first-order rate constants k_{obs} were determined at each pH by fitting the absorbance A at 343 nm versus time t to Equation 9,

$$A = a_1(1 - e^{-k_{obs}t}) + c \quad (\text{Eq. 9})$$

where a_1 is the burst magnitude, and c represents the value of the ordinate intercept.

The second-order kinetic constants k_2 were calculated by dividing the k_{obs} value by the concentration of 2PDS and then fitting to Equation 10,

$$k_2 = k_{min} + k' \left(\frac{1}{10^{(pK_{app} - pH)} + 1} \right) \quad (\text{Eq. 10})$$

in which k_{min} represents the second-order kinetic constant at pH 6, and k' represents the second-rate constant for the thiolate form. In the case of C67S protein, $k_{min} = 0$.

RESULTS

Production and Purification of the N-terminal Domain and of the Entire Form of PilB—The fact that PilB was shown to be secreted from the cytoplasm to the outer membrane of *N. gon-*

orrhoeae supported the presence of at least an N-terminal signal peptide. Therefore, the strategy to produce a soluble form of the N-terminal domain and of entire PilB should take in account this putative peptide. As indicated in Fig. 1, the N-terminal 31 amino acids are predicted to constitute a signal peptide using the SignalP software. Using the Network Protein Sequence Analysis web interface, an α helix between amino acids 4 and 22 is also predicted. Production of the N-terminal domain also required us to define where the truncation should be introduced between the C terminus of the N-terminal domain and the N terminus of the MsrA domain. For that purpose, the position of the truncation was based on the fact that 1) the MsrA domain was previously produced in a soluble form from a DNA construct in which a stop codon was introduced at a position corresponding to amino acid 195 (8), and 2) no secondary structural element is predicted after amino acid 171.

Production of a soluble form of PilB in good yield required not only to remove the peptide signal but also to change the AUG codon corresponding to Met-195 into an Ala one. Indeed, when overexpressed in *E. coli* two forms of PilB were shown to be produced in the cytoplasm of *E. coli* i.e. one form corresponding to amino acids 33–521 and the other, produced in a higher yield, corresponding to amino acids 196–521 (data and SDS-PAGE, not shown). This result strongly suggested that the AUG codon of Met-195 can be used as an internal initiation codon as already postulated for *N. gonorrhoeae* PilB (3). This is indeed the case. When an Ala codon was substituted for the Met-195 codon, only a form corresponding to amino acids 33–521 was produced and in a higher yield.

For N-terminal domain production, two plasmidic constructs under T7 promoter were therefore built that took in account all the information described above and then were tested for protein production in *E. coli*. Only one construct corresponding to truncations at the N terminus after position 32 and at the C terminus after position 175 gave a high overexpression of the N-terminal domain. The reason why truncation done at position 194 gave a small production of protein remains unknown. Therefore, only the plasmidic construct that coded the N-terminal domain from amino acids 33 to 175 was used to produce the domain. Purification was achieved by sequential ammonium sulfate precipitation, chromatography on ACA54, Q-Sepharose and phenyl-Sepharose. At the end of the purifica-




FIG. 2. Fluorescence emission and excitation spectra of the N-terminal domain under reduced and disulfide oxidized state. Excitation spectra of the reduced (●) and oxidized (○) N-terminal domains (emission followed at 347 nm) and the emission spectra of reduced (▲) and oxidized (△) N-terminal domains (excitation at 295 nm) were recorded with protein (10 μ M) at 25 °C in buffer A.

tion process, 150 mg of pure protein were obtained from 1 liter of culture. The elution profile on gel filtration under native conditions was in agreement with a monomeric state (profile not shown). As expected from the primary structure (Fig. 1), DTNB titration revealed two Cys under denaturing and native conditions. The process used for purification of the entire PilB was similar, and \sim 100 mg of entire PilB were obtained/liter of culture.

Characterization of the Redox Properties of the N-terminal Domain—The two cysteines in the N-terminal domain are separated by only two amino acids (Fig. 1). Therefore, the fact that they are included in a signature found in disulfide oxidoreductases suggested that these cysteines were redox sensitive. The addition of GSSG or DTNB led to the loss of the two cysteines as proved by DTNB titration. Addition of DTT, then followed by gel filtration restored the thiol titration. These results indicated that the Cys residues, Cys-67 and Cys-70 are oxidized by GSSG or DTNB and form an intramolecular disulfide bond that is sensitive to reduction by DTT.

It is known that proteins of the thiol-disulfide oxidoreductase family such as Trx show an increase in fluorescence emission intensity upon reduction of their active site disulfide bond. In the case of the Trx1 from *E. coli*, a 2.5-fold increase is observed, which was attributed essentially to the tryptophan located directly before the CXXC signature (24). In the N-terminal domain (amino acids 33–175), there exist five tryptophans of which one is situated at position 66 also before the CXXC motif (Fig. 1). Therefore, it was expected that the N-terminal domain behaved like Trx1. As shown in Fig. 2, when the oxidized N-terminal domain was excited at 295 nm for the selective excitation of tryptophan, an addition of 1 mM DTT led to a 2.0-fold increase in the fluorescence emission intensity at 347 nm. This result suggested that, as observed for *E. coli* Trx1, the fluorescence emission of Trp-66 of the N-terminal domain is quenched by formation of the disulfide bond.

By taking advantage of the fluorescence intensity change between oxidized and reduced N-terminal domains, the redox equilibrium constant K_{eq} of the domain was determined based on the variation of the fluorescence message at various GSH/GSSG ratios. Thus, the oxidized N-terminal domain was incubated in the presence of 100 μ M GSSG and increasing concentrations of GSH from 0 to 180 mM. The fraction of reduced N-terminal domain at equilibrium was measured by the intrinsic




FIG. 3. Redox equilibrium of the N-terminal domain with glutathione. The relative amount of reduced N-terminal domain at equilibrium (R) was measured using the N-terminal domain fluorescence message at 347 nm (excitation at 295 nm) after incubation of the oxidized form (1 μ M) for 15 h at 30 °C in buffer B containing 0.1 mM GSSG and various concentrations of GSH (0–180 mM). The data processing and the determination of the equilibrium constant are described under “Materials and Methods.” Using non-linear regression analyses, a value of $K_{eq} = 0.54 \pm 0.09$ M was determined for the N-terminal domain/glutathione system, corresponding to a redox potential of -232 ± 4 mV.

sic N-terminal domain fluorescence assuming no significant equilibrium concentration of N-terminal domain/glutathione mixed disulfide. After fitting the data by nonlinear regression according to Equation 4 (see “Materials and Methods”) (Fig. 3), an equilibrium constant for the N-terminal domain/glutathione system of 0.54 ± 0.09 M was determined. A standard redox potential of -0.232 ± 0.004 V for the active site cysteines of the N-terminal domain at 30 °C and pH 7.0 (E'_{ON}) was calculated from the Nernst equation (Equation 5) using a value of -0.240 V (22) for the glutathione standard redox potential (E'_{OG}).

To confirm the result, we used another method based on the analysis of the direct protein-protein redox equilibrium between the N-terminal domain and the *E. coli* Trx1, which has a well established redox potential E'_o of -0.270 V (23). To ensure that the redox equilibrium was indeed attained after incubation for 15 h, we checked with each protein pair that identical equilibrium constants were obtained irrespective of the redox state of the initial mixture. In both cases, the redox potential of the N-terminal domain (E'_{ON}) was determined to be -0.227 ± 0.005 V at pH 7. This is in good agreement with the value determined using the GSH/GSSG system as a reference.

Determination of the pK_{app} Values of the Cys Residues of the Redox Center of the N-terminal Domain—The pK_a values of the Cys were determined using 2PDS over a pH range of 6–10.5 under conditions where the wild-type, C67S, and C70S N-terminal domains are stable. The reaction of 2PDS with the N-terminal wild-type domain followed pseudo-first-order kinetics with formation of 2 mol of pyridine-2-thione/mol of the domain, as determined from the absorbance change at 343 nm (data not shown). The pH- k_2 curve between pH 6 and 10.5 fitted to a monosigmoidal profile with a pK_{app} value of 9.3 ± 0.2 and a k' value of 1.6×10^4 $M^{-1}\cdot s^{-1}$ (Fig. 4A). The fact that at pH 6 a k_2 value of 4.7×10^4 $M^{-1}\cdot s^{-1}$ was still observed indicated that at least one of the two cysteines has a pK_a with a value below 6. To assign the pK_a values of both Cys residues, their pK_a s were then determined in each of the two Cys mutated N-terminal domains.

As expected, the kinetics of 2PDS with the C67S and C70S




FIG. 4. pH dependence of the second-order rate constant k_2 for the reaction of the thiol group with 2PDS. Wild-type (A), C67S (B), and C70S (C) N-terminal domains. Reaction kinetics were performed at 25 °C over a pH-range of 6–10.5. The concentrations of the N-terminal domain and 2-PDS were 6.2 and 310 μ M, respectively. Values of k_{obs} were determined using non-linear regression analysis. Second-order rate constants k_2 were calculated dividing k_{obs} by the concentration of 2PDS and were fit to Equation 10 (solid line theoretical curve), which gave a $\text{p}K_{\text{app}}$ of 9.3 ± 0.2 with a k_{min} and a k' value of $(4.7 \pm 0.1) \cdot 10^4$ and $(1.6 \pm 0.1) \cdot 10^4$ M⁻¹.s⁻¹, respectively, for the wild-type, a $\text{p}K_{\text{app}}$ of 9.5 ± 0.2 with a k_{min} and a k' value of 0 and $(1.8 \pm 0.1) \cdot 10^4$ M⁻¹.s⁻¹, respectively, for the C67S, and a $\text{p}K_{\text{app}}$ of 9.1 ± 0.1 with a k_{min} and a k' value of $(1.4 \pm 0.1) \cdot 10^4$ and $(1.3 \pm 0.1) \cdot 10^4$ M⁻¹.s⁻¹, respectively, for the C70S N-terminal domains.

proteins followed pseudo-first-order but with formation of only 1 mol of pyridine-2-thione/mol of N-terminal domain (data not shown). The pH- k_2 curve with the C67S protein passed through the origin and fitted to a monosigmoidal profile with a $\text{p}K_{\text{app}}$ value of 9.5 ± 0.2 and a k' value of 1.8×10^4 M⁻¹.s⁻¹ (Fig. 4B). This result strongly suggested that the $\text{p}K_{\text{app}}$ of 9.3 observed in the wild-type N-terminal domain corresponded to that of Cys-

70. In the C70S protein, the Cys-67 reacted rapidly even at pH 6, with a k_2 constant of 1.4×10^4 M⁻¹.s⁻¹, indicating that the $\text{p}K_{\text{app}}$ value of Cys-67 was below 6 as expected (Fig. 4C). The fact that the k_2 value is 3-fold lower than that of the wild type remains to be explained. Another $\text{p}K_{\text{app}}$ value of 9.1 ± 0.1 with a k' value of 1.3×10^4 M⁻¹.s⁻¹ was also determined from the pH- k_2 curve, suggesting the presence of an amino acid not yet

TABLE I
Stoichiometry of Met formation/mol of Msrs and kinetic parameters of Msrs in the presence of the N-terminal domain

Stoichiometry of Met formation in the absence and presence of the N-terminal domain. The values indicated represent the average of three independent measurements of at least two N-terminal domain concentrations (S.D. range 10%). The quantity of Met formed/mol of Msr was determined by reverse phase chromatography, after 60 min of reaction between DL-Met-(*R,S*)-SO (100 mM) and MsrA/B (at two concentrations of 50 and 100 μM), in the absence or presence of stoichiometric concentrations of N-terminal domain (*i.e.* 50 and 100 μM , respectively). Kinetic parameters of Msrs in the presence of the N-terminal domain. Kinetic parameters were determined at a saturating concentration of Ac-L-Met-(*R,S*)-SO-NHMe (200 mM). Reactions were carried out in buffer A at 25 °C in the presence of MsrB (1 μM) or MsrA (2 μM) and various concentrations of N-terminal domain (10–800 μM). Data were obtained by fitting the experimental data to the Michaelis-Menten equation. ND, not determinable.

	Stoichiometry (mol Met/mol Msr)		Kinetic parameters of Msrs					
	Without N-terminal domain	With 1 eq of N-terminal domain	With the N-terminal domain			With Trx 1		
			k_{cat} s^{-1}	K_{M} μM	$k_{\text{cat}}/K_{\text{M}}$ $\mu\text{M}^{-1}\cdot\text{s}^{-1}$	k_{cat} s^{-1}	K_{M} μM	$k_{\text{cat}}/K_{\text{M}}$ $\mu\text{M}^{-1}\cdot\text{s}^{-1}$
MsrA	1	2	ND	ND	ND	3.4 ± 0.2^a	75 ± 15^a	0.045
MsrB	1	2	4.1 ± 0.2	280 ± 30	0.015	1.1 ± 0.2^b	58 ± 12^b	0.019

^a From Ref. 8.

^b From Ref. 15.

identified near Cys-67 whose deprotonation increases the reactivity of the thiolate of Cys-67 and which is likely involved in decreasing the $\text{p}K_{\text{a}}$ of Cys-67.

The N-terminal Domain Is Capable of Recycling Methionine Sulfoxide Reductase Activities—With the objective to get more information on the role of the N-terminal domain included in PilB, we first tested the ability of the N-terminal domain alone to reduce the oxidized MsrA and MsrB domains, produced independently as recently described (8). For that purpose, the stoichiometry of Met formation/mole of MsrA or MsrB was determined in the presence and absence of the N-terminal domain. Two mol of Met were formed/mol of MsrA or MsrB when an equimolar concentration of the N-terminal domain was added, whereas in its absence, only 1 mol of Met/mol of MsrA or MsrB was formed (Table I). Two mol of Met were also formed/mol of entire PilB in the presence of either L-Met-(*S*)-SO, which is selectively reduced by MsrA, or L-Met-(*R*)-SO which selectively reduced by MsrB (8). Altogether, these results clearly showed that the N-terminal domain, either as an isolated form or included in PilB, is able to reduce the disulfide bond in MsrA and in MsrB but did not give any information on the efficiency of the recycling process. Such data required the determination of the kinetic parameters, *i.e.* the k_{cat} value and the K_{m} for the N-terminal domain. This was done in the presence of saturating concentrations of Ac-L-Met-SO-NHMe (Table I). The rate was determined by following the formation of Ac-L-Met-NHMe. In the case of MsrB, a k_{cat} value of 4.1 s^{-1} and a K_{m} of 280 μM for the N-terminal domain were determined, which are in the range as those determined for Trx1 from *E. coli* (15) and from *N. meningitidis*.² In contrast, within the concentration range of 50–800 μM of the N-terminal domain tested, no saturating effect was observed for MsrA. At 800 μM , a k_{obs} value of 4.10^{-2} s^{-1} was determined that is 80-fold lower than the k_{cat} value determined with Trx1 from *E. coli* (8) and from *N. meningitidis*.²

DISCUSSION

PilB of *N. meningitidis* is composed of three domains. Although the central and the C-terminal domains display MsrA and MsrB activities, respectively, the function of the N-terminal domain is unknown. In this study, we have characterized the biochemical and enzymatic properties of a soluble form of the N-terminal domain.

The only two Cys of the N-terminal domain are located in a WCPLC motif that has been shown to form a disulfide redox-active site. The percentage of α helix and of β sheets predicted from CD spectra of the N-terminal domain and of the Trx1 from

E. coli are similar (data and CD spectra not shown). Altogether, these results suggested that the redox-active site is located in a Trx-like fold. Trx-like proteins are widely distributed and implicated in the control of the redox environment of subcellular compartments. The disulfide oxidoreductase activity of CXXC-containing proteins depends on various factors, among which the redox potential and the $\text{p}K_{\text{a}}$ of the thiol group of the N-terminal Cys in the motif. The redox potential ($E'^{\circ} = -230 \text{ mV}$) of the N-terminal domain is similar to those of different thiol reductants such as *E. coli* Trxs (-270 to -267 mV) (23), *E. coli* Grxs (-233 to -198 mV), and GSH (-240 mV) (22). This suggested that the N-terminal domain probably acts as a reductant. However, the $\text{p}K_{\text{a}}$ value of the N-terminal Cys in the WCPLC motif, which is below 6 corresponds to a value usually observed for an oxidant and not for a disulfide reductant. For instance, the $\text{p}K_{\text{a}}$ of the corresponding catalytic Cys in *E. coli* Trx1 is 7.5, whereas it is of 3.5 in the periplasmic oxidase DsbA (25, 26). The fact that the apparent second-order rate constant of reduction of the oxidized N-terminal domain by DTT ($5.10^3 \text{ M}^{-1}\cdot\text{s}^{-1}$) is in the same range as that measured on oxidized Trx1 from *E. coli* ($10^3 \text{ M}^{-1}\cdot\text{s}^{-1}$) and 3 orders of magnitude slower compared with that measured on oxidized DsbA ($10^6 \text{ M}^{-1}\cdot\text{s}^{-1}$) (27, 28) is another piece of data that supports a function of the N-terminal domain as a reductant.

This raises the question of the role of the N-terminal domain of PilB in the periplasm. As indicated in the Introduction, the fact that it is fused with MsrA and MsrB domains suggested a function of the N-terminal domain associated with the Msr activities. This is indeed the case. The N-terminal domain is able to recycle the oxidized form of MsrB from *N. meningitidis* with a catalytic efficiency, $k_{\text{cat}}/K_{\text{m}}$ of $0.015 \mu\text{M}^{-1}\cdot\text{s}^{-1}$, similar to that observed with Trx1 from *E. coli* and *N. meningitidis*. Therefore, the N-terminal domain can act as a disulfide reductase and is probably folded as a Trx. However, the fact that no saturating concentration effect for the *N. meningitidis* MsrA is observed supports subtle three-dimensional structural differences between the N-terminal domain and Trx1 from *E. coli* and *N. meningitidis*. This is confirmed by the fact that the *E. coli* Trx reductase is not able to recycle the oxidized N-terminal domain (data not shown) in contrast to that observed with oxidized Trx1. In that context, the knowledge of the x-ray structure of the N-terminal domain, the resolution of which is under progress, will be very informative.

The kinetic parameters obtained in the present study have been determined with separated and soluble domains. *In vivo*, PilB is localized in the periplasm on the outer membrane. Therefore, the question arises of whether the N-terminal domain in PilB is operative in recycling the MsrA and MsrB activities in the *in vivo* context. No data are presently avail-

² A. Olry, S. Boschi-Muller, and G. Branlant, unpublished results.

able. However, what is known from the present study done *in vitro* is that the disulfide bonds formed within PilB MsrA and MsrB domains are accessible and reduced by the N-terminal domain included in PilB. But this result does not give any indication of whether the recycling process by the N-terminal domain is intra or intermolecular. This question is of importance in the context of the localization of PilB *in vivo*. What can be concluded, however, is that whatever the mechanism, *i.e.* intra- or intermolecular, 1 mol of PilB is sufficient to reduce 1 mol of a mixture of a protein-Met-(R,S)-SO and to recycle the fractions of MsrA and MsrB domains in PilB, which have been oxidized under the disulfide state. Another point that has to be addressed is the nature of the proteins that are repaired in the periplasm by PilB and the possible relationship with the pathogenic character of *Neisseria*. Finally, the fact that the N-terminal domain displays a disulfide reductase activity suggests that in the periplasm of *N. meningitidis* and *N. gonorrhoeae* a disulfide oxidoreductase is present to recycle the N-terminal domain from the oxidized to the reduced form. Its nature remains to be identified.

The PilB organization is a specific human pathogen bacteria from the *Neisseria* genus. A similar organization only exists in *F. nucleatum*, a Gram-negative anaerobe, which is a human opportunistic pathogen (29). Therefore, the N-terminal domain which has 1) no homologue in other bacteria, 2) an outer membrane localization, and 3) a fold likely similar to Trx1 but with subtle structural differences, could be a good candidate as a drug target against pathogenic *Neisseria*.

Acknowledgments—We thank Dr. M. K. Taha for the gift of the *N. meningitidis* DNA and Dr. A. Van Dorsselear and G. Chevreux for mass determination. We also thank Dr. S. Azza and C. Gauthier for their efficient technical help, and A. Olry for helpful discussion.

REFERENCES

- Dehio, C., Gray-Owen, S. D., and Meyer, T. F. (2000) *Subcell. Biochem.* **33**, 61–96
- Boyle, E. C., and Finlay, B. B. (2003) *Curr. Opin. Cell Biol.* **15**, 633–639
- Skaar, E. P., Tobiason, D. M., Quick, J., Judd, R. C., Weissbach, H., Etienne, F., Brot, N., and Seifert, H. S. (2002) *Proc. Natl. Acad. Sci. U. S. A.* **99**, 10108–10113
- Moskovitz, J., Poston, J. M., Berlett, B. S., Nosworthy, N. J., Szczepanowski, R., and Stadtman, E. R. (2000) *J. Biol. Chem.* **275**, 14167–14172
- Sharov, V. S., Ferrington, D. A., Squier, T. C., and Schoneich, C. (1999) *FEBS Lett.* **455**, 247–250
- Moskovitz, J., Singh, V. K., Requena, J., Wilkinson, B. J., Jayaswal, R. K., and Stadtman, E. R. (2002) *Biochem. Biophys. Res. Commun.* **290**, 62–65
- Grimaud, R., Ezraty, B., Mitchell, J. K., Lafitte, D., Briand, C., Derrick, P. J., and Barras, F. (2001) *J. Biol. Chem.* **276**, 48915–48920
- Olry, A., Boschi-Muller, S., Marraud, M., Sanglier-Cianferani, S., Van Dorsselear, A., and Branlant, G. (2002) *J. Biol. Chem.* **277**, 12016–12022
- Tete-Favier, F., Cobessi, D., Boschi-Muller, S., Azza, S., Branlant, G., and Aubry, A. (2000) *Structure Fold. Des.* **8**, 1167–1178
- Lowther, W. T., Brot, N., Weissbach, H., and Matthews, B. W. (2000) *Biochemistry* **39**, 13307–13312
- Lowther, W. T., Weissbach, H., Etienne, F., Brot, N., and Matthews, B. W. (2002) *Nat. Struct. Biol.* **9**, 348–352
- Kauffmann, B., Favier, F., Olry, A., Boschi-Muller, S., Carpentier, P., Branlant, G., and Aubry, A. (2002) *Acta Crystallogr. Sect. D Biol. Crystallogr.* **58**, 1467–1469
- Taylor, A. B., Benglis, D. M., Jr., Dhandayuthapani, S., and Hart, P. J. (2003) *J. Bacteriol.* **185**, 4119–4126
- Boschi-Muller, S., Azza, S., Sanglier-Cianferani, S., Talfournier, F., Van Dorsselear, A., and Branlant, G. (2000) *J. Biol. Chem.* **275**, 35908–35913
- Boschi-Muller, S., Olry, A., Antoine, M., and Branlant, G. (2005) *Biochim. Biophys. Acta* **1703**, 231–238
- Antoine, M., Boschi-Muller, S., and Branlant, G. (2003) *J. Biol. Chem.* **278**, 45352–45357
- Olry, A., Boschi-Muller, S., and Branlant, G. (2004) *Biochemistry* **43**, 11616–11622
- Brot, N., and Weissbach, H. (2000) *Biopolymers* **55**, 288–296
- Stadtman, E. R., Moskovitz, J., and Levine, R. L. (2003) *Antioxid. Redox Signal.* **5**, 577–582
- Scopes, R. K. (1974) *Anal. Biochem.* **59**, 277–282
- Mossner, E., Huber-Wunderlich, M., and Glockshuber, R. (1998) *Protein Sci.* **7**, 1233–1244
- Aslund, F., Berndt, K. D., and Holmgren, A. (1997) *J. Biol. Chem.* **272**, 30780–30786
- Krause, G., Lundstrom, J., Barea, J. L., Pueyo de la Cuesta, C., and Holmgren, A. (1991) *J. Biol. Chem.* **266**, 9494–9500
- Slaby, I., Cerna, V., Jeng, M. F., Dyson, H. J., and Holmgren, A. (1996) *J. Biol. Chem.* **271**, 3091–3096
- Grauschopf, U., Winther, J. R., Korber, P., Zander, T., Dallinger, P., and Bardwell, J. C. (1995) *Cell* **83**, 947–955
- Chivers, P. T., Prehoda, K. E., Volkman, B. F., Kim, B. M., Markley, J. L., and Raines, R. T. (1997) *Biochemistry* **36**, 14985–14991
- Holmgren, A. (1979) *J. Biol. Chem.* **254**, 9627–9632
- Fabianek, R. A., Huber-Wunderlich, M., Glockshuber, R., Kunzler, P., Hennecke, H., and Thony-Meyer, L. (1997) *J. Biol. Chem.* **272**, 4467–4473
- Kapatral, V., Anderson, I., Ivanova, N., Reznik, G., Los, T., Lykidis, A., Bhattacharyya, A., Bartman, A., Gardner, W., Grechkin, G., Zhu, L., Vasieva, O., Chu, L., Kogan, Y., Chaga, O., Goltsman, E., Bernal, A., Larsen, N., D'Souza, M., Walunas, T., Pusch, G., Haselkorn, R., Fonstein, M., Kyrpides, N., and Overbeek, R. (2002) *J. Bacteriol.* **184**, 2005–2018

The N-terminal Domain of PILB from *Neisseria meningitidis* Is a Disulfide Reductase That Can Recycle Methionine Sulfoxide Reductases

Junzhu Wu, Fabrice Neiers, Sandrine Boschi-Muller and Guy Branlant

J. Biol. Chem. 2005, 280:12344-12350.

doi: 10.1074/jbc.M500385200 originally published online January 24, 2005

Access the most updated version of this article at doi: [10.1074/jbc.M500385200](https://doi.org/10.1074/jbc.M500385200)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 29 references, 13 of which can be accessed free at <http://www.jbc.org/content/280/13/12344.full.html#ref-list-1>