



HAL
open science

Quantification of human adenovirus and norovirus in river water in the north-east of France

Maryse Iris Sedji, Mihayl Varbanov, Marie Meo, Marius Colin, Laurence Mathieu, Isabelle Bertrand

► **To cite this version:**

Maryse Iris Sedji, Mihayl Varbanov, Marie Meo, Marius Colin, Laurence Mathieu, et al.. Quantification of human adenovirus and norovirus in river water in the north-east of France. *Environmental Science and Pollution Research*, 2018, 25 (30), pp.30497-30507. 10.1007/s11356-018-3045-4 . hal-01867900

HAL Id: hal-01867900

<https://hal.univ-lorraine.fr/hal-01867900>

Submitted on 29 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Quantification of human adenovirus and norovirus in river water in the north-east of**
2 **France**

3

4 Maryse Iris Sedji¹, Mihayl Varbanov², Marie Meo¹, Leslie Ogorzaly¹, Marius Colin²,
5 Laurence Mathieu^{1,3†}, Isabelle Bertrand^{1†*}

6

7 ¹Université de Lorraine, CNRS, LCPME, Nancy, F -54000, France

8 ²Université de Lorraine, CNRS, L2CM, Nancy, F-54000, France

9 ³ EPHE, PSL Research University, LCPME, UMR 7564, Vandoeuvre-lès-Nancy, F-54500,
10 France

11

12 †These authors contributed equally to this work

13

14 * **Corresponding author:** Isabelle Bertrand, Laboratoire de Chimie Physique et Microbiologie
15 pour les Matériaux et l'Environnement (LCPME), UMR 7564 CNRS Université de Lorraine,
16 Faculté de Pharmacie, 5 rue Albert Lebrun, BP 80403, F-54000 Nancy, France.

17 Tel: +33-3-72-74-73-52; Fax: +33-3-83-27-54-44; E-mail: isabelle.bertrand@univ-lorraine.fr

18

19 **Abstract**

20 Human adenoviruses (HAdVs) are a major cause of infection and have been proposed as viral
21 indicators of water quality. Human noroviruses (NoV) are the main cause of viral acute
22 gastroenteritis. Quantitative data on the environmental prevalence of both viruses are needed.
23 The genomes of HAdVs, enteric adenovirus type 41 (HAdV41) and noroviruses of
24 genogroups I and II (NoV GGI and GGII) were quantified over a six-month period in a river
25 located in north-eastern France. The samples were collected downstream from the discharge
26 of a wastewater treatment plant. The viruses were concentrated using a glass wool method
27 and the viral genomes were quantified using digital droplet PCR (ddPCR). All river water
28 samples (15/15) were positive for the genomes of HAdVs, HAdV41, NoV GGI and NoV
29 GGII. Concentrations of HAdVs, HAdV41 and NoV GII genomes were similar and HAdV41
30 represented ~80% of HAdVs. Infectious HAdVs were quantified in these samples using an
31 integrated cell culture - quantitative PCR method (ICC-qPCR); they were detected in 93%
32 (14/15) and quantified in 53% (8/15) of the samples. Thus, infectious HAdVs represented
33 0.3% to 12.2% of total HAdV particles detected by ddPCR. Infectious HAdV41 particles
34 were found in 73% (11/15) of the samples. This common presence of pathogenic enteric
35 viruses underlines the impact of wastewater discharge on quality of surface waters and may
36 constitute a threat for human health. The relative abundance of genome of HAdV41
37 underlines the need for studies focusing on the specific detection of its infectious forms along
38 water cycle.

39

40

41 **Keywords:** Adenovirus type 41; Norovirus; River water; Genome; Infectivity; ICC-qPCR;
42 Quantification

43

44 **Introduction**

45 HAdVs are a major cause of infections which can be fatal for immunocompromised
46 individuals. They are associated with a wide range of clinical illnesses including upper and
47 lower respiratory diseases, conjunctivitis, cystitis and gastroenteritis. All HAdVs are excreted
48 in faeces for an extended period of time even if diarrhoea is not present (Mena and Gerba
49 2009; Rodriguez-Lazaro et al. 2012). A recent study has shown the role of intestinal
50 lymphocytes as a reservoir for HAdV persistence and reactivation, whereas the intestinal
51 epithelium is the main site of viral proliferation preceding dissemination in
52 immunocompromised patients (Kosulin et al. 2016). HAdVs are non-enveloped viruses
53 characterized by a double-stranded DNA genome. They belong to the family *Adenoviridae* of
54 the genus *Mastadenovirus* and are divided into seven species (A to G). Ninety genotypes are
55 currently proposed by the Human Adenovirus Working Group (hadv.wg.gmu.edu; update July
56 2018).

57 HAdVs have been proposed as a viral indicator for water quality (Bofill-Mas et al.
58 2010; La Rosa et al. 2010; Wyn-Jones et al. 2011; Rusiñol et al. 2015; Rames et al. 2016;
59 Farkas et al. 2018). Indeed, these viruses reach at least the six following criteria: i) presence
60 from wastewater to surface water, ii) lack of seasonality in wastewater, iii) high resistance to
61 ultraviolet disinfection, iv) higher abundance relative to other enteric viruses, v) availability
62 of molecular methods and culture assays, and vi) human host specificity (Rames et al. 2016).
63 HAdVs seem to be highly persistent in the environment and may reflect the behaviour of
64 persistent enteric viruses (Farkas et al. 2018). Eight viral types of HAdVs belonging to A, C
65 and F species have been reported mainly in water (Ogorzaly et al. 2015). The most prevalent
66 HAdV type detected in wastewater and surface water is enteric type HAdV41 which causes
67 diarrhoea (Rames et al. 2016). Thus, HAdV41 genome was detected in 92% to 96% of
68 wastewater samples positive for HAdVs in Italy (Iaconelli et al. 2017a) and Japan (Haramoto
69 et al. 2007), and in 100% of the hospital wastewater samples in Tunisia (Ibrahim et al. 2018).

70 HAdV41 was also detected in 100% of the wastewater samples positive for HAdVs in
71 Luxembourg (n=8) with a relative abundance estimated to more than 80% using next-
72 generation sequencing (Ogorzaly et al. 2015). This enteric type is also highly prevalent in
73 surface water; it was detected in 94% to 100% of river water samples positive for HAdVs in
74 Luxembourg (Ogorzaly et al. 2015), Italy (La Rosa et al. 2017) and Japan (Haramoto et al.
75 2007). HAdV41 represented 12 to 77% of overall HAdVs in lake samples collected at
76 recreational beaches in the USA (Xagorarakis et al. 2007). The frequent presence of HAdV41
77 in recreational hot springs in Taiwan has also been recently described (Shih et al. 2017).

78 Environmental data on HAdVs have been mainly obtained using molecular methods
79 only; this led to the absence of information about the infectious status of HAdVs in
80 environmental samples (Girones et al. 2010; Ogorzaly et al. 2015). Various methods could be
81 combined with molecular methods to favour the detection of infectious enteric viruses. Most
82 of them are based on the increased permeability of the capsid to intercalating dyes such as
83 ethidium monoazide (EMA) or propidium monoazide (PMA) (Prevost et al. 2016).
84 Intercalating dyes may be useful for the discrimination between infectious and non-infectious
85 viruses such as HAdVs if the inactivation treatment alters the capsid with an increase in its
86 permeability (Leifels et al. 2016). This approach could also be applied for the monitoring of
87 viruses for which cell culture method could not be easily applied on environmental samples or
88 does not exist yet (Prevost et al. 2016).

89 Infectious HAdVs can be quantified using cell culture methods but long periods of time
90 are required for HAdVs to produce cytopathic effects (Rames et al. 2016). The integrated cell
91 culture-quantitative polymerase chain reaction (ICC-qPCR) approach combines cell culture
92 with qPCR. This approach does not require the observation of cytopathic effects and allows
93 the reduction of the incubation period down to one to three days even for environmental
94 samples (Rodriguez et al. 2013; Fongaro et al. 2015; Ogorzaly et al. 2015). Two analytical
95 approaches are described for HAdVs based on the detection of either viral DNA (Greening et

96 al. 2002) or viral mRNA (Ko et al. 2003) in host cells. Viral DNA detection is easily
97 performed, but it requires monitoring of the increase in initial genome copy numbers over
98 time thus reflecting viral replication upon infection (Ogorzaly et al. 2013). The presence of
99 viral mRNA in host cells is evidence of viral infection, but mRNA detection involves a
100 reverse transcription step and an enzymatic digestion of contaminating DNA, whose
101 efficiency has to be checked. Moreover, the variation in mRNA concentration during the virus
102 replication cycle and the susceptibility of such molecules to degradation under some
103 conditions require the use of housekeeping genes as cell-based positive control for mRNA
104 expression (Rodriguez et al. 2013). Another technical aspect of major importance in the use of
105 the ICC-qPCR method for the detection of infectious HAdVs in environmental samples
106 concerns the cell line used. Indeed, the 293 cell line, also called HEK293 or Graham 293
107 (G293), seems to be the most suitable for the detection of a wide diversity of HAdVs
108 occurring in water samples, especially HAdV41 (Ogorzaly et al. 2015). A549 and BGM cell
109 lines are also frequently used for the detection of infectious HAdVs using ICC-qPCR
110 approach (Greening et al. 2002; Lee et al. 2004; Lee et al. 2005; Wyn-Jones et al. 2011;
111 Fongaro et al. 2013; Fongaro et al. 2015), but their use may result in a lack of detection of
112 enteric adenoviruses types 40 and 41 (Wyn-Jones et al. 2011). The infection of permissive
113 cells with serial dilutions of a known HAdVs suspension in parallel with the sample allows
114 the quantification, and not only the detection, of infectious HAdVs. Whereas the ICC-qPCR
115 assay is currently used for quantifying infectious HAdVs in laboratory strains (Gerrity et al.
116 2008; Li et al. 2009; Ryu et al. 2015), very few studies have applied it for the quantification
117 of infectious HAdVs in environmental water samples (Fongaro et al. 2013; Rodriguez et al.,
118 2013; Fongaro et al., 2015).

119 Although river waters are commonly used for production of drinking water or culture
120 irrigation, data are still limited concerning the concentrations of HAdVs in such samples. The
121 occurrence and concentrations of HAdVs have been compared to those of other pathogenic

122 enteric viruses (*e.g.* noroviruses, enteroviruses) but mainly in wastewaters (Rames et al.
123 2016). Moreover, data are lacking on the relative abundance of infectious HAdVs.
124 Comparison between HAdVs and other pathogenic enteric viruses, in particular noroviruses,
125 in terms of total particles and data about concentrations of infectious HAdVs are needed for
126 their evaluation as viral indicators of water quality.

127 In this context, the present study aimed to improve the knowledge about HAdVs in
128 environmental samples. First, the genomes from all types of HAdVs, enteric type HAdV41,
129 NoV GGI and NoV GGII were quantified in 15 river water samples in order to give elements
130 in the evaluation of HAdVs as potential indicators of viral contamination. Second, a one-day
131 ICC-qPCR assay allowing the quantification of infectious HAdVs was applied on the same
132 river samples. This method allowed us to estimate the relative abundance of infectious
133 HAdVs among total HAdV particles. Third, the prevalence of infectious HAdV41 was also
134 investigated in these river water samples.

135

136 **Materials and methods**

137 **Cell line and virus**

138 Human embryonic kidney cell line 293A (Thermo Fisher Scientific, R705-07) was used in
139 this study between passages 15 and 30. Cell line 293A is a sub-clone of 293 cells with a rather
140 flat morphology which strongly adheres to conventional culture surfaces. They were cultured
141 in Dulbecco's modified Eagle's medium (DMEM) containing a high glucose concentration
142 and Glutamax (Gibco). This medium was supplemented with 5% of foetal bovine serum (FBS,
143 Gibco) and 1% of non-essential amino acids (NEAA, Gibco).

144 Human adenovirus type 2 (HAdV2; HPACC, NCPV#213) was chosen since it is a common
145 type in environmental water samples and one of the most studied types in lab-scale
146 experiments. HAdV2 was propagated in 293A cells after three days of culture. The culture
147 medium was supplemented with 2% FBS for HAdV2 replication. After three to four days,

148 FBS was added at a 20% final concentration just before three freeze/thaw cycles. A
149 centrifugation ($2,000 \times g$ for 10 min) was then performed to remove cell debris from the viral
150 lysate. The supernatant was supplemented with sterile glycerol at a 30% final concentration,
151 aliquoted (1.2 mL) and stored at -80°C .

152

153 **River water samples**

154 Over a six-month period (January-June 2016), a total of 15 river water samples (20 L) were
155 collected in the Meurthe River in Maxéville. The samples were collected downstream from
156 the metropolitan wastewater treatment plant (WWTP) (~250,000 inhabitants, Maxéville-
157 Nancy) (Fig. 1 in Supplementary material). The wastewater treatment is based on activated
158 sludge followed by removal of nitrogen and phosphorus. The metropolitan area is equipped
159 with rainwater collection systems with a retention capacity of $180,000 \text{ m}^3$ to reduce the effect
160 of heavy rain events on the operation of the WWTP. The average volume of treated
161 wastewater discharged by the WWTP is $\sim 100,000 \text{ m}^3/\text{day}$ and the mean flow of this river
162 ranges from $\sim 65.7 \text{ m}^3/\text{s}$ in January to $\sim 30.6 \text{ m}^3/\text{s}$ in June (mean of data collected from 1960 to
163 2018, <http://www.hydro.eaufrance.fr/stations/A6941020&procedure=synthese>).

164

165

166 **Quantification of infectious HAdVs in laboratory strain using the most probable** 167 **number assay**

168 The Most Probable Number (MPN) assay was performed to determine the infectious titre of
169 the laboratory strain that was then used for the standard of the ICC-qPCR method. This
170 method is used for the quantification of infectious enteric viruses (Gassilloud et al 2003;
171 Ogorzaly et al 2010; Brié et al 2017). In our case, the MPN assay was performed in 96-well
172 microplaques using 293A cells. Fifty-microlitre portions of successive logarithmic dilutions
173 of each sample were added to 200 μL of preparation containing 3×10^4 293A cells/mL in

174 DMEM containing 2% FBS. Each dilution was seeded in 40 wells. Plates were incubated in
175 5% CO₂ at 37°C for 11 days and then examined for cytopathic effects. The number of wells
176 with cytopathic effects were counted and the MPNCU (most probable number of cytopathic
177 units) was calculated. The final result of each sample analysed was expressed as the geometric
178 mean of the MPNCU per millilitre. The quantification limit of this method is of 1.5
179 MPNCU/mL.

180

181 **Virus concentration from river water samples**

182 A glass wool method was used for virus concentration, as previously described; this method is
183 well adapted to the detection of infectious HAdVs using cell culture methods (Wyn-Jones et
184 al 2011; Ogorzaly et al 2013). Briefly, river samples (20 L) were acidified with 1 M HCl to
185 pH 3.5 and 10 g of sterile glass wool was used for the filtration column. When the entire
186 sample had passed through the filter HAdVs were eluted from the glass wool by slow passage
187 of 200 mL 3% beef extract (w/v) at pH 9.5 in 0.05 M glycine buffer. The eluate was
188 flocculated by the addition of 1 M HCl to pH 3.5. The protein floc, containing viruses, was
189 recovered by centrifugation (7,000 × g for 30 min) and dissolved to a final volume of 20 mL
190 DMEM. The suspension was passed through a combination of pre-filter (5 µm, Sartorius
191 Minisart) and membranes (1.2 and 0.45 µm, Sartorius Minisart) before freezing at -80°C.

192

193 **Genome extraction in river water concentrates**

194 For the quantification of the viral genomes in the river water concentrates, a 1 mL aliquot was
195 mixed with 2 mL of lysis buffer and incubated at room temperature for 10 min. The sample
196 was then mixed with 3 mL of phenol/chloroform/isoamyl alcohol (24:5:1; Acros Organics)
197 and centrifuged (2,000 × g for 5 min). Viral DNA and RNA extraction was then performed on
198 the supernatant using a NucliSens Magnetic extraction kit (bioMérieux). The elution step was
199 carried out with 100 µL of the appropriate buffer and the extracted nucleic acids were

200 immediately stored at -80°C . This kit is well adapted for the extraction of viral genomes and
201 can be easily combined to the phenol/chloroform/isoamyl alcohol pre-treatment applied on
202 the water concentrates.

203

204 **Genome quantification in river water concentrates using digital droplet PCR**

205 The digital droplet PCR approach (dd-PCR) was performed for the quantification of the
206 genomes of HAdVs, HAdV41, NoV GGI and NoV GGII in river water concentrates. For
207 HAdVs, the assay described by Hernroth et al. (2002) using degenerated primers (AdF and
208 AdR) and a TaqMan probe (Ad/ACDEF) was used. This assay is based on the amplification
209 of the hexon gene and recognizes most HAdVs (Bofill-Mas et al. 2006). For HAdV41, the
210 assay based on the amplification of the fibre protein was designed in our laboratory. The
211 TaqMan MGB probe was: 5'-FAM CTT CGC CTT CAA AGT GC-MGB NFQ-3' (position
212 30415-30431 on AdV41 sequence, KY316161.1) and the primers were: forward 5'-CGC
213 CCC ACT TAA TGC TGA CA-3' (position 30387-30406), reverse primer 5'-TCC ACT
214 AGT CCA AGA GGT GCA-3' (position 30432-30452). Primer and probe sequences were
215 subsequently confirmed for specificity using BLAST. NoV GGI and NoV GGII genomes
216 were quantified using primers and probes described in ISO/TS Standard 15216-1:2013
217 (International Organization for Standardization, 2013). Amplifications were performed in a 20
218 μL reaction mixture containing 5 μL of template DNA for HAdVs and HAdV41 or RNA for
219 NoV GGI and GGII, and 15 μL of ddPCRTM Supermix for Probes (BIO-RAD) for HAdVs
220 and HAdV41 and One-Step RT-ddPCRTM Kit for Probes (BIO-RAD) for NoV GGI and GGII.
221 The reaction mix contained 0.9 μM of each primer, 0.225 μM of probe for HAdVs and 0.3
222 μM of probe for NoV. The samples were placed in the droplet generator using 70 μL of
223 generator oil. The resulting picolitre droplet emulsions (40 μL) were transferred to a Veriti
224 96-Well Thermal Cycler (Applied Biosystems). Viral DNA amplification was performed
225 using the following conditions: 10 min hold at 95°C , 40 cycles of 94°C for 30 s then 55°C for

226 60 s and finally a 10 min hold at 98°C with a ramp rate set at 70% between each step. For
227 viral RNA amplification, the reverse transcription was performed with 60 min hold at 50°C.
228 After amplification the plate was transferred to QX100™ Droplet Reader (BIO-RAD) using
229 QuantaSoft™ Software (BIO-RAD) to measure the number of positive and negative droplets
230 based on fluorescence amplitude. Each sample was analysed undiluted and after a 10-fold
231 dilution. Negative and positive controls were included in each experiment to ensure no
232 contamination and good reproducibility of the extractions and PCR assays. The genome
233 concentrations obtained with this method were given in genome copies (gc) per unit volume.

234

235 **Quantification of infectious HAdVs using ICC-qPCR method**

236 For the ICC-qPCR assay the river water samples were analysed concomitantly with a
237 10-fold dilution series of the laboratory strain HAdV2 used as standard in order to quantify
238 infectious HAdVs. The standard was made with concentrations ranging from 10^5 to 10
239 MPNCU/mL. A 1 mL volume of standard or river water sample was inoculated per well of
240 six-well plates (Nunc™) containing two-day-old 293A cells, with a growth area of 9.6 cm². A
241 negative control was included in all our experiments, consisting in the inoculation of cells
242 with 1 mL of DMEM medium. After inoculation, plates were incubated for 2 h at 37°C with
243 5% CO₂ for adsorption of viral particles. Afterwards, the inoculum was removed, and the
244 cells were washed twice with 2 mL of DMEM. Finally, 2 mL of DMEM containing 2% of
245 FBS and supplemented with a 1% final concentration of an antibiotic-antimycotic solution
246 (10,000 units/mL of penicillin, 10,000 units/mL of streptomycin and 25 µg/mL of
247 amphotericin B; Gibco) were added to each well. For the wells marked as T_0 viral DNA was
248 immediately extracted. The wells marked as T_{24h} were incubated at 37°C with 5% CO₂ for one
249 day, before viral DNA extraction which was performed using QIAamp viral RNA mini kit
250 (Qiagen). This kit allows the extraction of viral DNA in particular from infected cell cultures
251 (Ogorzaly et al. 2013, 2015). The culture medium was discarded from each well by aspiration

252 and the 293A cells - potentially infected by viruses - were covered with 1 mL of lysis buffer
253 (AVL buffer + Carrier RNA) for 15 min. Viral nucleic acids were eluted in 100 μ L of the
254 appropriate buffer, according to manufacturer's instructions. The extracts were immediately
255 stored at -80°C . Quantification of the HAdV and HAdV41 genomes in the cell culture
256 samples was performed by a generic real-time PCR system using the same primers and probes
257 as those described for ddPCR. Amplifications were performed in a 25 μ L reaction mixture
258 containing 10 μ L of template DNA and 15 μ L of TaqMan Universal MasterMix (Applied
259 Biosystems) containing 0.9 μ M of each primer and 0.225 μ M of TaqMan probe. Temperature
260 cycles were performed in CFX96 TouchTM Real-Time PCR Detection System (BIO-RAD).
261 PCR amplification was performed at 95°C for 10 min, followed by 45 cycles of 15 s at 95°C
262 and 60 s at 55°C . Negative and positive controls were included in each experiment to ensure
263 no contamination and good reproducibility of the extractions and real-time PCR steps.
264 Samples were considered positive for infectious HAdVs if a decrease in their quantification
265 cycle (Cq) values was observed between T_0 and T_{24h} . Three wells were used for each dilution
266 of the standard (one T_0 and two T_{24h}). For each environmental sample, four wells were each
267 infected with 1 mL of concentrate (one T_0 and three T_{24h}). The concentrations obtained with
268 this method were given in equivalent Most Probable Number of Cytopathic Unit (eqMPNCU)
269 per millilitre. The quantification limit of this method was determined prior to the analysis of
270 the river water samples.

271

272 **Statistical analysis**

273 The statistical analysis was performed using IBM SPSS Statistics version 23.0 Software. A
274 Wilcoxon signed rank test, a Spearman rank correlation test and linear regression tests were
275 performed to compare the viral genome concentrations of HAdVs, HAdV41, NoV GGI and
276 NoV GGII in the river water concentrates.

277

278 **Results**

279

280 **Quantification of viral genomes in the river water samples**

281 The pH and the temperature of the river water samples were measured prior to virus
282 concentration; the mean pH value was 8.2 ± 0.4 and the mean water temperature was
283 $12.0 \pm 3.5^\circ\text{C}$. The cumulative precipitation two days prior to sampling reached 4.1 ± 4.2 mm
284 (Table 1 in Supplementary material).

285 Prior to the quantification of infectious HAdVs in the river water samples, the genome
286 concentrations of HAdVs and HAdV41 were determined using ddPCR method (Fig. 1). The
287 15 samples were positive for these two targets with mean concentrations reaching
288 $3.6 \times 10^3 \pm 4.4 \times 10^3$ gc/L for HAdVs and $2.9 \times 10^3 \pm 2.9 \times 10^3$ gc/L for HAdV41. The
289 concentrations ranged from 1.1×10^2 gc/L (S-14 [Sample 14]) to 1.4×10^4 gc/L (S-11) for
290 HAdVs and from 1.3×10^2 gc/L (S-14) to 9×10^3 gc/L (S-9 and S-11) for HAdV41. The
291 genome concentrations of NoV GGI and NoV GGII were also evaluated in these samples
292 (Fig. 1). The mean concentrations reached $6.1 \times 10^2 \pm 7.4 \times 10^2$ gc/L for NoV GGI and
293 $3.7 \times 10^3 \pm 3.2 \times 10^3$ gc/L for NoV GGII. The lowest genome concentrations were
294 systematically observed for NoV GGI (P-value = 0.001, Wilcoxon signed rank test). The
295 genome concentrations obtained with undiluted and 10-fold diluted samples suggested that
296 our ddPCR assays were not impacted by inhibitory substances. These samples had been
297 collected between January and June, but no difference in the concentration was noticed
298 between the winter and spring periods for any virus. Similarly, no relation was observed
299 between the cumulative precipitation and the genome concentrations for any virus.

300 In order to assess correlation between these viruses, Spearman rank and linear
301 regression analyses were performed on these data (Table 2 in Supplementary material). The
302 strongest correlation was observed between HAdVs and HAdV41 with an r_s value of 0.943
303 (P-value < 0.01) and an R^2 value of 0.94 (Fig. 3a). The comparison between the genome

304 concentrations of HAdVs and HAdV41 showed that the enteric type HAdV41 represented
305 ~80% of HAdVs detected in these river water samples. A strong correlation was also
306 observed between NoV GGI and NoV GGII and between HAdV41 and NoV GGII with r_s
307 values reaching 0.844 (P-value < 0.01) and 0.808 (P-value < 0.01), respectively. Moreover, R^2
308 values of 0.59 and 0.61 were found between HAdV41 and NoV GGII (Fig. 3b) and between
309 NoV GGI and NoV GGII (Fig. 3c), respectively. The R^2 value reached 0.50 between HAdVs
310 and NoV GGII. These results suggested an interrelation between NoV GGII and HAdV41 in
311 these river water samples.

312

313 **Validation of the ICC-qPCR assay on HAdV2 suspension**

314 The ICC-qPCR assay was performed with 10-fold serial dilutions of the HAdV2 laboratory
315 strain (n = 17) ranging from 10^5 to 10 MPNCU/mL which were tested in duplicate in each
316 assay. At T_0 , all samples were positive (*i.e.* Cq value < 45) for concentrations ranging from
317 10^5 to 10^3 MPNCU/mL, whereas only four assays out of 17 were positive for
318 10^2 MPNCU/mL. For 10 MPNCU/mL, all tests were negative (*i.e.* Cq value \geq 45) at T_0 (data
319 not shown). The Cq value at T_0 showed the quantity of HAdVs bound to cells just after their
320 inoculation. Between T_0 and T_{24h} , a decrease in the Cq value was noticed for concentrations
321 ranging from 10^5 to 10 MPNCU/mL, reflecting the replication of viral DNA in the host cells.
322 When no infectious HAdVs were inoculated onto the cells, samples were negative at T_0 and
323 T_{24h} (data not shown). The Cq values obtained at T_{24h} were represented as a function of the
324 concentration of infectious HAdV2 inoculated onto cells (Fig. 1); a linear response was
325 obtained from 10^5 to 10 MPNCU/mL. The mean Cq values ranged from 17.3
326 (10^5 MPNCU/mL) to 30.3 (10 MPNCU/mL). The slope of this ICC-qPCR standard curve was
327 -3.33, with a squared correlation coefficient (R^2) of 0.9981. An increase in the Cq value was
328 observed between T_0 and T_{24h} in 15 assays out of 17 at a concentration of 10 MPNCU/mL
329 which evidenced that this ICC-qPCR method allowed the quantification of infectious HAdVs

330 with a quantification limit of 10 eqMPNCU/mL. Thus, we could expect a quantification limit
331 of 10 eqMPNCU/L of river water.

332 The ICC-qPCR method was then compared to the MPN method on the basis of HAdV2
333 suspensions exposed to different experimental conditions. A titration was performed with
334 both methods just after production for one suspension. Concentrations of infectious HAdVs
335 reached 1.1×10^{10} eqMPNCU/mL and 4.5×10^9 MPNCU/mL with the ICC-qPCR and MPN
336 methods, respectively. Thus, the concentration obtained with the ICC-qPCR method was
337 twice that obtained with the MPN assay. The titrations performed with two suspensions after
338 storage at -80°C for several weeks showed concentrations of 8.3×10^8 eqMPNCU/mL
339 (suspension 1) and 1.5×10^9 eqMPNCU/mL (suspension 2) using ICC-qPCR method and of
340 6.9×10^8 MPNCU/mL (suspension 1) and 2.0×10^9 MPNCU/mL (suspension 2) using the
341 MPN method. For these two suspensions, the concentrations detected with the ICC-qPCR
342 method reached 120% and 75% of those observed with the MPN method.

343

344 **Quantification of infectious HAdVs in the river water samples**

345 Infectious HAdVs were then quantified in the 15 river samples using the ICC-qPCR assay.
346 From the C_q values obtained at T_0 and T_{24h} , 14 samples (93%) were considered as positive for
347 infectious HAdVs. Among them, eight samples (53%) contained between 1.5×10^1 and
348 4.1×10^2 eqMPNCU/L and the other six were below the quantification limit (Fig. 4). When
349 quantifiable, infectious HAdVs represented between 0.3% and 12.2% of total HAdVs.

350 Infectious HAdV41 could not be specifically quantified using our ICC-qPCR assay
351 since it was based on a standard curve obtained with HAdV2. However, the quantification of
352 the HAdV41 genome in infected cells at T_{0h} and T_{24h} allowed us to determine the number of
353 samples positive for infectious HAdV41. Thus, an increase in the concentration of HAdV41
354 genome was observed in infected cells between T_0 and T_{24h} for 11 samples (73%); they were
355 marked with an asterisk in Fig. 4. Infectious HAdV41 particles were detected in the eight

356 samples in which infectious HAdVs were quantified and in three out of the six samples
357 containing infectious HAdVs below the quantification limit.

358

359 **Discussion**

360 The present work aimed to contribute to the evaluation of HAdVs as viral indicators of
361 water quality. For this purpose, comparison between the genome concentrations of human
362 NoV GGI and GGII, which are the main cause of viral acute gastroenteritis, and HAdVs were
363 performed in river water samples characterised by a recent faecal contamination. Infectious
364 HAdVs were also quantified in our river water samples using an ICC-qPCR method. Indeed,
365 the detection of viral genome in environmental samples does not provide information about
366 the infectivity (Girones et al. 2010), whereas this information is still of major importance in
367 the evaluation of the human health risk.

368 The virus particles were concentrated from river water using a glass wool method with
369 pre-acidification. This concentration method was chosen due to the relatively high volume of
370 river water analysed (20 L). Moreover, a mean recovery of 57.1% (range 34.2% - 78.2%)
371 (Wyn-Jones et al. 2011) has previously been obtained with surface water samples including
372 samples from the same river than our study. This concentration method has been previously
373 shown to allow the detection of infectious HAdVs (Wyn-Jones et al. 2011; Ogorzaly et al.
374 2013). Nevertheless, the pH range (pH 4-9) used in primary concentration methods based on
375 adsorption-elution using electronegative or electropositive filters such as glass wool method
376 and also in concentration methods based on organic flocculation and celite may interfere with
377 the detection of infectious viruses (Rames et al. 2016). Indeed, viral inactivation and
378 aggregation may occur with such concentration methods (Rames et al. 2016; Gerba and
379 Betancourt, 2017). Hollow fibre ultrafiltration (HFUF) is also adapted to large volume
380 samples and used for primary concentration, but entrapment of viruses on the filter surface

381 and inhibition of molecular methods from the use of sodium polyphosphate have been
382 described with this method (Rames et al. 2016).

383 The genomes of HAdVs and HAdV41 were detected in all 15 river water samples and
384 their concentrations ranged between 2 and 4 logs gc/L. Such a high proportion of positive
385 samples is in accordance with previous studies performed on the same river (Ogorzaly et al.
386 2009, 2013); prevalence of HAdVs about 90% has also been reported in surface water in other
387 countries (Hamza et al. 2009; Garcia et al. 2012; Ye et al. 2012; Calgua et al. 2013a; Vieira et
388 al. 2016; Farkas et al. 2018). The limitation of our sampling campaign to six months, the
389 exclusion of the fall and early winter season and the lack of heavy rain or snow event within
390 the two days prior to sampling may have contributed to the absence of strong variation
391 between the viral genome concentrations. In a recent study performed on the river Ruhr
392 (Germany) between July and September 2015, the authors observed that enhanced
393 precipitation in combination with an increased flow rate were associated with an increase in
394 the concentration of HAdVs (Mackowiak et al. 2018). HadVs are not known to show seasonal
395 differences in their excretion resulting in lack of seasonality in raw wastewater (Rames et al.
396 2016). In a study performed in the river Ruhr with 184 samples collected between May 2012
397 and June 2013, no significant difference in the percentage of positive samples for HAdVs
398 genome was observed between the bathing season (92.5%, n=80) and outside the bathing
399 season (88.5%, n=104), suggesting a lack of seasonality of this enteric virus (Timm et al.
400 2016). Nevertheless, seasonal occurrence has been described in environmental waters in some
401 cases (Rames et al. 2016). The discharge of treated wastewater from a large urban area
402 (~250,000 inhabitants) upstream from the sampling site had probably contributed to the high
403 prevalence and concentrations observed in our study. Overall, considerable variations in
404 prevalence and concentrations of HAdVs have been reported across studies. Indeed, 16% to
405 100% of surface water samples have been previously shown to be positive for HAdVs with
406 concentrations ranging from 1 to 7 logs gc/L (Bofill-Mas et al. 2013; Iaconelli et al. 2017a).

407 The high proportion of the enteric type HAdV41 (80%) observed in our river water samples is
408 in accordance with the few studies performed on surface water (Xagorarakis et al. 2007;
409 Haramoto et al. 2007; Bofill-Mas et al. 2010; Ogorzalay et al. 2015; La Rosa et al. 2017). The
410 use of the next-generation sequencing (NGS) technology which seems to be the most relevant
411 method to study the diversity of HAdVs in environmental waters has recently shown that
412 HAdV41 was the most abundant type in complex environmental matrices with high
413 concentrations of HAdVs such as wastewater (Ogorzalay et al. 2015; Iaconelli et al. 2017b).

414 All 15 river water samples were also positive for the genomes of NoV GGI and NoV
415 GGII and the concentrations of NoV GGII were similar to those observed for HAdVs and
416 HAdV41. In a recent study performed on the Seine River (France), HAdVs, NoV GGI and
417 NoV GGII genomes have been detected in 93%, 88% and 92% of the samples (n = 100),
418 respectively, and the impact of the wastewater treatment plant effluents on the viral
419 contamination of the river has been clearly shown (Prevost et al. 2015). Overall, considerable
420 variations between HAdVs and NoV have also been reported across studies. Highly different
421 prevalence has been described in some studies (Wyn-Jones et al. 2011; Corsi et al. 2014;
422 Vieira et al. 2016), whereas similar prevalence has been reported in others (Kishida et al.
423 2012; Prevost et al. 2015; Iaconelli et al. 2017a; Jones et al. 2017; La Rosa et al. 2017).
424 Differences in prevalence between sampling sites have also been reported in the same study
425 (Calgua et al., 2013a). Studies comparing the persistence of viral RNA and DNA are still
426 needed (Bertrand et al., 2012), but similar concentrations for HAdV and NoV GGII genomes
427 seem more likely to occur in case of a recent and / or high viral contamination.

428 The ICC-qPCR method used in the present work allowed the quantification of
429 infectious HAdVs in one day with a sensitivity of 10 MPNCU/L of river water. A linear
430 relationship between infectious dose and viral genome in infected cells at T_{24h} was observed
431 between 10^5 and 10^7 MPNCU/L. A few studies have quantified infectious HAdVs naturally
432 occurring in environmental waters and only a small part of them have used the ICC-qPCR

433 method in such a context. Two studies carried out in Brazil have observed high concentrations
434 of infectious HAdVs in groundwater, surface water (*i.e.* lagoon) and spring source water, but
435 also in chlorinated water from a network distribution system using ICC-RT-qPCR (Fongaro et
436 al. 2013, 2015). Thus, the authors have reported concentrations ranging from 10^2 to 10^3 gc/L
437 of infectious HAdVs in 60% (3/5) to 80% (4/5) of groundwater samples and from 10^2 to 10^5
438 gc/L of infectious HAdVs in 83% (5/6) of water samples from a network distribution system
439 (post chlorination). However, infectious HAdVs constituted $\leq 0.12\%$ of total HAdVs in all
440 these samples (Fongaro et al. 2015). Using ICC-RT-qPCR, Rodriguez et al. (2013) have
441 detected infectious HAdVs at concentrations ranging between 10^1 and 10^3 mRNA-Infectious
442 Units/L in sewage (n = 4) in the USA. The ratios between total and infectious HAdVs ranged
443 from 11/1 to 26/1 for three samples containing HAdV-F species only. One sample showed a
444 ratio of 381/1 and contained HAdVs belonging to A, E and F species. This meant that the first
445 three samples contained between 3.8% and 9.1% of infectious HAdVs, whereas the last
446 sample contained 0.2% of infectious HAdVs. Our concentrations (10^1 to 10^3 eqMPNCU/L)
447 and percentages (0.3% to 12.2%) of infectious HAdVs were similar to those observed in
448 sewage (Rodriguez et al. 2013) but different from those detected by Fongaro et al. (2013,
449 2015). The delay between viral contamination and sampling may significantly impact the
450 percentages of infectious HAdVs. The cell lines may also contribute to these discrepancies
451 concerning the percentages of infectious HAdVs. Indeed, Fongaro et al. (2013, 2015) have
452 used A549 cells, whereas Rodriguez et al. (2013) have used G293 cells. As discussed in the
453 introduction, several studies have shown that A549 cells do not allow the detection of
454 HAdV41 which is highly frequent in environmental samples. In the present study, infectious
455 HAdV41 were found in 73% of the river water samples (11/15). Only two studies have
456 described the detection or quantification of infectious HAdV41 in environmental waters and
457 were performed on sewage. Infectious HAdV41 have been detected in all sewage samples in
458 Luxembourg (n = 4) (Ogorzaly et al. 2015) and in the USA (n = 4) (Rodriguez et al. 2013).

459 Above, the possibility of an inactivation of virus particles during the concentration step has
460 been discussed. Rames et al. (2016) have reported in their review a comparison of HAdVs
461 concentrations in raw and treated wastewater using culture-based and qPCR methods for
462 different concentration methods; the percentages of infectious HAdVs ranged from <0.01% to
463 10%. Thus, our percentages of infectious HAdVs (0.3% to 12.2%) suggested that the glass
464 wool method does not impact more deeply the infectivity of virus particles than PEG
465 (polyethylene glycol) precipitation or ultracentrifugation. Overall, further studies on
466 concentration methods adapted to large volume of surface water samples and considering both
467 the recovery rates and the preservation of virus infectivity are still needed. These studies
468 should focus on HAdV41. Intercalating dye treatment prior to qPCR, in particular EMA-
469 qPCR, seems to be a valuable method to evaluate capsid integrity in water matrices. It could
470 be a rapid, relatively inexpensive and easy to use tool in the evaluation of the effect of
471 concentration methods on the virus integrity (Leifels et al. 2016; Prevost et al. 2016). For
472 ICC-qPCR approach, studies should take into account that specific cell lines such as 293
473 should be used and that HAdV41 might have a longer multiplication cycle than other HAdV
474 types (Siqueira-Silva et al. 2009).

475

476 **Conclusions**

477 Our work shows that not only HAdV genomes but also infectious particles can be frequent in
478 river water in Europe. Indeed, infectious HAdVs reached more than 10 infectious particles/L
479 in more than 50% of the 15 samples. HAdV41 genome was highly predominant in these river
480 samples and a correlation between HAdV41 and NoV GGII genomes was observed. The
481 presence of these two enteric viruses should be studied in relation with effluent discharge in
482 particular for rivers with a strong anthropic pressure. Infectious HAdV41 was detected with a
483 frequency higher than 70%. This predominance of HAdV41 genome over other HAdV types
484 and the high prevalence of infectious HAdV41 in environmental waters underline the need for

485 further studies focusing on the specific quantification of infectious particles of HAdV41. Such
486 data are needed to improve the knowledge on this enteric type and its potential as a viral
487 indicator of water quality.

488

489 **Acknowledgements**

490 The present work was financially supported by the Institut Jean Barriol (CNRS and Université
491 de Lorraine). Complementary financial support was obtained from Zone Atelier Moselle
492 (ZAM). The authors thank Romain Rivet for his excellent technical assistance in ddPCR
493 assays and Coline Wietrich for her contribution to the ddPCR analyses.

494

495 **Conflict of interest**

496 The authors declare that they have no conflict of interest.

497

498

499 **References**

- 500 Bertrand I, Schijven JF, Sánchez G, Wyn-Jones P, Ottoson J, Morin T, Muscillo M, Verani
501 M, Nasser A, de Roda Husman AM, Myrmet M, Sellwood J, Cook N, Gantzer C (2012)
502 The impact of temperature on the inactivation of enteric viruses in food and water: a
503 review. *J Appl Microbiol* 112:1059-7
- 504 Brié A, Razafimahefa R, Loutreul J, Robert A, Gantzer C, Boudaud N, Bertrand I (2017) The
505 Effect of heat and free chlorine treatments on the surface properties of murine
506 norovirus. *Food Environ Virol* 9:149-15
- 507 Bofill-Mas S, Albinana-Gimenez N, Clemente-Casares P, Hundesa A, Rodriguez-Manzano J,
508 Allard A, Calvo M, Girones R (2006) Quantification and stability of human
509 adenoviruses and polyomavirus JCPyV in wastewater matrices. *Appl Environ Microbiol*
510 72:7894-6
- 511 Bofill-Mas S, Calgua B, Clemente-Casares P, La Rosa G, Iaconelli M, Muscillo M, Rutjes S,
512 de Roda Husman AM, Grunert A, Gräber I, Verani M, Carducci A, Calvo M, Wyn-
513 Jones P, Girones R (2010) Quantification of human adenoviruses in European
514 recreational waters. *Food Environ Virol* 2:101-109
- 515 Bofill-Mas S, Rusiñol M, Fernandez-Cassi X, Carratalà A, Hundesa A, Girones R (2013)
516 Quantification of human and animal viruses to differentiate the origin of the fecal
517 contamination present in environmental samples. *Biomed Res Int*:192089
- 518 Calgua B, Fumian T, Rusiñol M, Rodriguez-Manzano J, Mbayed VA, Bofill-Mas S,
519 Miagostovich M, Girones R (2013a) Detection and quantification of classic and
520 emerging viruses by skimmed-milk flocculation and PCR in river water from two
521 geographical areas. *Water Res* 47:2797-810.
- 522 Corsi SR, Borchardt MA, Spencer SK, Hughes PE, Baldwin AK (2014) Human and bovine
523 viruses in the Milwaukee River watershed: hydrologically relevant representation and
524 relations with environmental variables. *Sci Total Environ* 490:849-60

525 Farkas K, Cooper DM, McDonald JE, Malham SK, de Rougemont A, Jones DL (2018)
526 Seasonal and spatial dynamics of enteric viruses in wastewater and in riverine and
527 estuarine receiving waters. *Sci Total Environ* 634:1174-1183

528 Fongaro G, Nascimento MA, Rigotto C, Ritterbusch G, da Silva AD, Esteves PA, Barardi CR
529 (2013) Evaluation and molecular characterization of human adenovirus in drinking
530 water supplies: viral integrity and viability assays. *Virology* 10:166

531 Fongaro G, Padilha J, Schissi CD, Nascimento MA, Bampi GB, Viancelli A, Barardi CR
532 (2015) Human and animal enteric virus in groundwater from deep wells, and
533 recreational and network water. *Environ Sci Pollut Res Int* 22:20060-6

534 Garcia LA, Viancelli A, Rigotto C, Pilotto MR, Esteves PA, Kunz A, Barardi CR (2012)
535 Surveillance of human and swine adenovirus, human norovirus and swine circovirus in
536 water samples in Santa Catarina, Brazil. *J Water Health* 10:445-52

537 Gassilloud B, Schwartzbrod L, Gantzer C (2003) Presence of viral genomes in mineral water:
538 a sufficient condition to assume infectious risk? *Appl Environ Microbiol* 69:3965-9

539 Gerba CP, Betancourt WQ (2017) Viral aggregation: impact on virus behavior in the
540 environment. *Environ Sci Technol* 51:7318-7325.

541 Gerrity D, Ryu H, Crittenden J, Abbaszadegan M (2008) UV inactivation of adenovirus type
542 4 measured by integrated cell culture qPCR. *J Environ Sci Health A Tox Hazard Subst*
543 *Environ Eng* 43:1628-38

544 Girones R, Ferrús MA, Alonso JL, Rodriguez-Manzano J, Calgua B, Corrêa Ade A, Hundesa
545 A, Carratala A, Bofill-Mas S (2010) Molecular detection of pathogens in water--the
546 pros and cons of molecular techniques. *Water Res* 44:4325-39

547 Greening GE, Hewitt J, Lewis GD (2002) Evaluation of integrated cell culture-PCR (C-PCR)
548 for virological analysis of environmental samples. *J Appl Microbiol* 93:745-50

549 Hamza IA, Jurzik L, Stang A, Sure K, Uberla K, Wilhelm M (2009) Detection of human
550 viruses in rivers of a densely-populated area in Germany using a virus adsorption elution
551 method optimized for PCR analyses. *Water Res* 43:2657-68

552 Haramoto E, Katayama H, Oguma K, Ohgaki S (2007) Quantitative analysis of human enteric
553 adenoviruses in aquatic environments. *J Appl Microbiol* 103:2153-9

554 Hernroth BE, Conden-Hansson AC, Rehnstam-Holm AS, Girones R, Allard AK (2002)
555 Environmental factors influencing human viral pathogens and their potential indicator
556 organisms in the blue mussel, *Mytilus edulis*: the first Scandinavian report. *Appl*
557 *Environ Microbiol* 68:4523-33

558 Iaconelli M, Muscillo M, Della Libera S, Fratini M, Meucci L, De Ceglia M, Giacosa D, La
559 Rosa G (2017a) One-year surveillance of human enteric viruses in raw and treated
560 wastewaters, downstream river waters, and drinking waters. *Food Environ Virol.* 9:79-
561 88

562 Iaconelli M, Valdazo-González B, Equestre M, Ciccaglione AR, Marcantonio C, Della Libera
563 S, La Rosa G (2017b) Molecular characterization of human adenoviruses in urban
564 wastewaters using next generation and Sanger sequencing. *Water Res* 121:240-247

565 Ibrahim C, Hassen A, Pothier P, Mejri S, Hammami S (2018) Molecular detection and
566 genotypic characterization of enteric adenoviruses in a hospital wastewater. *Environ Sci*
567 *Pollut Res Int* 25:10977-10987.

568 International Organization for Standardization. 2013. ISO/TS 152616-1:2013. Microbiology
569 of the food chain--Horizontal method for determination of hepatitis A virus and
570 norovirus using real-time RT-PCR--Part1: Method for quantification.
571 <https://www.iso.org/standard/65681.html>

572 Jones TH, Brassard J, Topp E, Wilkes G, Lapen DR (2017) Waterborne viruses and F-specific
573 coliphages in mixed-use watersheds: microbial associations, host specificities, and

574 affinities with environmental/land use factors. *Appl Environ Microbiol* 83, pii:
575 e02763-16

576 Kishida N, Morita H, Haramoto E, Asami M, Akiba M (2012) One-year weekly survey of
577 noroviruses and enteric adenoviruses in the Tone River water in Tokyo metropolitan
578 area, Japan. *Water Res* 46:2905-10

579 Ko G, Cromeans TL, Sobsey MD (2003) Detection of infectious adenovirus in cell culture by
580 mRNA reverse transcription-PCR. *Appl Environ Microbiol* 69:7377-84.

581 Kosulin K, Geiger E, Vécsei A, Huber WD, Rauch M, Brenner E, Wrba F, Hammer K,
582 Innerhofer A, Pötschger U, Lawitschka A, Matthes-Leodolter S, Fritsch G, Lion T
583 (2016) Persistence and reactivation of human adenoviruses in the gastrointestinal tract.
584 *Clin Microbiol Infect* 22:381.e1-381.e8

585 La Rosa G, Pourshaban M, Iaconelli M, Muscillo M (2010) Quantitative real-time PCR of
586 enteric viruses in influent and effluent samples from wastewater treatment plants in
587 Italy. *Ann Ist Super Sanita* 46:266-73

588 La Rosa G, Sanseverino I, Della Libera S, Iaconelli M, Ferrero VEV, Barra Caracciolo A,
589 Lettieri T (2017) The impact of anthropogenic pressure on the virological quality of
590 water from the Tiber River, Italy. *Lett Appl Microbiol.* 65:298-305.

591 Lee C, Lee SH, Han E, Kim SJ (2004) Use of cell culture-PCR assay based on combination of
592 A549 and BGMK cell lines and molecular identification as a tool to monitor infectious
593 adenoviruses and enteroviruses in river water. *Appl Environ Microbiol* 70:6695-705

594 Lee SH, Lee C, Lee KW, Cho HB, Kim SJ (2005) The simultaneous detection of both
595 enteroviruses and adenoviruses in environmental water samples including tap water
596 with an integrated cell culture-multiplex-nested PCR procedure. *J Appl Microbiol*
597 98:1020-9

598 Leifels M, Hamza IA, Krieger M, Wilhelm M, Mackowiak M, Jurzik L (2016) From lab to
599 lake - Evaluation of current molecular methods for the detection of infectious enteric
600 viruses in complex water matrices in an urban area. *PLoS One* 11:e0167105

601 Li F, Feng L, Liu Y, Zhen X, Chen L (2009) An integrated cell culture and quantitative
602 polymerase chain reaction technique for determining titers of functional and infectious
603 adenoviruses. *Anal Biochem* 391:157-9

604 Mena KD, Gerba CP (2009) Waterborne adenovirus. *Rev Environ Contam Toxicol*.
605 198:133-67

606 Ogorzaly L, Tissier A, Bertrand I, Maul A, Gantzer C (2009) Relationship between F-specific
607 RNA phage genogroups, faecal pollution indicators and human adenoviruses in river
608 water. *Water Res* 43:1257-64

609 Ogorzaly L1, Bertrand I, Paris M, Maul A, Gantzer C (2010) Occurrence, survival, and
610 persistence of human adenoviruses and F-specific RNA phages in raw groundwater.
611 *Appl Environ Microbiol* 76:8019-25

612 Ogorzaly L, Cauchie HM, Penny C, Perrin A, Gantzer C, Bertrand I (2013) Two-day
613 detection of infectious enteric and non-enteric adenoviruses by improved ICC-qPCR.
614 *Appl Microbiol Biotechnol* 97:4159-66

615 Ogorzaly L, Walczak C, Galloux M, Etienne S, Gassilloud B, Cauchie HM (2015) Human
616 Adenovirus diversity in water samples using a next-generation amplicon sequencing
617 approach. *Food Environ Virol* 7:112-121

618 Prevost B, Lucas FS, Goncalves A, Richard F, Moulin L, Wurtzer S (2015) Large scale
619 survey of enteric viruses in river and waste water underlines the health status of the
620 local population. *Environ Int* 79:42-50

621 Prevost B, Goulet M, Lucas FS, Joyeux M, Moulin L, Wurtzer S (2016) Viral persistence in
622 surface and drinking water: Suitability of PCR pre-treatment with intercalating dyes.
623 *Water Res* 91:68-76

624 Rames E, Roiko A, Stratton H, Macdonald J (2016) Technical aspects of using human
625 adenovirus as a viral water quality indicator. *Water Res* 96:308-26

626 Rodríguez RA, Polston PM, Wu MJ, Wu J, Sobsey MD (2013) An improved infectivity assay
627 combining cell culture with real-time PCR for rapid quantification of human
628 adenoviruses 41 and semi-quantification of human adenovirus in sewage. *Water Res*
629 47:3183-91

630 Rodríguez-Lázaro D1, Cook N, Ruggeri FM, Sellwood J, Nasser A, Nascimento MS,
631 D'Agostino M, Santos R, Saiz JC, Rzeżutka A, Bosch A, Gironés R, Carducci A,
632 Muscillo M, Kovač K, Diez-Valcarce M, Vantarakis A, von Bonsdorff CH, de Roda
633 Husman AM, Hernández M, van der Poel WH (2012) Virus hazards from food, water
634 and other contaminated environments. *FEMS Microbiol Rev* 36:786-814

635 Rusiñol M, Fernandez-Cassi X, Timoneda N, Carratalà A, Abril JF, Silvera C, Figueras MJ,
636 Gelati E, Rodó X, Kay D, Wyn-Jones P, Bofill-Mas S, Girones R (2015) Evidence of
637 viral dissemination and seasonality in a Mediterranean river catchment: Implications for
638 water pollution management. *J Environ Manage* 15:159:58-67

639 Ryu H, Cashdollar JL, Fout GS, Schrantz KA, Hayes S (2015) Applicability of integrated cell
640 culture quantitative PCR (ICC-qPCR) for the detection of infectious adenovirus type 2
641 in UV disinfection studies. *J Environ Sci Health A Tox Hazard Subst Environ Eng*
642 50:777-87

643 Shih YJ, Tao CW, Tsai HC, Huang WC, Huang TY, Chen JS, Chiu YC, Hsu TK, Hsu BM
644 (2017) First detection of enteric adenoviruses genotype 41 in recreation spring areas of
645 Taiwan. *Environ Sci Pollut Res Int* In Press

646 Siqueira-Silva J, Yeda FP, Favier AL, Mezin P, Silva ML, Barrella KM, Mehnert DU, Fender
647 P, Hársi CM (2009) Infection kinetics of human adenovirus serotype 41 in HEK 293
648 cells. *Mem Inst Oswaldo Cruz* 104:736-44

649 Timm C, Luther S, Jurzik L, Hamza IA, Kistemann T (2016) Applying QMRA and DALY to
650 assess health risks from river bathing. *Int J Hyg Environ Health* 9:681-692

651 Vieira CB, de Abreu Corrêa A, de Jesus MS, Luz SL, Wyn-Jones P, Kay D, Vargha M,
652 Miagostovich MP (2016) Viruses surveillance under different season scenarios of the
653 Negro river basin, Amazonia, Brazil. *Food Environ Virol* 8:57-69

654 Wyn-Jones AP, Carducci A, Cook N, D'Agostino M, Divizia M, Fleischer J, Gantzer C,
655 Gawler A, Girones R, Höller C, de Roda Husman AM, Kay D, Kozyra I, López-Pila J,
656 Muscillo M, Nascimento MS, Papageorgiou G, Rutjes S, Sellwood J, Szewzyk R, Wyer
657 M (2011) Surveillance of adenoviruses and noroviruses in European recreational waters.
658 *Water Res* 45:1025-38

659 Xagorarakis I, Kuo DH, Wong K, Wong M, Rose JB (2007) Occurrence of human
660 adenoviruses at two recreational beaches of the great lakes. *Appl Environ Microbiol.*
661 73:7874-81

662 Ye XY, Ming X, Zhang YL, Xiao WQ, Huang XN, Cao YG, Gu KD (2012) Real-time PCR
663 detection of enteric viruses in source water and treated drinking water in Wuhan, China.
664 *Curr Microbiol* 65:244-53

665

Figure legends

666

667

668 **Figure 1.** Genome concentrations of HAdVs, HAdV41, NoV GGI and NoV GGII in the river
669 water samples ($n = 15$) using ddPCR.

670

671 **Figure 2.** Standard curve of the ICC-qPCR assay generated from the range of dilutions of
672 HAdV2 ($n = 17 \times 2$).

673

674 **Figure 3.** Relation between the concentrations of (a) HAdVs and HAdV41, (b) NoV GGII
675 and HAdV41, and (c) NoV GGII and NoV GGI in the 15 river water samples.

676

677 **Figure 4.** Concentrations of HAdV genomes and infectious HAdVs in the river water samples
678 ($n = 15$) using ddPCR and ICC-qPCR, respectively. Samples positive for infectious HAdV41
679 are notified with an asterisk. ND: not detected. NQ: not quantified.*: water samples positive
680 for infectious HAdV41.

681

682

683 **Supplementary material**

684 **Figure 1.** Sampling point in the river Meurthe in the metropolitan area of Nancy. Black
685 arrow: discharge of treated wastewater in the river. Red star: sampling point in the river

686