The cephalic labial gland secretions of two socially parasitic bumblebees *Bombus hyperboreus* (*Alpinobombus*) and *Bombus inexpectatus* (*Thoracobombus*) question their inquiline strategy

Nicolas Brasero, Baptiste Martinet, Thomas Lecocq, Patrick Lhomme, Paolo Biella, Irena Valterova, Klára Urbanová, Maurizio Cornalba, Heather Hines, Pierre Rasmont

► To cite this version:

Nicolas Brasero, Baptiste Martinet, Thomas Lecocq, Patrick Lhomme, Paolo Biella, et al.. The cephalic labial gland secretions of two socially parasitic bumblebees *Bombus hyperboreus* (*Alpinobombus*) and *Bombus inexpectatus* (*Thoracobombus*) question their inquiline strategy. *Insect Science*, 2018, 25 (1), pp.75-86. 10.1111/1744-7917.12408 . hal-01972534

HAL Id: hal-01972534

<https://hal.univ-lorraine.fr/hal-01972534>

Submitted on 11 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The cephalic labial gland secretions of two socially parasitic bumblebees *Bombus hyperboreus* (*alpinobombus*) and *Bombus inexpectatus* (*thoracobombus*) question their inquiline strategy

Nicolas Brasero^{1*¶}, Baptiste Martinet^{1*¶}, Thomas Lecocq^{1,2}, Patrick Lhomme³, Paolo Biella⁴, Irena Valterova⁵, Klara Urbanova^{5,6}, Maurizio Cornalba⁷, Heather Hines³, Pierre Rasmont¹

1: University of Mons, Research institute of Biosciences, Laboratory of Zoology, Place du Parc 20, 7000 Mons, Belgium

2: Research Unit Animal and Functionalities of Animal Products (URAFPA), University of Lorraine – INRA, 2 Avenue de la Forêt de Haye, BP 172, 54505, Vandoeuvre-lès-Nancy, France

3: The Pennsylvania State University, Department of Biology, 208 Mueller Labs, University Park, PA 16802, United States of America (USA)

4: University of South Bohemia, Department of Zoology, Faculty of Science, Branišovská 31, 37005, České Budějovice, Czech Republic; Biology Centre of the Academy of Sciences of the Czech Republic, v.v.i., Institute of Entomology, Branišovská 31, 37005, České Budějovice, Czech Republic

5: Academy of Sciences of the Czech Republic, Institute of Organic Chemistry and Biochemistry, Flemingovo nám 2, CZ-166 10 Prague, Czech Republic

6: Czech University of Life Sciences, Faculty of Tropical AgriSciences, Department of Sustainable Technologies, Kamýcká 129, CZ-165 21 Prague, Czech Republic

7: University of Pavia, Department of Mathematics, via Ferrata 5, I-27100 Pavia, Italy

Short Title: Involvement of CLGS in the inquilinism of *B. hyperboreus* and *B. inexpectatus*

* Co-first authors and Corresponding authors; E-mail: nicolas.brasero@umons.ac.be

(NB) 0032(0)65373435; baptiste.martinet@umons.ac.be (BM) 0032(0)65373405

¶ These first co-authors contributed equally to this work

Running title: CLGS in social parasite

Abstract- Socially parasitic Hymenoptera have evolved morphological, chemical, and behavioral adaptations to overcome the sophisticated recognition and defense systems of their social host to invade host nests and exploit their worker force. In bumblebees, social parasitism appeared in at least three subgenera independently: in the subgenus *Psithyrus* consisting entirely of parasitic species, in the subgenus *Alpinobombus* with *B. hyperboreus*, and in the subgenus *Thoracobombus* with *B. inexpectatus*. Cuckoo bumblebee males utilize species-specific cephalic labial gland secretions for mating purposes which have been showed as acting in their inquiline strategy. We performed cephalic labial gland secretions comparative analyses of *B. hyperboreus*, *B. inexpectatus* and their hosts. Males of both parasitic species exhibit high species specific levels of cephalic gland secretions, including different main compounds. Our results show no chemical mimicry in the cephalic gland secretions between inquilines and their host and we did not identify the repellent compounds already known in other cuckoo bumblebees.

Key Words Social parasitism, bumblebees, *Bombus hyperboreus*, *Bombus inexpectatus*, cephalic labial gland secretions, inquiline strategy

INTRODUCTION

Brood care is an energy-consuming activity for social insects ranging from nest construction to offspring breeding (Smisteh et al. 2012). This high energetic cost has favored the evolution of cheating species that exploit brood care behavior of hetero-specifics for their own offspring rearing (Roldan & Soler 2011). This adaptive strategy involves the exploitation of host social behavior by specialized species (so called social parasites) (Lenoir et al. 2001). Social parasites have evolved morphological, chemical, and behavioral adaptations (usurpation behaviors) to overcome the sophisticated recognition system (based on species-specific pheromonal and cuticular signatures) and defense systems of their social host (Fisher & Sampson 1992; Martin et al. 2010).

Among bumblebees (Hymenoptera, Apidae, *Bombus*), the cuckoo species exemplify such a social parasitic strategy. Their life cycle relies on a non-mutual symbiotic relationship (called inquilinism) with their hosts (Lhomme et al. 2012). This interaction leads to a decrease or, in some cases, to a complete loss of host reproductive success (Lhomme et al. 2013). These “cheaters” infiltrate host colonies to take advantage of their food resources and workers (Nash & Boomsma 2008). In bumblebees, obligate inquilinism evolved in at least three subgenera independently (Hobbs 1967; Wilson 1971; Michener 1974): in the subgenus *Psithyrus* consisting entirely of parasitic species (Williams 1998), in the subgenus *Alpinobombus* with *B. hyperboreus* (Milliron & Oliver 1966; Richards 1973) (Fig. 1A), and in the subgenus *Thoracobombus* with *B. inexpectatus* (Yarrow 1970; Williams 1998; Müller 2006) (Fig. 1C).

All *Psithyrus* species are completely dependent on their host because these socially parasitic bumblebees do not produce a worker caste, do not have pollen collection baskets and produce little to no wax to build cells (Benton 2006; Sramkova & Ayasse 2008). In the *Alpinobombus* subgenus, the circum arctic *B. (Alpinobombus) hyperboreus* is mainly known

as an inquiline of *B. (Alpinobombus) polaris* (Fig. 1B) at least in some parts of its range (Milliron & Oliver 1966; LØken 1973; Richards 1973; Pape 1983; Stenström & Bergmann 1998; Rampini *et al.* 2012), but nests of *B. (Alpinobombus) balteatus* and *B. (Pyrobombus) jonellus* have been found to be infiltrated by *B. hyperboreus* in Norway (Bergwall 1970; Gjershaug 2009). In the Eurasian Arctic and in North America, some females were reported harvesting pollen (Gjershaug 2009), suggesting a facultative inquilinism for this species. In the *Thoracobombus* subgenus, Yarrow (1970) and Müller (2006) have highlighted a number of peculiar biological and morphological traits of *B. (Thoracobombus) inexpectatus* (i.e.: absence of worker caste, loss of wax production and reduction of the hair-like structures of the hind leg used to gather pollen) that likely are adaptations to parasitic life. *Bombus (Thoracobombus) ruderarius* (Fig. 1D) was described as the only host of *B. inexpectatus* (Müller 2006). This sub-alpine European species occurs in the Cantabrian Mountains and the Alps from 1200m to 2100m in altitude where it is uncommon (Tkalců 1963; Tkalců 1965; Yarrow 1970; Amiet 1996; Rasmont *et al.* 2015). Compared to their respective hosts, the geographical distribution of *B. inexpectatus* and *B. hyperboreus* and their ecological niches are restricted (Rasmont & Iserbyt 2014).

The infiltration strategies of parasitic females have been the focus of abundant research (Nash & Boomsma 2008): some bumblebee species produce very low amounts of cuticular hydrocarbons (chemical insignificance) and/or acquire their host's chemical profile (chemical mimicry) (Dronnet *et al.* 2005; Martin *et al.* 2010), whereas others use repellents (allomones) to defend themselves against the attacks of host workers (Zimma *et al.* 2003; Martin *et al.* 2010). More recently, few studies have started to investigate the strategies used by bumblebee inquiline offspring to be tolerated by host workers (Lhomme *et al.* 2012; Lhomme *et al.* 2015).

For male offspring of inquiline species, avoiding host recognition is especially challenging because of their early production of large amounts of the species-specific Cephalic Labial Gland Secretions (CLGS) involved in attracting conspecific females during the pre-mating behavior (Žáček *et al.* 2009; Lhomme *et al.* 2012; Ayasse and Jarau 2014). Indeed, CLGS are a complex mixture of (mainly aliphatic) compounds (Calam 1969) derived from saturated fatty acids (Luxová *et al.* 2003) involved in pre-mating behavior (Bergström *et al.* 1981). The species-specific CLGS are synthesized *de novo* by cephalic labial glands in the head of bumblebee males (Žáček *et al.* 2013). This highly concentrated specific signature are discriminated by host workers and potentially lead to male offspring destruction (Lhomme *et al.* 2012). Consequently, a strong selective pressure (beside the sexual selection) on the inquiline male CLGS should likely lead to adaptive changes. Similarly to alternative chemical infiltration strategies, one could expect three alternative adaptive patterns: (i) chemical insignificance meaning that males do not produce CLGS for pre-mating behavior or produce CLGS only after leaving the host nest or even males produce undetected compounds; (ii) chemical mimicry meaning that the inquiline actively produce the same compounds as its host; or (iii) chemical repellency meaning that specific repellent compounds are produced within the CLGS and act as "chemical shield" to avoid host worker aggressiveness. A previous study on the cuckoo bumblebee *B. (Psithyrus) vestalis* demonstrated a chemical repellent effect of CLGS on host workers triggered by tetradecyl acetate and (Z)-11-hexadecenyl acetate (Lhomme *et al.* 2015). Potential other chemical strategies used by other *Psithyrus* or non-*Psithyrus* inquiline species is unknown. Here, we performed (i) the first detailed description of the two non-*Psithyrus* inquiline CLGS and (ii) comparative analyses of CLGS composition between *B. hyperboreus*, *B. inexpectatus* and their hosts. We finally discuss the potential chemical strategies involved in these host-inquiline interactions.

Materials and Methods

Sampling

In total, 69 specimens collected belonging to two subgenera were used (Table 1; Supplementary material 1). The test group comprised *B. hyperboreus* (n=15) (Fennoscandia and Alaska), and its main host *B. polaris* (n=13) (Fennoscandia and Alaska), and *B. inexpectatus* (n=9) (Alps), and its host *B. ruderarius* (n=8) (Alps and Pyrenees). As a reference group, we used one phylogenetically close species of the host bumblebees of each parasite according to Cameron *et al.* [41] : *B. (Alpinobombus) alpinus* (n=14) (Fennoscandia) and *B. (Thoracobombus) sylvarum* (n=10) (Pyrenees and Massif Central) (Terzo *et al.* 2005). Concerning *B. alpinus*, the population in Alps have been excluded of the analysis to respect the geographical distribution of the host and parasite. For *Alpinobombus*, we chose to follow the taxonomy of Milliron (1971) and Løken (1973) as recent updates of Williams *et al.* (2015) still require further investigation. Exact localities are listed in (Supplementary material 1). All bumblebee specimens were killed by freezing at -20° C.

Extraction of Bumblebee Male Cephalic Labial Gland Secretions

We focused on the most studied reproductive trait involved in the bumblebee pre-mating recognition (Ayasse *et al.* 2001; Baer 2003; Ayasse & Jarau 2014; Lecocq *et al.* 2015b): the Cephalic Labial Gland Secretions (CLGS).

The CLGS were extracted from dissected whole cephalic labial glands using 400 µl of n-hexane -see protocol in (De Meulemeester *et al.* 2011). Samples were stored at - 40° C prior to the analyses. The composition of CLGS was compared to that established by (Žáček *et al.* 2013) with the same GC conditions to highlight potential common repellent compounds.

GC/MS Analysis- Identification

The composition was determined by Gas Chromatography - Mass Spectrometry (GC/MS) using a Focus GC (Thermo Scientific) with a non-polar DB-5 ms capillary column [5% phenyl (methyl) polysiloxane stationary phase; column length 30 m; inner diameter 0.25 mm; film thickness 0.25 μ m] coupled to DSQ II quadrupol mass analyzer (Thermo Scientific) with 70 eV electron impact ionization. We used a splitless injection mode (220°C) and helium as a carrier gas (one ml/min). The temperature program of the oven was set to 70°C for two minutes and then heated up at a rate of 10°C/min to 320°C. The temperature was then held at 320°C for five minutes. Compounds were identified in XcaliburTM using the retention times (t_r) and mass spectra of each peak, in comparison to those at National Institute of Standards and Technology library (NIST, U.S.A). Double bond positions (C=C) were determined by dimethyl disulfide (DMDS) derivatization (Cvacka *et al.* 2008).

GC-FID Analysis- Quantification

All samples were quantified (quantitative composition) with Gas Chromatography with a Flame Ionization Detector (GC-FID) Shimadzu GC-2010 system equipped with a non-polar SLB-5 ms capillary column [5% phenyl (methyl) polysiloxane stationary phase; column length 30 m; inner diameter 0.25 mm; film thickness 0.25 μ m]. The chromatographic conditions were the same as described above (cf. GC/MS Analysis). We quantified the peak areas of compounds in GC solution postrun (Shimadzu Corporation) with automatic peak detection and noise measurement. The relative areas (RAs, expressed in %) of compounds in each sample were calculated by dividing the peak areas of compounds by the total area of all compounds. All compounds for which RA were recorded as less than 0.1% for all specimens were excluded from the analysis (De Meulemeester *et al.* 2011). We identified the main compound as the compound that has the highest relative proportion (RA) among all compounds of CLGSs at least in one individual of the taxon. The data matrix (Supplementary material 2) for each taxa was based on the alignment of each relative proportion of compound

between all samples performed with GCAAligner 1.0 (Dellicour & Lecocq 2013a). To facilitate the alignment of compounds, before each sample injection, a standard (Kovats) were injected containing a mix of pure hydrocarbons (alkanes) from C 10 (decane) to C 40 (tetracontane). Kovats indices were calculated with GCKovats 1.0 according to the method described by (Dellicour and Lecocq 2013b).

Statistical Analysis

Each pair of parasite-host complex was considered independently. We performed statistical comparative analyses based on the CLGS matrix using R (R Development Core Team 2013) to detect chemical differentiations between hosts and parasites. A clustering method, computed with the unweighted pair-group method with average linkage (UPGMA) based on Pearson correlation distance matrices (RA of each compound) (Legendre & Legendre 1983; Paradis *et al.* 2004) was used to detect the divergence between taxa in the CLGS composition. We assessed the uncertainty in hierarchical cluster analysis using p-values calculated *via* multiscale bootstrap resampling with 100 000 bootstrap replications (R package pvclust) (Suzuki & Shimodaira 2011). Only bootstrap values greater than 90% were considered significant. To determine specific compounds of each taxon (indicator compounds - I.C.), the indicator-value (IndVal) method was used (Dufrêne & Legendre 1997). This value is the product of relative concentration and relative occurrence frequency of a compound within a group. The statistical significance of an indicator compound (threshold of 0.01) was evaluated with a randomization procedure.

RESULTS

Identified Compounds Within Cephalic Labial Gland Secretions

Alpinobombus

In total, we have detected 117 compounds for *Alpinobombus* subgenus (69 from *B. hyperboreus*, 72 from *B. polaris* and 53 from *B. alpinus*) in the CLGS (Supplementary material 2). The results showed that all species were differentiated in their CLGS composition (23 compounds shared by all species inside the *Alpinobombus* group). 34 compounds are shared between the parasite *B. hyperboreus* and the host *B. polaris*, 39 between *B. polaris* and *B. alpinus* and 27 between *B. hyperboreus* and *B. alpinus* (Supplementary material 2). In addition to qualitative differences in minor compounds, other differences are also present in main compounds. In *Alpinobombus* group, octadec-11-en-1-ol (23.90% - 56.21%, Table 2) is the main compound in the *B. hyperboreus* CLGS. While the CLGS of *B. polaris* is dominated by hexadec-9-enal (20.8% - 43.03%) and hexadec-9-en-1-ol (26.87% - 46.68%). The main compound of *B. alpinus* is hexadec-9-en-1-ol (63.95% - 89.43%) (Table 2). Thus the main compound of *B. alpinus* (hexadec-9-en-1-ol) is shared by its sister species *B. polaris* and not its social parasite (Table 2).

The IndVal method revealed several indicator compounds (I.C.) with high statistical significance (IndVal > 0.8) for these *Alpinobombus* taxa (Supplementary material 2): *B. alpinus* (I.C.= 8) excluding its main compound (hexadec-9-en-1-ol) which is shared with *B. polaris*; *B. hyperboreus* (I.C.= 34) including its main compound (octadec-11-en-1-ol); *B. polaris* (I.C.= 22) including one of its main compounds (hexadec-9-enal).

Thoracobombus

We detected 78 compounds in the CLGS of the species belonging to *Thoracobombus* subgenus (36 from *B. inexpectatus*, 52 from *B. ruderarius* and 39 from *B. sylvarum*) (Supplementary material 2). The results showed that all species were differentiated in their CLGS composition (12 compounds shared by all species inside *Thoracobombus* group). 16 compounds were shared between the parasite *B. inexpectatus* and the host *B. ruderarius*, 24 between *B. ruderarius* and *B. sylvarum* and 16 between *B. inexpectatus* and *B. sylvarum*

(Supplementary material 2). In addition to qualitative differences in minor compounds, other differences were also present in main compounds. In the *Thoracobombus* group, the main compounds of *B. inexpectatus* were hexadec-7-en-1-ol (17.23% - 36.14%), octadec-11-en-1-ol (12.25% - 32.57%) and octadec-11-enyl acetate (9.99% - 23.44%) while those of *B. ruderarius* and *B. sylvarum* were respectively hexadec-9-en-1-ol (16.49% - 70.64%) and hexadec-7-enyl acetate (48.74% - 65.34%) (Table 3).

The IndVal method revealed several indicator compounds (I.C.) with high statistical significance (IndVal > 0.8) for these *Thoracobombus* taxa (Supplementary material 2): *B. inexpectatus* (I.C.= 15) including the main compound octadec-11-enyl acetate; *B. ruderarius* (I.C.= 3) including the main compound hexadec-9-en-1-ol and *B. sylvarum* (I.C.= 9) including the main compound hexadec-7-enyl acetate.

Interpretation of Clusters

For both groups (*Alpinobombus* and *Thoracobombus*), the statistical analyses of the CLGS composition confirmed the interspecific differentiations and are supported by high values of multiscale bootstrap resampling (Fig. 2A, B).

DISCUSSION

Comparison of the Parasite CLGS Composition With Past

Identifications

Compared with Svensson and Bergström (1979), we have identified the same main compounds for *B. hyperboreus*. However, thanks to improved methods of chemical analyses, we identified many more compounds in the CLGS composition especially with double bounds information. Concerning *B. inexpectatus*, the description of CLGS composition is here reported for the first time.

Evidence for Host Evasion Strategies in Parasitic Males

Chemical Mimicry Hypothesis

In an older study investigating bumblebee taxonomic relationship based on their CLGS (Bellés *et al.* 1987), both cuckoos (*Bombus Psithyrus rupestris*) and hosts (*Bombus Melanobombus lapidarius*) appeared mixed together in the cluster analysis. According to the authors, there is a selection of the cuckoos for similarity to their hosts to improve their ability of parasitization. These data were in accordance with observations made by Cederberg (1979) showing that *B. rupestris* recognizes the specific marking secretion of its host *B. lapidarius*.

While chemical mimicry is a widespread strategy in social parasites (Dettner & Liepert 1994; Akino 2008), our CLGS composition-based comparative analyses seem to rule out the chemical mimicry hypothesis for both *B. hyperboreus* and *B. inexpectatus*. These results are convergent with previous studies on CLGS of other parasitic bumblebees (namely subgenus *Psithyrus*) that do not show similarities between inquiline and host species (Lecocq *et al.* 2011; Lecocq *et al.* 2013; Lecocq *et al.* 2015a; Lhomme *et al.* 2015).

Other previous studies have shown that CLGS play a major role the attractiveness of females during pre-nuptial behaviors (Lhomme *et al.* 2012; Ayasse & Jarau 2014). These secretions are very specific even in cryptic species (Bertsch *et al.* 2005; Brasero *et al.* 2015). Thus sharing the same recognition signals could affect the intrinsic role of CLGS in pre-nuptial behaviors. This may explain why chemical signals from these glands are maintained as distinct from their host species.

The interspecific CLGS comparisons (Rasmont *et al.* 2005; Lecocq *et al.* 2015a; Brasero *et al.* 2015) show higher composition similarities between phylogenetically closely related bumblebee species. For *Alpinobombus* group, the cluster analysis indicates that the CLGS composition of the hosts (*B. polaris*) is more similar to its sister-species (*B. alpinus*) than to their respective parasite (*B. hyperboreus*), which is a sister lineages to *B. alpinus* and *B. polaris* (Fig. 2A). In the *Thoracobombus* group, the composition of *B. inexpectatus* is

more similar to the reference species (*B. sylvarum*) than its own host (*B. ruderarius*) (Fig. 2B). Hines and Cameron (2010) showed that *B. inexpectatus* is phylogenetically closer to *B. sylvarum* than its host *B. ruderarius*.

Chemical Insignificance Hypothesis

Our results showed that mature male parasites emit a specific mixture of compounds from their cephalic labial glands so they are not using an absence of species-specific chemicals to avoid host detection (i.e., chemical insignificance).

Chemical Repellency Hypothesis

Since our results suggest that chemical insignificance and chemical mimicry strategies are unlikely in *B. hyperboreus* and *B. inexpectatus*, an alternative hypothesis seems to be the production of one or more repellent substances within CLGS by males. However, the repellent compounds identified by Lhomme *et al.* (2015) and Zimma *et al.* (2003) in other cuckoo bumblebees (dodecyl acetate, tetradecyl acetate and (Z)-11-hexadecenyl acetate) have not been found in the CLGS extraction of *B. hyperboreus* and *B. inexpectatus*. Other compounds in these glands or other glands that produce species-specific cuticular hydrocarbons could instead have a repellent effect.

Limitations and Future Prospects

Since we sampled only extra-nest males, we cannot exclude potential mimicry or chemical insignificance strategies during the early age of parasite males. However, we can underline that previous study showed that only the quantitative composition of the CLGS varies between intra-nest and mature emerged males (Žáček *et al.* 2009). Detailed assessment of these hypotheses requires *in natura* observations of nest infestations or bioassays on laboratory-reared specimens that will be challenging to obtain: wild nest infestation observations are uncommon (Lecocq *et al.* 2011) and species-specific year-round breeding

methods needed for bioassays (Lhomme *et al.* 2012; Lhomme *et al.* 2013) are available for only few species (Hasselrot 1960). Moreover, in recent decades, *B. hyperboreus* and *B. inexpectatus* have become scarce especially in Europe (Rasmont *et al.* 2015) increasing the difficulty of developing such a protocol with a suitable sampling. Indeed, as far as we know, just few people succeeded to breed *B. polaris* and *B. ruderarius* (Sladen 1912; Wojtowski 1963; Berezin 1995). Moreover, from a total of 988.187 data recorded in Rasmont *et al.* 2015, only 370 specimens of *B. hyperboreus* (0.03%) and 275 specimens of *B. inexpectatus* (0.02%) were recorded including them as the rarest West-Palearctic bumblebees.

In bumblebees, the chemical cue is not only due to the CLGS. Cuticular compounds should also play an important role in the recognition systems between host and parasite (Martin *et al.* 2010). Therefore, it would be interesting to investigate the composition of this cuticular signature to improve our knowledge on the intrusion and escape mechanisms of these non-*Psithyrus* parasites.

To test these hypotheses, neuro-physiological experiments could be useful (i.e. gas chromatography coupled with Electro-Antennographic Detector (Lhomme *et al.* 2015) or a glomerular function test (Devaud *et al.* 2015)). Unfortunately, the lack and the difficulty to breed these specimens make experiments simply impossible to realize for now.

CONCLUSIONS

In this study we find that the independently evolving social parasites *B. hyperboreus* and *B. inexpectatus* display a species-specific mating chemistry in their cephalic labial glands typical of bumblebees. Although these mating secretions could make them vulnerable to detection and exclusion from the host nest, our results do not support shifts in chemistry expected from major hypothesized host evasion strategies. These species are not devoid of sexual cephalic labial attractive compounds so are not chemically undetectable. These CLGS are also not more similar to their hosts than would be expected phylogenetically so are also

not involved in chemical mimicry. Finally, we have not identified the repellent compounds highlighted by Lhomme *et al.* (2015) in *B. vestalis*. Other repellent compounds in these or other glands could, however, be important for host evasion.

Acknowledgements

We acknowledge all people that helped us in our journey to the Abisko, Tarfala and Toolik stations: Tchad Singer, Mickael Abels, Peter Lakare and Gunhild Ninis Rosqvist (University of Stockholm), Magnus Augner and Linnea Wanhatalo (Abisko Station), Hannele Savela (Oulu University, Interact administration), Josefine Strand and Therese Wikstrom (Lansstyrelsen i Norrbottens län Naturvårdsenheten, Luleå). Financial support (IV and KU) was provided by the Institute of Organic Chemistry and Biochemistry of the Academy of Sciences of the Czech Republic (subvention for development of research organization RVO: 61388963). The research has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement no 244090, STEP Project (Status and Trends of European Pollinators, www.step-project.net) and from the INTERACT FP-7 EU project. Special thanks to Gilles Mahé (Nantes, France) for specimen of *B. inexpectatus*. PB contributes as PhD student funded by Czech Science Foundation (GAČR GP14-10035P). BM contributes as PhD student granted by Research Council of University of Mons and from the FRS-FNRS (Fonds de la Recherche Scientifique).

336

DISCLOSURE

337 The authors declare that they have no conflict of interest.

338

REFERENCES

339 Akino T. (2008) Chemical strategies to deal with ants: a review of mimicry, camouflage,
340 propaganda, and phytomimesis bu ants (Hymenoptera: Formicidae) and other arthropods.
341 Myrmecological News, 11, 173–181.

342 Amiet F. (1996) Hymenoptera Apidae, 1. Teil: Allgemeiner Teil, Gattungsschlüssel, die
343 Gattungen Apis, Bombus und Psithyrus. Insecta Helv Fauna, 12, 1–98.

344 Ayasse M. and Jarau S. (2014) Chemical ecology of bumble bees. Annu Rev Entomol, 59,
345 299–319, DOI: 10.1146/annurev-ento-011613-161949.

346 Ayasse M., Paxton R.J. and Tengö J. (2001) Mating behavior and chemical communication in
347 the order Hymenoptera. Annu Rev Entomol, 46, 31–78.

348 Baer B. (2003) Bumblebees as model organisms to study male sexual selection in social
349 insects. Behav Ecol Sociobiol, 54, 521–533.

350 Bellés X., Galofré A. and Ginebreda A. (1987) Taxonomic potential of the chemical
351 constituents in the cephalic marking secretions of Bombus and Psithyrus species
352 (Hymenoptera, Apidae): a numerical taxonomic study. Apidologie, 18, 231–242.

353 Benton T. (2006) Bumblebees, Harper Col. London. pp. 580.

354 Berezin M.V. (1995) Geographical diversity, species correlation, population structure and
355 cenotic interactions of Arctic bumble bees (Apidae, Bombus). In: Swedish-Russian Tundra

356 Ecology-expedition-94. Tundra Ecology-94. A Cruise Report. Swedish Polar Research
 357 Secretariat, Stockholm, pp. 205–215.

358 Bergström G., Svensson B.G., Appelgren M. and Groth I. (1981) Complexity of bumble bee
 359 marking pheromones: biochemical, ecological and systematical interpretations. Syst Assoc,
 360 special volume, 175–183.

361 Bergwall H.E. (1970) Ekologiska iakttagelser över några humlearter (*Bombus* Latr.) vid
 362 Staloluokta inom Padjelanta nationalpark, Lule lappmark. Entomol Tidskr, 91, 3–23.

363 Bertsch A., Schweer H., Titze A. and Tanaka H. (2005) Male labial gland secretions and
 364 mitochondrial DNA markers support species status of *Bombus cryptarum* and *B. magnus*
 365 (Hymenoptera, Apidae). Insectes Soc, 52, 45–54. DOI: 10.1007/s00040-004-0761-1.

366 Brasero N., Martinet B., Urbanová K., Valterová K., Torres A., Hoffmann W., Rasmont P.
 367 and Lecocq T. (2015) First Chemical Analysis and Characterization of the Male Species-
 368 Specific Cephalic Labial-Gland Secretions of South American Bumblebees. Chem Biodivers,
 369 12, 1535–1546. DOI: 10.1002/cbdv.201400375.

370 Calam D.H. (1969) Species and Sex-specific Compounds from the Heads of Male
 371 Bumblebees (*Bombus* spp.). Nature, 221, 856–857.

372 Cameron S.A., Hines H.M. and Williams P.H. (2007) A comprehensive phylogeny of the
 373 bumble bees (*Bombus*). Biol J Linn Soc, 91, 161–188.

374 Cederberg B. (1979) Odour guided host selection in *Psithyrus* (Hym., Apidae). Entomol
 375 Tidskr, 100, 128–129.

376 Cvacka J., Kofronová E., Vasícková S., et al. (2008) Unusual fatty acids in the fat body of the
 377 early nesting bumblebee, *Bombus pratorum*. *Lipids*, 43, 441–50. DOI: 10.1007/s11745-008-
 378 3174-5.

379 De Meulemeester T, Gerbaux P, Boulvin M, Coppée A. and Rasmont P. (2011) A simplified
 380 protocol for bumble bee species identification by cephalic secretion analysis. *Insectes Soc*, 58,
 381 227–236.

382 Dellicour S. and Lecocq T. (2013a) GCALIGNER 1.0: an alignment program to compute a
 383 multiple sample comparison data matrix from large eco-chemical datasets obtained by GC. *J*
 384 *Sep Sci*, 36, 3206–3209. DOI: 10.1002/jssc.201300388.

385 Dellicour S. and Lecocq T. (2013b) GCALIGNER 1.0 and GCKOVATS 1.0 - Manual of a
 386 software suite to compute a multiple sample comparison data matrix from eco-chemical
 387 datasets obtained by gas chromatography. University of Mons, Mons (Belgium).

388 Dettner K. and Liepert C. (1994) Chemical mimicry and camouflage. *Annu Rev Entomol*, 39,
 389 129–154.

390 Devaud J.M., Papouin T., Carcaud J., Sandoz J.C., Grünwald B. and Giurfa M. (2015)
 391 Neural substrate for higher-order learning in an insect: Mushroom bodies are necessary for
 392 configural discriminations. *Proc Natl Acad Sci, U S A*. pp 9. DOI: 10.1073/pnas.1508422112.

393 Dronnet S., Simon X., Verhaeghe J.C., Rasmont P. and Errard C. (2005) Bumblebee
 394 inquiline in *Bombus (Fernaldaepsithyrus) sylvestris* (Hymenoptera, Apidae): behavioural
 395 and chemical analyses of host-parasite interactions. *Apidologie*, 36, 59–70.

396 Dufrêne M. and Legendre P. (1997) Species assemblages and indicator species: The need for
 397 a flexible asymmetrical approach. *Ecol Monogr*, 67, 345–366.

398 Fisher R. and Sampson B. (1992) Morphological specializations of the bumble bee social
 399 parasite *Psithyrus ashtoni* (Hymenoptera, Apidae). *Canadian Entomol*, 124, 69-77.

400 Gjershaug J.O. (2009) The social parasite bumblebee *Bombus hyperboreus* Schönherr, 1809
 401 usurp nest of *Bombus balteatus* Dahlbom 1832 (Hymenoptera: Apidae) in Norway. *Nor J*
 402 *Entomol*, 56, 28–31.

403 Hasselrot (1960) Studies on Swedish bumblebees (genus *Bombus* Latr.), their domestication
 404 and biology. *Opusc Entomol Suppl*, 17,1–192.

405 Hines H.M., Cameron S.A. (2010) The phylogenetic position of the bumble bee inquiline
 406 *Bombus inexpectatus* and implications for the evolution of social parasitism. *Insectes Soc*,
 407 57, 379–383.

408 Hobbs R.M. (1967) Ecology of species of *Bombus* in southern Alberta. VI. Subgenus
 409 *Pyrobombus*. *Can Entomol*, 99,1271–1292.

410 Lecocq T., Brasero N., De Meulemeester T. et al (2015a) An integrative taxonomic approach
 411 to assess the status of Corsican bumblebees: implications for conservation. *Anim Conserv*, 18,
 412 236–148.

413 Lecocq T., Coppée A., Mathy T. et al (2015b) Subspecific differentiation in male
 414 reproductive traits and virgin queen preferences, in *Bombus terrestris*. *Apidologie*, 46,595–
 415 605. DOI: 10.1007/s13592-015-0349-y.

416 Lecocq T., Lhomme P., Michez D., Dellicour S., Valterová I. and Rasmont P. (2011)
 417 Molecular and chemical characters to evaluate species status of two cuckoo bumblebees:
 418 *Bombus barbutellus* and *Bombus maxillosus* (Hymenoptera, Apidae, Bombini). *Syst*
 419 *Entomol*, 36, 453–469. DOI: 10.1111/j.1365-3113.2011.00576.x.

420 Lecocq T., Vereecken N.J., Michez D. et al (2013) Patterns of genetic and reproductive traits
 421 differentiation in Mainland vs. Corsican populations of bumblebees. PLoS One, 8:e65642.
 422 DOI: 10.1371/journal.pone.0065642.

423 Legendre P. and Legendre L. (1983) Numerical Ecology. Developments in Environmental
 424 Modelling, 3. Elsevier Scient. Publ. Co., Amsterdam. pp. 419.

425 Lenoir A., D'Ettore P. and Errard C. (2001) Chemical Ecology and Social Parasitism in Ants.
 426 Annu Rev Entomol, 46, 573–599.

427 Lhomme P., Ayasse M., Valterová I., Lecocq T. and Rasmont P. (2012) Born in an alien nest:
 428 how do social parasite male offspring escape from host aggression? PLoS One, 7:e43053.
 429 DOI: 10.1371/journal.pone.0043053.

430 Lhomme P, Ayasse M, Valterová I, Lecocq T. and Rasmont P. (2015) A scent shield to
 431 survive: identification of the repellent compounds secreted by the male offspring of the
 432 cuckoo bumblebee *Bombus vestalis*. Entomol Exp Appl, 157, 263–270. DOI:
 433 10.1111/eea.12366.

434 Lhomme P, Sramkova A, Kreuter K, Lecocq T., Rasmont P. and Ayasse M. (2013) A method
 435 for year-round rearing of cuckoo bumblebees (*Psithyrus*). Ann la Société Entomol Fr, 49,
 436 117–125.

437 Løken A. (1973) Studies on Scandinavian bumble bees (Hymenoptera, Apidae). Nor Entomol
 438 Foren, 20, pp. 218.

439 Luxová A., Valterová I, Stránský K, Hovorka O. and Svatoš A. (2003) Biosynthetic studies
 440 on marking pheromones of bumblebee males. Chemoecology, 13, 81–87.

441 Martin S.J., Carruthers J.M., Williams P.H. and Drijfhout F.P. (2010) Host specific social
 442 parasites (*Psithyrus*) indicate chemical recognition system in bumblebees. *J Chem Ecol*, 36,
 443 855–863.

444 Michener C.D. (1974) The social behavior of the bees, a comparative study. Harvard
 445 University Press, Cambridge. pp. 408.

446 Milliron H.E. and Oliver D.R. (1966) Bumblebees From Northern Ellesmere Island, With
 447 Observations on Usurpation by *Megabombus hyperboreus* (Schönh.) (Hymenoptera: Apidae).
 448 *Can Entomol*, 98, 207–213.

449 Milliron H.E. (1971) A monograph of the western hemisphere bumblebees (Hymenoptera:
 450 Apidae; Bombinae). I. The genera *Bombus* and *Megabombus* subgenus *Bombias*. *Mem*
 451 *Entomol Soc Canada*, 82, 1–80.

452 Müller A. (2006) A scientific note on *Bombus inexpectatus* (Tkalcu, 1963): evidence for a
 453 social parasitic mode of life. *Apidologie*, 37, 408–409.

454 Nash D.R. and Boomsma J.J. (2008) Communication between hosts and social parasites. In:
 455 *Sociobiology of communication: an interdisciplinary perspective*. Hughes DP: Oxford
 456 University Press. pp. 55–79.

457 Pape T. (1983) Observations on nests of *Bombus polaris* Curtis usurped by *B. hyperboreus*
 458 Schönherr in Greenland. *Entomol Meddelelser*, 50, 145–150.

459 Paradis E., Claude J. and Strimmer K. (2004) APE: Analyses of phylogenetics and evolution
 460 in R language. *Bioinformatics*, 20, 289–290.

461 R Development Core Team (2013) R: A language and environment for statistical computing.

462 Rampini M., Russo C.D., Taylan M.S., Gelosa A. and Cobolli M. (2012) Four new species of
 463 Dolichopoda Bolivar, 1880 from Southern Sporades and Western Turkey (Orthoptera,
 464 Rhaphidophoridae, Dolichopodinae). Zookeys, 201, 43–58. DOI:
 465 10.3897/zookeys.201.2609.

466 Rasmont P., Franzen M., Lecocq T. et al. (2015) Climatic Risk and Distribution Atlas of
 467 European Bumblebees. BioRisk, 10, 1–236. DOI: 10.3897/biorisk.10.4749.

468 Rasmont P. and Iserbyt S. (2014) Atlas of the European Bees: genus *Bombus*. 3d Edition.
 469 STEP Project. In: Status Trends Eur. Pollinators, Atlas Hymenoptera, Mons, Gembloux.

470 Rasmont P., Terzo M, Aytekin AM et al. (2005) Cephalic secretions of the bumblebee
 471 subgenus *Sibiricobombus* Vogt suggest *Bombus niveatus* Kriechbaumer and *Bombus*
 472 *vorticatus* Gerstaecker are conspecific (Hymenoptera, Apidae, *Bombus*). Apidologie, 36,
 473 571–584. DOI: 10.1051/apido:2005047.

474 Richards K.W. (1973) Biology of *Bombus polaris* Curtis and *B. hyperboreus* Schonherr at
 475 lake Hazen, Northwest territories (Hymenoptera: Bombini). Quaest Entomol, 9, 115–157.

476 Roldan M. and Soler M. (2011) Parental-care parasitism: how do unrelated offspring attain
 477 acceptance by foster parents? Behav Ecol, 22, 679–691.

478 Sladen F.W.L. (1912) The Humble-bee. Its life-history and how to domesticate it with
 479 descriptions of all the British species of *Bombus* and *Psithyrus*, Macmillan . London; pp.
 480 1876-1921.

481 Smisteh T., Kölliker M. and Royle N.J. (2012) What is parental care? In: J. RN (ed) The
 482 Evolution of Parental Care. Oxford University Press, Oxford, pp. 355.

483 Sramkova A. and Ayasse M. (2008) Psithyrus females do possess wax glands. *Insectes Soc*,
484 55, 404–406. DOI: 10.1007/s00040-008-1020-7.

485 Stenström M. and Bergmann P. (1998) Bumblebees at an alpine site in northern Sweden:
486 temporal development, population size, and plant utilization. *Ecography*, 21, 306–316.

487 Suzuki R. and Shimodaira H. (2011) Pvcust: Hierarchical Clustering with P-Values via
488 Multiscale Bootstrap Resampling

489 Svesson B.O.G. and Bergström G. (1979) Marking pheromones of *Alpinobombus* males. *J*
490 *Chem Ecol*, 5, 603–615.

491 Terzo M., Urbanová K., Valterová I. and Rasmont P. (2005) Intra and interspecific variability
492 of the cephalic labial glands' secretions in male bumblebees: the case of *Bombus*
493 (*Thoracobombus*) *runderarius* and *B. (Thoracobombus) sylvarum* Hymenoptera, Apidae.
494 *Apidologie*, 36, 85–96.

495 Tkalců B. (1963) Eine neue Hummel-Art der Gattung *Agrobombus* Vogt aus dem
496 Alpengebiet. *Cas Ces Spol Entomol*, 60, 183–196.

497 Tkalců B. (1965) Über *Agrobombus inexpectatus* Tkalců. *Reichenbachia*, 5, 225–230.

498 Williams P.H., Byvaltsev A.M., Cederberg B. et al. (2015) Genes Suggest Ancerstral Colour
499 Polymorphisms Are Shared across Morphologically Cryptic Species in Artic Bumblebees.
500 *PLOS ONE*, 10(12), e0144544. DOI:10.1371/journal.pone.0144544.

501 Williams P.H. (1998) An annotated checklist of bumble bees with an analysis of patterns of
502 description (Hymenoptera: Apidae, Bombini). *Bull Nat Hist Museum*, 67, 79–152.

503 Wilson E.O. (1971) *The insect societies*. Harvard University Press, Cambridge. pp. 548.

504 Wojtowski F. (1963) Observations on the biology and reproduction of bumblebees
 505 (Bombinae). Zoologica Poloniae, 13, 137-152.

506 Yarrow I.H.H. (1970) Is *Bombus inexpectatus* (Tkalcu) a workless parasite? Insectes Soc, 17,
 507 95–112.

508 Žáček P., Kalinová B., Šobotník J. et al. (2009) Comparison of Age-dependent Quantitative
 509 Changes in the Male Labial Gland Secretion of *Bombus terrestris* and *Bombus lucorum*. J
 510 Chem Ecol, 35, 698–705.

511 Žáček P., Prchalová-Hornáková D., Tykva R. et al. (2013) De Novo Biosynthesis of Sexual
 512 Pheromone in the Labial Gland of Bumblebee Males. ChemBioChem, 14, 361–371.

513 Zimma B.O., Ayasse M., Tengö J., Ibarra F., Schulz C. and Francke W. (2003) Do social
 514 parasitic bumblebees use chemical weapons? (Hymenoptera, Apidae). J Comp Physiol A
 515 Neuroethol Sensory, Neural, Behav Physiol, 189, 769–775.

516 TABLES

517 *Table 1. Sampling sites and number (N) of specimens collected for B. hyperboreus , B.*
 518 *polaris, B. alpinus, B. inexpectatus, B. ruderarius, and B. sylvarum. All sampling*
 519 *information in supplementary material 1.*

Taxa	Collecting sites	Coordinates	N	Code
<i>Alpinobombus</i>				
<i>B. hyperboreus</i>	Sweden, Bjorklieden	68°24'22"N 18°40'26"E	1	BMAR054
	Sweden, Tarfala	67°53'55"N 18°37'57"E	1	PRAS0951
	Sweden, Tarfala	67°53'55"N 18°37'57"E	1	PRAS0954
	Sweden, Tarfala	67°54'42"N 18°36'36"E	1	PRAS0948
	Sweden, Tarfala	67°54'15"N 18°37'15"E	1	PRAS0952
	Norway, Narvik	68°06'23"N 17°29'20"E	2	PRAS0930 PRAS0931

	USA, Alaska, Galbraith lake	68°25'31"N 149°21'32"W	1	PRAS1067
	USA, Alaska, Galbraith lake	68°27'21"N 149°28'56"W	2	PRAS1062 BMAR0242
	USA, Alaska, Toolik lake	68°37'37"N 149°35'46"W	1	PRAS1078
	USA, Alaska, Toolik lake	68°38'22"N 149°34'35"W	2	BMAR0304 BMAR0305
	USA, Alaska, Toolik lake	68°37'37"N 149°35'46"W	1	BMAR0356
	USA, Alaska, Sag River	68°46'46"N 148°51'19"W	1	BMAR0377
<i>B. polaris</i>	USA, Alaska, Toolik Field station	68°37'40"N 149°35'07"W	10	BMAR0191 BMAR0157 BMAR0145 BMAR0190 BMAR0192 BMAR0195 BMAR0201 BMAR0206 BMAR0207 BMAR0221
	Norway, Narvik	68°06'23"N 17°29'20"E	1	PRAS0929
	Norway, Narvik	68°04'55"N 17°27'02"E	1	PRAS0936
	Sweden, Tarfala	67°54'42"N 18°36'36"E	1	PRAS0949
<i>B. alpinus</i>	Norway, Narvik	68°06'57"N 17°33'13"E	7	PRAS0939 PRAS0937 PRAS0938 PRAS0903 PRAS0904 PRAS0905 PRAS0906
	Norway, Narvik	68°09'06"N 17°29'45"E	1	PRAS0900
	Norway, Narvik	68°07'48"N 17°29'37"E	1	PRAS0935
	Norway, Narvik	68°07'59"N 17°28'52"E	1	PRAS0908
	Sweden, Tornehamn	68°26'28"N 18°37'45"E	1	PRAS0806
	Sweden, Kiruna	68°13'15"N 19°42'46"E	1	PRAS0885
	Sweden, Tarfala	67°54'42"N 18°36'36"E	2	PRAS0946 PRAS0947
<i>Thoracobombus</i>				
<i>B. inexpectatus</i>	Italy, Val Brembana	45°58'12"N 09°47'53"E	3	NBRA0918 NBRA0919 NBRA0920
	Italy, Val Brembana	45°58'13"N 09°47'58"E	1	NBRA0915
	Italy, Val Brembana	45°58'12"N 09°47'57"E	2	NBRA0916

	Italy, Val Brembana	45°58'11"N 09°47'52"E	3	NBRA0921 NBRA0922 NBRA0923
<i>B. ruderarius</i>	Switzerland, Valais, Motôt	46°05'21"N 07°24'13"E	4	NBRA0321 NBRA0322 NBRA0323 NBRA0324
	Switzerland, Valais, Mâssmatte	46°28'37.5"N 08°23'14.6"E	1	NBRA0365
	France, Oriental Pyrenees, Saillagouse	42°28'N 02°02'E	2	MT0015 MT0017
	France, Oriental Pyrenees, Err	42°25'N 02°03'E	1	MT0035
<i>B. sylvarum</i>	France, Oriental Pyrenees, Llo	42°27'N 02°04'E	1	MT0140
	France, Oriental Pyrenees, Dorres	42°28'N 01°45'E	3	MT0229 MT0231 MT0244
	France, Oriental Pyrenees, Eyne	42°29'N 02°05'E	4	MT0253 MT0258 MT0267 MT0277
	France, Lozère, Les Salces	44°33'N 03°07'E	1	MT0330
	France, Hérault, Romiguières	43°48'N 03°14'E	1	MT0346

520

521 Table 2. List of main compounds (**Bold**) identified in cephalic labial gland secretions of
522 *Bombus hyperboreus*, *B. polaris*, and *B. alpinus*. MW: the molecular weight, M: median of
523 relative concentration of compound (%). All identified compounds in Supplementary material
524 2.

Compounds	MW	<i>B. alpinus</i>			<i>B. polaris</i>			<i>B. hyperboreus</i>		
		Min	M	Max	Min	M	Max	Min	M	Max
Hexadec-9-enal	238	0.00	0.07	1.27	20.80	28.18	43.03	0.00	0.00	0.00
Hexadec-9-en-1-ol	240	63.95	83.16	89.43	26.87	36.33	46.68	0.00	0.00	0.00
Octadec-11-en-1-ol	268	0.00	0.03	0.19	0.00	0.00	0.00	23.90	38.39	56.21

525

526 Table 3. List of main compounds (**Bold**) identified in cephalic labial gland secretions of
 527 *Bombus inexpectatus*, *B. ruderarius*, and *B. sylvarum*. MW: the molecular weight; M:
 528 median of relative concentration of compound (%). All identified compounds in
 529 Supplementary material 2.

Compounds	MW	<i>B. inexpectatus</i>			<i>B. ruderarius</i>			<i>B. sylvarum</i>		
		Min	M	Max	Min	M	Max	Min	M	Max
Hexadec-7-en-1-ol	240	17.23	24.99	36.14	0.00	0.00	0.00	23.54	28.65	32.14
Hexadec-9-en-1-ol	240	4.94	7.47	8.32	16.49	37.26	70.64	0.00	0.00	0.00
Octadec-11-en-1-ol	268	12.25	17.66	32.57	3.97	7.57	11.25	0.00	0.00	0.00
Hexadec-7-enyl acetate	282	3.27	5.35	11.97	0.00	0.00	0.00	48.74	56.52	65.34

530

531

532

533

FIGURES

534

535 **Fig. 1:** Photos of the studied bumblebee species: A) *Bombus hyperboreus* male (PRAS1067)
536 from Alaska, USA (photo: P. Rasmont), B) *B. polaris* male (PRAS0936), from Norway (photo:
537 P. Rasmont), C) *B. inexpectatus* male from Alps, Italy (photo: P. Rasmont), D) *B. ruderarius*
538 male from Pyrenees, France (photo: P. Rasmont).

539

Fig. 2 Unweighted pair group method with arithmetic mean (UPGMA) cluster based on a correlation matrix calculated from the composition of cephalic labial gland secretions of *Bombus hyperboreus*, *B. polaris*, *B. alpinus* (A); *B. inexpectatus*, *B. ruderarius* and *B. sylvarum* (B). The values near nodes represent multiscale bootstrap resampling values (only values >90 are given). It= Italy; Fr= France and Ch= Switzerland.