

Sensitive response of sediment-grown Myriophyllum spicatum L. to arsenic pollution under different CO2 availability

Elisabeth M. Gross, Andréïna Nuttens, Paroshin Darya, Andreas Hussner

► To cite this version:

Elisabeth M. Gross, Andréina Nuttens, Paroshin Darya, Andreas Hussner. Sensitive response of sediment-grown Myriophyllum spicatum L. to arsenic pollution under different CO2 availability. Hydrobiologia, 2018, 812 (1), pp.177-191. 10.1007/s10750-016-2956-7. hal-02088981

HAL Id: hal-02088981 https://hal.univ-lorraine.fr/hal-02088981

Submitted on 30 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 2 3 4	Published in Hydrobiologia – DOI 10.1007/s10750-016-2956-7
5	Sensitive response of sediment-grown Myriophyllum spicatum L. to
6	arsenic pollution under different CO ₂ availability
7	
8	
9	Elisabeth M. Gross ^{1,2,*} , Andréïna Nuttens ^{1,2} , Darya Paroshin ^{1,2} & Andreas Hussner ³
10	
11	¹ Université de Lorraine, LIEC UMR 7360 CNRS, Rue Général Delestraint, Bâtiment IBISE,
12	F-57070 Metz, France.
13	² LTER-"Zone Atelier Moselle", LIEC UMR 7360 CNRS, Rue Général Delestraint, Bâtiment
14	IBISE, F-57070 Metz, France.
15	³ Institute of Botany, Heinrich-Heine-Universität Düsseldorf, Geb. 26.13.02, Universitätsstr.
16	1, D-40225 Düsseldorf, Germany
17	* Corresponding author: <u>gross5@univ-lorraine.fr</u> ; Tel.: +33 3 87 37 86 26; Fax +33 3 87 37
18	85 12

19 Abstract

Aquatic plants may face resource constraints or anthropogenic pollution, and effects might be 20 heightened under multiple stress conditions. We investigated if arsenate effects on 21 22 Myriophyllum spicatum L. would be stronger under CO₂ limitation and low phosphorus availability. In a factorial design, we exposed sediment-grown plants to either CO₂ (high 23 carbon or HC) or bicarbonate (low carbon or LC) and four levels of arsenate. We observed 24 25 strong effects of arsenate exposure on growth, biomass allocation (leaf, stem and root mass fractions), pigments and phenolic compounds. CO₂ availability strongly affected the content 26 in phenolic compounds and a few other response variables, yet overall effects were less 27 28 pronounced than expected. Strong interactive effects of CO₂ availability and arsenic concentration were only observed for carotenoids, the carotenoid/chlorophyll ratio and 29 phenolic compounds in leaves. Only the carbon content declined with increasing arsenic 30 concentration, otherwise leaf elemental content and stoichiometry were not affected by 31 arsenic or CO₂ availability, suggesting that plants strived to maintain leaf functions. The 32 33 observed effects on biomass allocation and plant quality, specifically elemental stoichiometry and phenolic compounds content of M. spicatum not only show direct changes in plant 34 performance but suggest also indirect effects on ecological interactions such as competition or 35 36 herbivory.

37

Keywords: Submerged macrophyte, Trace metal, Element pollution, OECD test guideline,
aquatic ecotoxicology, CO₂-based growth and physiology

41 Introduction

42

facing effects at the global, regional and local scale, strongly enhanced by anthropogenic 43 44 activities (Meybeck 2004; Segner et al. 2014). Since the response to a combination of stressors might be specific for a certain type or group of organisms (Segner et al. 2014), we 45 need to understand the reaction of key organisms, such as those used in environmental risk 46 47 assessment. The rooted submerged macrophyte M. spicatum is now used as OECD (Organisation for Economic Co-operation and Development) test organism because it allows 48 testing for both sediment and water exposure to anthropogenic pollutants (OECD 2014). 49 While normalized test systems in ecotoxicology strive for optimal growth (see details 50 on the OECD TG 239 for sediment grown *M. spicatum* below in materials and methods), 51 52 resource availability are known to affect plant growth and physiology (Hussner & Jahns 2015), and might thus also be involved in the response to pollution stress. Submerged 53 54 macrophytes, such as *M. spicatum*, can take up nutrients via roots or shoots, but root uptake is 55 the primary pathway (Grace & Wetzel 1978; Smith & Barko 1990). While excess or limiting nutrient amounts can be a constraint for submerged macrophytes, the availability of CO₂ can 56 even be more restraining, as CO₂ diffusion is much slower in stagnant water, and the 57 availability depends on the pH and total dissolved inorganic carbon (DIC) availability 58 (Hussner & Jahns 2015). Although the growth of *M. spicatum* is enhanced in alkaline lakes 59 (Smith & Barko 1990), CO₂ compensation points are lower than those for HCO₃⁻ (Grace & 60 Wetzel 1978). 61

Multiple stress scenarios are gaining focus both in ecology and ecotoxicology as we are

The genus *Myriophyllum* is known to accumulate trace metal elements, but this does
not exclude negative effects on plant growth and physiology (e.g., Harguinteguy et al. 2015;
Robinson et al. 2006). Arsenic is a trace metal element often found in aquatic systems

resulting from geochemical or anthropogenic processes. The New Zealand M. propinguum in 65 66 the Taupo Volcanic area hyperaccumulated 2101 mg [As] per kg dry mass (Robinson et al. 2006), and the North American M. sibiricum Kom. (syn. M. exalbescence Fernald) from a 67 lake polluted by mine tailings accumulated more than 140 mg kg⁻¹ [As] in its dry mass 68 (Dushenko et al. 1995). In freshwaters, arsenic occurs mainly as arsenate under oxygenic or 69 arsenite under anoxic conditions (Rahman et al. 2012), and the adsorption of arsenic to 70 71 sediments is influenced by metal oxides but also by carbonate and bicarbonate concentrations (Smedley & Kinniburgh 2002). 72

Various environmental factors influence the uptake of arsenic by plants. Plants take up 73 74 arsenate via the phosphate uptake transporter (Finnegan & Chen 2012). In laboratory experiments, high phosphate availability reduced the arsenate uptake by *Lemna minor* 75 (Mkandawire & Dudel 2012) and rice (Geng et al. 2005). However, no relationship between 76 in-situ phosphate availabiliy and arsenic uptake was observed for a range of submerged and 77 emergent macrophytes in Canadian lakes (Dushenko et al. 1995). Under alkaline pH, the fern 78 79 Pterris vittata contained less arsenic than under neutral pH, but bicarbonate concentration in natural ranges (up to 2 mM) showed little effect on arsenic uptake by the plant (Guo et al. 80 2012). Yet, eutrophic conditions reduced the arsenate uptake by the congeneric M. 81 82 alterniflorum, a species adapted to oligo-mesotrophic conditions (Krayem et al. 2016). The effect of alkalinity, and thus pH and CO₂ availability, on the uptake of arsenic has, as far as 83 we know, not yet been tested with submerged freshwater plants. In this context we performed 84 our study, knowing that *M. spicatum* prefers meso- to eutrophic and alkaline environments. 85

Environmental stressors and resource constraints can have a wide range of effects in plants. They may not only affect growth, but also morphological traits such as biomass allocation to leaves, shoots and roots or leaf dry matter content (LDMC), changes that may explain plant performance at the inter- and intraspecific level (Elger & Willby 2003; Fornoff

& Gross 2014; Hussner & Jahns 2015; Pérez-Harguindeguy et al. 2013) or physiological 90 91 traits, such as pigment composition, changes in elemental stoichiometry (i.e. carbon, nitrogen and phosphorus content), and the production of defensive substances such as phenolic 92 93 compounds (Fornoff & Gross 2014; Gross 2003; Hussner et al. 2015; Hussner & Jahns 2015). Phenolic compounds are widely involved in plant responses to abiotic and biotic stressors 94 (Dixon & Pavia 1995), and *M. spicatum* is a species well known for its high content in 95 bioactive polyphenols acting against competitors and herbivores (Fornoff & Gross 2014; Leu 96 et al. 2002) but also varying depending on resource availability (Gross 2003). High CO₂ 97 availability increased the production of phenolic compounds in terrestrial plants (Hartley et al. 98 99 2000), and trace metal elements affected the content of phenolic compounds (Lavid et al. 2001), possibly because of oxidative stress responses (Waśkiewicz et al. 2014). 100

Since the availability of CO₂ is a major constraint for submerged macrophytes, and 101 because of the potential impact of pH, bicarbonate and nutrient availability on the effect of 102 arsenic on aquatic plants, we used a factorial design experiment to test the effects of arsenate 103 104 under different CO₂ availability and at low phosphorus availability on *M. spicatum*. Our hypotheses were that 1) environmentally relevant arsenic concentrations would negatively 105 affect growth and physiological response variables (i.e. "endpoints" sensu ecotoxicological 106 test protocols, e.g., OECD 2014) in M. spicatum, 2) effects would be enhanced at high pH 107 reflecting low CO₂ availability, and 3) changes in morphological and physiological plant traits 108 of *M. spicatum* reflect specific responses towards a certain combination of stressors. We 109 included growth, pigment content, plant stoichiometry, and the content in phenolic 110 compounds as response variables to see if these "endpoints" are useful for predicting potential 111 112 ecological and ecosystem effects caused by multiple environmental stressors.

114 Material and Methods

115 Cultivation of plants and experimental set-up

We performed our experiment in a plant-water-sediment test system modified after the 116 currently accepted new OECD test guideline for the rooted aquatic plant M. spicatum (OECD 117 2014). This test system uses 2 L glass beakers, a sediment mixed from sand, kaolin, white 118 peat and calcium carbonate (74/20/5/1% on a weight basis), amended with rather high 119 120 concentrations of nutrients (each 200 mg of ammonium chloride and sodium phosphate per kg of dry sediment), and a water column based on the so called Smart & Barko medium (Smart 121 & Barko 1985), a mineral solution rich in calcium chloride, magnesium sulphate, potassium-122 and sodium bicarbonate, but free of nitrogen and phosphorus. In a typical set-up, three shoots 123 of *M. spicatum* per beaker grow without any aeration for 14 days. For our purposes, we 124 modified container size, shoot density, nutrient provision and CO₂ availability. 125

Approximately 4 cm long apical shoot sections of *M. spicatum* from axenic cultures 126 maintained in the laboratory of EMG (Gross 2003) were adapted during two weeks to Smart 127 & Barko medium (Smart & Barko 1985) supplemented initially with 143 µM N-NO₃ and 2.6 128 µM P-PO₄ as potassium salts, and after one week with half the dose of N and P. This resulted 129 in a molar N:P ratio of 55:1, as we intended to apply a slight phosphorus limitation (Güsewell 130 & Koerselman 2002) to enhance effects of arsenate. We applied the nutrients only to the 131 water column, and not in the sediment to avoid direct competition with arsenate. For the 132 sediment-based pre-culture lasting 5 days, shoots were planted in artificial sediment modified 133 slightly from the OECD 239 test protocol (OECD 2014), omitting CaCO₃ and nutrient 134 additions, and only using 75% sand, 20% kaolin 5% white peat. The water column consisted 135 of Smart & Barko (1985) solution amended with N and P as described above. 136

In the experiment, we applied 0, 1.44, 7.2 and 36 mg [As] in the form of Na₂HAsO₄ × 7 H₂O per kg dry sediment, mixed as aqueous solution with the wetted sediment to reach 20% water content. An equivalent of 200 g dry sediment was added to square plastic pots (250 ml; $11 \times 7 \times 3$ cm; w × d × h), and one each was then placed in a 3-L plastic container (15 × 11 × 19 cm; w × d × h). The total volume of medium added per container was 2.5 L.

We set up 32 containers, 16 dedicated for each of the two different DIC conditions 142 (HC / high CO₂ availability or LC / low CO₂ availability/ dominance of HCO₃⁻ treatment), 143 and four replicates for each [As] concentration (4 levels including control) and carbon 144 condition. Half of the containers were filled with the standard Smart & Barko solution at 0.85 145 mM DIC (LC treatment), the other half with only 0.45 mM DIC (HC treatment) but amended 146 with CO_2 aeration from a CO_2 tank (50 l; NE45 – 99.995%) via needle valves to yield final 147 DIC concentrations of 0.8 - 0.9 mM. The LC treatments were only bubbled with air to allow 148 gas exchange with the atmosphere. The addition of CO₂ to the HC treatments shifted the total 149 150 DIC concentration to 0.8 – 0.9 mM (Hussner & Jahns 2015). Thus, different proportions of CO₂ and HCO₃⁻ were available in the HC and LC treatments, as quantified by standard 151 methods (see detailed description in Hussner & Jahns, 2015, and yielding pH values of $6.8 \pm$ 152 0.2 and 8.2 ± 0.2 in the HC and LC treatments, respectively). All containers were placed in a 153 water bath at $22 \pm 2^{\circ}$ C and $120 (\pm 15\%)$ µmol photons m⁻² s⁻¹ illumination by OSRAM 154 SubstiTUBE LED light tubes (ST8-HV4-170-840) at a 16/8 h L/D photoperiod. 155

Each pot was planted with two shoots of *M. spicatum*. Shoots were selected from the pre-cultures to have a shoot length between 6 and 14 cm (9.0 ± 1.8 cm, mean ± 1 SD) and well developed roots of approximately 1 to 2 cm length. Because shoots differed much in length, we decided do use each one larger shoot (A; 8.7 - 14 cm) and one smaller shoot (B; 6 -11 cm) per container. Before planting, each shoot was carefully blotted dry to take the initial fresh weight (0.19 ± 0.06 g; equivalent to 0.014 ± 0.004 g dry mass based on dry-to-fresh mass calibration of 10 control shoots) before planting. Preliminary experiments had shown that carefully blotting-to-dry would not harm plant development. Care was taken to distribute shoots evenly among all containers. No difference in initial total length or fresh mass for the different treatments and containers existed (1-way ANOVA, F < 1.22; P > 0.33).

166

167 Maintenance and end of experiment

168 The experiment was run for 14 days, and pH and water level of all containers was controlled and eventually adjusted daily, or once every two days during the weekend. We 169 170 occasionally measured oxygen content around midday, and found all containers at or above 100% saturation, indicating ongoing photosynthetic activity of the plants. At the end of the 171 experiment, we carefully removed the two shoots per container, and measured main shoot 172 length, root length and eventual side shoot length, as well as total fresh mass. Each shoot was 173 photographed to document its appearance. Shoot A was dissected into root, shoot and leaf 174 biomass, each fraction was placed individually into pre-weighted tubes to determine fresh 175 mass, and shock frozen in liquid nitrogen. Shoot B was left intact and whole plants were shock 176 frozen using liquid nitrogen. All frozen samples were freeze-dried for 24 h, providing 177 individual total shoot or root, stem, and leaf dry mass. From leaves, we calculated for shoot A 178 the respective leaf mass fractions (LMF = leaf dry mass/total dry mass or LMF-a = leaf dry 179 mass/leaf+stem dry mass; the latter to account for aboveground biomass), as well as stem and 180 181 root mass fractions (SMF = stem dry mass/total dry mass; RMF = root dry mass/total dry mass). For all samples, we calculated also the respective dry matter content (DMC) for total 182 shoots or plant organs (leaf, stem and root dry matter content as LDMC, SDMC and RDMC, 183 respectively) based on the respective ratios between dry and fresh mass. 184

185 Relative growth rates were calculated for fresh and dry mass (RGR-fm, RGR-dm) and 186 length (RGR-L) based on the equation $RGR = (\ln W_f - \ln W_i)/d$ with W_f the end value, W_i the 187 initial value and d the time in days.

188

189 Chemical analyses

Inorganic nutrients in the water column (N-NO₃ and P-PO₄) were measured at the end
of the experiment using ion chromatography (ICS-1100, Thermo-Fisher Scientific) and
spectrophotometry following standard procedures (AFNOR 1990). Arsenic in the sediment
was analysed after mineralization in 35 % ultrapure HNO₃ for 24 h. Arsenic in the
mineralized sediment samples and the water samples were then quantified by atomic
absorption spectroscopy (AAS; Perkin-Elmer, AAnalyst 100).

Dry biomass samples were ground to a fine powder using a Fritsch pulverisette 23 ball 196 mill before use in chemical analyses. All sample mass data are approximate values, the exact 197 198 mass for each sample was determined with a Sartorius MC2 ultra-microbalance at a precision of 10 µg. Elemental content for carbon (C), nitrogen (N) and phosphorus (P) were determined 199 in leaves from plant A. The content in C and N were determined based on analyses of 200 201 subsamples of 1 mg encapsulated in tin containers in a Carlo-ERBA Na 2100 CE elemental analyser. P was determined according to AFNOR (1990) using alkaline digestion (1 N NaOH 202 203 and 0.45 M Na₂S₂O₈) of 1 mg samples for 2 h at 120 °C and 1 bar followed by spectrophotometric quantification in an automated analyser (Konelab 420-XTi, Thermo 204 Fischer Scientific). 205

Chlorophyll, anthocyanins and total phenolic contents were analysed based on
methods described in Fornoff & Gross (2014). All measurements were made in a Varian Cary
50 spectrophotometer. Chlorophyll was extracted from 0.5 mg samples extracted in 1 ml

209	acetone/TRIS-hydrochloric acid-buffer (0.1 M), 8/2 (v/v); pH 7.8, for 2 h at 16 °C. The
210	absorbance was measured at 470, 664 and 647 nm to calculate the carotenoid, the chlorophyll
211	a and b content per extracted biomass based on the equations provided by Lichtenthaler &
212	Buschmann (2001). Anthocyanin was extracted with acidified methanol
213	(methanol/water/hydrochloric acid 37 %, $80/20/1$ [v/v/v]) from 1 mg samples in 1 ml solvent
214	for 2 h at 16 °C. The absorbance at 532 and 653 nm was used to calculate the anthocyanin
215	content (cyanidin, the major anthocyanin found in Myriophyllum spp. (Reznik & Neuhausel
216	1959) based on the formula developed from Murray & Hackett (1991): Anthocyanin [mg/g] =
217	((A 532 nm) - (0.24 * A 653 nm)) / (30 / 287.24 * m); with A = absorbance at the respective
218	wave length and m the exact dry mass of the sample extracted in one ml solvent. Total
219	phenolic compounds (TPC) were extracted in aqueous acetone (50/50 $[v/v]$; 1 mg sample in 1
220	ml solvent) followed by the Folin-Ciocalteau assay with tannic acid (no. 403040, batch
221	06817CJ; Sigma-Aldrich) as standard, following methods described in Fornoff & Gross
222	(2014).

223

224 Statistical analyses

Data were analysed using R Studio (2015). Homoscedasticity and normality were 225 verified respectively by the Levene and Shapiro-Wilk tests, and if needed, data were log or 226 227 1/x transformed to ensure normality and equal variance. Two-way ANOVAs with carbon and arsenic condition as fixed factors were calculated for all response variables using a probability 228 threshold of $p \le 0.05$, followed by Tukey post-hoc tests in case of significant differences 229 among treatments. A Principal component analysis (PCA) was performed on the full set of 230 endpoints except global DMC to visualize differences according to the experimental 231 conditions using FactoMineR (Lê et al. 2008). The contribution of each variable (endpoint) 232

for the different axes (i.e. the quality of representation of endpoints upon axes) was computedand called column inertia.

235

236 **Results**

237 Sediment and water column

238 The pH in the high and low CO₂ treatments stayed constant at the values provided in 239 Material and Methods due to the continuous aeration (LC treatment) and CO₂ aeration (HC treatment). Oxygen concentrations remained near or slightly above 100%, also due to the 240 241 aeration-induced turbulence. No visual algal growth in the water column was observed, only occasionally we found some algae growing on the sediment, but not on the plants. Reference 242 plants which were visibly free from epiphytes are documented in Supplementary Figure A1. 243 Plants had absorbed about 36 % of the added N-NO3 and 22 % of the added P-PO4 (data not 244 shown). 245

246 At the end of the experiment, we typically recovered 75-150 % of the initially applied arsenic in the containers, when combining concentrations found in sediment and water (Table 247 1). Part of the arsenic had leached into the water column, and this effect increased with 248 249 increasing initial sediment arsenic concentrations. Controls exhibited low concentrations of arsenic, possibly leaching from background levels in sediments. In the low arsenic treatment 250 (1.44 ppm) we found about 50 % higher concentrations than intended, possibly caused by 251 sediment background levels. While the arsenic concentrations for the levels 7.2 and 36 ppm in 252 the HC treatments were close to the intended values with 110 and 100 % measured 253 254 concentrations, they were significantly lower in the LC treatments, with only 85 and 76 %. Thus, recovery of arsenic was significantly lower in the LC treatments (Tukey post-hoc test, p 255 < 0.05). 256

257

258 Plant responses

CO₂ availability and arsenic concentrations affected growth, plant traits, and plant 259 physiology of *M. spicatum*. In general, arsenic effects were pronounced at 7.2 and 36 ppm 260 [As] in sediment, while high or low CO₂ availability affected only a few response variables, 261 and almost no interactive effects between factors were observed (Table 2, referring also to all 262 response variables outlined in detail below). 263 264 A – Effects on the growth of shoots A and B 265 The growth of *M. spicatum* based on of length, fresh and dry weight was strongly 266 267 influenced by arsenic, while low CO₂ availability only significantly reduced length-based growth, with marginal effects on biomass-based growth parameters (RGR-L, RGR-fm and 268 RGR-dm). The dry matter content (DMC) increased by about 50% at arsenic concentrations 269 270 of 7.2 and 36 ppm, resulting in somewhat weaker effects on dry mass-related growth (Fig. 1 A-C). 271 272 B – Effects on biomass allocation of shoot A 273 274 Arsenic and to a lesser extend CO₂ availability affected the biomass allocation to 275 leaves, stems and roots in *M. spicatum* (Fig. 2). This resulted in differences in the respective mass fractions of leaves (LMF), stems (SMF) or roots (RMF). With increasing arsenic 276 277 concentrations, RMF declined while stems got thicker resulting in a higher SMF. Under low CO₂ availability, the allocation to leaves increased while that to stems decreased. For leaves, 278 we calculated two indices: One dividing leaf dry mass by total biomass (LMF), and one using 279

only the aboveground biomass (LMF-a). Since root biomass declined strongly with increasing
arsenic, the effect at the whole plant biomass on LMF was rather weak, and no differences
between treatments were found. When roots were omitted in the calculation, the LMF-a
declined with increasing arsenic concentrations, and that this effect was enhanced in LC
treatments. At the two highest arsenic concentrations, the dry matter content (DMC) increased
for all plant organs in shoot A (leaves - LDMC, stems - SDMC and roots - RDMC) while
high CO₂ availability only increased the RDMC.

287

288 C – Effects on pigment and phenolic compound content in leaves of shoot A

Effects on pigments in leaves are shown in Fig. 3A-D. Total chlorophyll content declined with increasing arsenic concentration, but the chlorophyll a/b ratio did not differ between treatments. A strong interaction between arsenic concentration and CO_2 availability was observed for the carotenoid concentration, which specifically declined at high CO_2 availability and higher arsenic concentrations. Subsequently, also the ratio of carotenoids to total chlorophyll exhibited a strong interaction effect between arsenic concentration and CO_2 availability.

Anthocyanins (Fig. 4A) declined at low CO₂ availability and higher arsenic concentrations in leaves, but we observed no arsenic-induced differences at high CO₂ availability. The content in total phenolic compounds (TPC; Fig. 4B) in leaves A was strongly affected by CO₂ availability and arsenic concentration, but with no clear pattern, thus resulting in strong interactive effects. While high CO₂ availability increased the TPC in leaves, we observed no consistent trend with increasing arsenic concentrations. Only the leaves from the LC treatment had higher TPC at higher arsenic concentrations.

303

304 D – Effects on elemental stoichiometry in leaves from shoot A (Fig. 5)

305	Both factors, arsenic concentration and CO ₂ availability, had only slight effects on the				
306	elemental composition of carbon, nitrogen and phosphorus in leaves. With higher arsenic				
307	concentrations, the C content declined in leaves, irrespective of CO ₂ availability. Only				
308	marginal effects were observed for nitrogen and phosphorus. Leaves in the LC treatment at				
309	higher arsenic concentrations had a slightly lower N content, and the phosphorus content was				
310	not affected by arsenic and only marginally reduced by low CO ₂ availability. Overall leaf				
311	stoichiometry was not affected, as neither CO2 availability nor arsenic concentration affected				
312	the C:N, C:P or N:P molar ratio.				
313					
314	E – Overall effects on plant performance (Fig. 6)				
315	The principal component analysis (PCA) on the whole set of endpoints and for the				
316	different treatment combinations showed distinct differences between treatment				
317	combinations. Axes 1 and 2 explained 37.2 and 15.7 % of the variance. The axes 3 and 4				
318	added another 9.9 and 7.8 % to explain in total more than 70 % of the total variance.				
319	Differences between low (0 and 1.44 ppm) or high (7.2 and 36 ppm) levels of [As] in				
320	sediment were visible along axis 1, while less pronounced differences based on CO ₂				
321	availability were related to axis 2. The correlation circle (Fig. 6A) shows the significant				
322	correlations for fresh mass based growth (RGR-fm), root mass fraction (RMF) and				
323	chlorophyll content (chl a , chl $a+b$), and in opposite direction stem mass fraction (SMF) and				
324	the dry matter content of all plant organs, leaves, roots and stems (LDMC, RDMC, SDMC)				
325	with axis 1. Axis 2 is mainly explained by phosphorus content (P) and the resulting C:P molar				
326	ratio, and to a lesser extent by the N:P molar ratio and pigment ratios (car/chl, chl <i>a/b</i>).				

327 Detailed correlation coefficients and associated P-values can be found in the Supplementary328 Material, Table A1.

329

330 **Discussion**

In a 14-day exposure to arsenate in combination with high or low CO₂ availability we observed treatment-specific effects on the growth, biomass allocation and plant metabolism in the rooted, submerged macrophyte *M. spicatum*. In the following we will discuss our initial hypotheses in detail.

335

Hypothesis 1: Environmentally relevant arsenic concentrations negatively affect growth andphysiological response variables in *M. spicatum*

Most selected endpoints measured in M. spicatum showed the effect of arsenic applied 338 to the sediment. Negative effects on growth and physiology started at a nominal concentration 339 of 7.2 ppm arsenic applied to the sediments. This is a rather low concentration, and in the 340 range of "normal", natural values found in lake or river sediments (Smedley & Kinniburgh 341 2002). However, the observed effects might have been higher in our experiment because the 342 applied arsenate is highly soluble, and diffused rapidly from the sediment to the overlying 343 water column, as our measurements show (Table 1). Also under natural conditions, sediments 344 and rocks release arsenic to the water column. Robinson et al. (2006) observed water 345 concentrations up to 180 µg l⁻¹ in the Taupo Volcanic Area, but in areas with no geochemical 346 or anthropogenic sources for arsenic, concentrations are mostly below 10 µg l⁻¹ (Smedley & 347 Kinniburgh 2002). 348

We could not assess the fate of arsenic accumulated in plant tissues or plant uptake as we only had limited biomass available (50 to 150 mg per replicate, depending on treatment

and used for all chemical analyses). Nevertheless, the negative effects already at 7.2 and 36 351 352 ppm nominal sediment concentrations suggest that at least part of the arsenic had been taken up by the plants, either from sediment or water. Laboratory studies showed growth reductions 353 in *M. propinguum* from the Taupo Volcanic Zone rich in arsenic at about 170 to 290 µg l⁻¹ 354 (Robinson et al. 2006). In our study, first effects occurred at measured final concentrations of 355 approximately 300 µg l⁻¹ in water and 3.25 ppm in sediment originally spiked with 7.2 ppm 356 [As]. No effects were observed in the control and the low arsenic treatment, although they 357 contained some arsenic in water and sediment. 358

A remarkable finding is the strong increase in the dry matter content (DMC) of the 359 360 whole shoots or in the different plant organs (leaves - LDMC, stems - SDMC, roots - RDMC) at higher arsenic concentrations. This indicates that plant physiology was altered, as plants 361 had reduced water content and/or changed some metabolites or components, such as increased 362 allocation to storage carbohydrates and starch (Hussner & Jahns 2015). Exposure to arsenic 363 also increased starch content in Ceratophyllum demersum due to negative effects on 364 365 photosynthesis (Mishra et al. 2014), thus it is likely that the observed increase in DMC here is caused by starch accumulation. 366

Plant leaves, stems, roots, and tissue biomass ratios also changed following exposure 367 to arsenic. Despite significant amounts of arsenic leaching into the water column, we 368 observed a strong reduction in root mass, both absolute and relative to total biomass (root 369 mass fraction / RMF), from above 15 % to around 5 % in high arsenic treatments. RMF 370 provides comparable results to the frequently used root/shoot ratio, with the difference that 371 the latter divides the root biomass only by the aboveground biomass, thus in our case the 372 root/shoot ratio in the controls and low arsenic treatments would be approximately 20 %, a 373 value comparable to that reported by Hussner & Jahns (2015). 374

The roots of arsenic exposed plants were thicker and darker than those found in 375 376 controls (Fig. A1). This is a well described phenomenon for macrophytes growing in metalpolluted sediments. The darkening results in part from iron plaques formed on the roots. 377 Many trace metal elements, including arsenic, bind to iron plaques on macrophyte roots 378 (Taggart et al. 2009), and the magnitude of iron plaque formation on roots influences the 379 accumulation of trace metal elements in aquatic plants (Tripathi et al. 2014). Stem appearance 380 381 did not differ visually (Fig. A1), but the biomass allocation to stems (SMF) rose sharply with higher arsenic exposure, increasing from 25 % up to 40 % of the total biomass. We assume 382 that plants used the stems to store arsenic and avoid negative effects on leaves. 383

The decline in chlorophyll *a* and *b* in leaves of plants exposed to higher arsenic concentrations is not unexpected, and has been observed in other plants exposed to arsenic or other trace metal elements. For example, negative effects on photosynthetic pigments have been observed in the floating fern *Azolla filiculoides* exposed to $60 \ \mu g \ l^{-1} As(V)$ (Sánchez-Viveros et al. 2011). Overall we can confirm our first hypothesis that arsenic exposure at environmentally realistic concentrations caused strong effects on different growth and physiology related parameters in *M. spicatum*.

391

392 Hypothesis 2: Effects of arsenic exposure are enhanced at high pH and low CO₂ availability

Low CO₂ availability caused much less, and usually less strong effects on the growth
 and physiology of *M. spicatum*, and only a few interactive effects between CO₂ availability

and arsenic exposure could be observed. High bicarbonate concentration in the medium

together with a high pH had affected arsenic uptake in the fern *Pteris vittata* (Guo et al.

397 2012). Yet, we observed only little differences between the two carbon treatments for arsenic

induced effects, and thus assume that the applied higher pH or lower CO₂ availability,

399	resulting in a higher bicarbonate concentration of the medium, might not have affected the
400	bioavailability, or negative effects of arsenic on M. spicatum. Only length growth was
401	significantly affected and mass-based growth rates declined only marginally. A much stronger
402	effect of CO ₂ availability was observed on biomass allocation to leaves, stems and roots. We
403	found an increase in RDMC in the high CO ₂ treatments, while Hussner & Jahns (2015) found
404	an increase in LDMC at high CO ₂ availability. Possibly the light intensity in our experiment
405	was too low, as the effects were also reduced or not visible in the low light treatments of
406	Hussner & Jahns (2015). Leaf biomass increased (LMF-a and LMF) in the low CO_2
407	treatments, and this can be seen in accompanying photographs (Fig. A1). We explicitly
408	calculated both the LMF based on total biomass and that based only on aboveground biomass
409	(LMF-a). While the former is commonly used as plant trait (Pérez-Harguindeguy et al. 2013),
410	the strong changes in root biomass resulted in a higher significance when considering only the
411	aboveground biomass for the calculation (LMF-a; Table 2). Our results indicate that plants in
412	the low CO ₂ treatments invested more in leaf biomass to improve the diffusional dissolved
413	inorganic carbon uptake (Eusebio Malheiro et al. 2013).

Except for a few cases, low CO₂ availability has not enhanced negative effects of arsenic exposure on *M. spicatum*. This might be due to the less severe carbon limitation in our set-up that provided still some aeration and kept the pH at about 8.2. If the higher pH of 9 as observed in the new OECD TG 239 (OECD 214) would have stronger effects needs to be shown. Yet, effects with the fern *Pteris vittata* (Guo et al. 2012) were also only visible at rather high bicarbonate levels, and not in environmentally relevant ranges as used in our experiment. We thus cannot confirm our second hypothesis.

422 Hypothesis 3: Changes in morphological and physiological traits of *M. spicatum* reflect
423 specific responses towards a certain combination of stressors.

Although we could not confirm our second hypothesis, we think that our extended 424 425 analysis of additional parameters provides a good picture of specific responses of *M. spicatum* 426 to certain stressors. To achieve this we used morphological plant traits, such as the dry matter content (DMC) and mass fractions for leaves, stems and roots (LMF, SMF, RMF), or 427 428 physiological traits such as the elemental composition and stoichiometry and the content in phenolic compounds. The principal component analysis shows (Fig. 6B) that for the two 429 higher arsenic concentrations CO₂ availability did make a difference in the overall plant 430 431 response, based on individual endpoints, or response variables.

Only arsenic increased the DMC in *M. spicatum*, but an increase in DMC has several 432 implications. From an ecological perspective, plants with a higher LDMC are less palatable to 433 herbivores (Elger & Willby 2003) and *M. spicatum* even increases the DMC of apical tissues 434 435 when attacked by herbivores (Fornoff & Gross 2014). With respect to common endpoints 436 used in ecotoxicology, an increased DMC can mask effects on dry mass based yield or relative growth rate calculations. Our results show that effects based on dry mass (RGR-dm) 437 were weaker than those based on fresh mass or total shoot length (RGR-fm; RGR-L). Arsenic 438 is not the only pollutant causing this effect, as the herbicides metsulfuron-methyl (Cedergreen 439 440 et al. 2004) and mesosulfuron-methyl previously showed no effects on dry-mass based growth estimations in *M. spicatum*, due to increases in DMC as we observed recently (Nuttens et al. 441 442 2016). From an ecotoxicological perspective, it seems thus prudent to select several growth endpoints to avoid the masking effect of increases in DMC on dry mass related results. 443

Changes in the proportion of leaf, stem and root biomass have ecological and
ecotoxicological implications. From an autecological perspective, stronger stems can provide

446 more stability and resistance to mechanical forces, but if this is at the expense of leaf biomass, 447 plants will have less photosynthetic tissue and thus risk slower growth rates. The new test 448 guideline for *M. spicatum* growing in a water-sediment set-up (OECD 2014) requires only 449 evaluation of root development. Though it seems unlikely that all plant traits used here could 450 be included for most normalized testing, the selection of additional endpoints would be quite 451 useful for understanding processes at the plant level and implications in an ecological context.

452 We found significant effects of both CO₂ availability and arsenic, and interaction effects on levels of phenolic compounds in *M. spicatum*, suggesting that this endpoint is 453 reacting strongly influenced by environmental conditions. A decline in phenolic compounds 454 455 was found in A. filiculoides exposed to arsenic (Sánchez-Viveros et al. 2011), yet we observed either no effect on TPC (HC for leaves) or an increase (LC for leaves exposed to 7.2 456 or 36 ppm [As]). Anthocyanin increased in Lemna exposed to water containing arsenic (Leao 457 et al. 2014), but we observed a decline in anthocyanins. Phenolic compounds are ubiquitous 458 459 in terrestrial and aquatic plants, but the relative content and composition can vary strongly 460 depending on environmental conditions, and in response to a wide range of abiotic and biotic 461 stressors (Dixon & Pavia 1995; Gross & Bakker 2012). We think that phenolic compounds are interesting indicators for environmental stressors acting on *M. spicatum* and other plants, 462 463 and propose to have a more detailed look how they are modified e.g. by trace metal elements in combination with environmental conditions, and how this affects plant performance and 464 biotic interactions. 465

Effects of CO₂ availability and arsenic on elemental composition and plant stoichiometry were relatively low in *M. spicatum* leaves. We expected that increased arsenic concentration would reduce the uptake of phosphate, as had been observed for *Lemna* (Mkandawire & Dudel 2012), but this effect was not found in leaves. The nitrogen content was slightly affected by arsenic (P = 0.066), and the significant lower carbon content due to

arsenic might be related to a lower photosynthetic activity (Mishra et al. 2014) and the lower 471 472 chlorophyll content. After two weeks exposure, we observed no effects on leaf stoichiometry. Apparently, N:P molar ratios are relatively robust, as Li et al. (2015) reported similar values 473 474 along a range of water depths for three common submerged macrophytes. Also other authors report a relative homeostasis of the N:P ratio in aquatic plants irrespective of environmental 475 nitrogen or phosphorus concentrations (Demars & Edwards 2007; Güsewell & Koerselman 476 477 2002). However, severe carbon stress due to very low CO_2 (and HCO_3^{-}) availability as it occurs in the standard set-up of the *M. spicatum* sediment-water OECD test system (OECD 478 2014) where no aeration occurs and resulting pH values exceed 9 (EMG, pers. observ.), might 479 480 modify plant stoichiometry. Nitrogen content in several Myriophyllum and other submerged macrophyte species was reduced under high CO₂ availability, with M. spicatum exhibiting the 481 482 lowest effects (Hussner et al., in press).

The principal component analysis revealed links between the different endpoints 483 chosen in our study. All different treatment combinations were rather well separated along 484 485 axes 1 and 2. The separation of treatments along axis 1 was influenced by differences in arsenic concentration, separating between the two low (0 and 1.44 ppm [As]) and the two 486 high [7.2 and 36 ppm [As]) concentrations. Differences based on CO₂ availability showed 487 488 along axis 2, but only for the two highest arsenic concentrations. Along axis 1, a high growth rate based on fresh mass corresponded to a high chlorophyll content, a high root mass and 489 aboveground leaf mass fraction (RMF, LMF-a), and was opposed to higher stem mass 490 fraction (SMF) and increases in all dry matter related endpoints (LDMC, RDMC, SDMC). 491 492 We thus consider endpoints pointing to the left in the circle of correlation (Fig. 6B) reflecting 493 active growth and development, while those pointing to the right as endpoints showing impaired growth. Interestingly, although we had not found many significant effects of CO₂ 494 availability on individual endpoints, treatments for one given arsenic level differed between 495

high and low CO₂ availability (HC, LC), with the strongest effect for 7.2 ppm [As] in
sediment. Thus, low CO₂ availability had not aggravated the impact of arsenic, but exerted
different effects on the plants. These were a slight reduction in phosphorus content in leaves,
modifications in the content of phenolic compounds (TPC) and the ratio of carotenoids to
chlorophyll (car/chl), the latter two possible indicators of oxidative stress (Waśkiewicz et al.
2014).

502

503 Conclusions

504 This laboratory experiment combined realistic CO₂ availability scenarios with presumed environmentally relevant arsenic exposure settings. Despite the high leaching of arsenic from 505 the sediment to the water column, the effective concentrations for *M. spicatum* were in the 506 507 range that can be found in areas with natural or anthropogenic high exposure to arsenic via sediment or water. We can thus partly confirm our first hypothesis that negative effects on 508 growth and physiology can be observed at environmentally relevant arsenic concentrations. 509 Our second hypothesis that low CO₂ availability would aggravate the effects of arsenic could 510 only partly be verified, but results might be different for a longer, chronic exposure or under 511 more severe CO₂ stress and pH values above 9, as caused by the complete absence of aeration 512 in the new OECD test guideline for *M. spicatum* (OECD 2014). With respect to our third 513 hypothesis, we think that our experiment shows the complex response of *M. spicatum* to 514 515 combined stressors. The combination of different endpoints allowed a rather clear distinction of treatment combinations. We suggest investigators consider whether a broader range of 516 response variables can help identifying responses to stress-induced changes. Additional 517 response variables may include plant traits such as biomass allocation, dry matter content, 518 stoichiometry and the content in secondary metabolites in addition to general "endpoints" 519

based on growth and pigment content. These additional endpoints might be useful in
predicting the plant performance in an ecological context, as for example increased leaf
biomass would result in better photosynthetic performance, or increased content in phenolic
compounds could explain the plant's capacity to be defended against competitors or
herbivores. Such effects on the "plant quality", i.e. the dry matter content, phenolic content or
elemental stoichiometry, have already been shown to affect herbivory (Fornoff & Gross
2014), and possibly other biotic interactions as allelopathy.

527

528 Acknowledgment

We greatly appreciate the help of P. Rousselle with the C:N analyses, J.-F. Poinsaint with 529 setting up the experimental infrastructure, S. Devin and L. Minguez for statistical advice. AH 530 benefited from an invitation by the UFR SciFa, Univ Lorraine for a visit of one month. AN 531 was funded by a PhD stipend from the MESR, Ministère de l'Enseignement Supérieur et de la 532 Recherche, France. The experiment was possible due to funding for the project "PICAI -533 pollutant induced changes in allelochemical interactions" in the program ECODYN / EC2CO, 534 INSU, CNRS to EMG and by support from the LTER – ZAM – Zone Atelier Moselle for AN, 535 536 DP and EMG. We thank three anonymous reviewers and the editors of the special issue for constructive comments on prior versions of this manuscript. 537

538

539 References

540 AFNOR, 1990. Eaux-Méthodes d'Essais. Association Française de Normalisation (Ed.). ISBN 2-12541 179041-1

542 Cedergreen, N., J. C. Streibig & N. H. Spliid, 2004. Sensitivity of aquatic plants to the herbicide
543 metsulfuron-methyl. Ecotoxicology and Environmental Safety 57: 153-161.

- Demars, B. O. L. & A. C. Edwards, 2007. Tissue nutrient concentrations in freshwater aquatic
 macrophytes: high inter-taxon differences and low phenotypic response to nutrient supply.
 Freshwater Biology 52: 2073-2086.
- 547 Dixon, R. A. & N. L. Pavia, 1995. Stress-induced phenylpropanoid metabolism. Plant Cell 7: 1085548 1097.
- 549 Dushenko, W. T., D. A. Bright & K. J. Reimer, 1995. Arsenic bioaccumulation and toxicity in aquatic
 550 macrophytes exposed to gold-mine effluent: relationships with environmental partitioning,
 551 metal uptake and nutrients. Aquatic Botany 50: 141-158.
- Elger. A. & N.J. Willby, 2003. Leaf dry matter content as an integrative expression of plant
 palatability: the case of freshwater macrophytes. Functional Ecology 17: 58-65.
- 554 Eusebio Malheiro, A. C., P. Jahns & A. Hussner, 2013. CO₂ availability rather than light and
- temperature determines growth and phenotypical responses in submerged *Myriophyllum aquaticum*. Aquatic Botany 110: 31-37.
- Finnegan, P. M. & W. Chen, 2012. Arsenic toxicity: The effects on plant metabolism. Frontiers in
 Physiology 3: 182
- Fornoff, F. & E. M. Gross, 2014. Induced defense mechanisms in an aquatic angiosperm to insect
 herbivory. Oecologia 175: 173-185.
- Geng, C. N., Y. G. Zhu, W. J. Liu & S. E. Smith, 2005. Arsenate uptake and translocation in seedlings
 of two genotypes of rice is affected by external phosphate concentrations. Aquatic Botany 83:
 321-331.
- Grace, J. B. & R. G. Wetzel, 1978. The production biology of Eurasian watermilfoil *(Myriophyllum spicatum* L.): a review. Joural of Aquatic Plant Management 16: 1-11.
- Gross, E. M., 2003. Differential response of tellimagrandin II and total bioactive hydrolysable tannins
 in an aquatic angiosperm to changes in light and nitrogen. Oikos 103: 497-504.
- Gross, E. M. & E. S. Bakker, 2012. The role of plant secondary metabolites in freshwater macrophyteherbivore interactions: Limited or unexplored chemical defences? In Iason, G. R., M. Dicke &

570	S. E. Hartley (eds.) The Integrative Role of Plant Secondary Metabolites in Ecological						
571	Systems. British Ecological Society / Cambridge University Press, Sussex, UK, pp. 154-169.						
572	Guo, H. M., Z. N. Zhong, M. Lei, X. L. Xue, X. M. Wan, J. Y. Zhao & T. B. Chen, 2012. Arsenic						
573	uptake from arsenic-contaminated water using hyperaccumulator Pteris vittata L.: Effect of						
574	chloride, bicarbonate, and arsenic species. Water Air and Soil Pollution 223: 4209-4220.						
575	Güsewell, S. & M. Koerselman, 2002. Variation in nitrogen and phosphorus concentrations of wetland						
576	plants. Perspectives in Plant Ecology Evolution and Systematics 5: 37-61.						
577	Harguinteguy, C. A., M. L. Pignata & A. Fernández-Cirelli, 2015. Nickel, lead and zinc accumulation						
578	and performance in relation to their use in phytoremediation of macrophytes Myriophyllum						
579	aquaticum and Egeria densa. Ecological Engineering 82: 512-516.						
580	Hartley, S. E., C. G. Jones, G. C. Couper & T. H. Jones, 2000. Biosynthesis of plant phenolic						
581	compounds in elevated atmospheric CO ₂ . Global Change Biology 6: 497-506.						
582	Hussner, A., D. Hofstra, P. Jahns & J. Clayton, 2015. Response capacity to CO ₂ depletion rather than						
583	temperature and light effects explain the growth success of three alien Hydrocharitaceae						
584	compared with native Myriophyllum triphyllum in New Zealand. Aquatic Botany 120: 205-						
585	211.						
586	Hussner, A. & P. Jahns, 2015. European native Myriophyllum spicatum showed a higher use capacity						
587	than alien invasive Myriophyllum heterophyllum. Hydrobiologia 746: 171-182.						
588	Hussner, A., T. Mettler-Altmann, A. P. M. Weber & K. Sand-Jensen, in press. Acclimation of						
589	photosynthesis to supersaturated CO ₂ in aquatic plant bicarbonate users. Freshwater Biology,						
590	accepted manuscript. doi:10.1111/fwb.12812						
591	Krayem M, V Deluchat, M. Rabiet, K. Cleries, J. F. Lenain, Z. Saad, V. Kazpard & P. Labrousse,						
592	2016. Effect of arsenate As(V) on the biomarkers of Myriophyllum alterniflorum in						
593	oligotrophic and eutrophic conditions. Chemosphere 147: 131-137.						
594	Lavid, N., A. Schwartz, O. Yarden & E. Tel-Or, 2001. The involvement of polyphenols and						
595	peroxidase activities in heavy-metal accumulation by epidermal glands of the waterlily						

596 (Nymphaeaceae). Planta 212: 323-331.

- 597 Lê, S., J. Josse & F. Husson, 2008. FactoMineR: An R Package for Multivariate Analysis. Journal of
 598 Statistical Software. 25: 1-18.
- Leao GA, JA de Oliveira, RTA Felipe, FS Farnese & GS Gusman, 2014. Anthocyanins, thiols, and
 antioxidant scavenging enzymes are involved in *Lemna gibba* tolerance to arsenic. Journal of
 Plant Interactions 9: 143-151.
- Leu, E., A. Krieger-Liszkay, C. Goussias & E. M. Gross, 2002. Polyphenolic allelochemicals from the
 aquatic angiosperm *Myriophyllum spicatum* L. inhibit photosystem II. Plant Physiology 130:
 2011-2018.
- Li, W., T. Cao, L. Y. Ni, G. R. Zhu, X. L. Zhang, H. Fu, X. Song & P. Xie, 2015. Size-dependent C, N
 and P stoichiometry of three submersed macrophytes along water depth gradients.

607 Environmental Earth Sciences 74: 3733-3738.

- 608 Lichtenthaler, H. K. & C. Buschmann, 2001. Chlorophylls and Carotenoids: Measurement and
- 609 Characterization by UV-VIS Spectroscopy Current Protocols in Food Analytical Chemistry.
 610 John Wiley & Sons, Inc.
- Meybeck, M., 2004. The global change of continental aquatic systems: Dominant impacts of human
 activities Water Science and Technology 49: 73-83.
- Mishra, S., H. J. Stark & H. Kupper, 2014. A different sequence of events than previously reported
 leads to arsenic-induced damage in *Ceratophyllum demersum* L. Metallomics 6: 444-454.
- Mkandawire, M. & E. G. Dudel, 2012. Homeostatic regulation of elemental stoichiometry by *Lemna gibba* L. G3 when nutrient interact with toxic metals. Ecotoxicology 21: 456-464.
- Murray, J. R. & W. P. Hackett, 1991. Dihydroflavonol reductase activity in relation to differential
 anthocyanin accumulation in juvenile and mature phase *Hedera helix* L. Plant Physiology
 97:343-351
- Nuttens, A., S. Chatellier, S. Devin, C. Guignard, A. Lenouvel & E. M. Gross, 2016. Does nitrate copollution affect biological responses of an aquatic plant to two common herbicides? Aquatic
 Toxicology 177: 355-364.

623 OECD, 2014. TG 239: Water-sediment *Myriophyllum spicatum* toxicity test. Test Guideline No. 239,
624 23 pp.

625	Pérez-Harguindeguy, N., S. Díaz, E. Garnier, S. Lavorel, H. Poorter, P. Jaureguiberry, M. S. Bret-
626	Harte, W. K. Cornwell, J. M. Craine, D. E. Gurvich, C. Urcelay, E. J. Veneklaas, P. B. Reich,
627	L. Poorter, I. J. Wright, P. Ray, L. Enrico, J. G. Pausas, A. C. de Vos, N. Buchmann, G.
628	Funes, F. Quétier, J. G. Hodgson, K. Thompson, H. D. Morgan, H. ter Steege, M. G. A. van
629	der Heijden, L. Sack, B. Blonder, P. Poschlod, M. V. Vaieretti, G. Conti, A. C. Staver, S.
630	Aquino & J. H. C. Cornelissen, 2013. New handbook for standardised measurement of plant
631	functional traits worldwide. Australian Journal of Botany 61: 167-234.
632	R Studio Team, 2015. RStudio: Integrated Development for R. RStudio, Inc., Boston, MA, USA.
633	Rahman, M. A., H. Hasegawa & R. P. Lim, 2012. Bioaccumulation, biotransformation and trophic
634	transfer of arsenic in the aquatic food chain. Environmental Research 116: 118-135.
635	Reznik, H. & R. Neuhausel, 1959. Farblose Anthocyane bei submersen Wasserpflanzen. Zeitschrift für
636	Botanik 47: 471-489.
637	Robinson, B., N. Kim, M. Marchetti, C. Moni, L. Schroeter, C. van den Dijssel, G. Milne & B.
638	Clothier, 2006. Arsenic hyperaccumulation by aquatic macrophytes in the Taupo Volcanic
639	Zone, New Zealand. Environmental and Experimental Botany 58: 206-215.
640	Sánchez-Viveros, G., R. Ferrera-Cerrato & A. Alarcón, 2011. Short-term effects of arsenate-induced
641	toxicity on growth, chlorophyll and carotenoid contents, and total content of phenolic
642	compounds of Azolla filiculoides. Water Air Soil Pollution 217: 455-462.
643	Segner, H., M. Schmitt-Jansen & S. Sabater, 2014. Assessing the impact of multiple stressors on
644	aquatic biota: The receptor's side matters. Environmental Science & Technology 48: 7690-
645	7696.
646	Smart, R. M. & J. W. Barko, 1985. Laboratory culture of submersed freshwater macrophytes on
647	natural sediments. Aquatic Botany 21: 251-263.
648	Smedley, P. L. & D. G. Kinniburgh, 2002. A review of the source, behaviour and distribution of
649	arsenic in natural waters. Applied Geochemistry 17: 517-568.

- 650 Smith, C. S. & J. W. Barko, 1990. Ecology of Eurasian watermilfoil. Journal of Aquatic Plant
 651 Management 28: 55-64.
- Taggart, M. A., R. Mateo, J. M. Charnock, F. Bahrami, A. J. Green & A. A. Meharg, 2009. Arsenic
 rich iron plaque on macrophyte roots an ecotoxicological risk? Environmental Pollution 157:
 946-954.
- Tripathi, R. D., P. Tripathi, S. Dwivedi, A. Kumar, A. Mishra, P. S. Chauhan, G. J. Norton & C. S.
 Nautiyal, 2014. Roles for root iron plaque in sequestration and uptake of heavy metals and
 metalloids in aquatic and wetland plants. Metallomics 6: 1789-1800.
- 658 Waśkiewicz, A., M. Beszterda & P. Goliński, 2014. Nonenzymatic Antioxidants in Plants. In:
- 659 Oxidative Damage to Plants: Antioxidant Networks and Signaling (ed. P. Ahmad). pp. 201-
- 660 234.

661 Table 1: Recovery of arsenic at the end of the experiment in the different treatments. Data represent means ± 1 SE; n = 4.

	Nominal sediment As concentration [mg kg ⁻¹]	Recovery		% of total recovery found in		% recovery of initially applied [As]	
Treat- ment		Sediment [mg kg ⁻¹]	Water [µg l ⁻¹]	Sediment Water			
HC	0	0.88 ± 0.14	2.75 ± 0.75	96.3	3.7		
	1.44	1.45 ± 0.12	50.50 ± 5.85	69.9	30.1	145.0 ± 6.6	
	7.20	3.60 ± 0.43	344.75 ± 25.96	45.6	54.4	110.0 ± 7.4	
	36.00	8.67 ± 0.60	2196.75 ± 226.01	24.5	75.5	100.4 ± 8.0	
LC	0	0.64 ± 0.79	16.50 ± 3.43	44.4	55.6		
	1.44	1.63 ± 0.16	52.0 ± 6.76	71.5	28.5	158.4 ± 10.4	
	7.20	2.89 ± 0.15	260.50 ± 25.01	47.4	52.6	85.4 ± 4.1	
	36.00	7.20 ± 0.67	1618.75 ± 344.10	28.1	71.9	76.2 ± 12.6	

Table 2: 2-way ANOVA results based on the fixed factors carbon and arsenic. Shown are the

- effect of CO₂ availability and arsenic pollution on different response variables measured in
- 666 *Myriophyllum spicatum*. Results are based a) on growth data for both shoots per microcosm,
- b) on plant traits and metabolism of leaves from shoot A and c) on the stoichiometry of the
- 668 whole shoot B. Degrees of freedom: df = 1 for carbon, df = 3 for arsenic; df = 3 for carbon x
- arsenic. * log transformed data. ** 1/x transformed data. Bold numbers indicate significant
- 670 results (P < 0.05). Marginally significant results (P < 0.1) are marked in italics.
- 671 RGR Relative growth rate based on dm dry mass, fm fresh mass or L length. DMC –
- 672 Dry matter content. LMF and LMF-a leaf mass fraction based on total biomass or
- aboveground biomass. SMF, RMF stem and root mass fractions. LDMC, SDMC and

674 RDMC – leaf, stem and root dry matter content. Chl – chlorophyll. Car – carotenoids. TPC –

total phenolic compounds. C – carbon. N – nitrogen. P – phosphorus.

	Carbon		Arsenic		Carbon x Arsenic		
	F	Р	F	Р	F	Р	
A) Growth rate and dry matter content of both shoots							
RGR-dm	3.494	0.074	5.366	0.006	0.607	0.617	
RGR-fm	3.235	0.085	42.95	< 0.001	1.683	0.197	
RGR-L	7.277	0.013	15.590	< 0.001	2.365	0.096	
DMC	0.015	0.902	27.831	< 0.001	1.779	0.178	
B1) Plant traits of shoot	A						
LMF**	5.415	0.029	3.530	0.030	0.396	0.757	
LMF-a	13.335	0.001	13.249	< 0.001	1.278	0.304	
SMF	12.329	0.002	32.794	< 0.001	1.065	0.382	
RMF	0.309	0.583	36.651	< 0.001	0.242	0.866	
LDMC	0.026	0.874	17.030	< 0.001	2.281	0.105	
SDMC	0.512	0.481	48.158	< 0.001	3.358	0.036	
RDMC*	24.906	< 0.001	49.897	< 0.001	1.478	0.246	
B2) Pigments in leaves of	of shoot A						
Chl a	1.064	0.313	10,755	< 0.001	2.206	0.114	
Chl b	3.375	0.079	9.630	< 0.001	1.870	0.162	
Chl <i>a</i> & <i>b</i>	1.768	0.196	10.744	< 0.001	2.109	0.126	
Chl <i>a/b</i> ratio	3.643	0.068	0.341	0.796	1.535	0.231	
Car	3.462	0.075	2.718	0.067	9.629	< 0.001	
Car/chl ratio	20.639	< 0.001	6.572	0.002	9.489	< 0.001	
Anthocyanins	1.977	0.173	4.991	0.008	2.782	0.063	
TPC	576.87	< 0.001	43.03	< 0.001	28.16	< 0.001	
C1) Plant stoichiometry	of leaves from s	shoot A					
C	0.934	0.346	6.610	0.002	0.997	0.411	
Ν	1.258	0.273	2.730	0.066	1.784	0.177	
Р	3.093	0.091	0.825	0.493	0.407	0.749	
CN	1.284	0.268	1.652	0.204	1.680	0.198	
СР	2.080	0.162	1.146	0.351	0.441	0.726	
NP	0.919	0.347	1.554	0.226	0.233	0.872	
C2) Plant stoichiometry	of shoot B						
C	5.848	0.024	0.573	0.638	7.242	0.001	
Ν	0.164	0.689	5.518	0.005	2.302	0.101	
Р	2.837	0.105	3.808	0.023	2.988	0.051	
CN	0.013	0.910	6.366	0.003	1.040	0.393	
СР	6.196	0.020	4.722	0.010	2.510	0.083	
NP	5.343	0.030	1.053	0.387	0.787	0.513	

676

678 Figure legends

Fig. 1. Effect of arsenic and CO_2 availability (HC – high CO_2 ; LC – low CO_2) on the relative

- 680 growth rate of *Myriophyllum spicatum* based on dry mass (RGR-dm; A) or total shoot length
- 681 (RGR-L; B) and the dry matter content (DMC; C). Results are based on data for both shoots
- A and B used per replicate, and represent means ± 1 SE from n = 4. Results from the Tukey
- post-hoc tests following the 2-way ANOVA are shown as small letter code next to each
- treatment. If no letters are shown for a response variable, treatments were not different to each
- other. Note that y-axes do not start with zero but are adapted to the data range.
- Fig. 2. Effect of arsenic and CO_2 availability (HC high CO_2 ; LC low CO_2) on plant traits,
- 687 here the biomass allocation to leaves, stems and roots (LMF, SMF, RMF). The proportion of
- leaves on the total aboveground biomass is shown as LMF-a. Results are based on data for
- shoots A, and represent means ± 1 SE from n = 4. Results from the Tukey post-hoc tests
- 690 following the 2-way ANOVA are shown as small letter code next to each treatment. If no
- 691 letters are shown for a response variable, treatments were not different to each other. Note that
- 692 y-axes do not start with zero but are adapted to the data range.
- Fig. 3. Effect of arsenic and CO_2 availability (HC high CO_2 ; LC low CO_2) on leaf
- 694 pigments and relevant pigment ratios. A Total chlorophyll content. B Carotenoid content.
- 695 C Chlorophyll a/b (chl a/b) ratio. D Carotenoid/total chlorophyll ratio (Car/chl ratio).
- Results are based on data for leaves from shoot A, and represent means ± 1 SE from n = 4.
- 697 Results from the Tukey post-hoc tests following the 2-way ANOVA are shown as small letter
- 698 code next to each treatment. If no letters are shown for a response variable, treatments were
- 699 not different to each other. Note that y-axes do not start with zero but are adapted to the data
- 701

700

range.

Fig. 4. Effect of arsenic and CO₂ availability (HC – high CO₂; LC – low CO₂) on leaf phenolic compounds, here content in anthocyanins (A) and total phenolic compounds (TPC; B). Results are based on data for leaves from shoot A, and represent means ± 1 SE from n = 4. Results from the Tukey post-hoc tests following the 2-way ANOVA are shown as small letter code next to each treatment. If no letters are shown for a response variable, treatments were not different to each other. Note that y-axes do not start with zero but are adapted to the data range.

709

Fig. 5. Effect of arsenic and CO₂ availability (HC – high CO₂; LC – low CO₂) on elemental 710 composition, (panels A - C: Content in carbon - C; nitrogen - N; phosphorus - P) and 711 resulting molar ratios (leaf stoichiometry; molar ratios of panels D – F: C:N, C:P and N:P). 712 713 Results are based on data for leaves from shoots A and represent means ± 1 SE from n = 4. Results from the Tukey post-hoc tests following the 2-way ANOVA are shown as small letter 714 715 code next to each treatment. If no letters are shown for a response variable, treatments were 716 not different to each other. Note that y-axes do not start with zero but are adapted to the data 717 range.

718

Fig. 6. Principal component analysis of the response of *Myriophyllum spicatum* to increasing
concentrations of arsenate in sediment (0, 1.44, 7 and 36 mg kg⁻¹), cultivated either in high or
low CO₂ availability (HC, LC), and after 14 days of exposure. A) Correlation circle showing
vectors for all measured endpoints (except DMC of whole shoots) as shown in Table 2 (see
list of abbreviations, aty - anthocyanins). B) PCA for all different treatment combinations.
Axes 1 and 2 explained 37.2 and 15.7 % of the total inertia.

















