



HAL
open science

**The 'Psychosocial Aspects in Hereditary Cancer'
questionnaire in women attending breast cancer genetic
clinics: Psychometric validation across French-,
German-and Spanish-language versions**

Anne Brédart, Amelie Anota, Julia Dick, Alejandra Cano, Antoine de Pauw,
Olivier Lareyre, Jean-Luc Kop, Neil Aaronson, Eveline Bleiker, Joan Brunet,
et al.

► **To cite this version:**

Anne Brédart, Amelie Anota, Julia Dick, Alejandra Cano, Antoine de Pauw, et al.. The 'Psychosocial Aspects in Hereditary Cancer' questionnaire in women attending breast cancer genetic clinics: Psychometric validation across French-, German-and Spanish-language versions. *European Journal of Cancer Care*, 2020, 29 (1), pp.e13173. 10.1111/ecc.13173 . hal-02275083

HAL Id: hal-02275083

<https://hal.univ-lorraine.fr/hal-02275083>

Submitted on 30 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The 'Psychosocial Aspects in Hereditary Cancer' questionnaire in women attending breast cancer genetic clinics: Psychometric validation across French-, German- and Spanish-language versions

Anne Brédart^{1,2}, Amélie Anota^{3,4}, Julia Dick⁵, Alejandra Cano⁶, Antoine De Pauw⁷, Olivier Lareyre¹, Jean-Luc Kop⁸, Neil K. Aaronson⁹, Eveline Bleiker⁹, Joan Brunet¹⁰, Peter Devilee¹¹, Dominique Stoppa-Lyonnet⁷, Rita Schmutzler⁵, Sylvie Dolbeault^{1,12}

¹Institut Curie, Supportive Care Department, Psycho-oncology Unit, 26 rue d'Ulm, 75005 Paris Cedex 05, Paris, France;

²University Paris Descartes, 71 avenue Edouard Vaillant, 92774 Boulogne-Billancourt, France;

³Methodology and Quality of Life in Oncology unit (INSERM UMR 1098), University Hospital of Besançon, France;

⁴French National Platform Quality of Life and Cancer, Besançon, France

⁵Familial Breast and Ovarian Cancer Centre, Cologne University Hospital and Faculty of Medicine, Kerpener Str. 34 | 50931, Cologne, Germany;

⁶University Autònoma of Barcelona, Clinical and Health Psychology Department, Carrer de Can'Altayó, s/n, 08193 Barcelona, Spain;

⁷Institut Curie, Cancer genetic clinic, 26 rue d'Ulm, 75005 Paris Cedex 05, France;

⁸Université de Lorraine, 2LPN (CEMA), 3 place Godefroy de Bouillon, 54 015 Nancy Cedex, Nancy, France;

⁹Division of Psychosocial Research and Epidemiology, The Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX, Amsterdam, The Netherlands;

¹⁰Catalan Institute of Oncology, Barcelona, Spain;

¹¹Netherlands Cancer Institute, Division of Molecular Pathology, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands ;

¹²CESP, University Paris-Sud, UVSQ, INSERM, University Paris-Saclay, 16 avenue Paul Vaillant-Couturier, 94807 Villejuif cedex, France.

Address for correspondence: Anne Brédart, PhD, Institut Curie, Psycho-Oncology Unit, 26 rue d'Ulm, 75.005 Paris Cedex 05, France. Telephone: +33-1-44 32 40 33, E-mail: anne.bredart@curie.fr

The 'Psychosocial Aspects in Hereditary Cancer' questionnaire in women attending breast cancer genetic clinics: Psychometric validation across French-, German- and Spanish-language versions

Abstract

Background

To design appropriate cancer genetic services for women at high breast cancer (BC) risk, it is essential to accurately assess their levels of genetic-specific psychosocial difficulties. We performed a comprehensive assessment of the psychometrics of the 'Psychosocial Aspects in Hereditary Cancer' (PAHC) questionnaire in French, German and Spanish.

Methods

Women consecutively approached in Cancer Genetic Clinics completed the PAHC, distress (HADS, IES-R) and satisfaction questionnaires at pre-testing (T1) and after the genetic test result disclosure (T2).

Results

Of 738 eligible counselees, 214 (90%) in France (Paris), 301 (92%) in Germany (Cologne) and 133 (77%) in Spain (Barcelona) completed the PAHC. A six-factor, revised PAHC model yielded acceptable CFA goodness-of-fit indexes and good all scales internal consistencies. PAHC scales demonstrated expected conceptual differences with distress and satisfaction with counselling. Different levels of psychosocial difficulties were evidenced between counselees' subgroups and over time (p values < 0.05). MID estimates based on changes in self-reported needs for additional counselling ranged from 8 to 15 for improvement and 9 to 21 for deterioration.

Conclusion

The PAHC French, German and Spanish versions are reliable and valid for evaluating the psychosocial difficulties of women at high BC risk attending genetic clinics.

Introduction

The hereditary breast and ovarian cancer (HBOC) syndrome is one of the most common hereditary cancer syndromes [Hiraki et al. 2014], suggested by a family history of breast or ovarian cancer, and caused by alterations in major BRCA1 or BRCA2 susceptibility genes [Couch et al. 2014]. Recently, other breast or ovarian cancer susceptibility genes have been identified [Taylor et al. 2018]. Simultaneous tests of multiple genes (gene panels) are now available and are currently being implemented in oncology practice [Domchek et al. 2013].

With these advances, genetic counselling has become increasingly complex. The information imparted may be difficult to interpret by counselees and potentially lead to false reassurance, inappropriate cancer risk management decisions and psychosocial distress [Bradbury et al. 2018]. There is a need to monitor the psychosocial difficulties that counselees may experience in this evolving genetic practice, identify their risk and protective factors, their impact on well-being, health behaviours and familial communication, and design and assess alternative models of genetic service delivery [Bradbury et al. 2018, Albada et al. 2012].

When tested for a highly penetrant cancer syndrome such as the HBOC and after careful pre-test counselling, most individuals do not appear to experience prolonged, clinically significant anxiety [Hamilton et al. 2009]. However, the majority of counselees experience psychosocial difficulties that are specific to the genetic context [Eijzenga et al. 2014a], which may then accentuate psychological distress [Farrelly et al. 2013] and elicit further counselling needs [Eijzenga et al. 2014a]. Addressing unmet psychosocial needs during the genetic consultation may reduce distress levels [Eijzenga et al. 2014b], and potentially result in better understanding of information provided, higher perceived control and satisfaction [Pieterse et al. 2005].

Based on the European Organisation for Research and Treatment of Cancer Quality of Life Group guidelines [Johnson et al. 2011], the “Psychosocial Aspects in Hereditary Cancer” (PAHC) questionnaire was recently developed to screen for the presence of genetic-specific psychosocial difficulties in clinical practice [Eijzenga et al. 2014a, 2014b]. Items of the PAHC address psychosocial concerns organized in six conceptually distinct domains. As this questionnaire may also be useful in observational studies and intervention trials, its dimensionality should also be demonstrated empirically.

The primary aim of the current study was to empirically test the PAHC questionnaire’s conceptual structure. In addition, we also assessed the questionnaire in terms of internal consistency, convergent and divergent validity, known-groups validity, responsiveness to change and interpretability (minimal important difference (MID)) [Mokkink et al. 2010].

This study was undertaken within the “Breast Cancer Risk after Diagnostic Gene Sequencing” (BRIDGES) consortium (<https://bridges-research.eu>), which provided the opportunity to investigate the PAHC’s psychometric performance in women attending genetics clinics for familial breast cancer in France, Germany and Spain.

Methods

This study protocol was approved in France by the Comité consultatif sur le traitement de l’information en matière de recherche dans le domaine de la santé (CCTIRS: Consultative committee for information management in health research – N°16.314) and by the Commission Nationale Informatique et Libertés (CNIL: French Information Technology and Privacy Commission), in Germany by the Ethics Committee of the University Hospital of Cologne (N°16-098) and, in Spain by the Ethics

Committee of the Instituto Catalán de Oncología of Barcelona (N° - PR111/16). All recruited women provided written informed consent.

Participants and procedure

From October 2016 to April 2018, women over the age of 18 years, unaffected or affected with a primary non-metastatic breast cancer, and eligible for BC gene panel or targeted testing were consecutively recruited in the Cancer Genetic Clinic of Institute Curie (France), the University Hospital of Cologne (Germany) and the Instituto Catalán de Oncología of Barcelona (Spain).

The primary analysis focused on evaluating the hypothesized scale structure of the 26-item PAHC. For confirmatory factor analysis, it is recommended to have a minimum of 5 cases per item in order to establish if there is a stable scale structure to the questionnaire [Tabachnik & Fidell, 2001]. Thus, we needed a minimum of 130 participants per country.

The study objectives were explained to the women on the day of the pre-test counselling visit (T1) and, when they agreed to participate, they were given questionnaires to complete at home, either on paper or online, and to return within the next two weeks. If necessary, one reminder was made by telephone call. Questionnaires not completed or received within 2 months after each genetic consultation were considered missing. Two months after the genetic test result disclosure consultation (T2) (i.e., between 1-8 months after T1), women were contacted again to complete the PAHC questionnaire following the same procedure.

Core measure

Psychosocial difficulties.

The PAHC questionnaire comprises six domains of psychosocial difficulties (i.e., hereditary predisposition, practical issues, family and social issues, living with cancer, general emotions, and children-related issues) [Eijzena et al. 2014a]. All items are scored on a four-point Likert scale with response options scored 1 (not at all), 2 (a little), 3 (quite a bit) and 4 (a lot). A non-applicable response option is also provided to items that may not be relevant in some circumstances (e.g., depending on the assessment time – pre- or post-testing, personal or familial actual cancer diagnosis). Additional PAHC items address counselees' needs for further counselling (yes/no) in relation to the domains and overall.

The PAHC was translated, adapted and pilot-tested into French, German and Spanish languages according to international guidelines [Kuliš et al. 2017]. A report on this process is available on the BRIDGES website (<https://bridges-research.eu/wp-content/uploads/2018/02/D52-PAHC-PU-v1.0.pdf>).

PAHC scale scores were transformed on a 0-100 scale, with higher scores for increasing difficulties. All questionnaire scale scores were computed replacing missing data by the mean score of the scale if a response was provided for more than 50% of the items in the scale; for children-related items, response frequencies were computed only on women with children.

Comparative measures

Psychological distress. The French [Razavi et al. 1990], German [Herrmann-Lingen et al. 2011] and Spanish [Herrero et al. 2003] versions of the Hospital Anxiety and Depression (HADS) were used to assess symptoms of anxiety and depression.

Specific distress. The Impact of Event Scale Revised (IES-R) [Weiss & Marmar, 1997], available in French [Brunet et al. 2003], German [Maerker & Schützwohl, 2000] and Spanish [Báguena et al. 2001] measures psychological reactions to a stressful event (i.e., in this case, referring to cancer risk).

Satisfaction with the genetic counselling visit [Pieterse et al. 2007] comprises eight items addressing expertise, communication, client-centeredness, information, responses to preferences, access, service organization and general satisfaction rated on a 10-point rating scale ranging from 1 (very bad) to 10 (excellent).

Needs for additional counselling were assessed by the specific PAHC items related to the six domains (as described above).

Additional data on sociodemographic characteristics were provided by the patient and clinical data were obtained from the medical record, including BC diagnosis status or type of genetic test result, i.e., positive pathogenic variant versus true negative or non-informative results.

Statistical analyses

Baseline sociodemographic and clinical characteristics of the participants were described using mean (standard deviation) or median (range) for continuous variables, and number (percentage) for qualitative variables. We used the Kruskal-Wallis test or Student's t-test for quantitative data and the Chi-square statistic for categorical. Effect sizes (Cohen's d) for group differences are given where values of 0.2 are considered small, 0.5 of moderate and 0.8 of large [Cohen, 1988].

Confirmatory factor analysis (CFA) was used to evaluate empirically the scale structure of the PAHC. We first tested a predefined conceptual model [Eijzena et al. 2014a]. Goodness of fit was evaluated using the χ^2 goodness-of-fit statistics and fit indexes: Root Mean Square Error of Approximation (RMSEA), Comparative Fit Index (CFI), Tucker-Lewis Index (TLI). Values <0.08 or <0.05 for the RMSEA, >0.90 or >0.95 for the CFI and TLI are indicative of acceptable or good fit between the hypothesised model and the observed data; the smallest likelihood ratio (χ^2/df), and Akaike Information criteria (AIC) was sought [Jackson et al. 2009]. Alternative measurement models were subsequently evaluated, where necessary.

Internal consistency reliability was calculated using Cronbach's coefficient alpha, with values >0.70 used as the criterion for acceptable internal consistency at the group level [Cronbach & Warrington, 1951].

Correlation analyses were performed for subscales of the PAHC with scores of HADS, IES-R scales and 'Satisfaction with the consultation' scales; correlations >0.50 were considered indicative of convergent validity and correlations $r < 0.35$ of divergent validity. Spearman or Pearson correlations were used, depending on the score distributions. We expected the distress scales to be highly correlated with the PAHC emotional scale [Farrelly et al. 2013, Brédart et al. 2005] (convergent validity) and moderately correlated (divergent validity) with other PAHC scales [Eijzena et al. 2015], and the PAHC scales and the 'satisfaction with the consultation' scale to be moderately correlated [Pieterse et al. 2005, Brédart et al. 2005] (divergent validity).

Known groups validity was evaluated based on hypothesized differences in PAHC scale scores as a function of sociodemographic and clinical characteristics. We hypothesized that younger counsees [Farrelly et al. 2013], those with a lower education, having children [Eijzena et al. 2015] or affected with BC [van Roosmalen et al. 2004] will present higher levels of difficulties; differences between

country settings were also expected due to case mix and counselling specificities (known-group differences).

Responsiveness was evaluated by calculating mean changes in PAHC scale scores between the two assessment-points and by genetic test result. We hypothesized that the PAHC scale scores would change significantly over time [Bennett et al. 2012] overall, and according to positive (BRCA1/2) or negative test results [Oberuggenberger et al. 2016, Lumish et al. 2017, Esteban et al. 2018] (responsiveness to change).

The minimal important difference (MID) was assessed for each PAHC scale based on an anchor-based counselee-derived approach [Jaeschke et al. 1989], using the six domain-specific counselee's self-reported needs for additional counselling at T1 and at T2 as anchors. A correlation of 0.30 or higher was considered as an acceptable association between the anchor and each scale at T1 [Revicki et al. 2008]. Each anchor was constructed considering three categories: no change (no need or persistent need at both measurement points), deteriorated (developed need at T2 not reported at T1), and improved (need reported at T1 not reported at T2). Subsequently, differences in PAHC scale scores were reported for each category of the anchor. To control for change observed for stable patients according to the anchor, MID estimates were obtained using the difference in two consecutive categories of the anchor, i.e. "improvement" versus "no change" and "no change" versus "deterioration" [Cella et al. 2002]. The 95% confidence intervals (CI) for the differences in mean change scores are reported. Anchor-based MID estimations were compared with the statistical distribution based method, based on fractions of 0.2, 0.3 and 0.5 of the PAHC scales' standard deviation (SD) at T1 and at T2, and where estimates <0.2 SD were considered too small [Maringwa et al. 2011].

Results

Patients' characteristics and completion rate

Among 239 counselees in France (Paris), 326 in Germany (Cologne) and 173 in Spain (Barcelona) consecutively approached, who accepted to participate in the study, 214 (90%), 301 (92%) and 133 (77%), respectively at T1, and 167 (70%), 221 (73%) and 67 (39%), respectively at T2, returned the PAHC questionnaire.

As shown in table 1, the mean age (standard deviation) of the French, German and Spanish respondents was 47.9 (11.9), 47.4 (10.7) and 47.7 (11.8) years, respectively and, 172 (80%), 253 (84%) and 86 (65%), respectively, were affected by BC.

Insert table 1 about here

There were no statistically significant differences observed between the 648 respondents and the 90 non-respondents in mean age (47.6 versus and 46.6 years), having children (75% versus 73%) or being affected by BC (79% versus 70%).

Questionnaire acceptability and item- descriptive statistics

Among the 648 respondents, 533 (83%) completed all PAHC items. The level of missing data per item was below 5% in all three countries. On a item scale score from 1 to 4, the mean item scores range from 1.17 to 3.29 (Supplementary material S1).

Confirmatory factor analyses

The data did not fit the originally hypothesized PAHC model well. Three successive, revised models were tested based on conceptual meaning and factor loadings. The best six-factor model yielded acceptable fit ($\chi^2/df = 4.64$; RMSEA = 0.075 [90%CI: 0.071-0.079]; CFI = 0.870, TLI = 0.848, AIC=34358.863) which could be improved (Supplementary material S2 and S3).

Examination of residuals evidenced pairs of items within the same factor with higher observed correlations than reproduced correlations. These pairs appeared semantically redundant [i.e. items 8 and 9 ($r_{\text{observed}}=0.65$; $r_{\text{reproduced}}=0.60$), 19 and 20 ($r_{\text{observed}}=0.56$; $r_{\text{reproduced}}=0.33$) and 23 and 26 ($r_{\text{observed}}=0.43$; $r_{\text{reproduced}}=0.30$)]. This model was thus modified allowing correlations between residuals of three redundant pairs of items within a same factor. This modified exploratory model provided an improved, more acceptable fit ($\chi^2/df = 3.64$, RMSEA = 0.061 [90%CI: 0.057-0.066], CFI = 0.91, TLI = 0.90).

Two new scales distinguish issues related to personal and familial cancer from the original 'Living with cancer' domain. The original 'Practical issues' domain was removed as one of the two items (item 7) had a low factor loadings (0.30). A table comparing the original hypothesized and newly proposed structure is provided in the supplementary material S2.

The six dimensions pertain to : 1) 'Hereditary predisposition' (items 1-4, 6); 2) 'Familial and social issues' (items 8-11, 13); 3) 'Emotions (items 14-18); 4) 'Familial cancer' (items 12, 19, 20, 23); 5) 'Personal cancer' (items 21, 22); 'Children-related issues' (items 24-26).

Reliability

The internal consistency of the PAHC scales was adequate, with Cronbach's alpha coefficients ranging from 0.71 ('Personal cancer') to 0.87 ('Emotions') (Supplementary material S1).

Convergent-divergent validity

As expected, the PAHC 'Emotions' correlated above 0.50 with the HADS and the IES-R scales. Correlations of other PAHC scales with HADS and IES-R ranged from 0.13 for 'Familial cancer' and HADS-Depression to 0.49 for 'Hereditary predisposition' and IES-R-Intrusion. All p values were below 0.001. The almost null correlations between the PAHC and satisfaction scales indicate that these scales address different concepts (Table 2).

Insert table 2 about here

Known-group differences

Except for 'Emotions', all scales of the PAHC were able to discriminate between counselees according to one or more of the clinical (BC diagnosis status), socio-demographic characteristics (age, level of education) or country criteria (p values < 0.05) (Table 3).

Insert table 3 about here

Responsiveness to change

Except for 'Familial and social issues' and 'Emotions', changes in mean scores of the PAHC scales over time were all statistically significant, ranging from -2.05 (p=0.034) ('Familial cancer') to -8.51 (p<0.0001) ('Hereditary predisposition') (Table 4). Analysis of change over time by genetic test result showed significant results only for ('Familial and social issues'), with a mean difference of 4.92 for women receiving a positive result compared to 0.20 for those with a negative result (p=0.04).

Insert table 4 about here

Minimal important difference (MID)

The mean change in scores of the PAHC scales by categories of the anchor is reported in Table 5. The deterioration category (i.e., occurrence of need for further counselling after the test result disclosure), comprised only few patients (ranging from 6 to 14). Differences in mean change between 'improvement' (i.e., resolution of need for further counselling after the test result disclosure) and 'no change' (i.e., no need or persistence of need at both time points) was statistically significant for each PAHC scale, with MID values varying from 8.42 for 'Personal cancer' to 14.86 for 'Emotions'. Differences in mean change between 'deterioration' and 'no change' were significant for 'Family and social issues' (21.05), 'Emotions' (9.07) and 'Personal cancer' (16.19).

Distribution of PAHC scores at T1 and T2 were similar (Supplementary material S4). The observed anchor-based MID values were close to 0.3 and 0.5 SD; the non-significant MID anchor-based values for deterioration in 'Familial cancer' and 'Children' was also reflected in more restricted score distribution, as below 0.2 SD.

Insert table 5 about here

Discussion

Our results provide evidence of adequate psychometric properties for the PAHC scales in women at high BC risk attending genetics clinics in France, Germany and Spain. The questionnaire proved to be acceptable to patients, with a low number of missing item responses across countries.

The PAHC conceptual model originally hypothesized was partially confirmed empirically in the current study. Our factor analysis led to make a distinction between items related to personal and familial cancer issues originally imbedded in the 'Living with cancer' domain; this reflects the difference between the impact of a hereditary cancer syndrome on those already affected by cancer versus on those who are unaffected at the time of genetic testing.

The new PAHC scale structure exhibits satisfactory index-of-fit and good internal consistency estimates. As expected [Farrelly et al. 2013, Eijzenga et al. 2015, Brédart et al. 2005], apart from the PAHC 'Emotions' scale, most PAHC scales exhibited only moderate correlations with the distress scales. All PAHC scales evidenced almost no correlations with satisfaction with the consultation. This may be explained, in our sample, as has been observed in other studies [Pieterse et al. 2005, Oberguggenberger et al. 2016], by a very high mean score (8.9 on a 1-10 scale) and little score variability (1.7) in the satisfaction scale scores.

Most PAHC scale scores were able to discriminate between counselees who differed in terms of age, education level, parental status, BC diagnosis, and country in the hypothesized direction. In addition, most PAHC scales evidenced significant change over time.

Differences in counselees' reported need for additional counselling at pre-test and after genetic test result disclosure served as the anchor to appraise the minimal important difference (MID). The mean change in PAHC scores was in the expected direction. Counselees classified as having improved by the anchor tended to present lower PAHC scores while those classified as having worsened by the anchor tended to present higher PAHC scores, and those classified as having not changed by the anchor generally presented PAHC scores close to 0. MID estimates varied between PAHC scales as has been reported for other patient-reported outcome measures [Maringwa et al. 2011]. The range of the anchor-based MID was 8 to 15 points for improvement, which is in line with the 5%-10% range of the instrument (5-10 points on 0-100 scale) [Maringwa et al. 2011]. For most PAHC scales, the anchor-based MID for improvement was close to 0.5 SD estimates, which reflects a 'moderate' effect size. Excluding values below 0.2 SD (too small effect size), anchor-based MID for deterioration ranged from 9 to 21 points; however further investigation is required considering the small number of counselees classified as having worsened by the anchor in this study.

Study limitations

We need to acknowledge a number of limitations of our study. First, although the PAHC was developed for any hereditary cancer syndrome, in this study, its psychometric performance was only addressed in women confronted with a HBOC syndrome. Moreover, the samples comprised primarily women opting for genetic testing and who were affected with BC. Although this reflects the current population of counselees in the participating centres (respondents did not differ on key available characteristics), our results may not generalize to broader populations.

Second, while we addressed most of the psychometric properties of the PAHC [Mokkink et al.

2010], it was not feasible in the context of this study to assess the temporal reliability (test-retest).

Third, MID was determined based on one anchor only. Although this anchor may be understandable and clinically relevant to clinicians and health managers in terms of estimating the need of personnel for genetic counselling service, a distinction can be made between reported difficulties and perceived need for psychosocial care [Brédart et al. 2013], and there are significant barriers to seeking psychosocial support [Sun et al. 2018].

Conclusions

The French, German and Spanish language versions of the PAHC questionnaire may be used to better understand counselees' psychosocial difficulties, highlight their needs, and potential gaps in genetic counselling services. This study provided an empirically derived PAHC scale structure, which are valid, reliable and responsive in assessing psychosocial difficulties in women attending genetic clinics for high BC risk.

References

- Albada, A., van Dulmen, S., Lindhout, D., Bensing, J. M., & Ausems, M. G. (2012). A pre-visit tailored website enhances counselees' realistic expectations and knowledge and fulfils information needs for breast cancer genetic counselling. *Familial Cancer, 11*(1), 85-95. doi:10.1007/s10689-011-9479-1
- Báguena, M. V., E & Beleña, María & Díaz Martinez, Amelia & Roldan, C & Reig, R. (2001). Psychometric properties of the Spanish version of the Impact of Event Scale-Revised (IES-R). *Analisis y Modificacion de Conducta, 27*, 581-604.(<https://www.uv.es/baguena/art-IES-R.pdf>)
- Bennett, P., Phelps, C., Hilgart, J., Hood, K., Brain, K., & Murray, A. (2012). Concerns and coping during cancer genetic risk assessment. *Psycho-Oncology, 21*(6), 611-617. doi:10.1002/pon.1938
- Bradbury, A. R., Patrick-Miller, L. J., Egleston, B. L., Hall, M. J., Domchek, S. M., Daly, M. B., . . . Yao, X. S. (2018). Randomized Noninferiority Trial of Telephone vs In-Person Disclosure of Germline Cancer Genetic Test Results. *Journal of the National Cancer Institute, 110*(9), 985-993. doi:10.1093/jnci/djy015
- Brédart, A., Kop, J. L., Griesser, A. C., Fiszer, C., Zaman, K., Panes-Ruedin, B., . . . Dolbeault, S. (2013). Assessment of needs, health-related quality of life, and satisfaction with care in breast cancer patients to better target supportive care. *Annals of Oncology, 24*(8), 2151-2158. doi:10.1093/annonc/mdt128
- Brunet, A., St-Hilaire, A., Jehel, L., & King, S. (2003). Validation of a French version of the impact of event scale-revised. *Canadian Journal of Psychiatry. Revue Canadienne de Psychiatrie, 48*(1), 56-61.
- Cella, D., Hughes, C., Peterman, A., Chang, C.-H., Peshkin, B. N., Schwartz, M. D., . . . Lerman, C. (2002). A brief assessment of concerns associated with genetic testing for cancer: The multidimensional impact of cancer risk assessment (MICRA) questionnaire. *Health Psychology, 21*(6), 564-572. doi:10.1037/0278-6133.21.6.564
- Cohen J. *Statistical power analysis for the behavioural sciences*. 2nd edn. New Jersey, Lawrence Earlbaum, 1988.
- Couch, F. J., Nathanson, K. L., & Offit, K. (2014). Two decades after BRCA: setting paradigms in personalized cancer care and prevention. *Science, 343*(6178), 1466-1470. doi:10.1126/science.1251827
- Cronbach, L. J., & Warrington, W. G. (1951). Time-limit tests: estimating their reliability and degree of speeding. *Psychometrika, 16*(2), 167-188.
- Domchek, S. M., Bradbury, A., Garber, J. E., Offit, K., & Robson, M. E. (2013). Multiplex genetic testing for cancer susceptibility: out on the high wire without a net? *Journal of Clinical Oncology, 31*(10), 1267-1270. doi:JCO.2012.46.9403 [pii]
- Eijzenga, W., Aaronson, N. K., Hahn, D. E., Sidharta, G. N., van der Kolk, L. E., Velthuisen, M. E., . . .
- Bleiker, E. M. (2014). Effect of Routine Assessment of Specific Psychosocial Problems on Personalized Communication, Counselors' Awareness, and Distress Levels in Cancer Genetic Counseling Practice: A Randomized Controlled Trial. *Journal of Clinical Oncology, 2*(27), 2998-

3004. doi:10.1200/JCO.2014.55.4576

Eijzenga, W., Bleiker, E. M., Hahn, D. E., Kluijdt, I., Sidharta, G. N., Gundy, C., & Aaronson, N. K. (2014). Psychosocial Aspects of Hereditary Cancer (PAHC) questionnaire: development and testing of a screening questionnaire for use in clinical cancer genetics. *Psycho-Oncology*, 23(8), 862-869. doi:10.1002/pon.3485

Eijzenga, W., Bleiker, E. M., Hahn, D. E., Van der Kolk, L. E., Sidharta, G. N., & Aaronson, N. K. (2015). Prevalence and detection of psychosocial problems in cancer genetic counseling. *Familial Cancer*, 14(4), 629-636. doi:10.1007/s10689-015-9809-9

Esteban, I., Vilaro, M., Adrover, E., Angulo, N., Carrasco, E., Gadea, N., . . . Balmana, J. (2018). Psychological impact of multi-gene cancer panel testing in patients with a clinical suspicion of hereditary cancer across Spain. *Psycho-Oncology*, 27(6), 1530-1537. doi:10.1002/pon.4686

Farrelly, A., White, V., Meiser, B., Jefford, M., Young, M. A., Ieropoli, S., . . . Duffy, J. (2013). Unmet support needs and distress among women with a BRCA1/2 mutation. *Familial Cancer*, 12(3), 509-518. doi:10.1007/s10689-012-9596-5

Herrero, M. J., Blanch, J., Peri, J. M., De Pablo, J., Pintor, L., & Bulbena, A. (2003). A validation study of the hospital anxiety and depression scale (HADS) in a Spanish population. *General Hospital Psychiatry*, 25(4), 277-283. doi:[https://doi.org/10.1016/S0163-8343\(03\)00043-4](https://doi.org/10.1016/S0163-8343(03)00043-4)

Hamilton, J. G., Lobel, M., & Moyer, A. (2009). Emotional distress following genetic testing for hereditary breast and ovarian cancer: a meta-analytic review. *Health Psychology*, 28(4), 510-518. doi:10.1037/a0014778

Hiraki, S., Rinella, E. S., Schnabel, F., Oratz, R., & Ostrer, H. (2014). Cancer risk assessment using genetic panel testing: considerations for clinical application. *J Genet Couns*, 23(4), 604-617. doi:10.1007/s10897-014-9695-6

Jackson, D. L., Gillaspay, J. A., & Purc-Stephenson, R. (2009). Reporting practices in confirmatory factor analysis: an overview and some recommendations. *Psychological Methods*, 14(1), 6-23. doi:10.1037/a0014694

Jaeschke, R., Singer, J., & Guyatt, G. H. (1989). Measurement of health status. Ascertaining the minimal clinically important difference. *Controlled Clinical Trials*, 10(4), 407-415.

Johnson, C., Blazeby, J.M., Bottomley, A., Fayers, P., Koller, K., Kuliš, D., JRamage, J., Sprangers, M., Velikova, G., Young, T.
Group Guidelines for developing questionnaire modules. EORTC Quality of Life Group, Bruxelles.

Kuliš, D., Bottomley, A., Velikova, G., Greimel, E., Koller, M. on behalf of the EORTC Quality of Life Group. (2017). EORTC Quality of Life Group Translation procedure. EORTC Quality of Life Group, Bruxelles.

Lee, A. J., Cunningham, A. P., Kuchenbaecker, K. B., Mavaddat, N., Easton, D. F., & Antoniou, A. C. (2014). BOADICEA breast cancer risk prediction model: updates to cancer incidences, tumour pathology and web interface. *British Journal of Cancer*, 110(2), 535-545. doi:10.1038/bjc.2013.730

Lumish, H. S., Steinfeld, H., Koval, C., Russo, D., Levinson, E., Wynn, J., . . . Chung, W. K. (2017). Impact of Panel Gene Testing for Hereditary Breast and Ovarian Cancer on Patients. *J Genet Couns*, 26(5), 1116-1129. doi:10.1007/s10897-017-0090-y

Maringwa, J., Quinten, C., King, M., Ringash, J., Osoba, D., Coens, C., . . . Brain Cancer, G. (2011). Minimal clinically meaningful differences for the EORTC QLQ-C30 and EORTC QLQ-BN20 scales in brain cancer patients. *Annals of Oncology*, 22(9), 2107-2112. doi:10.1093/annonc/mdq726

Mokkink LB, Terwee C, Patrick DL, Alonso J, Stratford PW, Knowl DL, Bouter LM, de Vet HCW. Cosmin checklist manual, February 2010, VU University Medical Center, Department of Epidemiology and Biostatistics, EMGO Institute for Health and Care Research, Amsterdam, The Netherlands (www.cosmin.nl

Oberguggenberger, A., Sztankay, M., Morscher, R. J., Sperner-Unterweger, B., Weber, I., Hubalek, M., . . . Meraner, V. (2016). Psychosocial outcomes and counselee satisfaction following genetic counseling for hereditary breast and ovarian cancer: A patient-reported outcome study. *Journal of Psychosomatic Research*, 89, 39-45. doi:10.1016/j.jpsychores.2016.08.005

Pieterse, A. H., Ausems, M. G., Van Dulmen, A. M., Beemer, F. A., & Bensing, J. M. (2005). Initial cancer genetic counseling consultation: change in counselees' cognitions and anxiety, and association with addressing their needs and preferences. *American Journal of Medical Genetics. Part A*, 137(1), 27-35. doi:10.1002/ajmg.a.30839

Pieterse, A. H., van Dulmen, A. M., Beemer, F. A., Bensing, J. M., & Ausems, M. G. (2007). Cancer genetic counseling: communication and counselees' post-visit satisfaction, cognitions, anxiety, and needs fulfillment. *J Genet Couns*, 16(1), 85-96. doi:10.1007/s10897-006-9048-1

Razavi, D., Delvaux, N., Farvacques, C., & Robaye, E. (1990). Screening for adjustment disorders and major depressive disorders in cancer in-patients. *British Journal of Psychiatry*, 156, 79-83.

Revicki D, Hays RD, Cella D, Sloan J (2008) Recommended methods for determining responsiveness and minimally important differences for patient-reported outcomes. *J Clin Epidemiol* 61:102– 109

Sun, K. S., Lam, T. P., Lam, K. F., Lo, T. L., Chao, D. V. K., & Lam, E. W. W. (2018). Enablers to seeking professional help for psychological distress—a study on Chinese primary care attenders. *Psychiatry Research*, 264, 9-14. doi:10.1016/j.psychres.2018.03.056

Taylor, A., Brady, A. F., Frayling, I. M., Hanson, H., Tischkowitz, M., Turnbull, C., . . . Group, U. K. C. G. (2018). Consensus for genes to be included on cancer panel tests offered by UK genetics services: guidelines of the UK Cancer Genetics Group. *Journal of Medical Genetics*. 55(6), 372-

377. doi:10.1136/jmedgenet-2017-105188

van Roosmalen, M. S., Stalmeier, P. F., Verhoef, L. C., Hoekstra-Weebers, J. E., Oosterwijk, J. C., Hoogerbrugge, N., . . . van Daal, W. A. (2004). Randomised trial of a decision aid and its timing for women being tested for a BRCA1/2 mutation. *British Journal of Cancer*, 90(2), 333-342. doi:10.1038/sj.bjc.6601525

Weiss DS, Marmar CR. The impact of event scale – revised. In: Wilson JP, Keane TM, editors. *Assessing psychological trauma and PTSD*. New York: Guilford Press; 1997. pp. 399–411.

Table 1. Socio-demographic and clinical characteristics across samples

	French Respondents (N=214)	German respondents (N=301)	Spanish respondents (N=133)
Age (years)			
Mean (SD)	47.9 (11.9)	47.4 (10.7)	47.7 (11.8)
Median (Q1-Q3)	48 (39 - 56)	48 (41 - 54)	48 (41 - 55)
Range	21-78	18-77	19-80
Education level (%)			
Medium education level or below	6 (2.83)	37 (12.37)	46 (34.59)
Secondary or superior education	206 (97.17)	262 (87.63)	87 (65.41)
Missing data	2	2	0
Marital status (%)			
Married/partnered	150 (70.4)	212 (71.1)	102 (77.3)
Others (widowed, separated/ divorced, single/never married)	63 (29.6)	86 (28.9)	30 (22.7)
Missing data	1	3	1
Having children (%)			
Yes	171 (79.9)	213 (70.8)	103 (77.4)
Personal breast cancer (%)			
Yes	172 (80.4)	253 (84.1)	86 (64.7)
Breast cancer lifetime risk			
BOADICEA estimates			
Mean (SD)	19.6 (11.9)	18.1 (9.2)	-
Median (range)	16.5 (0.7-81.1)	16.5 (0.7-81.1)	-

SD: standard deviation; Q1-Q3: first-third quartile.

Table 2. Correlation between the PAHC scales and, HADS, IES-R and Satisfaction with the consultation scales (N=648)

PAHC scales	HADS- Anxiety	HADS- Depression	IES- Hyperarousal	IES- Intrusion	IES- Avoidance	Satisfaction with consultation
Hereditary predisposition (HP)	0.42****	0.24****	0.44****	0.49****	0.40****	-0.01 NS
Familial & social issues (FSI)	0.45****	0.38****	0.41****	0.44****	0.35****	-0.04 NS
Emotions (E)	0.75****	0.57****	0.63****	0.66****	0.50****	0.0 NS
Familial cancer (FC)	0.31****	0.13***	0.24****	0.32****	0.27****	0.06 NS
Personal cancer (PC)	0.43****	0.36****	0.41****	0.47****	0.35****	0.03 NS
Children-related issues (CRI)	0.27****	0.17****	0.25****	0.30****	0.27****	0.06 NS

HADS: Hospital Anxiety and Depression Scale subscale scores; IES-R: Impact of Event Scale-Revised.
Note:

*correlations higher than |0.50| are in bold and lower than |0.35| in italic. NS=non-significant; *: p<0.05;*

*** : p<0.01; ***: p<0.001; ****: p<0.0001*

Table 3. Known-group comparisons

PAHC scales	Age			P value	Effect size
	Mean (SD)				
	<40, N=158	41-50, N=231	>50, N=259		
Hereditary predisposition (HP)	36.62 (21.65)	35.26 (25.17)	32.27 (24.45)	0.036	0.08
Familial & social issues (FSI)	15.13 (15.25)	14.91 (16.05)	15.93 (18.55)	0.809	0.009
Emotions (E)	35.01 (24.81)	30.05 (22.09)	30.18 (23.66)	0.102	0.08
Familial cancer (FC)	62.56 (27.05)	61.16 (25.90)	64.36 (25.70)	0.384	0.06
Personal cancer (PC)	55.84 (29.35)	59.49 (27.37)	56.47 (29.51)	0.429	0.05
Children-related issues (CRI)	50.24 (28.03)	50.78 (28.17)	45.80 (25.73)	0.234	0.09
	Level of education				
	Medium education level or below, N=89	Secondary or superior education, N=555			
Hereditary predisposition (HP)	39.57 (27.48)	33.63 (23.45)		0.056	0.23
Familial & social issues (FSI)	12.52 (16.42)	15.71 (16.67)		0.100	0.19
Emotions (E)	29.36 (23.78)	31.78 (23.71)		0.376	0.10
Familial cancer (FC)	68.36 (27.27)	61.80 (25.68)		0.029	0.25
Personal cancer (PC)	56.90 (32.60)	57.25 (28.15)		0.916	0.01
Children-related issues (CRI)	55.86 (27.09)	47.15 (26.95)		0.010	0.32
	Having children				
	Yes, N=487	No, N=161			
Hereditary predisposition (HP)	34.89 (24.79)	33.67 (22.12)		0.581	0.05
Familial & social issues (FSI)	16.02 (17.01)	13.70 (16.42)		0.134	0.14

Emotions (E)	31.07 (23.61)	33.19 (24.22)		0.329	0.09
Familial cancer (FC)	63.97 (25.86)	59.44 (26.12)		0.056	0.17
Personal cancer (PC)	57.90 (28.89)	55.42 (28.51)		0.344	0.09
Children-related issues (CRI)	48.64 (27.10)	--			
Affected/unaffected with BC					
	Yes, N=511	No, N=137			
Hereditary predisposition (HP)	33.27 (23.74)	39.51 (25.04)		0.007	0.26
Familial & social issues (FSI)	15.90 (17.06)	13.74 (16.16)		0.187	0.13
Emotions (E)	31.42 (23.78)	32.29 (23.79)		0.708	0.04
Familial cancer (FC)	60.63 (25.95)	71.15 (24.42)		<0.0001	0.42
Personal cancer (PC)	62.25 (26.28)	38.43 (30.19)		<0.0001	0.84

Children-related issues (CRI)	36.79 (30.92)	34.93 (33.94)		0.150	0.06
Country					
	German, N= 294	Spanish, N=129	French, N=210		
Hereditary predisposition (HP)	33.85 (24.12)	45.19 (26.56)	28.99 (20.26)	<0.0001	0.26
Familial & social issues (FSI)	16.76 (16.83)	11.84 (15.99)	15.84 (17.26)	0.0006	0.11
Emotions (E)	33.89 (24.27)	29.27 (23.23)	29.82 (23.17)	0.067	0.09
Familial cancer (FC)	51.94 (24.31)	79.99 (18.70)	67.65 (25.01)	<0.0001	0.48
Personal cancer (PC)	58.58 (28.65)	53.66 (31.72)	57.70 (26.97)	0.456	0.07
Children-related issues (CRI)	30.61 (29.96)	46.86 (33.68)	38.26 (30.69)	<.0001	0.25

issues (CRI)					
--------------	--	--	--	--	--

**Among different scales, sample sizes vary depending on the number of missing data; we provide: N = minimum sample size among the different scales; all PAHC scores are on a 0–100 scale; higher scores indicate higher expressed difficulties; values of 0.2, 0.5, 0.8 for effect size are considered small, moderate and large, respectively. P-value of T-tests for two sample or Kruskal-Wallis for three samples.*

Table 4. Responsiveness to change among patients over T1 and T2 assessment times overall and, by genetic test result

PAHC scales	T1	T2	T2-T1 [§]		Positive, N=90	Negative/Uninformative, N=405	
	Mean (SD)	Mean (SD)	Mean difference (SD)	P value	Mean difference (SD)	Mean difference (SD)	P value
Hereditary predisposition (HP)	33.03 (23.36)	24.52 (24.20)	-8.51 (24.52)	<0.0001	-3.94 (23.0)	-8.44 (23.35)	0.14
Familial & social issues (FSI)	14.72 (16.05)	15.05 (19.18)	0.34 (18.96)	0.712	4.92 (17.84)	0.20 (17.73)	0.04
Emotions (E)	31.34 (23.49)	29.51 (23.31)	-1.83 (20.52)	0.062	-0.12 (18.19)	-2.32 (20.10)	0.39
Familial cancer (FC)	62.17 (25.99)	60.12 (27.97)	-2.05 (20.45)	0.034	-0.34 (19.58)	-2.12 (20.25)	0.50
Personal cancer (PC)	57.90 (28.40)	53.73 (28.17)	-4.18 (24.88)	0.0004	0.68 (22.30)	-4.57 (25.67)	0.11
Children-related issues (CRI)	46.54 (26.40)	38.86 (26.86)	-5.76 (19.65)	<0.0001	-2.56 (19.35)	-8.26 (22.92)	0.10

[§]Patients responding at both assessment times; N range: 332 (Children) – 449 (Hereditary predisposition);

Positive test result N range: 50 (Children) – 73 (Familial Cancer, Personal Cancer);

Negative/uninformative test

result N range: 222 (Children)-296 (Hereditary predisposition). P-value of paired T-test.

Table 5. Change in need for help per PAHC scales over T1 and T2 assessment times and mean (SD) of PAHC change scores in three anchor-defined groups and the MID (95% CI) between no need and need

PAHC scales	Improved, no need anymore, N=48	No change, persistent need or no need, N=263	Deteriorate, developed need, N=6	MID (95% CI)* Difference in mean change	
				Improvement No need anymore – no change	Deterioration No change – developed need
Hereditary predisposition (HP)	-20.61 (28.15)	-6.09 (23.03)	1.07 (25.78)	-14.52 (-22.0; -7.04)	-7.16 (-20.88; 6.56)
Familial & social issues (FSI)	-8.24 (26.52)	0.71 (15.73)	21.76 (22.60)	-8.96 (-16.88; -1.04)	-21.05 (-31.62; -10.48)
Emotions (E)	-15.51 (24.30)	-0.65 (19.19)	8.42 (17.33)	-14.86 (-22.02; -7.70)	-9.07 (-17.12; -1.02)
Familial cancer (FC)	-10.91 (24.07)	-0.62 (19.58)	1.19 (20.37)	-10.29 (-16.38; -4.20)	-1.81 (-12.68; 9.06)
Personal cancer (PC)	-11.52 (29.82)	-3.10 (23.80)	13.10 (21.86)	-8.42 (-15.94; -0.90)	-16.19 (-27.92; -4.46)
Children- related issues (CRI)	-15.60 (31.89)	-5.92 (19.92)	-5.56 (18.26)	-9.68 (-19.11; -0.25)	-0.36 (-15.17; 14.45)

MID: Minimal clinical importance. 95% CI comprising 0 are statistically non-significant.