



## Massive gene flux in bacterial population fuels adaptation to forest soil ecosystem

Abdoul-Razak Tidjani, Jean-Noël Lorenzi, Olivier Lespinet, Cyril Bontemps, Pierre Leblond

### ► To cite this version:

Abdoul-Razak Tidjani, Jean-Noël Lorenzi, Olivier Lespinet, Cyril Bontemps, Pierre Leblond. Massive gene flux in bacterial population fuels adaptation to forest soil ecosystem. 15ème congrès de la Société Française de Microbiologie, Sep 2019, Paris 19ème, France. 2019. hal-02314235

**HAL Id: hal-02314235**

**<https://hal.univ-lorraine.fr/hal-02314235>**

Submitted on 11 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Massive gene flux in bacterial population fuels adaptation to forest soil ecosystem

Abdoul-Razak TIDJANI<sup>1</sup>, Jean-Noël LORENZI<sup>1/2</sup>, Olivier LESPINET<sup>2</sup>, Cyril BONTEMPS<sup>1</sup>, Pierre LEBLOND<sup>1</sup>

<sup>1</sup>Université de Lorraine, Inra, DynAMic, F-54000 Nancy, France, <sup>2</sup>Institut de Biologie Intégrative de la Cellule (I2BC), CEA, CNRS, Université Paris-Sud, UPSay, Gif-sur-Yvette cedex, France

## Introduction

As for plant or animal, **bacteria are naturally structured in populations**, *i.e.* individuals of the same species that interact together in the same environment.


The **strains of a population are not necessarily clones** and some can have specific genes that are not shared by all of the individuals of the population<sup>[1,2]</sup>. The real extent of this **population genome variation in soil bacteria remains poorly described**.

We aimed here to decipher by genome comparison **the proportion of the specific genes**, the **mechanisms shaping these variations** and their **potential roles** in a *Streptomyces* population.


***Streptomyces* are soil dwelling bacteria**. They have a linear chromosome and a prolific specialized metabolism (*e.g.* enzymes, antibiotics, antifungals...) often of high biotechnological and medical interests. They can play key roles in the plant rhizosphere by contributing to their growth, health and resilience.

## Strategy

**1. Sampling scheme:** To isolate a *Streptomyces* population we minimized the spatial scale by performing a micro-scale sampling (from grains of soil)



**2. Phylogenetic analyses** were then performed to select strains belonging to the same species (e.g. identical strains in 16S rDNA)



**3. The genomes of 11 strains of the population were sequenced and compared.**


The genomes are organized as linear chromosomes, some of them bearing large extrachromosomal elements

They are among the biggest bacterial genomes ever sequenced (ranging from **11.7 to 12.4 Mb**)

Phylogenomic analyses confirmed that the strains are closely related at the intra-specific level and derived from a recent common ancestor


Pairwise comparisons of genomes were performed to identify variable gene pools due to insertion or deletion events (**indels**)

### A large number of indels exists between the different strains of the population



The 11 sequenced genomes were compared with one reference chosen among the population (here RLB1\_8). Each bar represents a strain chromosome. Triangles above a chromosome represent an insertion of genes in the reference strain and those below, an insertion in the compared strain. The size of the triangle is proportional to the number of genes in the indel. The highlighted triangles show examples of diversity hotspots where different type of indels are present in the population.

### The population genetic diversity can have ecological impacts



## Conclusions

- This study revealed an unexpectedly genetic diversity in term of presence/absence of genes among a *Streptomyces* population.
- Massive gene fluxes can explain how such diversification can occur within short evolutionary time.
- The diversity observed between individuals can have a functional role and impacts the ecology of the whole population.

[1]Hao W., Golding G. B., (2006). The fate of laterally transferred genes: life in the fast lane to adaptation or death. *Genome research*, 16(5), 636-43.

[2]Wielgoss S., Didelot X., Chaudhuri R.R. et al. (2016); A barrier to homologous recombination between sympatric strains of the cooperative soil bacterium *Myxococcus xanthus*. *The ISME Journal*. 1-10.

[3]Vetsigian K., Jajoo R., and Kishony R., (2011). Structure and Evolution of *Streptomyces* Interaction Networks in Soil and In Silico. *PLoS Biol* 9 (10): e1001184. doi:10.1371/journal.pbio.1001184.