



HAL
open science

Serum and peritoneal exudate concentrations after high doses of β -lactams in critically ill patients with severe intra-abdominal infections

Lisa Leon, Philippe Guerci, Elise Pape, Nathalie Thilly, Amandine Luc, Adeline Germain, Anne-Lise Butin-Druoton, Marie-Reine Losser, Julien Birckener, Julien Scala-Bertola, et al.

► To cite this version:

Lisa Leon, Philippe Guerci, Elise Pape, Nathalie Thilly, Amandine Luc, et al.. Serum and peritoneal exudate concentrations after high doses of β -lactams in critically ill patients with severe intra-abdominal infections: an observational prospective study. *Journal of Antimicrobial Chemotherapy*, 2020, 75, pp.156-161. 10.1093/jac/dkz407 . hal-02350602

HAL Id: hal-02350602

<https://hal.univ-lorraine.fr/hal-02350602>

Submitted on 10 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Serum and peritoneal exudate concentrations after high doses of β -lactams in critically ill patients with severe intra-abdominal infections: an observational prospective study

Lisa Leon^{1,2}, Philippe Guerci^{1,2}, Elise Pape^{2,3}, Nathalie Thilly^{2,4}, Amandine Luc^{2,4}, Adeline Germain^{2,5}, Anne-Lise Butin-Druoton^{1,2}, Marie-Reine Losser^{1,2}, Julien Birckener^{1,2}, Julien Scala-Bertola^{2,3} and Emmanuel Novy^{1,2*}

¹Department of Anaesthesiology and Intensive Care Medicine, University Hospital of Nancy, Vandœuvre-Lès-Nancy, F-54511, France; ²University of Lorraine, F-54000, Nancy, France; ³Department of Clinical Pharmacology and Toxicology, University Hospital of Nancy, Nancy, F-54000, France; ⁴Plateforme d'aide à la recherche Clinique, University Hospital of Nancy, Vandœuvre-Lès-Nancy, F-54511, France; ⁵Department of Surgery, University Hospital of Nancy, Vandœuvre-Lès-Nancy, F-54511, France

*Corresponding author. E-mail: e.novy@chru-nancy.fr

Received 27 June 2019; returned 30 July 2019; revised 23 August 2019; accepted 24 August 2019

Background: Critically ill patients with severe intra-abdominal infections (IAIs) requiring surgery may undergo several pharmacokinetic (PK) alterations that can lead to β -lactam underdosage.

Objectives: To measure serum and peritoneal exudate concentrations of β -lactams after high doses and optimal administration schemes.

Methods: This observational prospective study included critically ill patients with suspicion of IAI who required surgery and a β -lactam antibiotic as empirical therapy. Serum and peritoneal exudate concentrations were measured during surgery and after a 24 h steady-state period. The PK/pharmacodynamic (PD) target was to obtain serum β -lactam concentrations of 100% $fT_{>4 \times MIC}$ based on a worst-case scenario (based on the EUCAST highest epidemiological cut-off values) before bacterial documentation (*a priori*) and redefined following determination of the MIC for the isolated bacteria (*a posteriori*). Registered with ClinicalTrials.gov (NCT03310606).

Results: Forty-eight patients were included with a median (IQR) age of 64 (53–74) years and a SAPS II of 40 (32–65). The main diagnosis was secondary nosocomial peritonitis. Piperacillin/tazobactam was the most administered β -lactam antibiotic (75%). The serum/peritoneal piperacillin/tazobactam ratio was 0.88 (0.64–0.97) after a 24 h steady-state period. Prior to bacterial documentation, 16 patients (33.3%) achieved the *a priori* PK/PD target. The identification of microorganisms was available for 34 patients (71%). Based on the MIC for isolated bacteria, 78% of the patients achieved the serum PK/PD target.

Conclusions: In severe IAIs, high doses of β -lactams ensured 100% $fT_{>4 \times MIC}$ in the serum for 78% of critically ill patients with severe IAIs within the first 24 h. In order to define optimal β -lactam dosing, the PK/PD target should take into account the tissue penetration and local ecology.

Introduction

Early surgery and appropriate antibiotic therapy are cornerstones of management of intra-abdominal infections (IAIs). Among the pivotal measures of optimal antibiotic therapy, adequate dosing and administration schemes are recommended to achieve pharmacokinetic (PK)/pharmacodynamic (PD) targets.¹ Insufficient serum antibiotic concentrations secondary to PK alterations are well described in critically ill patients with sepsis.^{2,3} In cases of IAIs,

impaired tissue penetration and presence of indwelling surgical drains may increase PK alterations.^{4–6}

β -Lactams are the first-line antimicrobial agents for IAIs.⁷ Current guidelines regarding the use of β -lactams for critically ill patients recommend higher doses at the onset of treatment and continuous infusion.⁸ There are few PK/PD studies focused on critically ill patients requiring emergent surgery, and data regarding peritoneal diffusion are scarce.^{9,10}

In the present study, we hypothesized that underdosage of β -lactams was likely to occur despite a high dosing regimen. The concentrations of β -lactams in the serum and in the peritoneal exudate of critically ill patients with IAI requiring surgical management were measured. The rates of PK/PD target attainment at the empirical phase and after bacterial documentation were assessed.

Methods

Study design and population

A single-centre observational prospective study was conducted in a surgical ICU at the University Hospital of Nancy, France. The study protocol was reviewed and approved by the Patient Protection Committee Sud Méditerranée (IDRCB: 2017-A00710-53) and was registered with ClinicalTrials.gov (NCT03310606). Written informed consent was obtained from the patients or their substitute decision-makers.

All consecutive patients aged ≥ 18 years admitted to the ICU with a suspected IAI were eligible if their empirical antibiotic treatment included a β -lactam (see the [Supplementary data](#) available at JAC Online) and if they underwent abdominal surgery.⁸ Patients (i) with inappropriate β -lactam administration, (ii) with a history of allergy to β -lactams or (iii) who were pregnant were excluded.

Clinical data collected from the electronic medical record included demographics, ICU hospitalization details, IAI characteristics and β -lactam exposure. Septic shock was defined according to Sepsis-3.¹¹

β -Lactam analysis in serum and peritoneal fluid

Serum and peritoneal samples were simultaneously collected at three different timepoints: peritoneal incision (T0); end of surgery (T1); and 24 h after the first sampling (T2). Peritoneal samples were sterilely collected after peritoneal incision with aspiration of 5 mL of peritoneal exudate (T0) and then drawn from intra-abdominal drains in the ICU (T1 and T2).

The total concentrations of β -lactams in the serum and peritoneal exudate were assessed using HPLC (see the [Supplementary data](#) available at JAC Online). The linearity ranged from 1 to 500 mg/L. The coefficients of variation for inter- and intra-assay precision were $<10\%$ and the accuracy was within 10% for all antibiotics. The free concentration was approximated using the literature estimates of the free unbound percentage of each β -lactam: 80% for cefotaxime, imipenem and piperacillin, and 10% and 100% for ceftriaxone and meropenem, respectively.⁸

PK/PD targets

In cases of undocumented IAI or empirical antibiotic therapy, the *a priori* PK/PD target was to achieve a serum concentration above $4 \times \text{MIC}$ [based on the EUCAST highest epidemiological cut-off (ECOFF) values].

In the case of documented IAI, the *a posteriori* PK/PD target was defined as a serum free concentration $>4 \times \text{MIC}$ for the isolated bacteria. The MIC of the β -lactam infused was determined using an automated system, i.e. the VITEK[®] 2 system (bioMérieux, Marcy-l'Étoile, France).

Overall, the PK/PD target was to obtain an adequate serum concentration at the three timepoints, corresponding to 100% of $T_{>4 \times \text{MIC}}$ in susceptible pathogens (see the [Supplementary data](#) available at JAC Online).

Outcomes

The primary outcome was to determine the percentage of patients who met the *a priori* and the *a posteriori* PK/PD targets.

Secondary outcomes included the impact of the factors associated with changes in PK/PD and the mortality at 28 days.

A subgroup analysis for piperacillin/tazobactam was performed to evaluate the correlation between serum and peritoneal concentrations and the concentration ratio.

Statistical analysis

The characteristics of the sample were described as percentages for categorical variables and as median (IQR) values for continuous variables. Parametric and non-parametric statistics were used to analyse risk factors of underdosage. Analysis of the explanatory factors of the T2 underdosing was performed using logistic regression models. Comparison tests between concentrations at T2 and T1 (on paired samples) were performed using the Wilcoxon signed rank test. Correlations were evaluated using the Spearman coefficient. The alpha risk was 5% for all analyses. These statistical analyses were performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA).

Results

Patients

From October 2017 to January 2019, 48 adult patients were prospectively included [Table 1 and Figure S1 (available as [Supplementary data](#) at JAC Online)]. All IAIs were secondary peritonitis. Thirty-nine (81%) were healthcare-associated IAIs. Piperacillin/tazobactam was the most administered β -lactam antibiotic (75%). The identification of microorganisms was available for 34 patients (71%). Among these, an MIC was available for 23 patients. The observed total serum and peritoneal piperacillin/tazobactam concentrations are presented in Figure 1 (those of other β -lactams are shown in Figure S2). Gram-negative species were the most identified bacteria. The MICs of piperacillin/tazobactam ranged from a minimum of 0.3 mg/L to a maximum of 8 mg/L. Overall, only 11% of bacteria isolated were MDR bacteria. Documented microorganisms are presented in Table S1.

Primary outcome

Overall, 16 patients (33%) achieved the *a priori* PK/PD target (Figure 1). In the 23 patients with documented infection and MIC determination, the *a posteriori* PK/PD target was achieved in 18 patients (78%). Among these 18 patients, 6 (33%) achieved both a *a priori* and a *a posteriori* PK/PD.

Secondary outcomes

In the bivariate analysis (Table S2), high creatinine clearance was associated with an underdosage ($P=0.01$). The presence of septic shock ($P=0.30$), the presence of acute renal failure ($P=0.40$), vascular filling ($P=0.61$) and BMI ($P=0.29$) were not associated with underdosage. For patients who were still treated with norepinephrine after the end of surgery, less underdosage at T2 was observed ($P=0.04$).

In the multivariate analysis, age ≤ 64 years increased the risk of underdosage at T2 by 9.82-fold (OR=9.82; 95% CI=1.84–52.39; $P=0.008$) (Table S2).

Correlation between piperacillin/tazobactam serum and peritoneal concentrations

At T1, a trend of correlation was observed, with the Spearman coefficient calculated at 0.73 ($P<0.0001$) (Figure S3). The median

Table 1. Demographics and clinical data for included patients; N=48

Demographic data and medical history	
age (years), median (IQR)	64 (54–74)
female, n (%)	12 (25)
BMI (kg/m ²), median (IQR)	25.3 (20.5–30.3)
arterial hypertension, n (%)	23 (48)
chronic renal failure ^a , n (%)	14 (29)
ischaemic heart disease, n (%)	10 (21)
IAI: type and localization, n (%)	
healthcare associated	39 (81)
localization	
colon	17 (35)
small intestine	14 (29)
gastric	6 (13)
hepato-biliary	6 (13)
multiple localizations	4 (8)
pancreatic	1 (2)
β-Lactams administered, n (%)	
piperacillin/tazobactam	36 (75)
carbapenem	3 (6)
imipenem	1 (2)
meropenem	2 (4)
third-generation cephalosporin	9 (19)
ceftriaxone	6 (13)
cefotaxime	3 (6)
β-Lactam total concentrations (mg/L), serum/peritoneal exudate, median (IQR)	
piperacillin concentrations	
T0	107.9 (62.2–179.9)/47.3 (19.1–84.5)
T1	70 (43.8–103.5)/69.3 (41.4–112.7)
T2	88 (51.5–116.3)/77.4 (56.8–108.7)
ceftriaxone concentrations	
T0	99.6 (78.3–135.7)/71.5 (33.9–112.8)
T1	58.8 (38.1–75.6)/60.1 (31.2–77.3)
T2	32.2 (19.2–70.6)/55.9 (27.8–91.4)
cefotaxime concentrations	
T0	–/–
T1	114.5 (55.4–135.9)/71.1 (36.8–124.5)
T2	49 (45.9–156.6)/74.7 (56–93.5)
imipenem concentrations	
T0	212
T1	160
T2	58
meropenem concentrations	
T0	26.8 (10.5–43.2)/23.6 (10.1–37.2)
T1	17.4 (5.7–29.2)/21.1 (8.8–33.5)
T2	10 (3.8–32)/11.9 (10.7–54.6)
Clinical and biological data prior to surgery	
leucocytes (×10 ⁹ /L), median (IQR)	13.8 (9.4–19)
protein at T2 (g/L), median (IQR)	50 (46–55)
C-reactive protein (mg/L), median (IQR)	193 (123–312)
norepinephrine infusion, n (%)	13 (27)
temperature (°C), median (IQR)	37.5 (37–38)
Intraoperative data	
open surgery, n (%)	44 (92)
vascular filling (mL), median (IQR)	2000 (1375–2250)

Continued

Table 1. Continued

norepinephrine maximal dose ($\mu\text{g}/\text{kg}/\text{min}$), median (IQR)	0.3 (0.1–0.8)
septic shock, <i>n</i> (%)	11 (23)
ICU characteristics	
mechanical ventilation at ICU admission, <i>n</i> (%)	24 (50)
SOFA score at ICU admission, median (IQR)	6 (2–10)
SAPS II, median (IQR)	40 (32–65)
measured creatinine clearance (mL/min) at 24 h ^b , median (IQR)	46.5 (20–87)
drain fluid output at T2 (mL), median (IQR)	200 (100–250)
mortality at day 28, <i>n</i> (%)	7 (15)
Microbiological data	
MIC of piperacillin/tazobactam, median (IQR)	4 (4–4)
microbiological identification in peritoneal exudate, <i>n</i> (%)	
monobacterial	7 (15)
polymicrobial	23 (48)
fungal	4 (8)
no identification	14 (29)

^aChronic renal failure was defined as glomerular filtration rate $<60 \text{ mL}/\text{min}/1.73 \text{ m}^2$ according to KDIGO recommendations (where KDIGO stands for Kidney Disease Improving Global Outcomes).

^bCreatinine clearance was measured using the formula (urinary creatinine \times urinary volume)/plasma creatinine.

ratio of serum to peritoneal exudate was 0.48 (0.24–0.82), 0.96 (0.88–1.4) and 0.88 (0.64–0.97) at T0, T1 and T2, respectively.

Discussion

Optimal β -lactam administration schemes ensured 100% $fT_{>4\times\text{MIC}}$ in 78% of critically ill patients with severe IAIs within the first 24 h. Considering solely a PK/PD target based on ECOFF values, this rate would have decreased to 33%. This therapeutic scheme resulted in a favourable concentration in the peritoneal exudate. Age ≤ 64 years and high creatinine clearance were identified as risk factors for target non-attainment.

Dhaese *et al.*¹² showed that in 253 critically ill patients, $<40\%$ of patients receiving continuous infusion of 16 g/day of piperacillin/tazobactam achieved the chosen PK/PD target. In their study, the PK/PD target was based on a worst-case scenario. One of the major findings of our study was the difference in target attainment between the *a priori* target (worst-case scenario) and the *a posteriori* target (documented MIC). The identification of microorganisms was available for 34 patients (71%) and the MIC for 23 patients. Consequently, the rate of target attainment in undocumented cases may be underestimated because most of the cases reached a concentration of unbound β -lactam $>16 \text{ mg}/\text{L}$ (corresponding to $4\times\text{MIC}$ of $4 \text{ mg}/\text{L}$). As one of the pivotal measures of antimicrobial stewardship, the consideration of local ecology and resistance patterns is crucial.¹³ In a recent survey, having access to local bacterial resistance data was reported in $>80\%$ of cases.¹⁴ Herein, the median MIC of piperacillin/tazobactam was calculated at 4 (4–4) mg/L and very few MDR bacteria were identified.

Another finding was the favourable diffusion of piperacillin/tazobactam in the peritoneal exudate, contrary to the findings

of a previous study.⁶ Indeed, while there was a large difference between serum and peritoneal concentrations (ratio of 0.48) at the onset, this difference was offset at T1 (ratio of 0.96) and T2 (ratio of 0.88) probably due to the use of continuous infusion. Further studies are needed to confirm the benefit from continuous infusion on peritoneal concentration. The other factors that could explain the favourable peritoneal diffusion were: (i) the type of IAI; (ii) a relatively low vascular filling; and (iii) a low drain fluid output.⁵

Limitations

First, because this study was a pragmatic and non-interventional study, we did not compare different dosing and/or administration schemes for the same antibiotic. Consequently, the rate of PK/PD target attainment observed here reflected real-life practices with an optimal dosing regimen. Second, free concentration was not measured but estimated. We acknowledge that the measurement of total antibiotic concentrations could lead to misinterpretation in the presence of hypoalbuminaemia, which is frequent in critically ill patients.^{15,16} Nevertheless, the most prescribed β -lactams were piperacillin/tazobactam and ceftriaxone. For piperacillin, the difference between estimated and measured free concentration was described in the case of low concentration (i.e. $<50 \text{ mg}/\text{L}$), which was not the case here. For ceftriaxone, which is highly protein bound, the estimated free concentration was thought to be underestimated.¹⁶ Consequently, the resulting bias related to the estimation of free concentrations in terms of PK/PD objectives can be considered low.

Conclusions

In severe IAIs, high doses of β -lactams ensured 100% $fT_{>4\times\text{MIC}}$ in the serum for 78% of critically ill patients within the first 24 h.




Figure 1. (a) Serum β -lactam PK/PD target attainment and total serum (b) and peritoneal (c) concentrations of piperacillin/tazobactam at each timepoint. (a) Difference in target attainment between *a priori* target (worst-case scenario) and *a posteriori* target (documented MIC, based on local ecology). ECOFF-based target: *a priori* PK/PD target attainment defined as a total concentration of β -lactams above $4 \times$ the MIC (based on ECOFF values and a worst-case scenario). MIC-based target: *a posteriori* PK/PD objective attainment defined as a free concentration above $4 \times$ the MIC for isolated bacteria. (b and c) The dotted line represents the ECOFF-based target (worst-case scenario) according to guidelines: the threshold was calculated at 80 mg/L corresponding to $4 \times$ the MIC for *Pseudomonas aeruginosa* (16 mg/L) and considering a free fraction of 80%. The dotted-dashed line represents the MIC-based target: the threshold was calculated at 20 mg/L corresponding to $4 \times$ the MIC of 4 mg/L and considering a free fraction of 80%. The boxplots represent median and IQR. C, serum concentration of β -lactam.

In order to define optimal β -lactam dosing, the PK/PD target should take into account the tissue penetration and local ecology.

Acknowledgements

We thank all operating room staff of the University Hospital of Nancy for their invaluable help in blood and fluid sampling.

Funding

This work was funded by the Department of Anaesthesiology and Intensive Care Medicine of the University Hospital of Nancy.

Transparency declarations

None to declare.

Supplementary data

Supplementary data, including Figures S1 to S3 and Tables S1 and S2, are available at JAC Online.

References

1 Roberts JA, Abdul-Aziz MH, Lipman J *et al.* Individualised antibiotic dosing for patients who are critically ill: challenges and potential solutions. *Lancet Infect Dis* 2014; **14**: 498–509.

- 2 Taccone FS, Laterre P-F, Dugernier T *et al.* Insufficient β -lactam concentrations in the early phase of severe sepsis and septic shock. *Crit Care* 2010; **14**: R126.
- 3 Dhaese SAM, Roberts JA, Carlier M *et al.* Population pharmacokinetics of continuous infusion of piperacillin in critically ill patients. *Int J Antimicrob Agents* 2018; **51**: 594–600.
- 4 Glatz T, Lederer A-K, Kulemann B *et al.* The degree of local inflammatory response after colonic resection depends on the surgical approach: an observational study in 61 patients. *BMC Surg* 2015; **15**: 108.
- 5 Adnan S, Paterson DL, Lipman J *et al.* Pharmacokinetics of β -lactam antibiotics in patients with intra-abdominal disease: a structured review. *Surg Infect (Larchmt)* 2012; **13**: 9–17.
- 6 Adnan S, Li JX, Wallis SC *et al.* Pharmacokinetics of meropenem and piperacillin in critically ill patients with indwelling surgical drains. *Int J Antimicrob Agents* 2013; **42**: 90–3.
- 7 Sartelli M, Weber DG, Ruppé E *et al.* Erratum to: antimicrobials: a global alliance for optimizing their rational use in intra-abdominal infections (AGORA). *World J Emerg Surg* 2017; **12**: 35.
- 8 Guilhaumou R, Benaboud S, Bennis Y *et al.* Optimization of the treatment with β -lactam antibiotics in critically ill patients—guidelines from the French Society of Pharmacology and Therapeutics (Société Française de Pharmacologie et Thérapeutique-SFPPT) and the French Society of Anaesthesia and Intensive Care Medicine (Société Française d'Anesthésie et Réanimation-SFAR). *Crit Care* 2019; **23**: 104.
- 9 Seguin P, Verdier MC, Chanavaz C *et al.* Plasma and peritoneal concentration following continuous infusion of cefotaxime in patients with secondary peritonitis. *J Antimicrob Chemother* 2009; **63**: 564–7.
- 10 Buijk S, Gyssens IC, Mouton JW *et al.* Pharmacokinetics of ceftazidime in serum and peritoneal exudate during continuous versus intermittent administration to patients with severe intra-abdominal infections. *J Antimicrob Chemother* 2002; **49**: 121–8.
- 11 Singer M, Deutschman CS, Seymour CW *et al.* The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016; **315**: 801–10.
- 12 Dhaese SAM, Thooft ADJ, Farkas A *et al.* Early target attainment of continuous infusion piperacillin/tazobactam and meropenem in critically ill patients: a prospective observational study. *J Crit Care* 2019; **52**: 75–9.
- 13 Timsit J-F, Bassetti M, Cremer O *et al.* Rationalizing antimicrobial therapy in the ICU: a narrative review. *Intensive Care Med* 2019; **45**: 172–89.
- 14 Delannoy M, Agrinier N, Charmillon A *et al.* Implementation of antibiotic stewardship programmes in French ICUs in 2018: a nationwide cross-sectional survey. *J Antimicrob Chemother* 2019; **74**: 2106–14.
- 15 Ulldemolins M, Roberts JA, Wallis SC *et al.* Flucloxacillin dosing in critically ill patients with hypoalbuminaemia: special emphasis on unbound pharmacokinetics. *J Antimicrob Chemother* 2010; **65**: 1771–8.
- 16 Wong G, Briscoe S, Adnan S *et al.* Protein binding of β -lactam antibiotics in critically ill patients: can we successfully predict unbound concentrations? *Antimicrob Agents Chemother* 2013; **57**: 6165–70.