



HAL
open science

Importance of digestive mucus and mucins for designing new functional food ingredients

Mounira Kebouchi, Zeeshan Hafeez, Yves Le Roux, Annie Dary-Mourot,
Magali Genay

► To cite this version:

Mounira Kebouchi, Zeeshan Hafeez, Yves Le Roux, Annie Dary-Mourot, Magali Genay. Importance of digestive mucus and mucins for designing new functional food ingredients. Food Research International, 2020, 131, pp.1-10. 10.1016/j.foodres.2019.108906 . hal-02450495

HAL Id: hal-02450495

<https://hal.univ-lorraine.fr/hal-02450495v1>

Submitted on 23 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Importance of digestive mucus and mucins for designing new**
2 **functional food ingredients**

3

4 Mounira Kebouchi ^(a, b), Zeeshan Hafeez ^(a), Yves Le Roux ^(b), Annie Dary-Mourot ^(a, c),

5 Magali Genay ^(a, c)*

6 ^(a) Université de Lorraine, CALBINOTOX, F-54000, Nancy, France

7 ^(b) Université de Lorraine, INRA, URAFPA, F-54000, Nancy, France

8 ^(c): Co-senior authors

9 ^(*) Corresponding author: Magali Genay. Composés Alimentaires : Biofonctionnalités et
10 Risques Neurotoxiques (CALBINOTOX), Faculté des Sciences et Technologies, B.P. 70239,
11 F-54506 Vandœuvre-lès-Nancy, France. *Phone: +33 (0) 3 72 74 51 90. E-mail:*
12 *magali.genay@univ-lorraine.fr*

13

14 **Abstract**

15 The mucus, mainly composed of the glycoproteins mucins, is a rheological substance
16 that covers the intestinal epithelium and acts as a protective barrier against a variety of
17 harmful molecules, microbial infection and varying lumen environment conditions.
18 Alterations in the composition or structure of the mucus could lead to various diseases such as
19 inflammatory bowel disease or colorectal cancer. Recent studies revealed that an exogenous
20 intake of probiotic bacteria or other dietary components (such as bioactive peptides and
21 probiotics) derived from food influence mucus layer properties as well as modulate gene
22 expression and secretion of mucins. Therefore, the use of such components for designing new
23 functional ingredients and then foods, could constitute a novel approach to preserve the
24 properties of mucus. After presenting some aspects of the mucus and mucins in the
25 gastrointestinal tract as well as mucus role in the gut health, this review will address role of
26 dietary ingredients in improving mucus / mucin production and provides new suggestions for
27 further investigations of how dietary ingredients / probiotics based functional foods can be
28 developed to maintain or improve the gut health.

29 **Keywords:** intestinal mucosa; mucins; digestive tract; probiotic bacteria; food ingredients;
30 functional foods

31

32 **1. Introduction**

33 The human digestive tract consists of a stable alliance among the epithelial barrier, the
34 immune system and the resident microbiota. Besides its implication in absorption and
35 assimilation of nutrients, exclusion of undigested food, protective role against harmful
36 molecules and pathogenic microorganisms, the contribution of the digestive tract to sensory
37 functions including the innate and adaptive immune responses have also been well
38 documented. The epithelium of the gastrointestinal tract (GIT) consists of enterocytes, goblet
39 cells, enteroendocrine cells, Paneth cells, microfold cells (M cells), cup and tuft cells, and is
40 coated with a gel layer, called mucus (Ahluwalia, Magnusson, & Lohman, 2017; McGuckin,
41 Thornton, & Whitsett, 2015). Furthermore, the human GIT also harbors more than 10^{13} of
42 resident bacteria per gram of feces colonizing the mucus layer (Sender, Fuchs, & Milo, 2016).
43 The number of microbial cells as well as the number of species varies throughout the GIT (Li
44 et al., 2015) with few resident bacteria in the stomach and small intestine (Bik et al., 2006),
45 and a dramatic increase in the distal ileum and the colon (Ermund, Schütte, Johansson,
46 Gustafsson, & Hansson, 2013; Li et al., 2015). The association of bacteria with the host have
47 led to co-evolution with mutual benefits and impacts (Bäckhed, Ley, Sonnenburg, Peterson, &
48 Gordon, 2005; Liang, Leung, Guan, & Au, 2018). Hence, the resident bacteria could influence
49 the composition and structuration of the mucus layer and finally the mucus properties
50 (Jakobsson et al., 2015).

51 The digestive mucus, mostly consisting of mucins, plays a major role in the good functioning
52 of the digestive tract and thereby the human health. Whatever the reasons, alterations in
53 composition or structure of the mucus result in a wide variety of diseases including
54 inflammatory bowel diseases or colorectal cancer (Johansson, Sjövall, & Hansson, 2013;
55 Velcich et al., 2002). Hence, mucins, such as MUC2 (“MUC” refers to human mucins

56 whereas “Muc” corresponds to mucins in mice and other animals), constitute a target for the
57 design of new therapeutic molecules because of their involvement in tumorigenesis (Aithal et
58 al., 2018; Li et al., 2018).

59 Scientific community and public are now more interested in natural products to maintain
60 consumer health. This is reflected by the ever-growing market of functional food products
61 containing health-promoting ingredients and / or bacteria (Douillard, & de Vos, 2019;
62 Vicentini, Liberatore, & Mastrocola, 2016). Therefore, it would be interesting to explore the
63 possibility of identifying functional ingredients able to modulate mucus production as the
64 drug treatments are usually associated with adverse effects. After a brief description of the
65 mucus, its beneficial role in maintaining human health and the control of this layer, this
66 review will address the effects of food ingredients on mucus or mucins production and the
67 possible ways to design / improve functional ingredients, a first step in developing functional
68 foods, summarizing thereby the recent insights in this domain.

69

70 **2. Overview of digestive mucus and its beneficial role**

71 Mucus, secreted by intestinal goblets cells, is characterized by its viscosity and
72 rheological properties. It consists in water ($\approx 95\%$), electrolytes, enzymes, salts, DNA, lipids,
73 growth factors, antibacterial peptides and proteins such as defensins, immunoglobulins,
74 lysozymes and trefoil factors (McGuckin, Thornton, & Whitsett, 2015), and especially
75 glycoproteins called mucins. It not only constitutes a lubricant facilitating the transport of the
76 chyme along the GIT (Corfield, 2015; Lai, Wang, Wirtz, & Hanes, 2009) but also serves as a
77 barrier that protects the epithelial mucosa against mechanical injuries as well as from harmful
78 enteric pathogenic bacteria, viruses and parasites (Birchenough, Johansson, Gustafsson,

79 Bergström, & Hansson, 2015; Ermund, Schütte, Johansson, Gustafsson, & Hansson, 2013;
80 Liévin-Le Moal & Servin, 2006; Pelaseyed et al., 2014). Moreover, it also plays a key role in
81 maintaining intestinal homeostasis (Faderl, Noti, Corazza, & Mueller, 2015).

82 Studies conducted in rodent models and humans have shown that the mucus organization and
83 composition differ along the GIT (Figure 1). In the small intestine, there is a single,
84 discontinuous and disorganized mucus layer which is continuously secreted from the crypt
85 openings (Atuma, Strugala, Allen, & Holm, 2001; Ermund, Schütte, Johansson, Gustafsson,
86 & Hansson, 2013; Pelaseyed et al., 2014). In the stomach and the colon, the mucus is thicker
87 and composed of two layers. The inner dense layer is strongly attached to the epithelium,
88 devoid of bacteria and is constantly renewed every 1-2 hours by surface goblet cells of the
89 colon (Ermund, Schütte, Johansson, Gustafsson, & Hansson, 2013; Gustafsson et al., 2012;
90 Johansson, 2012). The outer loose layer, facing the lumen, is disorganized, more hydrated and
91 thicker than the inner layer and is colonized by the commensal bacteria in the colon (Atuma,
92 Strugala, Allen, & Holm, 2001; Ermund, Schütte, Johansson, Gustafsson, & Hansson, 2013;
93 Hansson & Johansson, 2010; Johansson et al., 2008). It results from the inner layer probably
94 through the proteolytic cleavage occurring within the mucins (Johansson et al., 2008).

95 As major constituents of mucus, mucins represent large and highly glycosylated
96 proteins expressed in different organs. In the adult human digestive tract, a wide variety of
97 mucins are secreted at the epithelium surfaces (Figure 1). They are clustered into three distinct
98 subfamilies: i) secreted gel-forming mucins, ii) secreted non-gel forming mucins, iii)
99 membrane-bound mucins, and their physiological roles are described in Table 1 (Moniaux,
100 Escande, Porchet, Aubert, & Batra, 2001). Being highly glycosylated molecules, a high
101 proportion of mucin's molecular weight corresponds to carbohydrates (oligosaccharides or

102 glycan chains). Their structure described in Figure 2 is important for both viscoelasticity
103 (Demouveau, Gouyer, Gottrand, Narita, & Desseyn, 2018) and bioactivity of mucus (Petrou,
104 & Crouzier, 2018). Biosynthesis and secretion of all types of mucins present some similarities
105 such as N-glycosylation in the endoplasmic reticulum and O-glycosylation at PTS sequence
106 (proline, threonine and serine rich region) in the Golgi apparatus. Afterwards, membrane
107 bound mucins are inserted into the membrane *via* C-terminal transmembrane domain whereas
108 secreted mucins are oligomerized and packed into Goblet cell granules for secretion (Figure
109 3).

110 Mucins play a major role in the protection and functioning of different parts of the GIT. At
111 the oral sphere level, mucins of saliva, e.g. MUC5B and MUC7 are involved in maintenance
112 of its rheological properties and protect the floor of the mouth against mechanical forces
113 during eating and speaking. They also protect from different diseases by reducing bacterial
114 attachment and / or their biofilm formation capacity (Table 1 and Figure 1) (Amerongen,
115 Bolscher, & Veerman, 1995; Dawes et al., 2015; Plummer & Douglas, 2006; Stinson et al.,
116 1982; van't Hof, Veerman, Nieuw Amerongen, & Ligtenberg, 2014). Other mucins of oral
117 sphere (MUC1, MUC16 and MUC4) that line the duct of parotid, submandibular and minor
118 salivary glands are involved in cell signal transduction and binding of secreted mucins
119 MUC5B and MUC7 (Frenkel & Ribbeck, 2015b; Liu et al., 2002; O'Brien, Beard,
120 Underwood, & Shigemasa, 2002; Sengupta et al., 2001). In the rest of the GIT, the mucins
121 safeguard the cell surface epithelium from the endogenous toxic molecules and pathogenic
122 bacteria (Table 1 and Figure 1) (Allen & Flemström, 2005; Bevins, 2006). Consequently, they
123 prevent from different diseases such as ulcerative colitis and colon cancer (Johansson et al.,
124 2008; Sluis et al., 2006; Velcich et al., 2002). In addition, the alteration of mucus
125 glycosylation pattern impacts both the composition of intestinal microbiota and the gut

126 architecture in mouse models (Arike & Hansson, 2016; Sommer et al., 2014; Staubach et al.,
127 2012). For example, it has been demonstrated that the loss of mucin glycosylation especially
128 the Core 1–derived *O*-glycans (the predominant form of mucin *O*-glycans in mice) led to the
129 development of spontaneous colitis in a mouse model (Fu et al., 2011).

130

131 **3. Control of mucus layer, mucin synthesis and associated diseases**

132 Under normal physiological conditions, mucus is regularly renewed and this phenomenon
133 results from a balance between mucus degradation and synthesis. Intestinal bacteria take
134 important part in regenerating the mucus layer. The establishment of mucus-degrading
135 bacteria in the host intestine starts few months after birth and is completed gradually with age
136 (Palmer, Bik, DiGiulio, Relman, & Brown, 2007). This process is not only linked with the
137 individual age but also with the diet. Midtvedt et al. (1994) have observed that the
138 establishment of mucin degradation bacteria is delayed at least 4 months in breast-fed
139 children than those fed with formula milk (Midtvedt, Carlstedt-Duke, & Midtvedt, 1994).
140 Among intestinal bacteria known to have a mucolytic activity, *Akkermansia muciniphila* is
141 one of the bacteria having large enzymatic machinery for mucins oligosaccharides breakdown
142 (Derrien, Vaughan, Plugge, & de Vos, 2004; Derrien, Collado, Ben-Amor, Salminen, & de
143 Vos, 2008). In mouse, it is one of the first users of mucus (Berry et al., 2013). This bacterium
144 hydrolyzes mucins oligosaccharides and releases sugar residues such as N-acetylglucosamine,
145 N-acetylgalactosamine, fucose or sialic acid, which can be used by other bacteria such as
146 *Bifidobacterium infantis* (Derrien, Belzer, & de Vos, 2017; Garrido, Kim, German, Raybould,
147 & Mills, 2011) or *Bacteroides thetaiotaomicron* (For comprehensive review see Koropatkin,
148 Cameron, & Martens, 2012). Some other intestinal bacteria such as *B. bifidum*, *Ba. fragilis*

149 and *Ruminococcus gnavus* also present a mucolytic activity (Katoh et al., 2017; Macfarlane &
150 Gibson, 1991; Png et al., 2010).

151 The bacteria and / or their metabolites could modulate the expression and / or the secretion of
152 mucins (Liévin-Le Moal et al., 2002; Smirnova, Guo, Birchall, & Pearson, 2003), the
153 development of the mucus secreting goblet cells as well as the mucin glycosylation profile. It
154 has been shown that *Ba. thetaiotaomicron* influenced the development of goblet cells in
155 mono-associated rats by stimulating the expression of *klf4* gene encoding a transcription
156 factor involved in goblet cell differentiation. This study also reported that this bacterium
157 induced the expression of some genes encoding mucins (*muc2* and *muc4*) in the gnotobiotic
158 rats compared to germ free model. Furthermore, the presence of two metabolically
159 complementary bacteria, *Ba. thetaiotaomicron* and *Faecalibacterium prausnitzii* in
160 gnotobiotic rats modulates the mucin glycosylation and goblet cells development in the
161 colonic epithelium compared to germ free rats (Wrzosek et al., 2013).

162 Abnormal synthesis and / or processing of mucins leads to disruption of mucus structure and
163 imbalance of mucosal barrier function. This could favour the development of a pathogen
164 bacterium which becomes able to circumvent the protective function of the mucus barrier
165 (Navabi, Johansson, Raghavan, & Lindén, 2013). Likewise, the abnormal functioning of the
166 mucosal barrier leads to the development of intestinal diseases such as Inflammatory Bowel
167 Disease (Ulcerative colitis, Crohn's disease) and eventually colorectal cancer (Bergstrom et
168 al., 2017; Velcich et al., 2002; Wenzel et al., 2014). Patients with ulcerative colitis have
169 shown modifications in mucus layer properties, mucin gene expression, mucin proteins
170 synthesis as well as mucin's glycosylation structure (O-glycosylation, sulfation, and
171 silylation) leading thus to a deficient mucus barrier (Bergstrom et al., 2017; Yamamoto-

172 Furusho, Ascaño-Gutiérrez, Furuzawa-Carballeda, & Fonseca-Camarillo, 2015). For instance,
173 it has been reported that patients with active ulcerative colitis showed a significant decrease in
174 gene and protein expression and production of MUC20 in epithelial cells from mucosa as well
175 as submucosa (Yamamoto-Furusho, Ascaño-Gutiérrez, Furuzawa-Carballeda, & Fonseca-
176 Camarillo, 2015). In addition, in mice model lacking core 2 β 1,6-N-acetylglucosaminyl-
177 transferase enzyme which initiates core 2 type protein O-glycosylation in mucin protein, an
178 increase in intestinal permeability and susceptibility to experimental colitis (Dextran sulfate
179 sodium induced colitis) has been observed (Stone et al., 2009). Furthermore, in Muc2-
180 deficient mice, the bacterial contact with the epithelium explains the inflammation and the
181 development of the cancer in these animals (Johansson et al., 2008).

182

183 **4. Impact of food components on mucus and mucins**

184 As discussed before, the control of mucus layer plays an essential role in maintaining health
185 and should constitute a major target for conception of preventive strategies, including specific
186 diets. Although knowledge about the impact of food ingredients on the digestive mucus layer
187 remain relatively scarce. However, different dietary components such as milk proteins,
188 bacteria or their metabolites, bioactive molecules (released either by bacteria in certain
189 fermented foods or derived directly from food after digestion), fiber and short chain fatty
190 acids have been shown to modulate the mucin gene expression and secretion *via* interaction
191 with intestinal mucosa (Mack, Michail, Wei, McDougall, & Hollingsworth, 1999; Martinez-
192 Maqueda, Miralles, Cruz-Huerta, & Recio, 2013; Rokana, Singh, Mallappa, Batish, & Grover,
193 2016; Burger-van Paassen et al., 2009). Hence, the positive effects of different food

194 components on intestinal mucus properties particularly mucin modulation are discussed
195 hereafter.

196

197 **4.1. Effect of lactic acid and probiotic bacteria on mucin gene expression and** 198 **secretion**

199 Lactic Acid Bacteria (LAB) or probiotic bacteria present in fermented milk products can
200 modify mucus layer by regulating mucins expression. Such modulation has been shown to
201 occur either by direct contact of bacteria with mucosal surfaces or by bacterial products
202 (Liévin-Le Moal et al., 2002; Raja, Murali, Devaraj, & Devaraj, 2012; Smirnova, Guo,
203 Birchall, & Pearson, 2003). It has been demonstrated that *Lactobacillus. plantarum* 299v and
204 *Lb. rhamnosus* GG significantly increased the mucosal layer and mucosal integrity through
205 upregulated expression of MUC3 which inhibited adherence of the enterohaemorrhagic
206 *Escherichia coli* O157:H7 to intestinal epithelial cells HT29 (Mack, Ahrne, Hyde, Wei, &
207 Hollingsworth, 2003). In the same perspective, Matter and coworkers have also reported that
208 the expression of *MUC2* gene was induced at transcriptional and translational levels in Caco-
209 2 cell cultures by *Lb. rhamnosus* GG (Mattar et al., 2002). Similarly, intestinal integrity was
210 improved when mice were treated with *Lb. rhamnosus* CNCM I-3690. This was mainly due to
211 enhanced production of goblet cells and consequently increased expression of mucin MUC2
212 (Martín et al., 2019). The treatment of Caco-2 cells with a cow milk, initially fermented with
213 *Lb. paracasei* CBAL74 and heated to kill bacteria, led to increase of *MUC2* gene expression
214 and mucus layer thickness compared to non fermented milk (Paparo et al., 2018). Thus, this
215 study highlighted that the increased mucus production is not restricted to living probiotic
216 bacteria but their components could also preserve the mucus layer integrity. Similarly, it has
217 been shown that administration of milk fermented with *Lb. plantarum* MTCC 5690 enhanced

218 the intestinal barrier function by increasing expression of MUC2 in salmonella-infected mice
219 compared to those fed with unfermented probiotic milk (Rokana, Singh, Mallappa, Batish, &
220 Grover, 2016). Besides the use of individual probiotic strains or probiotic-fermented milk, the
221 oral administration of VSL#3 probiotic mixture (containing different species of
222 *Lactobacillus*, *Bifidobacterium*, and one species of *Streptococcus*) for seven days enhanced
223 by 60 % basal luminal mucin content in rats. In addition, VSL#3 directly inoculated into
224 colonic loops significantly enhanced Muc2 mRNA expression compared to Muc1 and Muc3
225 for which only a slight increase was observed (Caballero-Franco, Keller, De Simone, &
226 Chadee, 2007). Surprisingly, these results obtained *in vivo* were not reproducible *in vitro*
227 when authors incubated VSL#3 bacterial cultures with cultured LS 174T intestinal epithelial
228 cells suggesting that the cell culture conditions were not optimal for proper growth of
229 probiotic bacteria. On the other hand, the authors explored the fact that bacterial metabolites
230 could induce the mucin secretion in LS 174T cells. They reported that the bacterial secreted
231 products from VSL#3 contained in the conditioned media remarkably stimulated mucin
232 secretion and the stimulatory effect was similar to that obtained from the *in vivo* experiments
233 (Caballero-Franco, Keller, De Simone, & Chadee, 2007). More recently, this probiotic
234 mixture has been shown to reduce colonic inflammation and improve intestinal barrier
235 function in Muc2 mucin-deficient mice indicating that VSL#3 did not require an intact mucin
236 barrier to exert its beneficial effects (Kumar, Kisson-Singh, Coria, Moreau, & Chadee,
237 2016). Finally, it has been shown that despite poor mucus-adhesion, *S. thermophilus* promotes
238 mucus pathway in digestive tract of gnotobiotic rats by producing lactate which in turn
239 increased the number of goblet cells and amount of mucus inducer KLF4 transcription factor.
240 A significant increase in KLF4 protein due to lactate production by this bacterium was also
241 reported in HT29-MTX cells (Fernandez et al., 2018).

242

243 **4. 2. Impact of bioactive peptides on mucin expression and secretion**

244 In addition to bacteria, hydrolysates from milk proteins have been shown to affect mucin
245 expression and secretion in the intestinal mucosa (Table 2). Indeed, these proteins harbor
246 bioactive sequences released either after digestion (*in vivo* or *in vitro*) or under the effect of
247 proteolytic system of bacteria such as LAB during processing of fermented milks. These
248 proteins provide peptides with different positive health effects on the host, including blood
249 pressure lowering, immunomodulatory, antimicrobial, cholesterol-lowering and antioxidant
250 effects. Several studies performed either *in vitro* or *in vivo* have also reported their impacts on
251 the modulation of mucin secretion (Claustre et al., 2002; Han, Deglaire, Sengupta, &
252 Moughan, 2008 ; Plaisancié et al., 2013; Plaisancié et al., 2015; Trompette et al., 2003). It has
253 been shown that the luminal administration of casein and lactalbumin hydrolysates separately
254 in the isolated perfused rat jejunum induced release of mucins. Conversely, the administration
255 of native caseins did not significantly stimulate mucin secretion (Claustre et al., 2002).
256 Likewise, when rats were fed for 8 days with test diets enriched either with hydrolyzed casein
257 or a synthetic amino acid mixture as well as a protein-free diet, hydrolyzed casein enriched
258 diet significantly up-regulated the expression of Muc3 and Muc4 mRNA in the small
259 intestinal tissue and the colon, respectively, compared to other diets (Han, Deglaire, Sengupta,
260 & Moughan, 2008). Fernández-Tomé and colleagues also reported significantly higher fecal
261 mucin (*via* O-linked glycoproteins dosage in feces) content after 2- and 8-weeks
262 administration of casein hydrolysate in rats. Indeed, long term intake of casein hydrolysate
263 resulted in an increased expression of Muc2 and Muc3 mRNA in the ileum and the colon of
264 rats (Fernández-Tomé et al., 2017). Furthermore, mucin secretion was escalated when β -
265 casomorphin-7, an opioid peptide from bovine β -casein, was administered in the lumen of the

266 isolated vascularly perfused rat jejunum (Trompette et al., 2003). Further studies conducted
267 with β -casomorphin-7 in intestinal mucin-producing cells of rat (DHE) and human (HT29-
268 MTX) revealed an increased expression of different mucins in a time and a dose dependent
269 manner. In DHE cells, Muc2 and Muc3 expression was enhanced whereas in case of HT29-
270 MTX, an increase of transcription as well as secretion of MUC5AC was reported. The authors
271 concluded that this peptide contributed significantly to mucin secretion *via* a direct effect on
272 intestinal goblet cells and the activation of μ -opioid receptors (Zoghbi et al., 2006). Other
273 studies, performed with the human HT29-MTX cells, have also reported the modulation of
274 mucin gene expression and stimulation by the total peptide pool derived from a fermented
275 milk (yogurt) (Plaisancié et al., 2013; Plaisancié et al., 2015). Indeed, these peptides
276 stimulated *MUC2* and *MUC4* gene expression as well as mucin secretion in HT29-MTX cells
277 (Plaisancié et al., 2013). Interestingly, this study has identified peptide sequences carrying
278 biological activity for further tests in rat models. Thereby, the oral administration of the
279 peptide fraction (f94-123) of β -casein to pups enhanced the number of goblet cells and Paneth
280 cells along the small intestine which were associated with higher expression of intestinal
281 mucins (Muc2 and Muc4) and antibacterial factors (Plaisancié et al., 2013). Furthermore,
282 from the β -casein peptide sequence (f94-123), shorter peptides having different biological
283 activities were also identified. These peptides up-regulated MUC4 expression in HT29-MTX
284 cells whereas no effect was observed on the levels of secreted mucins MUC2 and MUC5AC.
285 In contrast, other identified shorter peptide sequences increased the gene expression of *MUC2*
286 and *MUC5AC* mucins (Plaisancié et al., 2015). In another study conducted in HT29-MTX
287 cells, the stimulatory effect of casein hydrolysate and isolated peptides on mucin production
288 was also investigated. Casein hydrolysate and casein derived peptides mainly α_{S1} -casein
289 (f143-149) (AYFYPEL) and α_{S1} -casein (f144-149) (YFYPEL) significantly enhanced the

290 expression of mucin gene encoding MUC5AC (over 1.7-fold basal level) after 4 h of
291 exposure. The authors reported that casein hydrolysate containing both above cited peptides
292 displayed a higher effect on mucin secretion as determined from mucin-like glycoproteins
293 quantification than individual peptides (Martinez-Maqueda, Miralles, Cruz-Huerta, & Recio,
294 2013). Moreover, whey-derived peptides, α -lactorphin and β -lactorphin with described or
295 probable opioid activity were also shown to induce secretion of MUC5AC mucin in HT29-
296 MTX cells. Concerning the gene expression level of this mucin, it was up-regulated by
297 α -lactorphin in these cells (Martinez-Maqueda et al., 2012). All these findings indicate that
298 peptides derived from yogurt may play an important role in protecting intestinal lumen from
299 damaging agents by modulating Paneth and mucus cells, which are known to play a crucial
300 role in intestinal protection (Plaisancié et al., 2013). Hence, the presence of bioactive peptides
301 including those probably released from LAB in dairy products from milk proteins could
302 thereby improve gut health in the host.

303 **4.3. Impact of other food ingredients on mucin expression and secretion**

304 Besides milk proteins, rye bran, oat bran and soybean hull (Table 2) were also shown
305 to increase goblet cell volume density in the proximal and distal small intestine of hamsters
306 (Lundin et al., 1993). Food fibers can also positively impact the mucus layer through the
307 production of SCFAs, such as acetate, propionate and butyrate products, which are the
308 metabolic products of the intestinal microbiota. Among SCFAs, butyrate appears to be the
309 most effective in stimulating mucus secretion in rat colon (Shimotoyodome, Meguro, Hase,
310 Tokimitsu, & Sakata, 2000). Moreover, the effect of propionate and butyrate on mucin gene
311 expression in human goblet cell-like LS 174T cells was also demonstrated. Butyrate at 1 mM,
312 as well as propionate induced an increase in MUC2 mRNA levels. This study has also
313 investigated the mechanisms that regulate butyrate-mediated effect on MUC2 synthesis. The

314 authors have revealed an active butyrate-responsive region within the MUC2 promoter and
315 have demonstrated that the regulation of MUC2 by butyrate is associated with increased
316 acetylation of histone H3 and H4 and methylation of H3 at the MUC2 promoter (Burger-van
317 Paassen et al., 2009). Similar results, i.e., stimulation of MUC2 mucin and increase in histone
318 3 acetylation were observed after treatment of LS 174T cells with 1-2 mM sodium butyrate.
319 The secreted mucin was detected by histological PAS staining of carbohydrate chains of
320 mucin and expression at the protein and mRNA levels was confirmed by immunoblotting
321 with anti-MUC2 antibody and real-time RT-PCR, respectively (Hatayama, Iwashita,
322 Kuwajima, & Abe, 2007). More recently, the variation of four mucin gene expression
323 (*MUC2*, *MUC3*, *MUC13*, and *MUC17*) in coculture of Caco-2/HT29-MTX cells was analyzed
324 by real-time RT-PCR in the presence of antioxidant compounds like chlorogenic acid,
325 epicatechin gallate, and quercetin (apple, tea, and coffee polyphenols, respectively). In
326 addition, the secretion level of MUC2 was also determined by enzyme-linked immunoassay
327 (ELISA) analysis. After exposing the cells to sub-lethal concentrations (10 μ M) of each of the
328 different polyphenolic compounds over a period of 48 h, it is reported that each phenolic
329 compound induces differential expression patterns of mucin genes and only a significant
330 increase in *MUC17* gene expression was observed especially with epicatechin gallate and
331 quercetin. However, results from ELISA analyses showed no significant differences in
332 secretion level of mucin MUC2 (Volstatova et al., 2019). In contrast, quercetin induced the
333 secretion of MUC2 and MUC5AC mucins in quercetin-treated LS 174T cells. Both mucins
334 were also detected at mRNA level. The secretion of these mucins in this cell line was
335 attributed to enhanced extracellular signal regulated kinase (ERK)1-2 activity as its inhibition
336 by the MEK inhibitor PD98059 reverted the stimulatory effects of quercetin. Besides LS
337 174T cells, quercetin treatment also enhanced the mRNA levels of MUC2 and MUC5AC *via*
338 ERK1-2 pathway in the human intestinal epithelial Caco-2 cells (Damiano et al., 2018).

339 **5. Research approaches to improve/design new functional ingredients/foods**

340 The increasing costs of health as well as the desire of numerous consumers to support a
341 preventive rather than a curative approach have created favourable conditions to develop
342 functional foods, which are claimed to display health specific properties because of the
343 presence of some specific functional and / or absence of some harmful components (Gupta &
344 Abu-Ghannam, 2012). Although the informations presented above remain scarce, the
345 evidence provided through the action of probiotics, bioactive peptides or some polyphenolic
346 compounds suggest that they may be valuable ingredients / tools for strengthening the
347 digestive mucus layer or mucus barrier that is central to consumer health. However, it must be
348 underlined that most of the work presented in this review focused on effects of tested
349 ingredients on mucins transcription (Table 2). Nevertheless, other parameters can also be
350 explored such as effects of food ingredients on i) the microbiota composition and equilibrium,
351 ii) the mucus producing cells and thus on the intestinal epithelium and / or iii) glycosylation
352 profile of mucins. The search of such ingredients constitutes a first step in designing
353 functional foods, i.e., before including them in a food matrix, study their bioavailability in this
354 matrix during digestion followed by their interactions with other constituents of the diet.

355 Two types of approaches can be used to identify functional ingredients directed towards
356 mucus and mucins. The first is to screen high throughput probiotic strains, bioactive peptides
357 or other dietary compounds for their adhesion capacity (for probiotics) and / or their ability to
358 induce the production of mucins (induction of transcription/translation/secretion) using cell
359 models. Indeed, several cell lines are known to exhibit the functional characteristics of
360 intestinal cells found in small and large intestines and express different mucins (Table 3).
361 Such screening can concern new probiotic strains or food ingredients, but also ingredients
362 already described for other properties such as the opioid bioactive peptides (see above). The

363 second approach consists of using animal models to explore the capacity of some ingredients
364 to increase the goblet cells and then the mucus production or modulate the microbiota and the
365 glycosylation profile of mucins, a path that seems insufficiently explored so far. This implies to
366 be able to characterize the microbiota rapidly and on cohorts, to determine which type of flore
367 is favored, how does it is maintained and what are the determinants. In addition; it could be
368 interesting to explore the relationship between food ingredients and the glycosylation state of
369 mucins.

370 To conclude, given the impact of diseases related to dysfunction of the digestive tract and
371 especially the amount of mucus and mucins produced, it is necessary to intensify research in
372 the field of functional ingredients with the prospect of exploiting them in the maintenance of
373 gut health.

374

375 **Acknowledgements/funding sources**

376 Mounira Kebouchi is the recipient of a PhD fellowship from the “Ministère de
377 l’Enseignement Supérieur et de la Recherche” of France.

378

379 **Declarations of interest:** none

380 **6. References**

381 Ahluwalia, B., Magnusson, M. K., & Lohman, L. (2017). Mucosal immune system of the
382 gastrointestinal tract: maintaining balance between the good and the bad. *Scandinavian*
383 *Journal of Gastroenterology* 52, 1185-1193.

384 Aithal, A., Rauth, S., Kshirsagar, P., Shah, A., Lakshmanan, I., Junker, W. M., ... & Batra, S.
385 K. (2018). MUC16 as a novel target for cancer therapy. *Expert Opinion on Therapeutic*
386 *Targets* 22, 675–686.

387 Albrecht, H., & Carraway, K. L. (2011). MUC1 and MUC4: Switching the Emphasis from
388 Large to Small. *Cancer Biotherapy and Radiopharmaceuticals* 26, 261–271.

389 Al-Hashimi, I., & Levine, M. J. (1989). Characterization of in vivo salivary-derived enamel
390 pellicle. *Archives of Oral Biology* 34, 289–295.

391 Allen, A., & Flemström, G. (2005). Gastroduodenal mucus bicarbonate barrier: protection
392 against acid and pepsin. *American Journal of Physiology-Cell Physiology* 288, C1-19.

393 Ambort, D., Johansson, M. E. V., Gustafsson, J. K., Ermund, A., & Hansson, G. C. (2012a).
394 Perspectives on Mucus Properties and Formation—Lessons from the Biochemical World.
395 *Cold Spring Harbor Perspectives in Medicine* 2.

396 Ambort, D., Johansson, M. E. V., Gustafsson, J. K., Nilsson, H. E., Ermund, A., Johansson,
397 B. R., ... Hansson, G. C. (2012b). Calcium and pH-dependent packing and release of the gel-
398 forming MUC2 mucin. *Proceedings of the National Academy of Sciences U.S.A.* 109, 5645–
399 5650.

400 Amerongen, A. V., Bolscher, J. G., & Veerman, E. C. (1995). Salivary mucins: protective
401 functions in relation to their diversity. *Glycobiology* 5, 733–740.

402 Arike, L., & Hansson, G. C. (2016). The Densely O-glycosylated MUC2 Mucin Protects the
403 Intestine and Provides Food for the Commensal Bacteria. *Journal of Molecular Biology* 428,
404 3221–3229.

405 Asker, N., Axelsson, M. A., Olofsson, S. O., & Hansson, G. C. (1998). Dimerization of the
406 human MUC2 mucin in the endoplasmic reticulum is followed by a N-glycosylation-
407 dependent transfer of the mono- and dimers to the Golgi apparatus. *Journal of Biological*
408 *Chemistry* 273, 18857–18863.

409 Atuma, C., Strugala, V., Allen, A., & Holm, L. (2001). The adherent gastrointestinal mucus
410 gel layer: thickness and physical state in vivo. *American Journal of Physiology-*
411 *Gastrointestinal and Liver Physiology* 280, G922-929.

412 Augeron, C., & Laboisse, C. L. (1984). Emergence of permanently differentiated cell clones
413 in a human colonic cancer cell line in culture after treatment with sodium butyrate. *Cancer*
414 *Research* 44, 3961–3969.

415 Bäckhed, F., Ley, R. E., Sonnenburg, J. L., Peterson, D. A., & Gordon, J. I. (2005). Host-
416 bacterial mutualism in the human intestine. *Science* 307, 1915–1920.

417 Bäckström, M., Ambort, D., Thomsson, E., Johansson, M. E. V., & Hansson, G. C. (2013).
418 Increased understanding of the biochemistry and biosynthesis of MUC2 and other gel-
419 forming mucins through the recombinant expression of their protein domains. *Molecular*
420 *Biotechnology* 54, 250–256.

421 Bafna, S., Kaur, S., & Batra, S. K. (2010). Membrane-bound mucins: the mechanistic basis
422 for alterations in the growth and survival of cancer cells. *Oncogene* 29, 2893–2904.

423 Bergstrom, K., Fu, J., Johansson, M. E., Liu, X., Gao, N., Wu, Q., ... Xia, L. (2017). Core 1-
424 and 3-derived O-glycans collectively maintain the colonic mucus barrier and protect against
425 spontaneous colitis in mice. *Mucosal Immunology* 10, 91-103.

426 Bernet-Camard, M. F., Coconnier, M. H., Hudault, S., & Servin, A. L. (1996). Differential
427 expression of complement proteins and regulatory decay accelerating factor in relation to
428 differentiation of cultured human colon adenocarcinoma cell lines. *Gut* 38, 248–253.

429 Berry, D., Stecher, B., Schintlmeister, A., Reichert, J., Brugiroux, S., Wild, B., ... Wagner,
430 M. (2013). Host-compound foraging by intestinal microbiota revealed by single-cell stable
431 isotope probing. *Proceedings of the National Academy of Sciences U.S.A.* 110, 4720–4725.

432 Bevins, C. L. (2006). Paneth cell defensins: key effector molecules of innate immunity.
433 *Biochemical Society Transactions* 34, 263–266.

434 Bik, E. M., Eckburg, P. B., Gill, S. R., Nelson, K. E., Purdom, E. A., Francois, F., ... Relman,
435 D. A. (2006). Molecular analysis of the bacterial microbiota in the human stomach.
436 *Proceedings of the National Academy of Sciences U.S.A.* 103, 732–737.

437 Birchenough, G. M. H., Johansson, M. E. V., Gustafsson, J. K., Bergström, J. H., & Hansson,
438 G. C. (2015). New developments in goblet cell mucus secretion and function. *Mucosal*
439 *Immunology* 8, 712–719.

440 Bobek, L. A., Tsai, H., Biesbrock, A. R., & Levine, M. J. (1993). Molecular cloning,
441 sequence, and specificity of expression of the gene encoding the low molecular weight human
442 salivary mucin (MUC7). *Journal of Biological Chemistry* 268, 20563–20569.

443 Brockhausen, I., Schachter, H., & Stanley, P. (2009). “O-GalNAc Glycans,” in *Essentials of*
444 *Glycobiology*, eds. A. Varki, R. D. Cummings, J. D. Esko, H. H. Freeze, P. Stanley, C. R.
445 Bertozzi, et al. (Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press). Available
446 at: <http://www.ncbi.nlm.nih.gov/books/NBK1896/> [Accessed June 20, 2018].

447 Bu, X.-D., Li, N., Tian, X.-Q., & Huang, P.-L. (2011). Caco-2 and LS174T cell lines provide
448 different models for studying mucin expression in colon cancer. *Tissue Cell* 43, 201–206.

449 Burger-van Paassen, N., Vincent, A., Puiman, P. J., van der Sluis, M., Bouma, J., Boehm, G.,
450 ... Renes, I. B. (2009). The regulation of intestinal mucin MUC2 expression by short-chain
451 fatty acids: implications for epithelial protection. *Biochemical Journal* 420, 211–219.

452 Caballero-Franco, C., Keller, K., De Simone, C., & Chadee, K. (2007). The VSL#3 probiotic
453 formula induces mucin gene expression and secretion in colonic epithelial cells.
454 *American Journal of Physiology-Gastrointestinal and Liver Physiology* 292, G315-322.

455 Claustre, J., Toumi, F., Trompette, A., Jourdan, G., Guignard, H., Chayvialle, J. A., &
456 Plaisancié, P. (2002). Effects of peptides derived from dietary proteins on mucus secretion in
457 rat jejunum. *American Journal of Physiology-Gastrointestinal and Liver Physiology* 283,
458 G521-528.

459 Corfield, A. P. (2015). Mucins: a biologically relevant glycan barrier in mucosal protection.
460 *Biochimica et Biophysica Acta* 1850(1), 236–252.

461 DDahiya, R., Lesuffleur, T., Kwak, K. S., Byrd, J. C., Barbat, A., Zweibaum, A., & Kim, Y.
462 S. (1992). Expression and characterization of mucins associated with the resistance to
463 methotrexate of human colonic adenocarcinoma cell line HT29. *Cancer Research* 52, 4655–
464 4662.

465 Damiano, S., Sasso, A., De Felice, B., Di Gregorio, I., La Rosa, G., Lupoli, G. A., ... Santillo,
466 M. (2018). Quercetin Increases MUC2 and MUC5AC Gene Expression and Secretion in
467 Intestinal Goblet Cell-Like LS174T *via* PLC/PKC α /ERK1-2 Pathway. *Frontiers in*
468 *Physiology* 9, 357.

469 Dawes, C., Pedersen, A. M. L., Villa, A., Ekström, J., Proctor, G. B., Vissink, A., ... Wolff,
470 A. (2015). The functions of human saliva: A review sponsored by the World Workshop on
471 Oral Medicine VI. *Archives of Oral Biology* 60, 863–874.

472 Demouveau, B., Gouyer, V., Gottrand, F., Narita, T., & Desseyn, J. L. (2018). Gel-forming
473 mucin interactome drives mucus viscoelasticity. *Advances in Colloid and Interface Science*
474 252, 69-82.

475 Derrien, M., Vaughan, E. E., Plugge, C. M., & de Vos, W. M. (2004). Akkermansia
476 muciniphila gen. nov., sp. nov., a human intestinal mucin-degrading bacterium. *International*
477 *Journal of Systematic and Evolutionary Microbiology* 54, 1469–1476.

478 Derrien, M., Collado, M. C., Ben-Amor, K., Salminen, S., & de Vos, W. M. (2008). The
479 Mucin degrader Akkermansia muciniphila is an abundant resident of the human intestinal
480 tract. *Applied and Environmental Microbiology* 74, 1646–1648.

481 Derrien, M., Belzer, C., & de Vos, W. M. (2017). Akkermansia muciniphila and its role in
482 regulating host functions. *Microbial Pathogenesis* 106, 171–181.

483 Desseyn, J.-L., Tetaert, D., & Gouyer, V. (2008). Architecture of the large membrane-bound
484 mucins. *Gene*, 410(2), 215–222.

485 Douillard, F. P., & de Vos, W. M. (2019). Biotechnology of health-promoting bacteria.
486 *Biotechnology Advances* 37, 107369.

487 Ermund, A., Schütte, A., Johansson, M. E. V., Gustafsson, J. K., & Hansson, G. C. (2013).
488 Studies of mucus in mouse stomach, small intestine, and colon. I. Gastrointestinal mucus
489 layers have different properties depending on location as well as over the Peyer's patches.
490 *American Journal of Physiology-Gastrointestinal and Liver Physiology* 305, G341-347.

491 Faderl, M., Noti, M., Corazza, N., & Mueller, C. (2015). Keeping bugs in check: The mucus
492 layer as a critical component in maintaining intestinal homeostasis. *IUBMB Life* 67, 275–285.

493 Fernández-Tomé, S., Martínez-Maqueda, D., Tabernero, M., Largo, C., Recio, I., & Miralles,
494 B. (2017). Effect of the long-term intake of a casein hydrolysate on mucin secretion and gene
495 expression in the rat intestine. *Journal of Functional Foods*. 33, 176–180.

496 Fernandez, N., Wrzosek, L., Radziwill-Bienkowska, J.M., Ringot-Destrez, B., Duviau, M.P.,
497 Noordine, M.L., ... Mercier-Bonin, M. (2018). Characterization of Mucus-Related Properties
498 of *Streptococcus thermophilus*: From Adhesion to Induction. *Frontiers in Physiology*, 9, 980

499 Fogh, J., Fogh, J. M., & Orfeo, T. (1977). One hundred and twenty-seven cultured human
500 tumor cell lines producing tumors in nude mice. *Journal of the National Cancer Institute* 59,
501 221–226.

502 Frenkel, E. S., & Ribbeck, K. (2015a). Salivary mucins in host defense and disease
503 prevention. *Journal of Oral Microbiology* 7.

504 Frenkel, E. S., & Ribbeck, K. (2015b). Salivary Mucins Protect Surfaces from Colonization
505 by Cariogenic Bacteria. *Applied and Environmental Microbiology* 81, 332–338.

506 Fu, J., Wei, B., Wen, T., Johansson, M. E., Liu, X., Bradford, E., ... Xia, L. (2011). Loss of
507 intestinal core 1-derived O-glycans causes spontaneous colitis in mice. *Journal of Clinical*
508 *Investigation* 121, 1657-1666.

509 Garrido, D., Kim, J. H., German, J. B., Raybould, H. E., & Mills, D. A. (2011).
510 Oligosaccharide Binding Proteins from *Bifidobacterium longum* subsp. *infantis* Reveal a
511 Preference for Host Glycans. *PLOS ONE* 6, e17315.

512 Godl, K., Johansson, M. E. V., Lidell, M. E., Mörgelin, M., Karlsson, H., Olson, F. J., ...
513 Hansson, G. C. (2002). The N Terminus of the MUC2 Mucin Forms Trimers That Are Held
514 Together within a Trypsin-resistant Core Fragment. *Journal of Biological Chemistry* 277,
515 47248–47256.

516 Göttke, M. U., Keller, K., Belley, A., Garcia, R. M., Hollingsworth, M. A., Mack, D. R., &
517 Chadee, K. (1998). Functional heterogeneity of colonic adenocarcinoma mucins for inhibition
518 of *Entamoeba histolytica* adherence to target cells. *Journal of Eukaryotic Microbiology* 45,
519 17S-23S.

520 Gum, J. R., Crawley, S. C., Hicks, J. W., Szymkowski, D. E., & Kim, Y. S. (2002). MUC17,
521 a Novel Membrane-Tethered Mucin. *Biochemical and Biophysical Research Communications*
522 291, 466–475.

523 Gupta, S., & Abu-Ghannam, N. (2012). Probiotic fermentation of plant based products:
524 possibilities and opportunities. *Critical Reviews in Food Science and Nutrition* 52, 183-199.

525 Gururaja, T. L., Ramasubbu, N., Venugopalan, P., Reddy, M. S., Ramalingam, K., & Levine,
526 M. J. (1998). Structural features of the human salivary mucin, MUC7. *Glycoconjugate*
527 *Journal* 15, 457-467.

528 Gustafsson, J. K., Ermund, A., Johansson, M. E. V., Schütte, A., Hansson, G. C., & Sjövall,
529 H. (2012). An ex vivo method for studying mucus formation, properties, and thickness in
530 human colonic biopsies and mouse small and large intestinal explants. *American Journal of*
531 *Physiology-Gastrointestinal and Liver Physiology* 302, G430-438.

532 Han, K. S., Deglaire, A., Sengupta, R., & Moughan, P. J. (2008). Hydrolyzed casein
533 influences intestinal mucin gene expression in the rat. *J. Agric. Food Chem.* 56, 5572-6.

534 Hansson, G. C., & Johansson, M. E. (2010). The inner of the two Muc2 mucin-dependent
535 mucus layers in colon is devoid of bacteria. *Gut Microbes* 1, 51-54.

536 Hatayama, H., Iwashita, J., Kuwajima, A., & Abe, T. (2007). The short chain fatty acid,
537 butyrate, stimulates MUC2 mucin production in the human colon cancer cell line, LS174T.
538 *Biochemical and Biophysical Research Communications* 356, 599-603.

539 Hattrup, C. L., & Gendler, S. J. (2008). Structure and function of the cell surface (tethered)
540 mucins. *Annual Review of Physiology* 70, 431-457.

541 Ho, S. B., Takamura, K., Anway, R., Shekels, L. L., Toribara, N. W., & Ota, H. (2004). The
542 adherent gastric mucous layer is composed of alternating layers of MUC5AC and MUC6
543 mucin proteins. *Digestive Diseases and Sciences* 49, 1598-1606.

544 Huet, G., Kim, I., de Bolos, C., Lo-Guidice, J. M., Moreau, O., Hemon, B., ... Degand, P.
545 (1995). Characterization of mucins and proteoglycans synthesized by a mucin-secreting HT-
546 29 cell subpopulation. *Journal of Cell Science* 108, 1275–1285.

547 Jakobsson, H. E., Rodríguez-Piñeiro, A. M., Schütte, A., Ermund, A., Boysen, P., Bemark,
548 M., ... Johansson, M. E. (2015). The composition of the gut microbiota shapes the colon
549 mucus barrier. *EMBO Reports* 16, 164–177.

550 Jarry, A., Merlin, D., Velcich, A., Hopfer, U., Augenlicht, L. H., & Laboisse, C. L. (1994).
551 Interferon-gamma modulates cAMP-induced mucin exocytosis without affecting mucin gene
552 expression in a human colonic goblet cell line. *European Journal of Pharmacology* 267, 95–
553 103.

554 Johansson, M. E. V., Phillipson, M., Petersson, J., Velcich, A., Holm, L., & Hansson, G. C.
555 (2008). The inner of the two Muc2 mucin-dependent mucus layers in colon is devoid of
556 bacteria. *Proceedings of the National Academy of Sciences U. S. A.* 105, 15064–15069.

557 Johansson, M. E. V. (2012). Fast renewal of the distal colonic mucus layers by the surface
558 goblet cells as measured by in vivo labeling of mucin glycoproteins. *PloS One*, 7, e41009.

559 Johansson, M. E. V., Sjövall, H., & Hansson, G. C. (2013). The gastrointestinal mucus system
560 in health and disease. *Nature Reviews Gastroenterology & Hepatology* 10, 352–361.

561 Jonckheere, N., Skrypek, N., Frénois, F., & Van Seuning, I. (2013). Membrane-bound
562 mucin modular domains: from structure to function. *Biochimie* 95, 1077–1086.

563 Juge, N. (2012). Microbial adhesins to gastrointestinal mucus. *Trends Microbiology* 20, 30–
564 39.

565 Katoh, T., Maeshibu, T., Kikkawa, K.-I., Gotoh, A., Tomabechi, Y., Nakamura, M., ...
566 Katayama, T. (2017). Identification and characterization of a sulfoglycosidase from
567 *Bifidobacterium bifidum* implicated in mucin glycan utilization. *Bioscience, Biotechnology,*
568 *and Biochemistry* 81, 2018–2027.

569 Kawashima, H. (2012). Roles of the gel-forming MUC2 mucin and its O-glycosylation in the
570 protection against colitis and colorectal cancer. *Biological and Pharmaceutical Bulletin* 35,
571 1637–1641.

572 Koropatkin, N. M., Cameron, E. A., & Martens, E. C. (2012). How glycan metabolism shapes
573 the human gut microbiota. *Nature Reviews Microbiology* 10, 323–335.

574 Kumar, M., Kisson-Singh, V., Coria, A. L., Moreau, F., & Chadee, K. (2016). Probiotic
575 mixture VSL#3 reduces colonic inflammation and improves intestinal barrier function in
576 *Muc2* mucin-deficient mice. *American Journal of Physiology-Gastrointestinal and Liver*
577 *Physiology* 312, G34–G45.

578 Lai, S. K., Wang, Y. Y., Wirtz, D., & Hanes, J. (2009). Micro- and macrorheology of mucus.
579 *Advanced Drug Delivery Reviews* 61, 86–100.

580 Lesuffleur, T., Barbat, A., Dussaulx, E., & Zweibaum, A. (1990). Growth adaptation to
581 methotrexate of HT-29 human colon carcinoma cells is associated with their ability to
582 differentiate into columnar absorptive and mucus-secreting cells. *Cancer Research* 50, 6334–
583 6343.

584 Lesuffleur, T., Kornowski, A., Augeron, C., Dussaulx, E., Barbat, A., Labois, C., &
585 Zweibaum, A. (1991). Increased growth adaptability to 5-fluorouracil and methotrexate of

586 HT-29 sub-populations selected for their commitment to differentiation. *International Journal*
587 *of Cancer* 49, 731–737.

588 Lesuffleur, T., Porchet, N., Aubert, J. P., Swallow, D., Gum, J. R., Kim, Y. S., ... Zweibaum,
589 A. (1993). Differential expression of the human mucin genes MUC1 to MUC5 in relation to
590 growth and differentiation of different mucus-secreting HT-29 cell subpopulations. *Journal of*
591 *Cell Science* 106, 771–783.

592 Lesuffleur, T., Roche, F., Hill, A. S., Lacasa, M., Fox, M., Swallow, D. M., ... Real, F. X.
593 (1995). Characterization of a mucin cDNA clone isolated from HT-29 mucus-secreting cells.
594 The 3' end of MUC5AC? *Journal of Biological Chemistry* 270, 13665–13673.

595 Li, H., Limenitakis, J. P., Fuhrer, T., Geuking, M. B., Lawson, M. A., Wyss, M., ...
596 Macpherson, A. J. (2015). The outer mucus layer hosts a distinct intestinal microbial niche.
597 *Nature Communications* 6, 8292.

598 Li, C., Zuo, D., Yin, L., Lin, Y., Li, C., Liu, T., & Wang, L. (2018). Prognostic value of
599 MUC2 expression in colorectal cancer: A systematic review and meta-analysis.
600 *Gastroenterology Research and Practice* 2018, 6986870

601 Liang, D., Leung, R. K.-K., Guan, W., & Au, W. W. (2018). Involvement of gut microbiome
602 in human health and disease: brief overview, knowledge gaps and research opportunities. *Gut*
603 *Pathogens* 10.

604 Liévin-Le Moal, V., Huet, G., Aubert, J.-P., Bara, J., Forgue-Lafitte, M.-E., Servin, A. L., &
605 Coconnier, M. H. (2002). Activation of mucin exocytosis and upregulation of MUC genes in

606 polarized human intestinal mucin-secreting cells by the thiol-activated exotoxin listeriolysin
607 O. *Cellular Microbiology* 4, 515–529.

608 Liévin-Le Moal, V., & Servin, A. L. (2006). The front line of enteric host defense against
609 unwelcome intrusion of harmful microorganisms: mucins, antimicrobial peptides, and
610 microbiota. *Clinical Microbiology Reviews* 19(2):315-337.

611 Liu, B., Lague, J. R., Nunes, D. P., Toselli, P., Oppenheim, F. G., Soares, R. V., ... Offner, G.
612 D. (2002). Expression of membrane-associated mucins MUC1 and MUC4 in major human
613 salivary glands. *Journal of Histochemistry & Cytochemistry* 50, 811–820.

614 Lundin, E., Zhang, J. X., Huang, C. B., Reuterving, C. O., Hallmans, G., Nygren, C., &
615 Stenling, R. (1993). Oat bran, rye bran, and soybean hull increase goblet cell volume density
616 in the small intestine of the golden hamster. A histochemical and stereologic light-microscopy
617 study. *Scandinavian Journal of Gastroenterology* 28(1):15-22.

618 Macao, B., Johansson, D. G. A., Hansson, G. C., & Härd, T. (2006). Autoproteolysis coupled
619 to protein folding in the SEA domain of the membrane-bound MUC1 mucin. *Nature*
620 *Structural & Molecular Biology* 13, 71–76.

621 Macfarlane, G. T., & Gibson, G. R. (1991). Formation of glycoprotein degrading enzymes by
622 *Bacteroides fragilis*. *FEMS Microbiology Letters* 61, 289–293.

623 Mack, D. R., Michail, S., Wei, S., McDougall, L., & Hollingsworth, M. A. (1999). Probiotics
624 inhibit enteropathogenic *E. coli* adherence in vitro by inducing intestinal mucin gene
625 expression. *American Journal of Physiology* 276, G941-950.

626 Mack, D. R., Ahrne, S., Hyde, L., Wei, S., & Hollingsworth, M. A. (2003). Extracellular
627 MUC3 mucin secretion follows adherence of *Lactobacillus* strains to intestinal epithelial cells
628 in vitro. *Gut* 52, 827–833.

629 Maher, D. M., Gupta, B. K., Nagata, S., Jaggi, M., & Chauhan, S. C. (2011). Mucin 13:
630 structure, function, and potential roles in cancer pathogenesis. *Molecular Cancer Research* 9,
631 531–537.

632 Maoret, J. J., Font, J., Augeron, C., Codogno, P., Bauvy, C., Aubery, M., & Laboisse, C. L.
633 (1989). A mucus-secreting human colonic cancer cell line. Purification and partial
634 characterization of the secreted mucins. *Biochemical Journal* 258, 793–799.

635 Martín, R., Chamignon, C., Mhedbi-Hajri, N., Chain, F., Derrien, M., Escribano-Vázquez, U.,
636 ... Langella, P. (2019). The potential probiotic *Lactobacillus rhamnosus* CNCM I-3690 strain
637 protects the intestinal barrier by stimulating both mucus production and cytoprotective
638 response. *Scientific Reports*, 9, 5398.

639 Martínez-Maqueda, D., Miralles, B., De Pascual-Teresa, S., Reverón, I., Muñoz, R., & Recio,
640 I. (2012). Food-derived peptides stimulate mucin secretion and gene expression in intestinal
641 cells. *Journal of Agricultural and Food Chemistry* 60, 8600–8605.

642 Martínez-Maqueda, D., Miralles, B., Cruz-Huerta, E., & Recio, I. (2013). Casein hydrolysate
643 and derived peptides stimulate mucin secretion and gene expression in human intestinal cells.
644 *International Dairy Journal* 32, 13-19.

645 Mattar, A. F., Teitelbaum, D. H., Drongowski, R. A., Yongyi, F., Harmon, C. M., & Coran,
646 A. G. (2002). Probiotics up-regulate MUC-2 mucin gene expression in a Caco-2 cell-culture
647 model. *Pediatric Surgery International* 18, 586–590.

648 McGuckin, M. A., Thornton, D. J., & Whitsett, J. A. (2015). "Chapter 14 - Mucins and
649 Mucus," in *Mucosal Immunology (Fourth Edition)*, eds. J. Mestecky, W. Strober, M. W.
650 Russell, B. L. Kelsall, H. Cheroutre, and B. N. Lambrecht (Boston: Academic Press), 231–
651 250.

652 Mehrotra, R., Thornton, D. J., & Sheehan, J. K. (1998). Isolation and physical
653 characterization of the MUC7 (MG2) mucin from saliva: evidence for self-association.
654 *Biochemical Journal* 334, 415–422.

655 Midtvedt, A. C., Carlstedt-Duke, B., & Midtvedt, T. (1994). Establishment of a mucin-
656 degrading intestinal microflora during the first two years of human life. *Journal of Pediatric*
657 *Gastroenterology and Nutrition* 18, 321–326.

658 Moniaux, N., Nollet, S., Porchet, N., Degand, P., Laine, A., & Aubert, J. P. (1999). Complete
659 sequence of the human mucin MUC4: a putative cell membrane-associated mucin.
660 *Biochemical Journal* 338, 325–333.

661 Moniaux, N., Escande, F., Porchet, N., Aubert, J. P., & Batra, S. K. (2001). Structural
662 organization and classification of the human mucin genes. *Frontiers in Bioscience* 1;6:D1192-
663 206.

664 Navabi, N., Johansson, M. E. V., Raghavan, S., & Lindén, S. K. (2013). Helicobacter pylori
665 infection impairs the mucin production rate and turnover in the murine gastric mucosa.
666 *Infection and Immunity* 81, 829–837.

667 Nilsson, H. E., Ambort, D., Bäckström, M., Thomsson, E., Koeck, P. J. B., Hansson, G. C., &
668 Hebert, H. (2014). Intestinal MUC2 mucin supramolecular topology by packing and release
669 resting on D3 domain assembly. *Journal of Molecular Biology* 426, 2567–2579.

670 O’Brien, T. J., Beard, J. B., Underwood, L. J., & Shigemasa, K. (2002). The CA 125 gene: a
671 newly discovered extension of the glycosylated N-terminal domain doubles the size of this
672 extracellular superstructure. *Tumour biology: the journal of the International Society for*
673 *Oncodevelopmental Biology and Medicine* 23, 154–169.

674 Ouellette, A. J. (2010). Paneth cells and innate mucosal immunity. *Current Opinion in*
675 *Gastroenterology* 26, 547–553.

676 Palmer, C., Bik, E. M., DiGiulio, D. B., Relman, D. A., & Brown, P. O. (2007). Development
677 of the human infant intestinal microbiota. *PLOS Biology* 5, e177.

678 Paparo, L., Aitoro, R., Nocerino, R., Fierro, C., Bruno, C., & Canani, R. B. (2018). Direct
679 effects of fermented cow’s milk product with *Lactobacillus paracasei* CBA L74 on human
680 enterocytes. *Beneficial Microbes* 9, 165–172. doi:10.3920/BM2017.0038.

681 Pelaseyed, T., Bergström, J. H., Gustafsson, J. K., Ermund, A., Birchenough, G. M. H.,
682 Schütte, A., ... Hansson, G. C. (2014). The mucus and mucins of the goblet cells and
683 enterocytes provide the first defense line of the gastrointestinal tract and interact with the
684 immune system. *Immunological Reviews* 260, 8–20.

685 Perez-Vilar, J., & Hill, R. L. (1999). The structure and assembly of secreted mucins. *Journal*
686 *of Biological Chemistry* 274, 31751–31754.

687 Perez-Vilar, J., & Mabolo, R. (2007). Gel-forming mucins. Notions from in vitro studies.
688 *Histology & Histopathology* 22, 455–464.

689 Petrou, G., & Crouzier, T. (2018). Mucins as multifunctional building blocks of biomaterials.
690 *Biomaterials Science* 6, 2282-2297.

691 Plaisancié, P., Claustre, J., Estienne, M., Henry, G., Boutrou, R., Paquet, A., & Léonil, J.
692 (2013). A novel bioactive peptide from yoghurts modulates expression of the gel-forming
693 MUC2 mucin as well as population of goblet cells and Paneth cells along the small intestine.
694 *Journal of Nutritional Biochemistry* 24, 213–221.

695 Plaisancié, P., Boutrou, R., Estienne, M., Henry, G., Jardin, J., Paquet, A., & Léonil, J.
696 (2015). β -Casein(94-123)-derived peptides differently modulate production of mucins in
697 intestinal goblet cells. *Journal of Dairy Research* 82, 36–46.

698 Plummer, C., & Douglas, C. W. I. (2006). Relationship between the ability of oral
699 streptococci to interact with platelet glycoprotein Iba α and with the salivary low-
700 molecular-weight mucin, MG2. *FEMS Immunology and Medical Microbiology* 48, 390–399.

701 Png, C. W., Lindén, S. K., Gilshenan, K. S., Zoetendal, E. G., McSweeney, C. S., Sly, L. I.,
702 ... Florin, T. H. (2010). Mucolytic bacteria with increased prevalence in IBD mucosa
703 augment in vitro utilization of mucin by other bacteria. *The American Journal of*
704 *Gastroenterology* 105, 2420–2428.

705 Raja, S. B., Murali, M. R., Devaraj, H., & Devaraj, S. N. (2012). Differential expression of
706 gastric MUC5AC in colonic epithelial cells: TFF3-wired IL1 β /Akt crosstalk-induced

707 mucosal immune response against *Shigella dysenteriae* infection. *Journal of Cell Science* 125,
708 703–713.

709 Rokana, N., Singh, R., Mallappa, R. H., Batish, V. K., & Grover, S. (2016). Modulation of
710 intestinal barrier function to ameliorate *Salmonella* infection in mice by oral administration of
711 fermented milks produced with *Lactobacillus plantarum* MTCC 5690-a probiotic strain of
712 Indian gut origin. *Journal of Medical Microbiology* 65, 1482–1493.

713 Rong, M., Rossi, E. A., Zhang, J., McNeer, R. R., van den Brande, J. M. H., Yasin, M., ...
714 Carraway, K. L. (2005). Expression and localization of Muc4/sialomucin complex (SMC) in
715 the adult and developing rat intestine: implications for Muc4/SMC function. *Journal of*
716 *Cellular Physiology* 202, 275–284.

717 Sender, R., Fuchs, S., & Milo, R. (2016). Revised estimates for the number of human and
718 bacteria cells in the body. *PLOS Biology* 14:e1002533

719 Sengupta, A., Valdramidou, D., Huntley, S., Hicks, S. J., Carrington, S. D., & Corfield, A. P.
720 (2001). Distribution of MUC1 in the normal human oral cavity is localized to the ducts of
721 minor salivary glands. *Archives of Oral Biology* 46, 529–538.

722 Sheng, Y. H., Lourie, R., Lindén, S. K., Jeffery, P. L., Roche, D., Tran, T. V., ... McGuckin,
723 M. A. (2011). The MUC13 cell-surface mucin protects against intestinal inflammation by
724 inhibiting epithelial cell apoptosis. *Gut* 60, 1661–1670.

725 Shimotoyodome, A., Meguro, S., Hase, T., Tokimitsu, I. & Sakata, T. (2000). Short chain
726 fatty acids but not lactate or succinate stimulate mucus release in the rat colon. *Comparative*
727 *Biochemistry and Physiology Part A Molecular and Integrative Physiology* 125, 525-31.

728 Sluis, M. V. der, Koning, B. A. E. D., Bruijn, A. C. J. M. D., Velcich, A., Meijerink, J. P. P.,
729 Goudoever, J. B. V., ... Einerhand, A. W. (2006). Muc2-Deficient Mice Spontaneously
730 Develop Colitis, Indicating That MUC2 Is Critical for Colonic Protection. *Gastroenterology*
731 131, 117–129.

732 Smirnova, M. G., Guo, L., Birchall, J. P., & Pearson, J. P. (2003). LPS up-regulates mucin
733 and cytokine mRNA expression and stimulates mucin and cytokine secretion in goblet cells.
734 *Cellular Immunology* 221, 42–49.

735 Sommer, F., Adam, N., Johansson, M. E. V., Xia, L., Hansson, G. C., & Bäckhed, F. (2014).
736 Altered Mucus Glycosylation in Core 1 O-Glycan-Deficient Mice Affects Microbiota
737 Composition and Intestinal Architecture. *PLOS ONE* 9, e85254.

738 Sonnenburg, J. L., Xu, J., Leip, D. D., Chen, C.-H., Westover, B. P., Weatherford, J., ...
739 Gordon, J. I. (2005). Glycan foraging in vivo by an intestine-adapted bacterial symbiont.
740 *Science* 307, 1955–1959.

741 Staubach, F., Künzel, S., Baines, A. C., Yee, A., McGee, B. M., Bäckhed, F., ... Johnsen, J.
742 M. (2012). Expression of the blood-group-related glycosyltransferase *B4galnt2* influences the
743 intestinal microbiota in mice. *The ISME Journal* 6, 1345–1355.

744 Stinson, M. W., Levine, M. J., Cavese, J. M., Prakobphol, A., Murray, P. A., Tabak, L. A., &
745 Reddy, M. S. (1982). Adherence of *Streptococcus sanguis* to salivary mucin bound to glass.
746 *Journal of Dental Research* 61, 1390–1393.

747 Stone, E. L., Ismail, M. N., Lee, S. H., Luu, Y., Ramirez, K., Haslam, S. M., ... Marth, J. D.
748 (2009). Glycosyltransferase Function in Core 2-Type Protein O Glycosylation. *Molecular and*
749 *Cellular Biology* 29(13), 3770-3782.

750 Tailford, L. E., Crost, E. H., Kavanaugh, D., & Juge, N. (2015). Mucin glycan foraging in the
751 human gut microbiome. *Frontiers in Genetics* 6, 81.

752 Toribara, N. W., Robertson, A. M., Ho, S. B., Kuo, W. L., Gum, E., Hicks, J. W., ... Kim, Y.
753 S. (1993). Human gastric mucin. Identification of a unique species by expression cloning.
754 *Journal of Biological Chemistry* 268, 5879–5885.

755 Trompette, A., Claustre, J., Caillon, F., Jourdan, G., Chayvialle, J. A., & Plaisancié, P. (2003).
756 Milk bioactive peptides and beta-casomorphins induce mucus release in rat jejunum. *Journal*
757 *of Nutrition* 133, 3499–3503.

758 van Klinken, B. J., Oussoren, E., Weenink, J. J., Strous, G. J., Büller, H. A., Dekker, J., &
759 Einerhand, A. W. (1996). The human intestinal cell lines Caco-2 and LS174T as models to
760 study cell-type specific mucin expression. *Glycoconjugate Journal* 13, 757–768.

761 van't Hof, W., Veerman, E. C. I., Nieuw Amerongen, A. V., & Ligtenberg, A. J. M. (2014).
762 Antimicrobial defense systems in saliva. *Monographs in Oral Science* 24, 40–51.

763 Velcich, A., Yang, W., Heyer, J., Fragale, A., Nicholas, C., Viani, S., ... Augenlicht, L.
764 (2002). Colorectal cancer in mice genetically deficient in the mucin Muc2. *Science* 295,
765 1726–1729.

766 Vicentini, A., Liberatore, L., & Mastrocola, D. (2016). Functional foods: Trends and
767 development of the global market. *Italian Journal of Food Science* 28, 338-351.

768 Volstatova, T., Marchica, A., Hroncova, Z., Bernardi, R., Doskocil, I., & Havlik, J. (2019).
769 Effects of chlorogenic acid, epicatechin gallate, and quercetin on mucin expression and
770 secretion in the Caco-2/HT29-MTX cell model. *Food Science & Nutrition* 7, 492–498.

771 Wenzel, U. A., Magnusson, M. K., Rydström, A., Jonstrand, C., Hengst, J., Johansson, M. E.,
772 ... Wick, M. J. (2014). Spontaneous colitis in Muc2-deficient mice reflects clinical and
773 cellular features of active ulcerative colitis. *PLoS One* 19;9(6):e100217.

774 Williams, S. J., McGuckin, M. A., Gotley, D. C., Eyre, H. J., Sutherland, G. R., & Antalis, T.
775 M. (1999). Two Novel Mucin Genes Down-Regulated in Colorectal Cancer Identified by
776 Differential Display. *Cancer Research* 59, 4083–4089.

777 Williams, S. J., Wreschner, D. H., Tran, M., Eyre, H. J., Sutherland, G. R., & McGuckin, M.
778 A. (2001). Muc13, a novel human cell surface mucin expressed by epithelial and hemopoietic
779 cells. *Journal of Biological Chemistry* 276, 18327–18336.

780 Wreschner, D. H., McGuckin, M. A., Williams, S. J., Baruch, A., Yoeli, M., Ziv, R., ...
781 Gordon, S. (2002). Generation of ligand-receptor alliances by “SEA” module-mediated
782 cleavage of membrane-associated mucin proteins. *Protein Science : A Publication of the*
783 *Protein Society* 11, 698–706.

784 Wrzosek, L., Miquel, S., Noordine, M.-L., Bouet, S., Joncquel Chevalier-Curt, M., Robert,
785 V., ... Thomas, M. (2013). Bacteroides thetaiotaomicron and Faecalibacterium prausnitzii
786 influence the production of mucus glycans and the development of goblet cells in the colonic
787 epithelium of a gnotobiotic model rodent. *BMC Biology* 11, 61.

788 Yamamoto-Furusho, J. K., Ascaño-Gutiérrez, I., Furuzawa-Carballeda, J., & Fonseca-
789 Camarillo, G. (2015). Differential expression of MUC12, MUC16, and MUC20 in patients
790 with active and remission ulcerative colitis. *Mediators of Inflammation* 659018.

791 Zoghbi, S., Trompette, A., Claustre, J., El Homsí, M., Garzón, J., Jourdan, G., ... Plaisancié,
792 P. (2006). β -Casomorphin-7 regulates the secretion and expression of gastrointestinal mucins
793 through a μ -opioid pathway. *American Journal of Physiology-Gastrointestinal and Liver*
794 *Physiology* 290, 1105-13.

795

Table 1 Mucins in the digestive tract, their localization, cell expression and physiological functions

796

Mucins	Expression in each digestive tract segment	Cell type expression (Johansson, Sjövall, & Hansson., 2013)	Function	References
Secreted mucins				
- Gel-forming mucins				
MUC2	Small intestine (jejunum and ileum), colon	Goblet cells, Paneth cells	Formation of a protective barrier, lubrication, entrapment	Bäckström, Ambort, Thomsson, Johansson, Sjövall & Hansson, 2013 Nilsson et al., 2014 Kawashima, 2012
MUC5AC	Stomach	Mucous cells	Formation of a protective barrier, lubrication, entrapment	Allen & Flemström, 2005 Ho et al., 2004
MUC5B	Salivary glands, submandibular glands (mouth)	Mucous cells, Goblet cells	Lubrication, pathogens entrapment	Al-Hashimi & Levine, 1989 Frenkel & Ribbeck, 2015a, 2015b
MUC6	Stomach, ileum	Mucous cells	Protection, lubrication, entrapment	Toribara et al., 1993 Ho et al., 2004
- Non gel-forming mucins				
MUC7	Salivary glands (mouth)	Mucous cells	Clearance of harmful substances and pathogenic bacteria	Bobek, Tsai, Biesbrock, & Levine, 1993; Frenkel & Ribbeck, 2015b; Mehrotra, Thornton, & Sheehan, 1998; Plummer & Douglas, 2006; Stinson et al., 1982
Cell-membrane bound mucins				
MUC1	All intestinal segments	Epithelial cells	Cellular signal transduction, adhesion properties to bacteria, barrier activity	Albrecht & Carraway, 2011 Bafna, Kaur, & Batra, 2010
MUC3A/B	Small intestine, colon	Enterocytes	Adhesion modulation and cell signaling	Bafna, Kaur, & Batra, 2010 Desseyn, Tetaert, & Gouyer, 2008
MUC4	Salivary glands, small intestine, colon	Epithelial cells, Goblet cells	Lubrication, epithelial cell differentiation	Albrecht & Carraway, 2011 Moniaux et al., 1999 Rong et al., 2005
MUC12	Colon	Enterocytes	Epithelial cell regulation	Desseyn, Tetaert, & Gouyer, 2008 Williams et al., 1999
MUC13	Small intestine, colon	Enterocytes	Epithelial barrier function	Maher, Gupta, Nagata, Jaggi, & Chauhan, 2011 Sheng et al., 2011 Williams et al., 2001
MUC16	Mouth	Epithelial cells	Apical surface protection	Desseyn Tetaert, & Gouyer, 2008 O'Brien, Beard, Underwood, & Shigemasa, 2002
MUC17	Stomach, duodenum, colon	Enterocytes	Extracellular matrix constituent, apical surface protection	Desseyn, Tetaert, & Gouyer, 2008 Gum, Crawley, Hicks, Szymkowski, & Kim, 2002

Food ingredients	Enzyme hydrolyzed /	Increased mucin expression / secretion	Study model	References
Casein hydrolysate	Pancreatin	Muc3, Muc4 mRNA	Male rats	Han, Deglaire, Sengupta, & Moughan 2008
Casein hydrolysate	Pepsin	Muc2, Muc3 mRNA	Male rats	Fernández-Tomé et al., 2017
β -casomorphin-7	Purified / synthesized	Mucins	Male rats	Trompette et al., 2003
β -casomorphin-7	Purified / synthesized	Muc2, Muc3 mRNA	Rat DHE cells	Zoghbi et al., 2006
β -casomorphin-7	Purified / synthesized	MUC5AC	Human HT29-MTX cells	Zoghbi et al., 2006
Fermented milk (Yogurt)	Commercial product	MUC2, MUC4	Human HT29-MTX cells	Plaisancié et al., 2013
β -casein (f94-123)	Chemically synthesized	MUC2, MUC4	Human HT29-MTX cells	Plaisancié et al., 2013
β -casein (f94-123)	Chemically synthesized	Muc2, Muc4 mRNA	Rat pups	Plaisancié et al., 2013
β -casein (f108-113), β -casein (f114-119)	Purified	MUC4	Human HT29-MTX cells	Plaisancié et al., 2015
β -casein (f94-108)	Purified	MUC2, MUC5AC mRNA	Human HT29-MTX cells	Plaisancié et al., 2015
Casein hydrolysate	Pepsin	MUC5AC mRNA	Human HT29-MTX cells	Martínez-Maqueda, Miralles, Cruz-Huerta, & Recio, 2013
α_{s1} -casein (f143-149) α_{s1} -casein (f144-149)	Synthesized	MUC5AC mRNA	Human HT29-MTX cells	Martínez-Maqueda, Miralles, Cruz-Huerta, & Recio, 2013
α -lactorphin	Synthesized	MUC5AC	Human HT29-MTX cells	Martinez-Maqueda et al., 2012
Sodium butyrate	Commercial	MUC2 mRNA	Human LS 174T	Burger-Van Paassen et al., 2009; Hatayama, Iwashita, Kuwajima, & Abe., 2007
Quercetin, Epicatechin	Commercial	MUC17 mRNA	Caco-2/HT29-MTX cells co-culture	Volstatova et al., 2019
Quercetin	Commercial	MUC2, MUC5AC	Human LS 174T, Human Caco-2	Damiano et al., 2018

799 **Table 3** Summary of most cell lines *in vitro* models used in bacterial adhesion studies

Cell lines	Origin and characteristics	<i>MUC</i> gene expressed	References
Caco-2 TC7	Human colorectal adenocarcinoma cell line which differentiate spontaneously <i>in vitro</i> forming a polarized cell monolayer. The cell monolayer is devoid of mucus	<i>MUC1, MUC3, MUC4, MUC2, MUC5AC, MUC6</i> and <i>MUC13</i>	Bernet-Camard, Coconnier, Hudault, & Servin, 1996; Bu, Li, Tian, & Huang, 2011; Fogh, Fogh, & Orfeo, 1977; Götcke et al., 1998; van Klinken et al., 1996
HT-29	Human colorectal adenocarcinoma cell line forming polarized cell monolayer in culture (enterocytes + goblet cells)	<i>MUC1, MUC2, MUC3</i>	Augeron & Laboisse, 1984; Dahiya et al., 1992
HT29-MTX	Human colorectal adenocarcinoma cell line deriving from HT-29 cell line after growth adaptation to methotrexate thus acquiring the goblet cell-like phenotype with a high capacity of mucus production	<i>MUC1, MUC2, MUC3, MUC5AC</i>	Huet et al., 1995; Lesuffleur, Barbat, Dussaulx, & Zweibaum, 1990; Lesuffleur et al., 1991; Lesuffleur et al., 1993; Lesuffleur et al., 1995
HT29-Cl.16E	Human colorectal adenocarcinoma cell line deriving from HT-29 cell line after growth adaptation to sodium butyrate thus acquiring the goblet cell-like phenotype with a high capacity of mucus production	<i>MUC3, MUC2</i>	Augeron & Laboisse, 1984; Jarry et al., 1994; Maoret et al., 1989

800

801

Mucus layer

Mucins

No mucus layer

Oral cavity

MUC7, MUC5B,
MUC1, MUC16

Inner mucus layer:
attached and organized
Outer mucus layer:
unattached and mobile

Stomach

MUC5AC, MUC6,
MUC1

Outer mucus layer:
unattached and discontinuous

Small intestine

MUC2, MUC6, MUC1,
MUC3, MUC4,
MUC12, MUC13,
MUC17


Inner mucus layer:
attached and organized
Outer mucus layer:
unattached and mobile

Large intestine


MUC2, MUC1, MUC3,
MUC4, MUC12,
MUC13, MUC17

a) Secreted mucins: Gel-forming mucin

MUC2




MUC5AC




b) Secreted mucins: Non gel-forming mucin

MUC7



c) Cell membrane bound mucin

MUC1



Peptide signal

Von Willebrand D1-D4

Cysteine-rich domain

Tandem repeat (TR) sequence

Histidin like-sequence

Serine-Threonine rich region

Von Willebrand B

Von Willebrand C

Cysteine knot

SEA domain

Cytoplasmic domain

Leucine-zipper

