

On gaussian curvature equation in R^2 with prescribed nonpositive curvature

Huyuan Chen, Dong Ye, Feng Zhou

▶ To cite this version:

Huyuan Chen, Dong Ye, Feng Zhou. On gaussian curvature equation in R^2 with prescribed non-positive curvature. Discrete & Continuous Dynamical Systems - A, 2020, 40 (6), pp.3201-3214. 10.3934/dcds.2020125 . hal-02894440

HAL Id: hal-02894440 https://hal.univ-lorraine.fr/hal-02894440

Submitted on 8 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On Gaussian curvature equation in \mathbb{R}^2 with prescribed nonpositive curvature

HUYUAN CHEN¹

Department of Mathematics, Jiangxi Normal University, Nanchang, Jiangxi 330022, PR China

Dong Ye^2

Center for PDEs, School of Mathematical Sciences, East China Normal University, Shanghai Key Laboratory of PMMP, Shanghai 200062, PR China IECL, UMR 7502, University of Lorraine, 57050 Metz, France

Feng $ZHOU^3$

Center for PDEs, School of Mathematical Sciences, East China Normal University, Shanghai Key Laboratory of PMMP, Shanghai 200062, PR China

Dedicated to Professor Wei-Ming Ni's seventieth birthday

Abstract

The purpose of this paper is to study the solutions of

 $\Delta u + K(x)e^{2u} = 0 \quad \text{in } \mathbb{R}^2$

with $K \leq 0$. We introduce the following quantity:

$$\alpha_p(K) = \sup\left\{\alpha \in \mathbb{R} : \int_{\mathbb{R}^2} |K(x)|^p (1+|x|)^{2\alpha p + 2(p-1)} dx < +\infty\right\}, \quad \forall \ p \ge 1.$$

Under the assumption (\mathbb{H}_1) : $\alpha_p(K) > -\infty$ for some p > 1 and $\alpha_1(K) > 0$, we show that for any $0 < \alpha < \alpha_1(K)$, there is a unique solution u_{α} with $u_{\alpha}(x) = \alpha \ln |x| + c_{\alpha} + o(|x|^{-\frac{2\beta}{1+2\beta}})$ at infinity and $\beta \in (0, \alpha_1(K) - \alpha)$. Furthermore, we show an example $K_0 \leq 0$ such that $\alpha_p(K_0) = -\infty$ for any p > 1 and $\alpha_1(K_0) > 0$, for which we study the asymptotic behavior of solutions. In particular, we prove the existence of a solution u_{α_*} such that $u_{\alpha_*} - \alpha_* \ln |x| = O(1)$ at infinity for some $\alpha_* > 0$, but who does not converge to a constant at infinity. This example exhibits a new phenomenon of solution with logarithmic growth, finite total curvature, and non-uniform asymptotic behavior at infinity.

Key words: Gaussian curvature; Conformal metrics; Asymptotic behavior. MSC 2010: 35R06, 35A01, 35J66.

1 Introduction

In this paper, we study the solutions of

$$\Delta u + K(x)e^{2u} = 0 \quad \text{in } \mathbb{R}^2, \tag{1.1}$$

where K is a nonpositive, non trivial and locally bounded function in \mathbb{R}^2 .

Equation (1.1) finds its origin from conformal geometry: Given a Riemannian surface (\mathcal{M}, g) and a function K on \mathcal{M} , we may ask whether there is a new Riemannian metric g_1 , conformal to

¹chenhuyuan@yeah.net

²dye@math.ecnu.edu.cn, dong.ye@univ-lorraine.fr

³fzhou@math.ecnu.edu.cn

g, such that K is the Gaussian curvature of g_1 . In other words, we need to find a function u on \mathcal{M} such that $g_1 = e^{2u}g$ and u is a solution of the elliptic equation

$$\Delta_q u - k_q + K e^{2u} = 0 \quad \text{on } \mathcal{M}.$$
(1.2)

Here Δ_g is the Laplace-Beltrami operator, k_g is the Gaussian curvature on (\mathcal{M}, g) . In the Euclidean case $\mathcal{M} = \mathbb{R}^2$ and $g = |dx|^2$, (1.2) reduces to equation (1.1).

The equation (1.1) has been studied extensively, especially for the nonpositive, nonconstant curvature case, see [5, 9, 8, 6, 3] and the references therein. The nonexistence of entire solution in \mathbb{R}^2 with $K \equiv -1$ was given by Ahlfors [1] in 1938, and then improved by Sattinger [9], Ni [8] for $K \leq 0$ in \mathbb{R}^2 and $K(x) \leq -C|x|^{-2}$ at infinity with C > 0. The first general existence result for (1.1) is due to Ni [8], he proved that if $K \leq 0$, $K \neq 0$ in \mathbb{R}^2 with $|K(x)| \leq C|x|^{-\ell}$ at infinity for some $\ell > 2$, then for any $\alpha > 0$ but small enough, there exists a solution u satisfying $u(x) = \alpha \ln |x| + O(1)$ at infinity. Later on, it was proved by McOwen [6], Cheng-Ni [3] that if $K \leq 0$ in \mathbb{R}^2 , $K(x) \sim -|x|^{-\ell}$ at infinity with $\ell > 2$, then for any $\alpha \in (0, \frac{\ell-2}{2})$, (1.1) has a unique solution satisfying

$$u_{\alpha}(x) = \alpha \ln |x| + c_{\alpha} + o(1) \text{ as } |x| \to \infty, \text{ with } c_{\alpha} \in \mathbb{R}.$$
(1.3)

The solutions set has layer structure, that is $u_{\beta} < u_{\alpha}$ in \mathbb{R}^2 , for any $0 < \beta < \alpha < \alpha_1(K)$. Moreover, a unique maximal solution U exists such that $U(x) = \frac{\ell-2}{2} \ln |x| - \ln \ln |x| + O(1)$ at infinity, see [3].

For general curvature functions K, the following quantity was introduced by Cheng-Ni in [4],

$$\alpha_1(K) = \sup\left\{\alpha \in \mathbb{R} : \int_{\mathbb{R}^2} |K(x)| (1+|x|)^{2\alpha} dx < +\infty\right\}$$
(1.4)

with the convention $\sup \emptyset = -\infty$. When K is controlled polynomially at infinity, using $\alpha_1(K)$, Cheng-Ni characterized the existence to problem (1.1): Assume that $K \leq 0$ satisfies

$$(\mathbb{H}_0) \qquad \qquad \alpha_1(K) > 0 \text{ and } |K(x)| < C(|x|^m + 1) \text{ in } \mathbb{R}^2 \text{ for some } m > 0.$$

Then for any $\alpha \in (0, \alpha_1(K))$, (1.1) has a unique solution such that

$$u_{\alpha} = \alpha \ln |x| + O(1) \quad \text{as} \quad |x| \to \infty.$$
(1.5)

Later on, by employing a variational method with weighted Sobolev spaces (firstly applied by McOwen in [7]), Cheng-Lin [2] succeeded to handle the existence of problem (1.1) without any growth control for K: Assume that $\alpha_1(K) > 0$ for a nonpositive, locally bounded function K.

• For any $\alpha \in (0, \alpha_1(K))$, equation (1.1) possesses a unique solution u_α satisfying

$$\int_{\mathbb{R}^2} K e^{2u} dx = -2\alpha\pi \quad \text{and} \quad \int_{\mathbb{R}^2 \setminus B_1} \left| \nabla (u - \alpha \ln |x|) \right|^2 dx < +\infty.$$
(1.6)

- Moreover, for any $0 < \beta < \alpha < \alpha_1(K)$, there holds $u_\beta < u_\alpha$ in \mathbb{R}^2 .
- If $\alpha_1(K) < \infty$, then $\lim_{\alpha \to \alpha_1(K)} u_\alpha = u_{\alpha_1}$ is a solution to (1.1) satisfying (1.6) with $\alpha = \alpha_1(K)$.

The quantity

$$\mathcal{C}(u) := \int_{\mathbb{R}^2} K e^{2u} dx$$

is called the total curvature of the conformal metric $g_u = e^{2u} |dx|^2$. It plays an important role in the analysis of (1.1) and has a notable meaning in geometry. There are many interesting discussions

in [2]. For example, by Theorem 1.4 in [2], when $K \leq 0$, a solution u to (1.1) is logarithmically up bounded at infinity if and only if $\mathcal{C}(u)$ is finite and the metric $e^{2u}|dx|^2$ is complete.

However, to claim the asymptotic behavior like (1.5), Cheng-Lin need additional assumptions like (\mathbb{H}_0). A naturel question is to understand the asymptotic behavior of solutions provided by variational method in [2]. We will point out that any solution satisfying (1.5) is just a Cheng-Lin type solution, see Proposition 2.1 below. Moreover, we want to prove the existence of solutions satisfying (1.5) without any growth assumption on the curvature function K.

Another naturel question is to ask if there exist solutions with logarithmic growth but anisotropic behavior at infinity, more precisely, solutions of (1.1) satisfying (1.5) but not (1.3). Our purpose here is to answer these questions.

In fact, we propose to replace the pointwise growth assumption (\mathbb{H}_0) by an integral estimate. To this end, consider the following quantity related to K, which generalizes $\alpha_1(K)$:

$$\alpha_p(K) = \sup\left\{\alpha \in \mathbb{R} : \int_{\mathbb{R}^2} |K(x)|^p (1+|x|)^{2\alpha p + 2(p-1)} dx < +\infty\right\}, \quad \forall \ p \ge 1.$$
(1.7)

We use always the convention $\sup \emptyset = -\infty$. Our first main result states as follows.

Theorem 1.1. Let $K \leq 0$ be locally bounded in \mathbb{R}^2 . Assume that

$$(\mathbb{H}_1) \qquad \qquad \alpha_1(K) > 0 \text{ and there exists } p > 1 \text{ such that } \alpha_p(K) > -\infty.$$

Then for any $\alpha \in (0, \alpha_1(K))$, problem (1.1) possesses a unique solution u_α satisfying (1.5). Moreover, the total curvature of u_α is $-2\alpha\pi$; and there exists $c_\alpha \in \mathbb{R}$ such that for any $\beta \in (0, \alpha_1(K) - \alpha)$,

$$u_{\alpha}(x) = \alpha \ln |x| + c_{\alpha} + o\left(|x|^{-\frac{2\beta}{1+2\beta}}\right) \quad \text{as } |x| \to \infty.$$
(1.8)

Remark 1.1. Combining with Proposition 2.1 below, Theorem 1.1 means that under (\mathbb{H}_1) , for any $\alpha \in (0, \alpha_1(K))$, the unique solution satisfying (1.6) given by [2] has the asymptotic behavior (1.8).

Remark 1.2. Assumption (\mathbb{H}_1) is weaker than (\mathbb{H}_0) , see Lemma 2.2 below. So we obtain more precise asymptotic behavior (1.8) also for Cheng-Ni's solutions in [3].

Our construction is based on the understanding of $\alpha_p(K)$. Indeed, under (\mathbb{H}_1) , we can prove that $\alpha_p(K)$ are positive for p > 1 but nearby. This observation allows us to construct suitable super and sub solutions of (1.1). Let w_0 be a smooth radial function such that

$$w_0(x) = \ln |x| \quad \text{in } \mathbb{R}^2 \setminus B_1, \tag{1.9}$$

where $B_1 = \{x \in \mathbb{R}^2, |x| < 1\}$. In the following, $B_r(x)$ stands for the standard open ball with center $x \in \mathbb{R}^2$ and the radius r; B_r denotes always $B_r(0)$. We will look for a solution u_{α} to (1.1) in the form

$$u_{\alpha} = \alpha w_0 + v_{\alpha}$$

where v_{α} resolves

$$-\Delta v = K e^{2\alpha w_0} e^{2v} + \alpha \Delta w_0 \quad \text{in } \mathbb{R}^2$$

The crucial point is to know whether the above equation has a bounded solution, or equivalently, whether $Ke^{2\alpha w_0}e^{2v} + \alpha \Delta w_0$ belongs to the Kato's class. If the answer is affirmative, the Perron's method can be employed to get a solution of (1.1).

Our second concern is to understand solutions to (1.1) with an exotic $K_0 \leq 0$ such that

$$\alpha_p(K_0) = -\infty \text{ for any } p > 1 \text{ and } \alpha_1(K_0) > 0.$$
 (1.10)

We construct here an explicit model K_0 as follows. Consider the sequence $a_n := (n, 0)_{n \ge 2} \in \mathbb{R}^2$. Let

$$K_0(x) = -\sum_{n=2}^{\infty} r_n^{-2} n^{-\ell} \eta_0 \left(\frac{x - a_n}{r_n}\right) \quad \text{in } \mathbb{R}^2,$$
(1.11)

where

 $\ell > q > 1$, $r_n = e^{-n^q}$ for all $n \ge 2$

and η_0 is a smooth radial cut-off function satisfying

$$\chi_{B_{\frac{1}{2}}} \le \eta_0 \le \chi_{B_1}.$$
(1.12)

Here χ_{Ω} means the usual characteristic function of Ω . Observe that $r_n \leq e^{-2} < \frac{1}{4}$ for any $n \geq 2$, so

$$\int_{\mathbb{R}^2} |K_0(x)|^p (1+|x|)^{2\alpha p+2(p-1)} dx = \sum_{n=2}^\infty \int_{B_{r_n}(a_n)} |K_0(x)|^p (1+|x|)^{2\alpha p+2(p-1)} dx =: \sum_{n=2}^\infty I_n.$$

Direct computation implies then: for $p \ge 1$, $\alpha \in \mathbb{R}$, (as n goes to ∞)

$$\begin{split} I_n &:= r_n^{-2p} n^{-\ell p} \int_{B_{r_n}(a_n)} (1+|x|)^{2\alpha p+2(p-1)} \eta_0 \left(\frac{x-a_n}{r_n}\right)^{2\alpha p+2(p-1)} dx \\ &\sim r_n^{-2p} n^{2\alpha p+2(p-1)-\ell p} \int_{B_{r_n}(a_n)} \eta_0 \left(\frac{x-a_n}{r_n}\right)^{2\alpha p+2(p-1)} dx \\ &= M_p r_n^{2-2p} n^{2\alpha p+2(p-1)-\ell p} \\ &= M_p e^{2(p-1)n^q} n^{(2\alpha+2-\ell)p-2}, \end{split}$$

with $M_p = \int_{\mathbb{R}^2} \eta_0^p(z) dz > 0$. We can verify easily (1.10) and $\alpha_1(K_0) = \frac{\ell - 1}{2}$.

The following result exhibits clearly the asymptotic behavior of Cheng-Lin's solutions to (1.1) with $K = K_0$.

Theorem 1.2. Assume that $\ell > q > 1$ and $K = K_0$ given by (1.11). By [2], for any $0 < \alpha < \frac{\ell-1}{2}$, there is a unique solution of (1.1) u_{α} satisfying (1.6). Denote $\alpha_* = \frac{\ell-q}{2}$, we have:

- (i) For any $0 < \alpha < \alpha_*$, u_α satisfying (1.3).
- (ii) For $\alpha = \alpha_*$, the solution u_{α_*} satisfies (1.5), but the remainder term $u_{\alpha_*} \alpha_* \ln |x|$ does not converge to any constant as $|x| \to \infty$.
- (iii) For any $\alpha_* < \alpha < \frac{\ell-1}{2}$, the remainder term $u_{\alpha} \alpha \ln |x|$ is unbounded at infinity.

To our best knowledge, this is a first example of K_0 without any symmetry or polynomial growth assumption, where we show a complete picture of asymptotic behavior for solutions provided in [2]. It shows that under the only condition $\alpha_1(K) > 0$, the solution with finite total curvature to (1.1) can have very different asymptotic behavior at infinity. This is equally a first example of solution to (1.1) with nonpositive Gaussian curvature, logarithmic growth but non uniform behavior at infinity, it is also a first example of infinitely many solutions with finite total curvature but unbounded remained behavior to $u_{\alpha} - \alpha \ln |x|$.

The rest of our paper is organized as follows. In Section 2, we show the link between solutions satisfying (1.5) and that provided by Cheng-Lin, we show also some properties of $\alpha_p(K)$ and the relationship between (\mathbb{H}_0) and (\mathbb{H}_1) . In Section 3, we prove Theorem 1.1. Section 4 is addressed to the study of solutions with K_0 . In the following, C or C_i denote always generic positive constants, their values could be changed from one line to another.

2 Preliminaries

First, we indicate the relationship between solutions satisfying (1.5) and solutions given by [2].

Proposition 2.1. Let u be a continuous solution to (1.1) satisfying (1.5) with $\alpha < \alpha_1(K)$. Then u is the Cheng-Lin's solution, i.e. u satisfies (1.6).

Proof. We involve function w_0 given by (1.9). So $\Delta w_0(x) = 0$ for |x| > 1 and

$$\int_{\mathbb{R}^2} \Delta w_0 dx = 2\pi. \tag{2.1}$$

We claim that

$$\int_{\mathbb{R}^2} |\nabla(u - \alpha w_0)|^2 dx < \infty \quad \text{and} \quad \int_{\mathbb{R}^2} K e^{2u} dx = -2\alpha \pi.$$
(2.2)

Denote $v = u - \alpha w_0$, we prove first $v \in L^2(\mathbb{R}^2)$. To this end, set $\eta_\rho(t) = \eta_0(t/\rho)$ for $\rho > 0$, where η_0 is a smooth function satisfying (1.12). Observe that

$$-\Delta v = K e^{2\alpha w_0} e^{2v} + \alpha \Delta w_0 \quad \text{in } \mathbb{R}^2.$$
(2.3)

As v is bounded in \mathbb{R}^2 and $\alpha < \alpha_1(K)$, there holds

J

$$2\int_{\mathbb{R}^2} \eta_{\rho}^2 |\nabla v|^2 dx = \int_{\mathbb{R}^2} \eta_{\rho}^2 \left(\Delta v^2 - 2v\Delta v\right) dx$$

$$= \int_{\mathbb{R}^2} v^2 \Delta \eta_{\rho}^2 dx - 2 \int_{\mathbb{R}^2} \eta_{\rho}^2 v \Delta v dx$$

$$\leq \|v\|_{\infty}^2 \int_{\mathbb{R}^2} |\Delta \eta_{\rho}^2| dx + 2\|v\|_{\infty} \int_{\mathbb{R}^2} |\Delta v| dx$$

$$\leq C_1 + 2\|v\|_{\infty} \int_{\mathbb{R}^2} \left(|K|e^{2\alpha w_0 + 2\|v\|_{\infty}} + |\alpha\Delta w_0|\right) dx \leq C_2.$$

Here C_i are positive constants independent on ρ . Passing $\rho \to \infty$, we see that $\nabla v \in L^2(\mathbb{R}^2)$.

Furthermore,

$$\begin{split} \left| \int_{\mathbb{R}^2} \eta_{\rho} \Delta v dx \right| &= \left| \int_{\mathbb{R}^2} \nabla \eta_{\rho} \cdot \nabla v dx \right| \\ &\leq \left(\int_{B_{2\rho} \setminus B_{\rho}} |\nabla v|^2 dx \right)^{\frac{1}{2}} \left(\int_{B_{2\rho} \setminus B_{\rho}} |\nabla \eta_{\rho}|^2 dx \right)^{\frac{1}{2}} \\ &= \| \nabla \eta_0 \|_{L^2(\mathbb{R}^2)} \left(\int_{B_{2\rho} \setminus B_{\rho}} |\nabla v|^2 dx \right)^{\frac{1}{2}}. \end{split}$$

Remark that $\Delta v \in L^1(\mathbb{R}^2)$ and $\nabla v \in L^2(\mathbb{R}^2)$, we obtain, taking $\rho \to \infty$,

$$\int_{\mathbb{R}^2} \Delta v dx = 0.$$

Combining (2.3) and (2.1), we obtain the second part of (2.2).

We give some elementary consequences of (\mathbb{H}_1) and (\mathbb{H}_0) .

Lemma 2.1. Assume that (\mathbb{H}_1) holds true with $p_0 > 1$. Then for any $p \in [1, p_0)$, $\alpha_p(K) \ge \alpha_{p_0}(K)$ and $\alpha_p(K) \to \alpha_1(K)$ as $p \to 1^+$.

Proof. For simplicity, we denote $\alpha_p(K)$ by α_p . Take any $\beta < \alpha < \alpha_{p_0}$ and $p \in [1, p_0)$. Denote $\tau_{\alpha} = -2 + \frac{2p_0p}{p_0-p}(\beta - \alpha) < -2$, then

$$\tau_{\alpha} \left(1 - \frac{p}{p_0} \right) + \left[2\alpha p_0 + 2(p_0 - 1) \right] \frac{p}{p_0} = 2\beta p + 2(p - 1)$$

By Hölder's inequality, as $\alpha < \alpha_{p_0}$,

$$\int_{\mathbb{R}^2} |K(x)|^p (1+|x|)^{2\beta p+2(p-1)} dx$$

$$\leq \left(\int_{\mathbb{R}^2} |K(x)|^{p_0} (1+|x|)^{2\alpha p_0+2(p_0-1)} dx \right)^{\frac{p}{p_0}} \left(\int_{\mathbb{R}^2} (1+|x|)^{\tau_\alpha} dx \right)^{1-\frac{p}{p_0}} < \infty.$$

It means that $\alpha_p \ge \alpha_{p_0}$ for any $p \in [1, p_0)$. We see also that the function $p \mapsto \alpha_p$ is non-increasing in $[1, p_0]$.

If $\alpha_{p_0} = \alpha_1$, then $\alpha_p \equiv \alpha_{p_0}$ for $p \in [1, p_0]$ and we are done. Now assume that $\alpha_{p_0} < \alpha_1$. Let $\varepsilon \in (0, \alpha_1 - \alpha_{p_0})$, take

$$\lambda = \alpha_1 - \varepsilon, \quad \beta_2 < \alpha_{p_0} < \lambda < \beta_1 < \alpha_1,$$

and consider

$$h(p) := \beta_1(p_0 - p) + \beta_2 p_0(p - 1) - \lambda p(p_0 - 1)$$
 in $[1, p_0]$.

Readily, $h(1) = (\beta_1 - \lambda)(p_0 - 1) > 0$ and $h(p_0) = (\beta_2 - \lambda)(p_0 - 1)p_0 < 0$. Hence, there exists $p_* \in (1, p_0)$ such that $h(p_*) = 0$, or equivalently

$$2\beta_1\theta + \Big[2\beta_2p_0 + 2(p_0 - 1)\Big](1 - \theta) = 2\lambda p_* + 2(p_* - 1) \quad \text{where} \quad \theta := \frac{p_0 - p_*}{p_0 - 1} \in (0, 1).$$

As $\theta + p_0(1 - \theta) = p_*$, there holds

$$\begin{split} &\int_{\mathbb{R}^2} |K(x)|^{p_*} (1+|x|)^{2\lambda p_*+2(p_*-1)} dx \\ &\leq \left(\int_{\mathbb{R}^2} |K(x)| (1+|x|)^{2\beta_1} dx \right)^{\theta} \left(\int_{\mathbb{R}^2} |K(x)|^{p_0} (1+|x|)^{2\beta_2 p_0+2(p_0-1)} dx \right)^{1-\theta} < \infty. \end{split}$$

Thus we get $\alpha_{p_*} \ge \lambda$, so $\alpha_1 - \varepsilon \le \alpha_p \le \alpha_1$ for $p \in [1, p_*]$, which means $\lim_{p \to 1^+} \alpha_p = \alpha_1$.

Remark 2.1. By Lemmas 2.1, we see that (\mathbb{H}_1) is equivalent to (\mathbb{H}'_1) : There exists p > 1 such that $\alpha_p(K) > 0$.

Lemma 2.2. If K satisfies (\mathbb{H}_0) , then there exists p > 1 such that $\alpha_p(K) > 0$.

Proof. Fix some $\alpha \in (0, \alpha_1)$ and choose p > 1 but nearby such that

$$\beta := \alpha + \left(\frac{m}{2} + \alpha + 1\right)(p-1) < \alpha_1.$$

Then

$$|K(x)|^{p-1}(1+|x|)^{2(\alpha+1)(p-1)} \le C(1+|x|)^{m(p-1)+2(\alpha+1)(p-1)} = C(1+|x|)^{2\beta-2\alpha},$$

so that

$$\begin{split} \int_{\mathbb{R}^2} |K(x)|^p (1+|x|)^{2\alpha p+2(p-1)} dx &= \int_{\mathbb{R}^2} |K(x)| (1+|x|)^{2\alpha} \Big[|K(x)|^{p-1} (1+|x|)^{2(\alpha+1)(p-1)} \Big] dx \\ &\leq C \int_{\mathbb{R}^2} |K(x)| (1+|x|)^{2\beta} dx < \infty. \end{split}$$

The proof is completed.

We end this section by a simple fact of uniqueness, which can be derived from Theorem 4.1 in [3]. For the convenience of readers, we give a short proof here.

Lemma 2.3. Let $K \leq 0$, then for any $\alpha \in \mathbb{R}$, there exists at most one continuous solution to (1.1) satisfying (1.5).

Let \tilde{u}_{α} and u_{α} be two solutions of (1.1) satisfying (1.5) with $K \leq 0$. Denote $v = \tilde{u}_{\alpha} - u_{\alpha}$, then

$$-\Delta v^{2} = -2v\Delta v - 2|\nabla v|^{2} \le -2v\Delta v = 2Ke^{2u_{\alpha}}v\left(e^{2v}-1\right) \le 0,$$

that is, v^2 is bounded and subharmonic in \mathbb{R}^2 , hence v^2 is a constant. So $0 = -\Delta v = Ke^{2u_\alpha}(e^{2v}-1)$ in \mathbb{R}^2 , which implies that $v \equiv 0$ and the uniqueness follows.

3 Proof of Theorem 1.1

We begin with a result which plays an important role to obtain the asymptotic behavior (1.8).

Proposition 3.1. Let

$$f \in \dot{L}^1(\mathbb{R}^2) := \left\{ f \in L^1(\mathbb{R}^2), \ \int_{\mathbb{R}^2} f(x) dx = 0 \right\}.$$

Assume that $\alpha_{p_0}(f) > 0$ with $p_0 > 1$, then there exists a unique ground state solution to $-\Delta w = f$ in \mathbb{R}^2 . Moreover, $w \in C(\mathbb{R}^2)$ and

$$\lim_{|x|\to\infty} w(x)|x|^{\frac{2\beta}{1+2\beta}} = 0 \quad \text{for any } \beta \in (0,\alpha_1(f)).$$
(3.1)

Proof. The uniqueness of the ground state solution is a trivial consequence of Liouville's Theorem. So we prove only the existence of a solution satisfying (3.1). By $\alpha_{p_0}(f) > 0$ and Lemma 2.1, we get $\alpha_1(f) > 0$ and $f(z) \ln(2 + |z|) \in L^1(\mathbb{R}^2)$. As $f \in \dot{L}^1 \cap L^{p_0}_{loc}$, consider

$$w(x) = -\frac{1}{2\pi} \int_{\mathbb{R}^2} \ln |x - y| f(y) dy \quad \text{in } \mathbb{R}^2.$$

Obviously, $-\Delta w = f$ in \mathbb{R}^2 , $w \in C(\mathbb{R}^2)$ and for any $x \neq 0$,

$$2\pi w(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \ln \frac{|x|}{|x-y|} f(y) dy = \sum_{i=1}^3 \int_{\Omega_i} \ln \frac{|x|}{|x-y|} f(y) dy =: \sum_{i=1}^3 J_i(x),$$

where $\Omega_1 = B_R$, $R \ge 2$, $\Omega_2 = B_{\frac{|x|}{2}}(x)$ and $\Omega_3 = \mathbb{R}^2 \setminus (\Omega_1 \cup \Omega_2)$. For $|x| \ge 4R > 0$,

$$|J_1(x)| \le C \int_{B_R} \frac{|y|}{|x|} |f(y)| dy \le \frac{CR}{|x|} ||f||_{L^1(\mathbb{R}^2)}.$$

Let $e_x = \frac{x}{|x|}$, $p'_0 = \frac{p_0}{p_0-1}$ and fix $0 < \beta < \alpha_{p_0}(f)$, we get

$$\begin{aligned} |J_{2}(x)| &= \left| \int_{\Omega_{2}} \ln \frac{|x|}{|x-y|} f(y) dy \right| \\ &\leq \left(\int_{\Omega_{2}} |f(y)|^{p_{0}} (1+|y|)^{2\beta p_{0}+2(p_{0}-1)} dy \right)^{\frac{1}{p_{0}}} \left(\int_{\Omega_{2}} \left| \ln \frac{|x|}{|x-y|} \right|^{p_{0}'} (1+|y|)^{-2-2\beta p_{0}'} dy \right)^{\frac{1}{p_{0}'}} \\ &\leq C \left(\int_{B_{\frac{1}{2}}(e_{x})} \left| \ln |e_{x}-z| \right|^{p_{0}'} |z|^{-2-2\beta p_{0}'} |x|^{-2\beta p_{0}'} dz \right)^{\frac{1}{p_{0}'}} \\ &= C|x|^{-2\beta}. \end{aligned}$$

Let $|x| \ge 4R \ge 8$. In Ω_3 , as min $(|x - y|, |x|, |y|) \ge R \ge 2$, we can check that

$$\forall y \in \Omega_3, \qquad \left| \ln \frac{|x|}{|x-y|} \right| \le \ln |y|,$$

since $\frac{1}{|y|} \leq \frac{|x|}{|x-y|} \leq |y|$. Then there holds

$$\begin{aligned} |J_{3}(x)| &\leq \int_{\Omega_{3}} |f(y)| \ln |y| dy \\ &\leq \left(\int_{\Omega_{3}} |f(y)|^{p_{0}} (1+|y|)^{2\beta p_{0}+2(p_{0}-1)} dy \right)^{\frac{1}{p_{0}}} \left(\int_{\Omega_{3}} (\ln |y|)^{p_{0}'} |y|^{-2-2\beta p_{0}'} dy \right)^{\frac{1}{p_{0}'}} \\ &\leq C \left(\int_{\mathbb{R}^{2} \setminus B_{R}} (\ln |y|)^{p_{0}'} |y|^{-2-2\beta p_{0}'} dy \right)^{\frac{1}{p_{0}'}} \\ &\leq C R^{-2\beta} \ln R. \end{aligned}$$

Notice that all the constants C involved are independent of $|x| \ge 4R$ and $R \ge 2$. Taking $R = |x|^{\frac{1}{1+2\beta}}$, there holds

$$|w(x)| \le C|x|^{-\frac{2\beta}{1+2\beta}} \ln|x|$$
 for $|x|$ large. (3.2)

This ends the proof of (3.1), because we can get (3.2) with any $p \in (1, p_0)$ and any $\beta < \alpha_p(f)$. \Box

We are now in position to prove Theorem 1.1.

Proof of Theorem 1.1. Let $\alpha \in (0, \alpha_1(K))$ and w_0 be given by (1.9), there exists then a unique $t_{\alpha} \in \mathbb{R}$ such that

$$g_{\alpha} := K e^{2\alpha w_0} e^{2t_{\alpha}} + \alpha \Delta w_0 \in \dot{L}^1(\mathbb{R}^2).$$

Indeed, as $K \leq 0$ in \mathbb{R}^2 and $\alpha > 0$, the unique choice is given by

$$t_{\alpha} = \frac{1}{2} \ln \left(-\frac{2\alpha\pi}{\int_{\mathbb{R}^2} K e^{2\alpha w_0} dx} \right).$$

On the other hand, using (\mathbb{H}_1) and Lemma 2.1, there exists p > 1 such that $\alpha < \alpha_p(K)$. As $\alpha \Delta w_0$ is compactly supported, it's not difficult to see that

$$\alpha_p(g_\alpha) = \alpha_p(K) - \alpha > 0.$$

Using Proposition 3.1, we get a continuous ground state solution to $-\Delta \tilde{w} = g_{\alpha}$ in \mathbb{R}^2 .

We conclude by super-sub solution method. Let $\overline{v} = \alpha w_0 + t_\alpha + \widetilde{w} + \|\widetilde{w}\|_{\infty}$. As $K \leq 0$, it's easy to see that

$$-\Delta \overline{v} = -\alpha \Delta w_0 - \Delta \widetilde{w} = K e^{2(\alpha w_0 + t_\alpha)} \ge K e^{2\overline{v}}.$$

Hence \overline{v} is a supersolution to equation (1.1). Similarly, $\underline{v} = \alpha w_0 + t_\alpha + \widetilde{w} - \|\widetilde{w}\|_\infty$ is a subsolution to (1.1). Obviously $\overline{v} \geq \underline{v}$ in \mathbb{R}^2 . Using the standard Perron's method, there exists a solution u to (1.1) such that $\underline{v} \leq u \leq \overline{v}$ in \mathbb{R}^2 . Clearly, u satisfies (1.5).

By Proposition 2.1, there holds $Ke^{2u} + \alpha \Delta w_0 \in \dot{L}^1(\mathbb{R}^2)$. Applying Lemma 2.1, there exists p > 1 such that

$$\alpha_p \left(K e^{2u} + \alpha \Delta w_0 \right) = \alpha_p(K) - \alpha > 0.$$

By Proposition 3.1, we get a continuous solution of $-\Delta w = Ke^{2u} + \alpha \Delta w_0$ in \mathbb{R}^2 , which satisfies (3.1) for any $0 < \beta < \alpha_1(K) - \alpha$. Therefore, $v = u - \alpha w_0 - w$ is a bounded harmonic function over \mathbb{R}^2 , hence v is a constant c_{α} . Finally $u = \alpha w_0 + c_{\alpha} + w$, so (1.8) holds trues for any $0 < \beta < \alpha_1(K) - \alpha$. \Box

Similarly, using the super-sub solution method and Proposition 2.1, we can claim

Proposition 3.2. Assume that for $K \leq 0$, $\alpha > 0$ and $t \in \mathbb{R}$, there exists a bounded solution to $-\Delta v = Ke^{2\alpha w_0+2t} + \alpha \Delta w_0$ in \mathbb{R}^2 , then the equation (1.1) admits a unique solution u_α satisfying (1.5). Moreover, the total curvature of u_α is $-2\alpha\pi$.

4 Exotic curvature K_0 and anisotropic behavior solutions

Here we prove Theorem 1.2. We divide our study by several steps.

4.1 Solution satisfying (1.5) for $\alpha \leq \alpha_*$.

Remark that we cannot apply Theorem 1.1 in [2] or its proof, since they don't provide the asymptotic behavior. Indeed, we will construct directly solutions satisfying (1.5), and Proposition 2.1 ensures then they are solutions given by [2]. Remark also that we cannot apply the results of Section 3, since $\alpha_p(K_0) = -\infty$ for any p > 1.

Recall that $\alpha_1(K_0) = \frac{\ell-1}{2} > 0$ and w_0 is given by (1.9). Let $0 < \alpha \leq \frac{\ell-q}{2}$ and

$$v_{\alpha}(x) = -\frac{1}{2\pi} \int_{\mathbb{R}^2} K_0(y) e^{2\alpha w_0(y)} \ln |x - y| dy.$$

It is easy to see that v_{α} is well defined, continuous and locally bounded in \mathbb{R}^2 . Let

$$n \ge 2$$
 and $x \in B_{n+\frac{1}{2}} \setminus \overline{B}_{n-\frac{1}{2}}.$ (4.1)

We decompose $v_{\alpha}(x)$ as follows,

$$2\pi v_{\alpha}(x) = \sum_{k=2, k\neq n}^{\infty} r_{k}^{-2} k^{-\ell} \int_{B_{r_{k}}(a_{k})} |y|^{2\alpha} \eta_{0} \left(\frac{y-a_{k}}{r_{k}}\right) \ln|x-y| dy$$
$$+ r_{n}^{-2} n^{-\ell} \int_{B_{r_{n}}(a_{n})} |y|^{2\alpha} \eta_{0} \left(\frac{y-a_{n}}{r_{n}}\right) \ln|x-y| dy$$
$$=: E_{1}(x) + E_{2}(x).$$

Denote

$$s_k := \int_{B_{r_k}(a_k)} |y|^{2\alpha} \eta_0\left(\frac{y - a_k}{r_k}\right) dy \le C(k+1)^{2\alpha} r_k^2, \quad \forall \ k \ge 2.$$

Consider first $k \leq \frac{n}{2}$ and $k \geq 2$. Notice that there exists C > 0 such that

$$\left| \ln |x - y| - \ln |x| \right| \le C \frac{|y|}{|x|}, \quad \forall |x| \ge 2|y| > 0.$$

For x satisfying (4.1), there holds then

$$\begin{split} \left| \int_{B_{r_k}(a_k)} |y|^{2\alpha} \eta_0\left(\frac{y-a_k}{r_k}\right) \ln|x-y| \, dy - s_k \ln|x| \right| &= \left| \int_{B_{r_k}(a_k)} |y|^{2\alpha} \eta_0\left(\frac{y-a_k}{r_k}\right) \ln\frac{|x-y|}{|x|} dy \right| \\ &\leq \frac{C}{n} (k+1)^{2\alpha+1} \int_{B_{r_k}(a_k)} \eta_0\left(\frac{y-a_k}{r_k}\right) dy \\ &= \frac{C}{n} k^{2\alpha+1} r_k^2. \end{split}$$

Let now $k \neq n, k > \frac{n}{2}$. As $|y| \ge \frac{7}{4}$ and $|x - y| \ge \frac{1}{4}$ in $B_{r_k}(a_k)$, there exists C > 1 (independent of $k, n \ge 2, k \neq n$) such that

$$\forall y \in B_{r_k}(a_k), \quad |y|^{-C} \le \frac{\frac{1}{4}}{|y| + \frac{1}{4}} \le \frac{|x - y|}{|y| + |x - y|} \le \frac{|x - y|}{|x|} \le 1 + |y| \le |y|^C.$$

We obtain

$$\begin{split} \left| \int_{B_{r_k}(a_k)} |y|^{2\alpha} \eta_0 \left(\frac{y - a_k}{r_k} \right) \ln |x - y| \, dy - s_k \ln |x| \right| &= \left| \int_{B_{r_k}(a_k)} |y|^{2\alpha} \eta_0 \left(\frac{y - a_k}{r_k} \right) \ln \frac{|x - y|}{|x|} dy \right| \\ &\leq C \left| \int_{B_{r_k}(a_k)} |y|^{2\alpha} \eta_0 \left(\frac{y - a_k}{r_k} \right) \ln |y| dy \right| \\ &\leq C(k + 1)^{2\alpha} r_k^2 \ln(k + 1). \end{split}$$

Therefore, for x in (4.1),

$$\begin{aligned} & \left| E_1(x) - \left(\sum_{k=2}^{\infty} r_k^{-2} k^{-\ell} s_k \right) \ln |x| \right| \\ & \leq \frac{C}{n} \sum_{k=2}^{\left[\frac{n}{2}\right]} k^{2\alpha - \ell + 1} + C \sum_{k > \left[\frac{n}{2}\right], k \neq n} k^{2\alpha - \ell} \ln k + r_n^{-2} n^{-\ell} s_n \ln |x| \\ & \leq \frac{C}{n} \sum_{k=2}^{\left[\frac{n}{2}\right]} k^{2\alpha - \ell + 1} + C \sum_{k=\left[\frac{n}{2}\right] + 1}^{\infty} k^{2\alpha - \ell} \ln k. \end{aligned}$$

Here $\left[\frac{n}{2}\right]$ is the integer part of $\frac{n}{2}$. Finally, as $2\alpha - \ell + 1 \leq 1 - q < 0$, there holds

$$\lim_{|x| \to \infty} \left| E_1(x) - A_\alpha \ln |x| \right| = 0, \quad \text{where} \quad A_\alpha = \sum_{k=2}^\infty r_k^{-2} k^{-\ell} s_k \in (0,\infty).$$
(4.2)

On the other hand, for x satisfying (4.1),

$$\begin{aligned} |E_{2}(x)| &\leq Cr_{n}^{-2}n^{2\alpha-\ell}\int_{B_{r_{n}}(a_{n})}\eta_{0}\left(\frac{y-a_{n}}{r_{n}}\right) \times (-\ln|x-y|)dy \\ &= -Cn^{2\alpha-\ell}\int_{B_{1}}\eta_{0}(z)\ln|x-a_{n}-r_{n}z|dz \\ &\leq -Cn^{2\alpha-\ell}\int_{B_{1}}\eta_{0}(z)\ln|r_{n}z|dz \\ &= Cn^{2\alpha-\ell}(|\ln r_{n}|+1) \\ &\leq Cn^{2\alpha-\ell+q}. \end{aligned}$$
(4.3)

For the third line, we used the symmetric decreasing rearrangement argument, since η_0 and $-\ln |r_n z|$ are radial non-increasing positive functions in B_1 .

Combining (4.2)–(4.3), v_{α} is a solution to $-\Delta v_{\alpha} = K_0 e^{2\alpha w_0}$ in \mathbb{R}^2 such that

$$v_{\alpha}(x) = \frac{A_{\alpha}}{2\pi} \ln |x| + O(1), \text{ as } |x| \to \infty.$$

Taking

$$t_{\alpha} = \frac{1}{2} \ln \frac{2\alpha \pi}{A_{\alpha}},$$

 $h_{\alpha} = e^{2t_{\alpha}}v_{\alpha} - \alpha w_0$ is clearly a bounded solution to

$$-\Delta h = K_0 e^{2\alpha w_0 + 2t_\alpha} + \alpha \Delta w_0 \quad \text{in } \ \mathbb{R}^2.$$

By Propositions 3.2, for any $\alpha \in (0, \frac{\ell-q}{2}]$, equation (1.1) with $K = K_0$ admits a unique solution satisfying (1.5), and whose total curvature is equal to $-2\alpha\pi$.

4.2 Uniform behavior for $0 < \alpha < \alpha_*$.

For $\alpha \in (0, \frac{\ell-q}{2})$, we have a continuous solution u_{α} to (1.1) satisfying (1.5). Rewrite $u_{\alpha} = \alpha w_0 + \xi_{\alpha}$ with $\xi_{\alpha} \in L^{\infty}(\mathbb{R}^2)$. Let

$$\widetilde{v}_{\alpha}(x) = -\frac{1}{2\pi} \int_{\mathbb{R}^2} K_0(y) e^{2\alpha w_0(y) + 2\xi_{\alpha}(y)} \ln|x - y| dy.$$

By decomposing \tilde{v}_{α} as above and using

$$\widetilde{s}_k := \int_{B_{r_k}(a_k)} e^{2\xi_\alpha(y)} |y|^{2\alpha} \eta_0\left(\frac{y-a_k}{r_k}\right) dy \quad \text{for } k \ge 2,$$

the similar estimates to (4.2)–(4.3) hold true for \tilde{v}_{α} , which yield that (as now $2\alpha - \ell + q < 0$)

$$\lim_{|x|\to\infty} \left[\widetilde{v}_{\alpha}(x) - \frac{\widetilde{A}_{\alpha}}{2\pi} \ln |x| \right] = 0, \quad \text{where} \quad \widetilde{A}_{\alpha} = \sum_{k=2}^{\infty} r_k^{-2} k^{-\ell} \widetilde{s}_k \in (0, +\infty).$$
(4.4)

So $u_{\alpha} - \tilde{v}_{\alpha}$ is a harmonic function in \mathbb{R}^2 and $(u_{\alpha} - \tilde{v}_{\alpha})(x) = O(\ln |x|)$ at infinity, hence $u_{\alpha} - \tilde{v}_{\alpha}$ is a constant in \mathbb{R}^2 . The behavior (4.4) means that u_{α} satisfies (1.3).

4.3 Anisotropic behavior for $\alpha = \alpha_*$.

Let $\alpha = \alpha_* = \frac{\ell-q}{2}$, and u_{α_*} be the unique solution to (1.1) satisfying (1.5), hence $\xi_* := u_{\alpha_*} - \alpha_* w_0$ is uniformly bounded in \mathbb{R}^2 . However, we will show that ξ_* does not converge to a constant at infinity. Define

$$\begin{aligned} 2\pi v_*(x) &= -\int_{\mathbb{R}^2} K_0(y) e^{2\alpha_* w_0(y) + 2\xi_*(y)} \ln |x - y| dy \\ &= \sum_{k=2, \, k \neq n}^{\infty} r_k^{-2} k^{-\ell} \int_{B_{r_k}(a_k)} e^{2\xi_*(y)} |y|^{2\alpha} \eta_0 \left(\frac{y - a_k}{r_k}\right) \ln |x - y| \, dy \\ &+ r_n^{-2} n^{-\ell} \int_{B_{r_n}(a_n)} e^{2\xi_*(y)} |y|^{2\alpha} \eta_0 \left(\frac{y - a_n}{r_n}\right) \ln |x - y| \, dy \\ &=: F_1(x) + F_2(x). \end{aligned}$$

We see that

$$-\Delta v_* = K_0 e^{2\alpha_* w_0 + 2\xi_*} = K_0 e^{2u_{\alpha_*}} \text{ in } \mathbb{R}^2 \text{ and } v_*(x) = O(\ln|x|) \text{ as } |x| \to \infty.$$

The logarithmic control of v_* at infinity can be obtained as above, using $\xi_* \in L^{\infty}(\mathbb{R}^2)$.

We shall estimate the difference between $v_*(a_n)$ and $v_*(-a_n)$. The estimation of F_1 is very similar to E_1 in the above. Let $n \ge 2$. If $k \le \frac{n}{2}$ and $k \ge 2$, there holds

$$\begin{split} & \left| \int_{B_{r_k}(a_k)} e^{2\xi_*(y)} |y|^{2\alpha_*} \eta_0 \left(\frac{y - a_k}{r_k} \right) \left[\ln |a_n - y| - \ln |a_n + y| \right] dy \right| \\ & \leq \frac{C}{n} e^{2\|\xi_*\|_{\infty}} (k+1)^{2\alpha_*} \int_{B_{r_k}(a_k)} \eta_0 \left(\frac{y - a_k}{r_k} \right) |y| \, dy \\ & \leq \frac{C}{n} k^{2\alpha_* + 1} r_k^2. \end{split}$$

Let $k > \frac{n}{2}$ and $k \neq n$, we have

$$\begin{split} & \left| \int_{B_{r_k}(a_k)} e^{2\xi_*(y)} |y|^{2\alpha_*} \eta_0\left(\frac{y-a_k}{r_k}\right) \left[\ln|a_n-y| - \ln|a_n+y| \right] dy \right| \\ & \leq C(k+1)^{2\alpha_*} (\ln n + \ln k) \int_{B_{r_k}(a_k)} \eta_0\left(\frac{y-a_k}{r_k}\right) \, dy \\ & \leq Ck^{2\alpha_*} r_k^2 \ln k. \end{split}$$

As $2\alpha_* - \ell = -q < -1$, we get then

$$|F_1(a_n) - F_1(-a_n)| \le \frac{C}{n} \sum_{k=2}^{\left\lfloor \frac{n}{2} \right\rfloor} k^{2\alpha_* - \ell + 1} + C \sum_{k=\left\lfloor \frac{n}{2} \right\rfloor + 1}^{\infty} k^{2\alpha_* - \ell} \ln k \to 0, \quad \text{as } n \to \infty.$$
(4.5)

Now we estimate $F_2(\pm a_n)$:

$$0 \le F_2(-a_n) \le Cr_n^{-2}n^{-\ell+2\alpha_*} \int_{B_{r_n}(a_n)} \eta_0\left(\frac{y-a_n}{r_n}\right) \ln|a_n+y| dy \le Cn^{-\ell+2\alpha} \ln(2n+1)$$

and

$$\begin{aligned} -F_{2}(a_{n}) &= -r_{n}^{-2}n^{-\ell}\int_{B_{r_{n}}(a_{n})}e^{2\xi_{*}(y)}|y|^{2\alpha_{*}}\eta_{0}\left(\frac{y-a_{n}}{r_{n}}\right)\ln|a_{n}-y|dy\\ &\geq -e^{-2||\xi_{*}||_{\infty}}r_{n}^{-2}n^{-\ell}(n-1)^{2\alpha_{*}}\int_{B_{r_{n}}(a_{n})}\eta_{0}\left(\frac{y-a_{n}}{r_{n}}\right)\ln|a_{n}-y|dy\\ &\geq Cn^{-\ell+2\alpha_{*}}\int_{B_{1}}\eta_{0}(z)\Big[-\ln r_{n}-\ln|z|\Big]dz\\ &\geq -Cn^{-\ell+2\alpha_{*}}\ln r_{n}\\ &= Cn^{-\ell+2\alpha_{*}+q}=C>0. \end{aligned}$$

Therefore $\liminf_{n\to\infty} |F_2(a_n) - F_2(-a_n)| \ge C > 0$. Combining with (4.5), we get

$$\liminf_{n \to \infty} |v_*(a_n) - v_*(-a_n)| \ge \frac{C}{2\pi} > 0.$$

Furthermore, as $-\Delta(v_* - u_{\alpha_*}) = 0$ in \mathbb{R}^2 and $v_* - u_{\alpha_*}$ is $O(\ln |x|)$ at infinity, the standard Liouville's theorem implies that $v_* - u_{\alpha_*}$ is constant in \mathbb{R}^2 . Consequently,

$$\liminf_{n \to \infty} |\xi_*(a_n) - \xi_*(-a_n)| = \liminf_{n \to \infty} |v_*(a_n) - v_*(-a_n)| \ge \frac{C}{2\pi} > 0.$$

The above estimate means that ξ_* does not converge to a constant at infinity.

4.4 Unbounded remainder term for $\alpha_* < \alpha < \alpha_1(K_0)$.

Recall that by Theorem 1.1 in [2], for any $\alpha \in \left(\frac{\ell-q}{2}, \frac{\ell-1}{2}\right)$, problem (1.1) possesses a unique solution u_{α} satisfying (1.6). We shall prove that $\xi_{\alpha} := u_{\alpha} - \alpha w_0$ is unbounded.

Suppose the contrary, assume that ξ_{α} is bounded in \mathbb{R}^2 . Let

$$\eta_{\alpha}(x) = -\frac{1}{2\pi} \int_{\mathbb{R}^2} K_0(y) e^{2u_{\alpha}(y)} \ln |x - y| dy \text{ in } \mathbb{R}^2,$$

then we have

$$2\pi\eta_{\alpha}(x) = \sum_{k=2, k\neq n}^{\infty} r_{k}^{-2} k^{-\ell} \int_{B_{r_{k}}(a_{k})} |y|^{2\alpha} e^{2\xi_{\alpha}} \eta_{0} \left(\frac{y-a_{k}}{r_{k}}\right) \ln|x-y| dy$$
$$+ r_{n}^{2} n^{-\ell} \int_{B_{r_{n}}(a_{n})} |y|^{2\alpha} e^{2\xi_{\alpha}} \eta_{0} \left(\frac{y-a_{n}}{r_{n}}\right) \ln|x-y| dy$$
$$=: G_{1}(x) + G_{2}(x).$$

We will estimate $\eta_{\alpha}(a_n)$. For $k \neq n, k \geq 2$,

$$\begin{split} & \left| \int_{B_{r_k}(a_k)} |y|^{2\alpha} e^{2\xi_\alpha} \eta_0 \left(\frac{y - a_k}{r_k} \right) \ln |a_n - y| \, dy \right| \\ & \leq C e^{2\|\xi_\alpha\|_\infty} (k+1)^{2\alpha} \int_{B_{r_k}(a_k)} \eta_0 \left(\frac{y - a_k}{r_k} \right) \left(\ln n + \ln |y| \right) dy \\ & \leq C k^{2\alpha} r_k^2 \left(\ln n + \ln k \right). \end{split}$$

Since $\alpha < \frac{\ell-1}{2}$, there exists C > 0 such that $|G_1(a_n)| \leq C \ln n$ for n large. On the other hand,

$$-G_{2}(a_{n}) \geq -e^{-2\|\xi_{\alpha}\|_{\infty}} r_{n}^{-2} n^{-\ell} (n-1)^{2\alpha} \int_{B_{r_{n}}(a_{n})} \eta_{0} \left(\frac{y-a_{n}}{r_{n}}\right) \ln|a_{n}-y| dy$$

$$\geq -Cn^{-\ell+2\alpha} \ln r_{n}$$

$$= C_{3} n^{-\ell+2\alpha+q}.$$

Combining the estimations of $G_1(a_n)$ and $G_2(a_n)$, we get, as $\alpha > \frac{\ell - q}{2}$,

$$\lim_{n \to \infty} \frac{\eta_{\alpha}(a_n)}{\ln n} = -\infty.$$
(4.6)

Moreover, remark that for x satisfying (4.1) and $y \in B_{r_n}(a_n)$, there holds $|x-y| \leq 3|x|$. Similar estimations as above yield that there exists C > 0 such that

$$\eta_{\alpha}(x) \leq C \ln(2+|x|)$$
 in \mathbb{R}^2 .

As $\tilde{v} := \eta_{\alpha} - u_{\alpha} = \eta_{\alpha} - \alpha w_0 - \xi_{\alpha}$ is harmonic over \mathbb{R}^2 , and $\tilde{v}(x) \leq C \ln(2 + |x|)$ in \mathbb{R}^2 , \tilde{v} is constant by Liouville's theorem. However, $\eta_{\alpha} = \alpha w_0 + \xi_{\alpha} + C$ with $\xi_{\alpha} \in L^{\infty}$ contradicts the estimate (4.6). This means that ξ_{α} cannot be bounded in \mathbb{R}^2 .

Acknowledgements. H.C. is supported by NSFC (No. 11726614 and 11661045), and Jiangxi Provincial Natural Science Foundation (No. 20161ACB20007). F.Z. and D.Y. are supported by Science and Technology Commission of Shanghai Municipality (STCSM), grant No. 18dz2271000. F.Z. is also supported by NSFC (No. 11726613 and 11431005).

References

- [1] L.V. Ahlfors, An extension of Schwartz's lemma, Trans. Amer. Math. Soc. 43, 359-364 (1938).
- [2] K.-S. Cheng and C.-S. Lin, Conformal metrics with prescribed nonpositive Gaussian on ℝ², Calc. Var. PDE 11, 203-231 (2000).
- [3] K.-S. Cheng and W.-M. Ni, On the structure of the conformal Gaussian curvature equation on ℝ², Duke Math. J. 62, 721-737 (1991).
- [4] K.-S. Cheng and W.-M. Ni, On the structure of the conformal Gaussian curvature equation on ℝ² II, Math. Ann. 290, 671-680 (1991).
- [5] J. Kazdan and F. Warner, Curvature functions for open 2-manifolds, Ann. Math. 99, 203-219 (1974).
- [6] R. McOwen, On the equation $\Delta u + Ke^{2u} = f$ and prescribed negative curvature in \mathbb{R}^2 , J. Math. Anal. Appl. 103, 365-370 (1984).
- [7] R. McOwen, Conformal metrics in ℝ² with prescribed Gaussian curvature and positive total curvature, *Indiana Univ. Math. J.* 34, 97-104 (1985).
- [8] W.-M. Ni, On the elliptic equation $\Delta u + K(x)e^{2u} = 0$ and conformal metric with prescribed Gaussian curvatures, *Invent. Math.* 66, 343-352 (1982).
- [9] D. Sattinger, Conformal metrics in \mathbb{R}^2 with prescribed curvature, *Indiana Univ. Math. J.* 22, 1-4 (1972).