

Westward migration of oceanic ridges and related asymmetric upper mantle differentiation

Françoise Chalot-Prat, Carlo Doglioni, Trevor Falloon

▶ To cite this version:

Françoise Chalot-Prat, Carlo Doglioni, Trevor Falloon. Westward migration of oceanic ridges and related asymmetric upper mantle differentiation. Lithos, 2017, 268-271, pp.163-173. 10.1016/j.lithos.2016.10.036 . hal-02936297

HAL Id: hal-02936297 https://hal.univ-lorraine.fr/hal-02936297

Submitted on 15 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. 1 2

Westward migration of oceanic ridges and related asymmetric upper mantle differentiation

3

4 Françoise Chalot-Prat^a, Carlo Doglioni^b, Trevor Falloon^c

- ^aCentre de Recherches Petrographiques et Geochimiques Lorraine University, BP20, 15 rue Notre
 Dame des Pauvres, F-54501 Vandoeuvre-les-Nancy Cedex, France.
- 7 ^bDipartimento di Scienze della Terra, Sapienza University, P.le A. Moro, 5, 00185 Roma, Italy and
- 8 Istituto Nazionale di Geofisica e Vulcanologia, Via di Vigna Murata 605, 00143 Roma, Italy
- 9 ^cSchool of Physical Sciences, Discipline of Earth Sciences, University of Tasmania, Hobart, Tasmania
- 10 7001, Australia
- 11
- 12 Corresponding author: Françoise Chalot-Prat françoise.chalot-prat@univ-lorraine.fr
- 1314 Highlights:

15 1. Westward ridge migration induces higher mantle refertilisation of the western plate ...88

- Upwelling residual mantle is abandoned eastward consistent with seismological asymmetry ...
 91
- 18 3. Ridge migration provides permanently MORB mantle source from the west ...73

19 Abstract

20 Combining geophysical, petrological and structural data on oceanic mantle lithosphere, underlying 21 asthenosphere and oceanic basalts, an alternative oceanic plate spreading model is proposed in the 22 framework of the westward migration of oceanic spreading ridges relative to the underlying 23 asthenosphere. This model suggests that evolution of both the composition and internal structure of 24 oceanic plates and underlying upper mantle strongly depends at all scales on plate kinematics. We 25 show that the asymmetric features of lithospheric plates and underlying upper asthenosphere on both 26 sides of oceanic spreading ridges, as shown by geophysical data (seismic velocities, density, 27 thickness, and plate geometry), reflect somewhat different mantle compositions, themselves related to 28 various mantle differentiation processes (incipient to high partial melting degree, percolation/reaction 29 and refertilisation) at different depths (down to 300 km) below and laterally to the ridge axis. The 30 fundamental difference between western and eastern plates is linked to the westward ridge migration 31 inducing continuing mantle refertilisation of the western plate by percolation-reaction with ascending 32 melts, whereas the eastern plate preserves a barely refertilized harzburgitic residue. Plate thickness on 33 both sides of the ridge is controlled both by cooling of the asthenospheric residue and by the 34 instability of pargasitic amphibole producing a sharp depression of the water-undersaturated solidus, 35 its intersection with the geotherm at ~90 km, and incipient melt production right underneath the 36 lithosphere-asthenosphere boundary (LAB). Thus the intersection of the geotherm with the water-37 undersaturated lherzolite solidus explains the existence of a low-velocity zone (LVZ). As oceanic 38 lithosphere is moving westward relative to asthenospheric mantle, this partially molten upper 39 asthenosphere facilitates the decoupling between lower asthenosphere and lithosphere. Thereby the 40 westward drift of the lithosphere is necessarily slowed down, top to down, inducing a progressive 41 decoupling within the mantle lithosphere itself. This intra-mantle decoupling could be at the origin of 42 asymmetric detachment faults allowing mantle exhumation along slow-spreading ridges. Taking into 43 account the asymmetric features of the LVZ, migration of incipient melt fractions and upwelling paths

44 from the lower asthenosphere through the upper asthenosphere are oblique, upward and eastward.

45 MORB are sourced from an eastward and oblique, near-adiabatic mantle upwelling from the lower 46 asthenosphere. This unidirectional mantle transfer is induced by isostatic suction of the migrating 47 spreading ridge.

48 ³

49 Key words

Asymmetric oceanic lithosphere, Westward drift of the lithosphere, Migrating ridge, Mantle melting,
 Percolation-reaction and refertilization, Detachment fault and mantle core complex, Secondary
 Iherzolite and upper mantle differentiation

53 1. Introduction

54 The theory of plate tectonics, the post-1960's version of continental drift (Wegener, 1915), 55 proposes that the lithosphere (crust + uppermost mantle) is divided into a number of plates that can 56 move over, as well as descend into, the underlying asthenosphere and even deeper. However, since the 57 development of plate tectonics theory, the link between geodynamics and the compositional 58 differentiation of lithospheric and asthenospheric mantles has received relatively little attention. Most 59 commonly, both plates and underlying asthenosphere are treated in geophysical modelling as closed 60 systems evolving separately with little mutual interaction. Petrological and structural studies of 61 exposed mantle lithologies and experimental petrology demonstrate significant interactions between 62 lithospheric and asthenospheric mantles at mid-oceanic spreading ridges. Moreover in recent years, 63 geophysical observations of oceanic plates have demonstrated a significant asymmetry of the oceanic 64 lithosphere on both sides of oceanic spreading ridge axis (Doglioni et al., 2003; Muller et al., 2008; 65 Panza et al., 2010 and references therein) in terms of seismic velocities, density, thickness, and plate 66 geometry (Fig. 1). These observations are fundamental and require an explanation that integrates the 67 whole data in order to improve our understanding of how the lithosphere and asthenosphere interact at 68 and near spreading ridges.

69 Since the 1960s, plate tectonics is conceived as driven either by active mantle upwelling ('bottom 70 up' or alternatively 'ridge push') or alternatively by the negative buoyancy of slabs ('top down' or 71 alternatively 'slab pull'). It is clear that creation of oceanic lithosphere and its subsequent subduction 72 are fundamental mechanisms in governing the cooling of the Earth. In other words, the surface 73 dissipation of both the Earth's internal primordial and radiogenic heat is tied with the re-entrance of 74 the cold lithosphere along subduction zones to determine the cooling of the planet. However, the 75 mechanisms determining mantle drag and mantle convection remain a matter of debate. The 76 asymmetry of subduction zones (those directed to the west are steeper and faster, e.g., Doglioni et al., 77 2007; Riguzzi et al., 2010) and plate motion reconstructions relative to the mantle reference frame 78 support a westerly-directed drift of the lithosphere relative to the underlying mantle, also called net 79 rotation (as a mean value) or westward drift (Wegener, 1915; Holmes, 1944; Carey, 1958; Le Pichon, 80 1968; Bostrom, 1971; Moore; 1973; Ricard et al., 1991; Gripp and Gordon, 2002; Crespi et al., 2007; 81 Cuffaro and Doglioni, 2007). The net rotation or the westward drift of the lithosphere requires both a 82 decoupling at the lithosphere-asthenosphere interface and a mechanism driving this rotation (Doglioni 83 et al., 2011; Doglioni and Anderson, 2015; Doglioni and Panza, 2015). The motion is not east-west, 84 but along an undulated flow that varies from WNW to E-W to ENE trends, along the so-called 85 'tectonic equator' (Crespi et al., 2007). Proposed mechanisms involve either the negative buoyancy of 86 the lithosphere (Ricard et al., 1991), or the astronomical drag induced by the Earth's rotation

combined with tidal friction (Riguzzi et al., 2010). Potentially astronomical forces may interact with
mantle convection resulting from the cooling of the Earth (Holmes, 1944; Carey, 1958; Scoppola et
al., 2006).

90 The aim of our paper is to build an oceanic spreading model integrating existing compositional, 91 structural and geophysical data across oceanic ridges and the underlying upper mantle. We show the 92 interdependence between the oceanic spreading process, the upper mantle differentiation and plate 93 kinematics driven by the westward drift of the lithosphere. Our model enables to integrate, in space 94 and time, natural and experimental petrology results on upper mantle differenciation (down to 300 km 95 depth) and oceanic, tholeiitic and alkalic, basalt genesis (Green et al., 2014; Green 2015; Green and 96 Falloon, 2015 and references therein), and provides a mechanism for upper mantle geophysical 97 stratification with lateral heterogeneity as evidenced by Thybo (2006).

We will use the well accepted terminology of geophysicists, i.e., upper asthenosphere as the equivalent of "petrological' asthenosphere" corresponding to the Low-Velocity Zone (LVZ), and lower asthenosphere as the equivalent of "petrological' sub-asthenosphere" that reaches about 410 km of depth (Green and Falloon, 1998). They are defined in terms of depth, thickness and composition later in this paper.

103 2. Asymmetry on both sides of oceanic spreading axis and preliminary model

104 Since the recognition of magnetic anomalies on both sides of oceanic ridges (Vine and Matthews, 105 1963), oceanic rift zones were considered as having symmetric features, and plate spreading as a 106 symmetric geological process. As shown by Doglioni et al. (2003), observations on the bathymetry of ridge flanks (shallower eastward) of Pacific and Atlantic oceans and on variable rates of oceanic 107 108 spreading (Muller et al., 1997; 2008; Mallows and Searle, 2012) contradict the concept of full 109 symmetry. Panza et al. (2010) and Panza and Romanelli (2014) provided evidence for asymmetry in 110 both shear wave velocity and thickness within mantle lithosphere and asthenosphere when comparing 111 the two sides of most spreading ridges. In general, the old (>60 Ma) western lithospheric plates have a 112 faster shear wave velocity and are thicker (≈ 100 km versus ≈ 80 km in the eastern flank). Seismic 113 tomography (Thybo, 2006; Panza et al., 2010; Schmerr, 2012; Panza and Romanelli, 2014) provides 114 robust evidence of an asymmetric LVZ at depths from about 80-100 km down to 180-225 km. The 115 average Vs within the western LVZ is slower than the eastern one. The LVZ shape is itself 116 asymmetric, being thicker and wider on the western side of ridges. Also passive continental margins 117 and the related continent-ocean transition support a systematic asymmetry of rift zones (Lavier and 118 Manatschal, 2006; Brune et al., 2014).

All these asymmetric features appear to be independent from the age of the oceanic lithosphere, although they become more prominent moving toward older lithospheric ages (Panza et al., 2010). These observations suggest that mantle compositions and related physical parameters such as rigidity μ , density ρ and their ratio μ/ρ , are systematically different from one side to the other of a spreading ridge within both the lithosphere and the upper asthenosphere, i.e., the LVZ.

Panza et al. (2010) explain these observations as a result of the depletion of the upper asthenosphere, that evolves from pre-melting lherzolite into post-melting harzburgite composition below the oceanic ridge, while the ridge is moving westward relative to the asthenospheric mantle, i.e., in the hot-spot reference frame (Crespi et al., 2007). The role of ridge migration (Scheirer et al., Small and Danyushevsky, 2003; Doglioni and Panza, 2015) in producing an oceanic plate asymmetry is a key parameter of the model. To the east of the ridge, the lithospheric mantle (named also LID) would represent the residual product from partial melting of upwelling asthenosphere,

- 131 abandoned and in the process of cooling after the ridge migration to the "west". Since ridges are
- shifting above a fertile asthenospheric mantle, there is a continuous supply of MORB source mantle
- beneath the migrating spreading ridge (Panza et al., 2010). This model, deduced from geophysical and
- kinematic data (Fig. 1), suggests that the westward drift of the lithosphere relative to the underlying
- 135 asthenosphere is a global phenomenon, a concept that implies decoupling between lithosphere and
- 136 asthenosphere occurring within the LVZ, i.e., in the upper asthenosphere.
- 137 Important questions that arise from the above observations are as follows:
- 138 i) How does oceanic lithospheric mantle form?
- 139 ii) How can the mantle lithosphere composition evolve differently on each side of the ridge?
- 140 iii) How to combine oceanic spreading and westward drift of the lithosphere? Could the detachment-
- 141 mode of seafloor spreading (Maffione et al., 2013; Whitney et al. 2013 and references therein), be the 142 visible major effect in the field?
- 143 iv) How does asthenosphere evolve in relation with oceanic lithosphere during plate spreading? What
- 144 controls the differentiation between lower and upper asthenosphere and the rather homogeneous
- 145 thickness (\approx 125km) of the upper part?
- 146 v) How to combine the ascent of MORB source mantle from the lower asthenosphere according to
- experimental P-T-depth conditions (Green et al., 2014) with oceanic spreading and westward drift of
- 148 the lithosphere?
- 149 vi) Is the oceanic lithosphere denser than the underlying asthenosphere?

150 3. Deciphering the compositional and physical evolution of the oceanic lithospheric mantle on 151 both sides of a spreading ocean ridge

- Lithospheric plates of slow spreading oceans (up to 4cm/yr, e.g., Atlantic, Arctic, southwest Indian and Antarctic) are known to be mostly composed of mantle rocks. Volumes of extrusive and intrusive magmatic rocks, representative of the magmatic oceanic crust, remain minor (Cannat, 1993; Chalot-Prat, 2005; Smith et al., 2008, 2012). Lithospheric mantle is thus the main component of the oceanic plate, which reinforces the major role of oceanic mantle for understanding the nature and construction of the ocean floor. For this reason the following discussion (below) will focus firstly on the physical then secondly on the petrological features of the lithospheric mantle.
- 158 then secondly on the petrological features of the hthospheric mainter
- 159 *3.1. General physical features and composition of oceanic lithosphere*
- 160 The oceanic lithosphere is deeply fractured at the ridge axis. This major fracture network reaches 161 the top of the ductile layer in the lithosphere, and two plates, A and B, form and spread apart. The 162 western plate A always moves westerly faster than the eastern plate B (Fig. 2) as shown by Crespi et 163 al. (2007). In our model, due to the westward drift of the lithosphere (Le Pichon, 1968; Doglioni et al., 164 2003), both plates move to the west, the western plate A at VA and the eastern plate B at VB, being 165 VA>VB (Fig. 2). The oceanic ridge moves west at Vr = (VA + VB)/2. Plate growth rate is due to the 166 difference between VA and VB. Thus the separation between points on each plate increases at the half 167 spreading rate (hS) which is given by (VA-VB)/2. Vr is always faster than hS.
- 168 The thickness of the oceanic lithosphere can be estimated from geophysical data including seismic, 169 gravity, and electrical conductivity It can also be inferred from geotherm/lherzolite solidus 170 intersection based on experimental studies at controlled P, T and water contents.
- 171 Geophysical data show lithospheric thickness decreasing towards ridge axes above shallowing 172 asthenosphere. However we do not think that the lithosphere-asthenosphere boundary (LAB) reaches 173 the base of the newly formed basaltic crust at fast spreading ridges, nor approaches the sea floor at

174 magma-starved slow-spreading ridges. This view is based on the petrology (phase assemblages), 175 chemical compositions and isotopic compositions of peridotites sampled from rifted margins, 176 ridge/transform intersections and slow-spreading ridges. Only lithospheric (secondary harzburgite or 177 secondary lherzolite; see Piccardo et al., 2007b), but not asthenospheric mantle compositions have 178 been identified either at the bottom of rift axis in slow-spreading active and fossil oceans (e.g., Seyler 179 and Bonatti, 1997; Dijkstra et al., 2001; Takahashi, 2001; Tartarotti et al., 2002; Müntener et al., 180 2004; Piccardo et al., 2004, 2007a b, 2014; Kaczmarek and Müntener, 2008, 2010; Rampone et al., 181 1997, 2005, 2008; Rampone and Borghini, 2008; Warren and Shimizu, 2010; Muntener et al., 2010; 182 Piccardo and Garnieri, 2010; Rampone and Hofmann, 2012; Warren, 2016), or below oceanic crust in 183 fast-spreading active and fossil oceans (Dick and Natland, 1996; Abily and Ceuleneer, 2013). The 184 depth of the LAB under the ridge may be inferred in two ways: 1. estimating the depth of melt 185 segregation for parental MOR picrites or basalts and 2. by the P, T conditions recorded in ridge 186 peridotites (including their P, T decompression paths in some cases). Indeed experimental results (Fig. 187 3) demonstrate that parental MOR picrite segregation from residual peridotite occurs mostly between 188 2 to 1.6 GPa (up to 2.2 GPa; down to 1.1 GPa), thus between 55 to 45 km depth (down to 65 km; up 189 to 30 km), around 1400°C (up to 1430°C) from a trace element depleted lherzolite (Eggins, 1992a,b; 190 Falloon et al., 2007a,b; Green et al., 2001; Green and Falloon, 2005; Green et al., 2014; Green and 191 Falloon, 2015). Magma genesis produces mantle residual peridotite less dense than the asthenospheric 192 protolith, accreting above its asthenospheric source and potentially subject to percolation-reaction of 193 new magma ascents, giving birth to a refertilized constantly growing oceanic lithospheric mantle. 194 From experimental results on liquid compositions in equilibrium with spinel and/or plagioclase 195 lherzolite assemblages (Falloon et al., 2007a, b; Chalot-Prat et al., 2010; 2013), analogous to natural 196 oceanic mantle samples, the oceanic lithospheric mantle thickness at the ridge is at least 20 km and up 197 to 40 km in thickness. Note that density differences are due to chemical/mineralogical differences but 198 also to temperature differences (e.g., residual mantle temperature has been reduced by the lost of the 199 latent heat of melting). Thus determining density differences is subject to significant uncertainties.

200 Laterally on both sides of the ridge axis, seismic data show a progressive thickening of oceanic 201 mantle lithosphere until around 80-100 km maximum, equivalent to 3 GPa of pressure. This 202 thickening is assumed to derive first from the addition of the cooled uppermost part of asthenosphere 203 (below 1100°C; Green and Falloon, 2005; Green et al., 2014). Green and Falloon (2005) and Green et 204 al. (2010, 2011, 2014) have shown experimentally that the lithospheric mantle thickness depends on 205 its water storage capacity (Fig. 3). Indeed at pressures greater than 2.5 to 3 GPa (80 to 100 km depth), 206 instability of pargasite causes a sharp drop in the water storage capacity of fertile lherzolite, and for 207 water content greater than ~ 200 ppm in the lithosphere, the mantle solidus is depressed at 2.5-3 GPa 208 along the geotherm. In both cases partial melting can explain both the thickness of the oceanic 209 lithosphere and the occurrence of a LVZ just below the LAB. Therefore from a petrological point of 210 view, the LAB detected by the slowing of seismic waves is the intersection of the geotherm with the 211 vapour-undersaturated lherzolite solidus, and the LVZ is a layer in which there is incipient melt 212 (probably <1-2 % melt).

As far as it concerns the depth/temperature of the base of the oceanic lithosphere outside the ridge (>500 km), the model of Afonso et al. (2007) predicts 1300°C at 105 km depth. This temperature is somewhat too high according to experimental data of Green et al. (2010, 2014), showing that a "wet" (>200 ppm H₂O) oceanic lithospheric mantle is above its solidus below 80-100 km if temperature is higher than 1000 °C (Fig. 3). Below this depth, this is the LVZ, thus the upper asthenosphere. An incipient melt is present, the melt fraction being determined primarily by the water and CO₂ contents. This melt has low permeability (i.e. can only move slowly by porous flow), but has a large effect on seismic and rheological properties.

3.2. How oceanic mantle lithosphere of the western limb of the ridge axis becomes thicker and densercompared to the eastern limb?

223 *3.2.1 Intra-lithospheric mantle decoupling*

224 In this section we will focus on the modalities of mantle mass transfer upwards and laterally within 225 both spreading plates in the framework of the westward drift of the lithosphere. Oceanic mantle 226 lithosphere is created at the ridge through accretion of residues above the asthenospheric partial 227 melting area. These residues are successively transferred upwards and laterally at hS rate within the 228 mantle lithosphere on each side of the ridge (Fig. 2). As Vr is higher than hS, the transfer rate of 229 successive residues above the melting area is slower than the westward velocity of the melting area 230 itself. Accordingly, mantle exhumed at the bottom of any active axial ridge is always older than the 231 basaltic volcanoes emplaced at the axial ridge (Bonatti et al., 2003). It is the same for intrusive gabbro 232 bodies which move upwards and laterally, driven by and with the upwelling of the host-mantle: for 233 instance, for hS=2cm/yr, Vr=4cm/y and a mantle residue initial depth of Di=40 km, the mantle residue 234 takes t1 2 Ma (t1=Di/hS) to ascend and to be exhumed at the surface. Thus active volcanoes at the 235 ridge axis are about 2 Ma younger than the mantle on which they are formed. Besides, while this 236 mantle residue is rising up to the ocean floor, the ridge itself with the uppermost part of the 237 lithosphere will have moved (t1*Vr) 80 km westward. However it may be possible that due to 238 'second-stage melting', the extra buoyancy added to the residue may accelerate the upwelling – in 239 which case some residue may be younger than 2Ma.

240 Therefore, besides the notion that asthenosphere is moving "eastward" relative to the lithosphere 241 (or more precisely along the flow lines parallel to the undulate tectonic equator of Crespi et al., 2007 242 and Cuffaro and Doglioni, 2007), the westward drift of the lithosphere is necessarily slowed down, 243 top to down, inducing a decoupling within the mantle lithosphere itself. This is visualized on the 244 model (Fig. 2) with the progressive top-down eastward shifting of the "half-spreading residues" 245 relative to each other within both plates. This decoupling would be progressive such that the top of the 246 melting area would be somewhat delayed relative to the top of the axial ridge, even if they are 247 interdependent.

248 Normal faults are the natural feature associated with rift zones and as indicated by rock mechanics, 249 their mean dip is around 60°. Lower dips may naturally result from a decrease of friction due to the 250 high heat flow and moving into shear zones within the underlying ductile layer. Besides faults and 251 fractures may dip predominantly in one direction if spreading is asymmetric. Indeed the half spreading 252 rate is often slightly asymmetric (Mueller et al., 2008), but this is related not to the velocity of plates 253 relative to the mantle, but to the velocity of the ridge which sometimes is not exactly (VA+VB)/2; it 254 may be (VA+VB)/1.95 or (VA+VB)/2.05 etc. Therefore faster and larger spreading may occur in one 255 flank without varying the relative velocity of the two plates with respect to the mantle. This can be 256 explained by the lateral viscosity variations in the flanks right beneath the ridge. Therefore, ridge-257 related faults and associated fractures would dip mainly "eastward" and be convex upwards since the 258 westward drift is faster upwards, but their dip may also be opposite due to the half-spreading process. 259 This is the geometry of detachment faults, i.e., large offset low-angle faults capping the eastern (rarely 260 western) side of oceanic mantle core complexes (OCC; Smith et al., 2008, 2012; MacLeod et al., 261 2009; Reston and Ranero, 2011; Mallows and Searle, 2012; Whitney et al., 2013; Maffione et al., 262 2013) and extending along steep faults probably rooted in a melt-rich zone. They are responsible for

7

exhumation of mantle and plutonic rocks onto the seafloor in the footwall of normal faults, whereasextrusion of basalt occurs on the hanging wall.

265 This intra-mantle decoupling within the lithosphere means also that even a mid-oceanic ridge and 266 its related fracture network shift probably westward within the lithosphere, moving itself westward 267 relative to the asthenosphere. This cannot be visible from the surface, and this is why general 268 symmetry of magnetic stripes on the oceanic sea floor is preserved. However Carbotte et al. (2004) 269 and Mallows and Searle (2012) note a westward shift of volcanic activity with time along the active 270 Mid-Atlantic ridge axis. The geological maps of Mallows and Searle (Fig. 4; 2012) and Smith et al. 271 (Fig. 4; 2012) show that most OCC outcrops on the western side of the ridge and that spreading rates 272 are asymmetric with faster spreading on western plates. According to Whitney et al. (2013 and 273 references therein), OCC account for 60% to 100% of the total plate spreading, showing that tectonic 274 process is the main driver of mantle exhumation in the oceans and of heat and mass transfer in the 275 Earth. Furthermore, Maffione et al. (2013) introduce the concept of detachment-mode of seafloor 276 spreading. Moreover, the usual sill shape of gabbro intrusive bodies both at multimetric and 277 multikilometric scale (Henstock et al., 1993 and references therein; Chalot-Prat, 2005) testifies sub-278 horizontal magma injections within weakness zones of lithospheric mantle, supporting intra-mantle 279 decoupling during ocean spreading. Volcanoes grow on the topmost part of the hangingwall of these 280 large detachment faults, structuring hangingwall rider blocks (Whitney et al., 2013; Maffione et al., 281 2013 and references therein). Small-scale decoupling occurs between the base of volcanoes just 282 emplaced and underlying mantle or/and gabbro bodies in the process of exhumation. This decoupling 283 occurs on both sides of relief where eruptions take place, underlining a pseudo-symmetric spreading 284 (Chalot-Prat, 2005).

3.2.2. Asymmetric evolution of lithospheric mantle compositions and physical parameters on each side of the ridge

287 The variation of Vs between the two sides of the oceanic ridge should have an explanation in terms 288 of composition related to the different kinematics. As the melting area moves westward with the 289 migrating spreading axis, the eastern plate B incorporates successively the abandoned depleted eastern 290 parts of the ridge. It undergoes a relatively brief time of refertilization by ascending MORB and its 291 composition remains close to a harzburgitic residue and is relatively Fe-poor. At the same time, 292 harzburgitic residues are also continuously added to the western plate A, but, in contrast to the eastern 293 plate, they are also continuously percolated by MORB as the ridge is moving westward above the 294 underlying melting area linked to the ridge location. Thus the composition of the western plate A is 295 that of a strongly refertilized harzburgite, thus a secondary lherzolite. An outcropping example could 296 be the lherzolites from the Lanzo oceanic mantle area (Piccardo et al., 2007, and references therein). 297 Therefore lithospheric mantle of the plate A is continuously re-enriched in Fe and Ca, whereas 298 lithospheric mantle of the plate B remains a harzburgitic residue barely re-enriched in Fe and Ca.

Nevertheless as Vr>hS, the axial ridge is moving westwar within the lithosphere itself, and the western plate A is progressively incorporated within the eastern plate B. This means that the plate B composition will become somewhat more refertilized than presented above. However the overall consequence is that the western plate A is thicker, always richer in Fe and thus denser than the eastern plate B, resulting in a first order asymmetry in composition between the two plates.

In addition, if the westward drift is producing intra-lithospheric deformation preferentially on the western plate – then we might expect easier and more abundant pathways for off-axis magmatism and hence off-axis melt impregnation and channelling. Some of the upwelling diapirs may lose part of their melt fraction before arriving directly under the ridge axis. Evidence for this could be an 308 asymmetry in abundance of off-axis seamounts (i.e. volcanic chains on western plate as in Shen et al.,

- 309 1993, White et al., 1998). If this off-axis process occurs then there will be less melt arriving directly
- 310 under ridge axis and hence the eastern flank will receive long-term less fertilised residue than the western flank.
- is higher than ρB (previous paragraph), μA must be much higher than μB . Since the modulus of rigidity of Fe is around 3 to 4 times higher than that of Mg and Si, the rigidity of rocks is expected to be enhanced if they are richer in Fe-rich minerals (Fyfe, 1960; Jordan, 1979). Thus higher μA compared to μB is an expected natural consequence of enhanced melt refetilization under the western
- 469 plate A during westward migration of the spreading ridge.

470 4. Deciphering the physical and compositional evolution of the asthenosphere below a spreading471 oceanic lithosphere

472 *4.1. Lower asthenosphere*

473 The lower asthenosphere is subsolidus garnet lherzolite, about 180 km thick (230 to 410 km depth) 474 overlying the Transitional Zone beginning at ~ 410 km (Anderson, 2007; 2010; Schmerr, 2012). From 475 the top to the bottom of this layer, seismic waves gradually increase their speed with respect to the 476 LVZ. Vs is much higher than in the upper asthenosphere and more or less similar to that of the 477 lithosphere (Panza and Romanelli 2014). Its mineralogical and chemical composition is inferred from 478 experimental petrology on MORB genesis (Green et al., 2014; Green, 2015), being the N-MORB 479 mantle source located below 230 km. It is a fertile garnet lherzolite, depleted in the most incompatible 480 trace elements, with \geq 200 ppm H₂O and with incipient melt at \leq 7 GPa/~230 km, 1490°C. As shown 481 in Fig. 3, an oceanic intraplate geotherm joins the mantle adiabat at about 230 km depth defining the 482 base of the upper asthenosphere. The lower asthenosphere may migrate upwards as diapirs within the 483 230-100 km including very small, relatively immobile, near-solidus melts (0.05–0.1 % of melting).

484 *4.2. Upper asthenosphere*

485 Within the upper part (80-220 km depth) of the asthenosphere, i.e., the LVZ, Vs is slower in the 486 western than in the eastern side of the ridge, the reverse of what it is observed in the oceanic 487 lithosphere (Panza et al., 2010). We infer that during its transfer from west to east below the oceanic 488 lithosphere, the eastern part of upper asthenosphere becomes somewhat Fe-depleted by incipient 489 melting (Fig. 2) inducing a density decrease for a potentially similar value on rigidity from one side to 490 the other of the axis. Moreover as shown by experiments of Conder and Wiens (2006), Hammond and 491 Toomey (2003) and Green et al. (2014), the western part of the upper asthenosphere is hotter and/or 492 richer in H₂O+CO₂ than its eastern part and includes higher melt fractions inducing a rigidity decrease 493 in the western side. It follows that to the "west" of a spreading ridge, both the oceanic lithospheric 494 mantle (down to 90-100 km) and the upper asthenosphere (down to 230 km) should be more fertile 495 and denser than the "eastern" part. Let us remember that in both mantle layers, density is determined 496 not only by mantle fertility and Fe-enrichment, but also by the pargasite content in the lithospheric 497 mantle and by percentage of melt within the upper asthenospheric mantle.

The LVZ has been seismically recognized below all oceans, both beneath the "western" and the "eastern" plates. It has a very small aspect ratio (thickness: 130 km; length in spreading direction: $6*10^3$ km below the Atlantic to $11*10^3$ km below the Pacific). Experimental results (Fig. 3) show that the upper asthenosphere corresponds petrologically to a mantle layer including small interstitial melt fractions (i.e. 0.05 to 0.1 %). According to Doglioni et al. (2005), the degree of partial melting would be increased (up to 1.5 %) by shear heating (more than 100°C) generated by decoupling between lithosphere and lower asthenosphere (Fig. 3) and with largest contribution i.e. largest E-W relative movement at ~230 km depth. It may be alternatively due to higher H₂O content in the mantle (Bonatti, 1990; Ligi et al., 2005).

507 Moreover, seismic images from Panza and Romanelli (2014) show an asymmetric shape of the 508 LVZ, thicker and more elongated westward (2/3 in volume) than eastward (1/3 in volume) of ridge 509 axis in all oceans. This spectacular and permanent asymmetry of the LVZ shape raises the question 510 not only of the origin of partial melting, but also of its much greater development below the western 511 part of the oceans. From experimental results (Fig. 3), the presence of these very small melt fractions 512 have two distinct origins depending on the H₂O content of the mantle sources: 1- below the 513 lithosphere-asthenosphere boundary (LAB) and far away from the ridge (>500 km): because of 514 destabilization of amphibole, below 2.5-3 GPa/ 80-100 km/ 1000-1100°C, when "wet" (>200 ppm 515 H2O) oceanic mantle lithosphere, i.e. a subsolidus pargasite-bearing garnet lherzolite cools, subsides 516 and becomes asthenosphere, i.e. a pargasite-out garnet lherzolite with incipient melting. 2- above the 517 lower-upper asthenosphere boundary: during adiabatic ($\approx 1490-1450^{\circ}$ C) "dry" (<200 ppm H2O) ascent 518 of the lower asthenosphere from \geq 230 km/ \geq 7 GPa towards the MORB genesis area, i.e. 65-45 km/2-519 1.6 GPa below the ridge axis. Let us note that if the first partial melting origin corresponds to a rather 520 static process linked only to the permanent thickening of the lithospheric mantle, the second is 521 synchronous with a dynamic process of partially molten transfer of matter, both being imaged by the 522 LVZ.

In parallel, the LVZ asymmetry suggests that the on-going ascent of partially molten mantle from lower asthenosphere is oblique, and not vertical, as already proposed by Carbotte et al. (2004). Taking into account its dimension, the ascent would be initiated around some thousands kilometers "west" from the ridge (Fig. 2), and driven by a suction effect towards the shallow area below the spreading ridge moving "westward", due to the removal of the overlying lithosphere and the consequent asthenospheric isostatic compensation.

529 Nevertheless according to the entrance angle of lower asthenosphere into upper asthenosphere, this 530 lateral and upward mass transfer is more or less counteracted by the relative "eastward" horizontal 531 mantle flow linked to the net rotation of the lithosphere, whose origin can be ascribed either to lateral 532 viscosity variation (e.g., Ricard et al., 1991) or to the Earth's rotation (Scoppola et al., 2006; Riguzzi 533 et al., 2010). Two different paths of transfer would exist. The main path involving the largest mantle 534 volume follows a very low angle trajectory, barely deviated in passing below the spreading ridge; it 535 would concern the mantle flux emerging rather far laterally (more than 500 km?) from the spreading 536 centre. From experimental results (Fig. 3), this mantle flux crosscuts the intraplate geotherm, modified 537 by shear heating, and undergoes incipient to low degrees of partial melting (up to 4% down to 80-100 538 km deep / 2.5-3 GPa, Green and Falloon, 2005). The other path is single inasmuch as, all things 539 (composition, flowing velocity, temperature, pressure) being equal, a more opened entrance angle of 540 mantle flux at a shorter distance from the ridge (500 km or less?) enables it to be mainly monitored by 541 the suction effect coming from the strongly depressurized shallow area below the spreading ridge. 542 This single path is assumed to be the one producing MORB after 15-20% melting between 65 and 45 543 km depth (2 to 1.6 GPa) (see section 5.1 below). Indeed at small-scale, we suggest that convective 544 eddies form by peeling off the upper surface of the lower asthenosphere and ascend driven by 545 buoyancy and rheology contrast with the upper asthenosphere. These ascending eddies would become 546 'isolated' or encapsulated by a cooling rind/skin as the temperature contrast increases between eddy 547 core and ambient mantle, forming diapirs. These eddies/diapirs move obliquely towards the shallow 548 area below the spreading ridge where they are tapped by fracturing at the rift.

549 "Eastward" of the ridge, because of the "westward" drift of the lithosphere, the suction effect fades 550 and mantle melting progressively mitigates top to bottom to become quite insignificant at more than 551 1000 km from the ridge. Accordingly in this model, off-axis melt lenses on the eastern side of the East 552 Pacific Ridge (Canales et al., 2012; Toomey, 2002, 2012) witness the asthenospheric partial melting 553 zone underlying the ridge just before eastward abandonment due to the westward migration of the 554 ridge.

According to this new scenario, the westward drift of the lithosphere and the related oceanic plate spreading have a strong mechanical effect on lateral (several thousands kilometers long) and upward (from 230 to 45 km depth) mantle mass transfer below the western plate A. This effect appears as extremely relevant in terms of mass transfer and is synchronous with partial melting and percolation/reaction processes, and thus mantle differentiation.

- 560 Therefore, as already inferred by Green et al. (2014) and Green (2015), the upper asthenosphere 561 mantle composition is heterogeneous at a small scale, being influenced from below by the ascending 562 lower asthenosphere and from above by the oceanic lithosphere during its journey and related process 563 of thickening. In other terms, from 230 km to 90-100 km, the upper asthenosphere should be a mantle 564 layer including incipient melt fractions coming from two distinct mantle sources: the lower 565 asthenosphere, the "dry" lherzolitic source of MORB on one hand; the lowermost part of lithosphere, 566 made of harzburgitic residues produced below the ridge axis, more (westward) or less (eastward) 567 refertilized and "wet" once accreted to the lithosphere.
- 568 All these data confirm that the asthenosphere forms a whole, 300-320 km thick, with a lower 569 "nearly dry" (~200 ppm H₂O) solid part (\approx 185 km thick; 2/3 in volume) and an upper "relatively wet 570 (> 200 ppm H₂O) solid + incipient melt" part (\approx 125 km thick; 1/3 in volume), the latter interacting 571 with the overlying lower lithosphere. The existence of the asymmetric LVZ below the oceans 572 indicates that below divergent plates, upwards and "eastward" transfer of lower asthenosphere is more 573 active than elsewhere, accelerating its replacement by eastward lateral mantle flow coming from 574 below the adjoining continents and related passive continental margin located to the "west" of a given 575 ocean. So before flowing below and interacting with oceanic lithosphere, the upper asthenosphere 576 underwent an earlier history of interaction with continental lithosphere, hence adding further chemical 577 heterogeneity to intraplate mantle sources.

578 5. How to understand, within the framework of plate tectonics, the coexistence of mid-oceanic 579 basalt and oceanic intraplate basalt mantle sources within the asthenosphere?

580 5.1. Location of mid-oceanic basalt mantle sources

Experimentally (Fig. 3), MORB are shown to be sourced from a roughly adiabatic (1490 to 1430°C) upwelling of the uppermost part (250-230 km) of lower asthenosphere with melt fraction increasing dramatically above its anhydrous solidus (3 GPa/~100 km; 1450°C). MORB melt segregation occurs at 15-20% melting between 2 and 1.6 GPa (65 to 45 km; Green and Falloon, 2005; Falloon et al., 2007b) within the so called "melting area" below the ridge axis in our model (Fig. 2).

The MORB mantle source composition is lherzolitic and fertile in terms of mineral assemblage and major element contents (Green and Falloon, 1998; 2005). However both trace element contents and Nd-Sr-Pb isotopic signatures are those of a mantle depleted in the most incompatible trace elements, *interpreted* as representative of a residual mantle composition. This decoupling between major and trace element composition interpretations is explained considering that the MORB source is itself a secondary lherzolite coming from the refertilization of a mantle residue by melts extracted from an already trace element depleted lherzolitic mantle (Frey and Green, 1974; Piccardo et al., 2007), or an eclogitic magmatic crust (Eller et al., 2000), or recycled components (e.g. oceanic crust, Eiler et al., 2000; or metasomatized oceanic lithosphere, Niu et al. 2002; Shimizu et al., 2015), or a mix (Rosenthal et al., 2014). So our model assumes that the uppermost part of the lower asthenosphere is a secondary lherzolite, being the refertilization mostly provided from the recycling of W-directed subducting oceanic lithospheric slabs (Rosenthal et al., 2014; Doglioni and Anderson, 2015; Green, 2015).

599 As shown by our model, the unique and unidirectional path of transfer of a mantle with a MORB 600 source signature, combined with the westward drift of the lithosphere and of the spreading ridge itself, 601 highlights the systematic and permanent identical replacement of the MORB mantle source within the 602 same P-T conditions of ascent and partial melting, which fits with the observed relative compositional 603 homogeneity of MORB (dominant N-MORB) through geological times. This compositional 604 homogeneity testifies itself for an absence of significant interactions of this oblique succession of 605 mantle diapirs (Green, 1971) crosscutting the surrounding upper asthenosphere, and therefore for the 606 relatively high velocity of such mantle diapirs driven by the suction effect of the spreading ridge. 607 Diapir buoyancy driven by density, rheology and isolation from wall-rock reaction, is probably more a 608 consequence of increasing temperature contrast if the diapir is near adiabatic i.e garnet lherzolite with 609 incipient melt into a rind of spinel lherzolite to plagioclase lherzolite (foliated to mylonitic).

610 5.2. Location of oceanic intraplate basalt mantle sources

611 As summarized by Green and Falloon (1998; 2005), Green et al. (2014), Green (2015) and Green 612 and Falloon (2015), intraplate magmas, including oceanic intraplate basalts (or "hot spots" basalts), 613 range from olivine melilitites and nephelinites, to olivine-rich basanites and finally to alkali basalts 614 and olivine tholeites. Experimentally, these basalts represent low degree (2 to 10%) melts from 615 mantle sources located within the uppermost asthenosphere on 50 km thickness (\approx 110-160 km depth; 616 \approx 3.3 to 5 GPa) or only 20 km thickness (\approx 105-125 km depth; \approx 3 to 4 GPa) depending on the 617 considered geotherm ("intraplate" or "intraplate+ shear heating effect"; Fig. 3). This upper 618 asthenospheric source mantle is lherzolitic as the MORB mantle source and thus fertile in terms of 619 mineral assemblage and major element contents, but above all rather enriched, and not depleted as the 620 MORB source, in water (> 0.02%) and in the most incompatible trace elements. As inferred in the 621 previous chapter, they are secondary lherzolites in which refertilization of mantle residues can have 622 two distinct origins. On one hand the lherzolites could derive from destabilization (\geq 90-100 km 623 depth) of the lowermost part of the subsiding oceanic lithosphere far from the axial ridge. This result 624 validates the hypothesis of an origin of some intraplate basalts from the lower lithospheric mantle in 625 terms of composition (Anderson, 2010, and references therein). In that case, mantle metasomatism 626 occurred below the spreading ridge by percolation/reaction of lithospheric mantle by MORB a rather 627 long time ago. On the other hand, these secondary lherzolites could derive from metasomatism of 628 lower asthenospheric material ascending throughout the upper asthenosphere along the main path with 629 a very low angle trajectory (Fig. 2; see § 4.2 for explanations). In that case, metasomatism results 630 from successive percolation/reaction of incipient melts with mantle constantly on the move laterally. 631 Nevertheless the Vs variations in the LVZ from one side to another of the ridge axis (cf. § 4.2) 632 suggests that the uppermost mantle fertility is somewhat reduced eastward.

Besides the Nd-Sr-Pb isotopic signatures of intraplate basalts attest for mantle sources also metasomatized at some point in their long history by continental lithospheric products (Pilet et al., 2005; 2008; 2011). Notice that most Pacific, but also Indian "hot spots" basalts (dominant alkali intraplate basalts) are on the western or south-western plates relative to their respective axial ridges, thus are sourced within the upper asthenosphere having previously interacted with western adjoining and overlying continental plates. Such an interaction may be a candidate for explaining this
 continental imprint on intraplate basalt mantle sources, coherently with the relative "eastward" mantle
 flow.

641 6. Towards an alternative oceanic plate spreading model for a polarized plate tectonics

642 To date, a number of geophysical data at different scales demonstrate asymmetry of oceanic plates, 643 whereas mantle petrologists still conceive mantle differentiation as a symmetric process on both sides 644 of divergent plate boundaries and the underlying asthenosphere. To resolve this inconsistency, 645 combination of updated geophysical, structural and petrological data on oceanic upper mantle (down 646 to about 300 km) and also related concepts on mantle and magma genesis and plate kinematics, leads 647 to the development of an oceanic spreading model where both the growing oceanic lithospheric 648 mantle and the underlying upper asthenospheric mantle are to varying degrees asymmetric in 649 composition. Our model takes place in the frame of the westward drift of the lithosphere, which 650 means that oceanic ridges migrate laterally to the "west" relative to the asthenospheric mantle, and 651 thus are always moving over a fertile mantle.

From the geophysical data, the asymmetry concerns bathymetry of the ridge flanks (less steep eastward), plate velocity rates (slower for the eastern one), thickness of plates (thinner eastward), and faster shear wave velocity in the western lithosphere. It comes that plates on each side of the ridge differ in density as well as rigidity, thus in composition, and thereby in growing conditions of mantle lithosphere at the axial ridge.

To understand from a petrological but also kinematic point of view the aforementioned asymmetry, mantle lithosphere and mantle asthenosphere were examined separately, even if they are interdependent.

660 From natural and experimental data, three processes, two just below and one outside the spreading 661 ridge, are known to generate compositional heterogeneity of oceanic mantle lithosphere of both plates: 662 1) sub-ridge accretion of partial melting residues formed by melt segregation from mantle 663 asthenosphere upwelling; mantle lithosphere thickness at the axial ridge is estimated, via experimental 664 results on mantle phase equilibrium, between 30 to 80 km; 2) melt-rock reactions or mantle 665 metasomatism by basaltic melt percolation through previously accreted residues below the ridge, 666 inducing refertilisation and secondary lherzolite formation; 3) capture and addition of asthenospheric 667 mantle to the base of mantle lithosphere, because of cooling of the uppermost asthenosphere far from 668 the ridge. This last process occurs when the geotherm drops below 1050°C at 3 GPa which yields a 669 LAB at ~90km below which neither pargasite or hydrous carbonate-bearing silicate melt could exist. 670 It gives a thickened lithosphere with the growing layer of subsolidus phlogopite-bearing garnet-671 lherzolite with carbonatite, or graphite+ H₂O+CH₄ fluid.

672 Structural data released by mapping numerous mantle core complexes, mostly immediately 673 westward of active slow-spreading ridges, give strong structural constraints for understanding 674 lithospheric mantle dynamics during plate spreading. Indeed whether in the west or east, these core 675 complexes represent outcrops of mantle and related intrusive gabbro bodies exhumed along the 676 footwall of convex up-wards detachment faults dipping eastward and rooted down to 7-8 km. We 677 interpret these detachment faults as effects of top to bottom asymmetric lithospheric mantle shear, 678 linked to the eastward flowing of asthenosphere delaying the base of the lithosphere moving 679 westward. The usual sill shape of gabbro intrusive bodies is another observation supporting intra-680 mantle decoupling during ocean spreading. Furthermore, at the topmost part of the hangingwall of 681 these large detachment faults, a small-scale decoupling occurs between the base of volcanoes and 682 underlying mantle or/and gabbro bodies in process of exhumation, underlining this time a pseudo683 symmetric spreading. It follows that whatever the scale, the top to bottom decoupling dynamics 684 enables the exhumation of deep rocks (mantle with or without intrusive gabbro), thus the creation of 685 new surfaces, giving all its meaning to the concept of "detachment-mode of seafloor spreading" 686 developed by Maffione et al (2013). *This top to bottom mantle decoupling preserves totally magnetic* 687 *anomalies recorded at the surface on each side of the axis.*

688 Considering all these results, the mantle lithosphere composition can evolve differently on each 689 side of the ridge, coming from a permanent refertilisation of the western plate whereas the eastern 690 plate just preserves the barely refertilized feature of a harzburgitic residue generated during 691 asthenosphere partial melting below axial ridge. So rigidity, density to a lesser extent and thickness of 692 the western plate become much higher, whereas the shallower bathymetry of the east flank of the 693 ridge is explained by isostatic adjustment because of a somewhat lower density. Also as the eastern 694 plate mantle is lighter and more viscous, there is a higher coupling at the lithosphere-asthenosphere 695 boundary, determining a slower velocity of the eastern plate during ridge migration. The top to bottom 696 and west to east internal decoupling of the mantle lithosphere, linked to its westward drift above the 697 asthenospheric mantle flowing eastward, explains detachment faults and related mantle core complex 698 exhumation.

699 The petrology of asthenospheric mantle is known experimentally since asthenosphere is the mantle 700 reservoir, at distinct P-T-depths, of MORB (lower asthenosphere) and intraplate basalts (middle to 701 upper asthenosphere). An asymmetric "incipient melt + solid" mantle zone, the LVZ, exists between 702 230 and 100 km, being much more developped and more fertile westward. So MORB are sourced 703 from a lherzolitic mantle in equilibrium at $\approx 250-230$ km / 7 GPa, the ascent of which up to 65-45 km 704 / 2-1.6 GPa just below the ridge is roughly adiabatic, thus rather fast and without any significant 705 interaction with surrounding crosscut upper asthenosphere. According to the relative "eastward" 706 direction of the asthenospheric mantle flow lines, the diapiric rise of MORB mantle source from lower 707 asthenosphere is significantly oblique and eastward. This mantle transfer is triggered and checked by 708 the suction effect of the spreading ridge, and the westward ridge migration determines a self-709 perpetuating mechanism for permanently renewing the MORB source mantle, which explains in 710 return the great homogeneity of MORB through time. Synchronously, major mantle transfers within 711 middle to upper asthenosphere come from the lower asthenosphere farther westward and follow the 712 eastward mantle flow lines, slightly slantwise but barely deviated by the suction effect of the 713 spreading ridge. Mantle undergoes incipient melting, except that the interstitial melts percolate 714 laterally and upwards, react with and enrich the upper asthenosphere in water and the most 715 incompatible elements, a possible mantle reservoir for intraplate basalts. This incipient melting zone 716 represents also the lithosphere-asthenosphere decoupling zone affected by shear heating, reducing the 717 P/depth of intraplate magma genesis. Also the higher shear wave velocity eastward means a density 718 decrease in relation with a somewhat residual feature of the eastern asthenospheric mantle and 719 reflecting a mixing with residual mantle coming from the high degree melting area below the ridge 720 and not accreted to the lithosphere.

721 7. Conclusions

The proposed oceanic plate spreading model shows that numerous asymmetric features observed with geophysics and structural analysis in the first 300 km below the oceans are consistent with petrological data on upper mantle and basaltic compositions. Our multi-disciplinary study emphasizes the major role of physical and chemical dynamics of the upper mantle for understanding oceanic plate tectonics, and in turn how evolution of both the composition and internal structure of oceanic plates strongly depends at all scales on plate kinematics. Our model suggests that both plate composition and

- renewal, by mass transfer at all scales, particularly in the shallow 300 km of the Earth's mantle below
- the whole width of the oceans. Another major concept is that all petrological processes, occurring
- during different types of solid or/and liquid mantle mass transfers, ultimately lead to the genesis of a
- more or less fertile mantle composition, i.e. a lherzolite but secondary in nature (residual harzburgite
- refertilized by melt interaction), which is the main composition of the first 300 km of the Earth below
- oceans and most likely the continents as well. Incompatible trace elements abundances and isotopic
- values of mantle sources will necessarily vary from one specific mantle site to another reflecting the
- 737 detailed individual geological histories of 'lherzolites'.

738 Acknowledgments

739 We are grateful for fruitful and helpful discussions to David H. Green, Don Anderson, Eugenio 740 Carminati, Marco Cuffaro and Giuliano Panza. F. Chalot-Prat is grateful to the CRPG for its financial 741 support to allow brainstorming meetings between the authors of this paper. We gratefully 742 acknowledge David H. Green for his thorough, very constructive and interesting review that helped us 743 to deepen our observations and our model, and to an anonymous reviewer for his numerous pertinent 744 remarks and his careful and patient work on the form of the text, usefully clarifying the content and 745 form of this multi-disciplinary study. Our warm thanks go to Marco Scambelluri for his extensive 746 work during the reviewing process and all his attention as editor of this paper. The data used are listed

747 in the references. This is the CRPG-CNRS contribution no. 2433.

748 **References**

- Abily, B., Ceuleneer G. 2013. The dunitic mantle-crust transition zone in the Oman ophiolite: Residue
 of melt-rock interaction, cumulates from high-MgO melts, or both? Geology, 41, 67-70.
 Doi :10.1130/G33351.1
- Afonso, J. C., Fernandez M., Ranalli G., Griffin W. L., Connolly J. A. D. 2008. Integrated geophysical-petrological modeling of the lithosphere and sublithospheric upper mantle: Methodology and applications, Geochem. Geophys. Geosyst., 9, Q05008, doi:10.1029/2007GC001834.
- Anderson, D.L., 2007. New Theory of the Earth, Cambridge University Press.
- Anderson, D.L., 2010. Hawaii, boundary layers and ambient mantle-geophysical constraints. J.
 Petrology, doi: 10.1093/petrology/egq068.
- Bonatti, E., Ligi M., Brunelli D., Cipriani A., Fabretti P., Ferrante V., Ottolini L. 2003. Mantle
 thermal pulses below the Mid Atlantic Ridge and temporal variations in the oceanic
 lithosphere, Nature, 423, 499–505.
- 762 Bostrom, R.C., 1971. Westward displacement of the lithosphere. Nature 234, 536–538.
- Brune S., Heine, C., Pérez-Gussinye M., Sobolev S.V. 2014. Rift migration explains continental
 margin asymmetry and crustal hyper-extension. Nature Communications, 5:4014,
 doi:10.1038/ncomms5014.
- Canales, J.P., Carton, H., Carbotte, S. M., Mutter, J.C., Nedimovi'c, M.R., Xu, M., Aghaei, O.,
 Marjanovi'c, M., Newman, K., 2012. Network of off-axis melt bodies at the East Pacific
 Rise. Nature Geoscience, 5, 279-283.
- Cannat, M., 1993. Emplacement of mantle rocks in the seafloor at mid-ocean ridges. Journal of
 Geophysical Research, Solid Earth, 98, B3, 4163–4172.

- Carbotte, S.M., Small C. Donnelly K. 2004. The influence of ridge migration on the magmatic
 segmentation of mid-ocean ridges, Nature, 429, 743–746.
- Carey, S.W., 1958. Continental Drift—A symposium (Univ. of Tasmania, Hobart, March 1956), 177–
 363.
- Chalot-Prat, F, 2005. An undeformed ophiolite in the Alps: field and geochemical evidences for a link
 between volcanism and shallow plate tectonic processes. In Plates, Plumes & Paradigms,
 edited by G.R. Foulger, D.L. Anderson, J.H. Natland and D.C. Presnall, Geological Society
 of America, Special Paper 388, 751–780.
- Chalot-Prat, F., Falloon T. J., Green D. H. Hibberson W. O., 2010. An experimental study of liquid
 compositions in equilibrium with plagioclase+spinel lherzolite at low pressures (0.75GPa).
 Journal of Petrology, 51(11), 23–49-2376.
- Chalot-Prat, F., Falloon T. J., Green D. H., Hibberson W. O., 2013. Melting of plagioclase + spinel
 lherzolite at low pressures (0.5 GPa): An experimental approach to the evolution of basaltic
 melt during mantle refertilisation at shallow depths. Lithos, 172-173, 61–80.
- Conder, J.A., Wiens, D.A., 2006. Seismic structure beneath the Tonga arc and Lau backarc basin
 determined from joint Vp, Vp/Vs tomography. Geochemistry, Geophysics, Geosystems 7,
 Q03018.
- 788 Crespi, M., Cuffaro, M., Doglioni, Giannone, C., Riguzzi, F., 2007. Space geodesy validation of the
 789 global lithospheric flow. Geophysical Journal International, 168, 491-506, doi:
 790 10.1111/j.1365-246X.2006.03226.x.
- Cuffaro, M., Doglioni, C., 2007. Global kinematics in deep versus shallow hotspot reference frames,
 In: Foulger, G.R., and Jurdy, D.M. (Eds.), Plates, plumes, and planetary processes:
 Geological Society of America Special Paper 430, 359–374, doi: 10.1130/2007.2430(18).
- Dijkstra, A.H., Drury, M.R., Visser, L.R.M., 2001. Structural petrology of plagioclase peridotite in the
 West Othris mountains (Grece): melt impregnation in mantle lithosphere. Journal of
 Petrology 42, 5–24.
- Dick, H.J.B., Natland, J.H. 1996. Late-stage melt evolution and transport in the shallow mantle
 beneath the East Pacific Rise, in Mével, C., et al., eds. Proceedings of the Ocean Drilling
 Program, Scientific Results, 147:College Station, Texas, Ocean Drilling Program, 103–134.
- Boglioni, C., Carminati, E., Bonatti, E., 2003. Rift asymmetry and continental uplift. Tectonics, 22,
 1024-1037, doi:10.1029/2002TC001459
- Boglioni, C., Carminati, E., Cuffaro, M., Scrocca, D., 2007. Subduction kinematics and dynamic
 constraints. Earth Science Reviews, 83, 125-175, doi:10.1016/j.earscirev.2007.04.001.
- Boglioni, C., Green, D.H., Mongelli, F., 2005. On the shallow origin of hotspots and the westward
 drift of the lithosphere. In: Foulger, G.R., Natland, J.H., Presnall, D.C., and Anderson, D.L.
 (Eds.), Plates, plumes, and paradigms: Geological Society of America Special Paper 388,
 735–749, doi: 10.1130/2005.2388(42).
- Boglioni, C., Ismail-Zadeh, A., Panza, G., Riguzzi, F. 2011. Lithosphere-asthenosphere viscosity
 contrast and decoupling, Physics of the Earth and Planetary Interiors, 189, 1-8.
- Boglioni, C., Anderson, D.L., 2015. Top driven asymmetric mantle convection. In The
 Interdisciplinary Earth: A volume in honor of Don L. Anderson. Geological Society of
 America Special Paper 514, American Geophysical Union Special Publication 71,
 doi:10.1130/2015.2514(05).
- Doglioni, C., Panza, G.F., 2015. Polarized plate tectonics. Advances in Geophysics, 56, 3, 1-167,
 http://dx.doi.org/10.1016/bs.agph.2014.12.001

- 816 Eggins, S. M., 1992a. Petrogenesis of Hawaiian tholeiites: 1. Phase equilibria constraints.
 817 Contributions to Mineralogy and Petrology 110, 387-397.
- Eggins, S. M., 1992b. Petrogenesis of Hawaiian tholeiites: 2. Aspects of dynamic melt segregation.
 Contributions to Mineralogy and Petrology110, 398-410.
- Eiler J.M., Schiano, P., Kitchen, N., Stolper E.M., 2000. Oxygen-isotope evidence for recycled crust
 in the sources of mid-ocean-ridge basalts. nature, 403, 530-534.
- Falloon, T. J., Danyushevsky, L. V., Ariskin, A., Green, D. H., 2007a. The application of olivine
 geothermometry to infer crystallization temperatures of parental liquids: implications for the
 temperature of MORB magmas. Chemical Geology 241, 207^233.
- Falloon, T.J., Green, D.H. Danyushevsky, L.V. 2007b. Crystallization temperatures of tholeiite
 parental liquids: Implications for the existence of thermally driven mantle plumes, in
 Foulger, G.R., and Jurdy, D.M., eds., Plates, plumes, and planetary processes: Geological
 Society of America Special Paper 430, 235–260, doi:10.1130/2007.2430(12)
- Frey, F.A., Green, D. H., 1974. The mineralogy, geochemistry and origin of lherzolite inclusions in
 Victorian basanites. Geochimica et Cosmochimica Acta, 38, 1023-1059.
- Fyfe, W.S., 1960. The possibility of *d*-electron coupling in olivine at high pressures. Geochimica et
 Cosmochimica Acta, 19, 141-143.
- Green, D. H., 1971. Composition of basaltic magmas as indicators of conditions of origin: application
 to oceanic volcanism. Philosophical Transactions of the Royal Society of London, A 268,
 707-725.
- Green, D. H., Falloon, T. J., 1998. Pyrolite: A Ringwood concept and its current expression. In:
 Jackson INS (ed) The Earth's mantle: composition, structure and evolution. Cambridge
 University Press, Cambridge, pp 311–380.
- Green, D. H., Falloon, T. J., Eggins, S. M., Yaxley, G. M., 2001. Primary magmas and mantle
 temperatures. European Journal of Mineralogy 13(3), 437-451.
- Green, D. H., Falloon, T. J., 2005. Primary magmas at mid-ocean ridges, 'hotspots', and other intraplate settings: Constraints on mantle potential temperature. In: Foulger, G. R., Natland, J. H., Presnall, D. C. & Anderson, D. L. (eds) Plates, Plumes, and Paradigms. Geological Society of America, Special Papers 388, 217-247.
- Green, D. H., Falloon, T. J., (2015), Mantle-derived magmas: intraplate, hot-spots and mid-ocean
 ridges, Sci. Bull., doi:10.1007/s11434-015-0920-y
- Green, D.H., Hibberson, W.O., Kovacs, I., Rosenthal, A., 2010. Water and its influence on the
 lithosphere asthenosphere boundary. Nature 467, 448–451. doi:10.1038/nature09369.
- Green, D.H., Hibberson, W.O., Kovacs, I., Rosenthal, A., 2011. Water and its influence on the
 lithosphere-asthenosphere boundary, 467, 448-451, 2010). [Addendum.]. Nature 472(7344),
 504.
- Green, D.H., Hibberson, W.O., Rosenthal, A., Kovacs, I., Yaxley, G.M., Falloon, T.J., Brink, F. 2014.
 Experimental study of the influence of water on melting and phase assemblages in the upper
 mantle. Journal of Petrology, 55, 10, 2067-2096 doi: 10.1093/petrology/egu050.
- Green, D.H., 2015. Experimental petrology of peridotites, including effects of water and carbon on
 melting in the Earth's uppermantle. Phys Chem Miner 42, 95–102. doi: 10.1007/s00269014-0729-2.
- Gripp, A.E., Gordon, R.G., 2002. Young tracks of hotspots and current plate velocities: Geophysical
 Journal International, v. 150, p. 321–364, doi: 10.1046/j.1365-246X.2002.01627.x.
- Hammond, W. C., Toomey, D. R., 2003. Seismic velocity anisotropy and heterogeneity beneath the
 Mantle Electromagnetic and Tomography Experiment (MELT) region of the East Pacific

- Rise from analysis of P and S body waves, Journal of Geophysical Research, 108, 2176, doi:
 10.1029/2002JB001789, B4.
- Henstock, T. J., Woods, A. W., White, R. S., 1993. The accretion of oceanic crust by episodic sill
 intrusion. Journal of Geophysical Research, Solid Earth, 98, 4143–4161.
- Harper, G. D., 1985. Tectonics of slow spreading mid-ocean ridges and consequences of a variable
 depth to the brittle/ductile transition. Tectonics, 4(4), 395-409. doi:
 10.1029/TC004i004p00395.
- Holmes, A., 1944. Principles of physical geology. In Thomas Nelson and Sons LTD, Edinburgh (ed.).
- Jordan, T. H., 1979. Mineralogies, densities and seismic velocities of garnet lherzolites and their
 geophysical implications, in The Mantle Sample: Inclusion in Kimberlites and Other
 Volcanics (eds F.R. Boyd and H. O.A. Meyer), American Geophysical Union, Washington
 D.C. 1-13. doi: 10.1029/SP016p0001
- Kaczmarek, M.A., Müntener, O., 2008. Juxtaposition of melt impregnation and high temperature
 shear zones in the upper mantle; field and petrological constraints from the Lanzo peridotite
 (Northern Italy). Journal of Petrology 49, 2187–2220.
- Kaczmarek, M.A., Müntener, O., 2010. The variability of peridotite composition across a mantle shear
 zone (Lanzo massif, Italy): interplay of melt focusing and deformation. Contribution to
 Mineralogy and Petrology 160, 663–679
- Lavier, L. L., Manatschal, G., 2006. A mechanism to thin the continental lithosphere at magma-poor
 margins. Nature 440, 324–328.
- Le Pichon, X., 1968. Sea-floor spreading and continental drift. Journal of Geophysical Research, 73,
 12, 3661-3697.
- Ligi, M., Bonatti, E., Cipriani, A., Ottolini, L., 2005. Water-rich basalts at mid-ocean-ridge cold spots.
 Nature, 434, 7029, 66-69.
- MacLeod, C.J., Searle, R.C., Murton, B.J., Casey, J.F., Mallows, C., Unsworth, S.C., Achenbach,
 K.L., Harris, M., 2009. Life cycle of oceanic core complexes. Earth and Planetary Science
 Letters 287,333–344
- Maffione, M., Morris, A., Anderson, M.W., 2013. Recognizing detachment-mode seafloor spreading
 in the deep geological past. Scientific report 3, 2336, doi: 10.1038/srep02336.
- Mallows, C., Searle, R. C., 2012. A geophysical study of oceanic core complexes and surrounding
 terrain, Mid-Atlantic Ridge 13_N-14_N, Geochem. Geophys. Geosyst., 13, Q0AG08,
 doi:10.1029/2012GC004075.
- Moore, G.W., 1973. Westward tidal lag as the driving force of plate tectonics. Geology, 1, 99–100.
- Müller, R.D., Roest, W.R., Royer, J.Y., Gahagan, L.M., Sclater, J.G., 1997. Digital isochrons of the
 world's ocean floor, Journal of Geophysical Research, 102, 3211–3214.
- Müller, R.D., Sdrolias, M., Gaina, C., Roest, W.R., 2008. Age, spreading rates, and spreading asymmetry of the world's ocean crust, Geochemistry, Geophysics, Geosystems, 9, Q04006, doi:10.1029/2007GC001743.
- Müntener, O., Pettke, T., Desmurs, L., Meier, M., Schaltegger, U., 2004. Refertilization of mantle
 peridotite in embryonic ocean basins: trace element and Nd-isotope evidence and
 implications for crust-mantle relationships, Earth and Planetary Science Letters 221, 293–
 308.
- Muntener, O., Manatschal, G., Desmurs, L., Pettke T., 2010. Plagioclase peridotites in ocean continent transitions: Refertilized mantle domains generated by melt stagnation in the
 shallow mantle lithosphere, Journal of Petrology, 51, 255-294.

- Niu, Y. L., Regelous, M., Wendt, I. J., Batiza, R. and O'Hara, M. J., 2002. Geochemistry of near-EPR
 seamounts: importance of source vs. process and the origin of enriched mantle component.
 Earth Planetary Science Letters, 199, 327–345.
- Panza, G.F., Doglioni, C., Levshin, A., 2010. Asymmetric ocean basins, Geology, 38, 1, 59-62. doi:
 10.1130/G30570.1.
- Panza, G.F., Romanelli, F., 2014. Seismic waves in 3-D: from mantle asymmetries to reliable seismic
 hazard assessment. Earthq Sci 27(5), 567–576, doi:10.1007/s11589-014-0091-y.
- Piccardo, G.B., Müntener, O., Zanetti, A., Pettke, T., 2004. Ophiolitic peridotites of the Alpine Apennine system: mantle processes and geodynamic relevance, International Geology
 Review 46, 1119–1159
- 917 Piccardo, G.B., Poggi, E., Vissers, R.L.M., 2007a. The pre-oceanic evolution of the Erro–Tobbio
 918 peridotite (Voltri Massif, Ligurian Alps, Italy), Journal of Geodynamics 43, 417–449.
- Piccardo, G.B., Zanetti, A., Müntener, O., 2007b. Melt/peridotite interaction in the Southern Lanzo
 peridotite: field, textural and geochemical evidence. Lithos 94, 181–209.
- Piccardo, G.B., Guarnieri, L. 2010. The Monte Maggiore peridotite (Corsica, France): a case study of
 mantle evolution in the Ligurian Tethys. Geological Society, London, Special Publications,
 337, 7-45, doi:10.1144/SP337.2
- Piccardo, G.B., M. Padovano, and L. Guarnieri (2014), The ligurian tethys: mantle processes and
 geodynamics. Earth-Science Reviews, 138, 409–434, doi:10.1016/j.earscirev.2014.07.002.
- Pilet, S., Baker, M.B., and Stolper, E.M., 2008. Metasomatized lithosphere and the origin of alkaline
 lavas, Science, 320, 916-919.
- Pilet, S., Baker, M.B., Muntener, O. Stolper, E. M., 2011. Monte Carlo simulations of metasomatic
 enrichment in the lithosphere and implications for the source of alkaline basalts, Journal of
 Petrology, 52, 1415-1442.
- Pilet, S., Hernandez, J., Sylvester, P., Poujol, M., 2005. The metasomatic alternative for ocean island
 basalt chemical heterogeneity, Earth and Planetary Science Letters, 236, p. 148-166.
- Rampone, E., Borghini, G., 2008. The melt intrusion/interaction history of the Erro–Tobbio
 peridotites (Ligurian Alps, Italy): insights on mantle processes at nonvolcanic passive
 margins. European Journal of Mineralogy 20, 573–585.
- Rampone, E., Piccardo, G.B., Hofmann, A.W., 2008. Multi-stage melt-rock interaction in the Mt.
 Maggiore (Corsica, France) ophiolitic peridotites: microstructural and geochemical records.
 Contribution to Mineralogy and Petrology 156, 453–475.
- Rampone, E., Piccardo, G.B., Vannucci, R., Bottazzi, P., 1997. Chemistry and origin of trapped melts
 in ophiolitic peridotites. Geochimica et Cosmochimica Acta 61, 4557–4569.
- Rampone, E., Romairone, A., Abouchami, W., Piccardo, G.B., Hofmann, W., 2005. Chronology,
 petrology and isotope geochemistry of the Erro–Tobbio peridotites (Ligurian Alps, Italy):
 records of late Palaeozoic lithospheric extension, Journal of Petrology 46, 799–827.
- Rampone, E., Hofmann, A.W., 2012. A global overview of isotopic heterogeneities in the oceanic
 mantle, Lithos 148, 247–261.
- Reston, T. J., Ranero, C. R., 2011. The 3-D geometry of detachment faulting at mid-ocean ridges,
 Geochem. Geophys. Geosyst., 12, Q0AG05, doi:10.1029/2011GC003666.
- Ricard, Y., Doglioni, C., Sabadini, R., 1991. Differential rotation between lithosphere and mantle: A
 consequence of lateral mantle viscosity variations, Journal of Geophysical
 Research, 96(B5), 8407–8415, doi:10.1029/91JB00204.
- Riguzzi, F., Panza, G., Varga, P. Doglioni, C., 2010. Can Earth's rotation and tidal despinning drive
 plate tectonics? Tectonophysics, 484, 60-73. doi:10.1016/j.tecto.2009.06.012.

- Scoppola, B., Boccaletti, D., Bevis, M., Carminati, E., Doglioni, C., 2006. The westward drift of the
 lithosphere: a rotational drag? Geological Society of America Bulletin, 118, 199–209,
 doi:10.1029/2004TC001634.
- Rosenthal, A., Yaxley, G.M., Green, D. H., Hermann, J., Kovacs, I., Spandler, C., 2014. Continuous
 eclogite melting and variable refertilisation in upwelling heterogeneous mantle, Scientific
 Reports 4, 6099, 10.1038/srep06099.
- Scheirer, D.S., Forsyth, D.W., Cormier, M.H., Macdonald, K.C., 1998. Shipboard geophysical
 indications of asymmetry and melt production beneath the East Pacific Rise near the MELT
 experiment: Science, 280, 1221–1224, doi: 10.1126/science.280.5367.1221.
- Shen, Y., Forsyth, D.W., Scheirer, D.S., Macdonald, K.C., 1993. Two forms of volcanism:
 implications for mantle flow and off-axis crustal production on the west flank of the
 southern east pacific rise. Journal of Geophysical Research 98, B10, 17,875-17,889.
- Shimizu, K., Saal, A.E., Myers, C.E., Nagle, A.N., Hauri, E.H., Forsyth, D.W., Kamenetsky, V.S.,
 Niu, Y., 2016. Two-component mantle melting-mixing model for the generation of midocean ridge basalts: Implications for the volatile content of the Pacific upper mantle.
 Geochimica et Cosmochimica Acta, 176, 44–80.
- Schmerr, N., 2012. The Gutenberg discontinuity: Melt at the lithosphere-asthenosphere boundary,
 Science 335, 14801483.
- Seyler, M., Bonatti, E., 1997. Regional-scale melt-rock interaction in lherzolitic mantle in the
 Romanche fracture zone (Atlantic Ocean), Earth and Planetary Science Letters 146, 273–
 287.
- Small, C., Danyushevsky L. 2003. A plate kinematic explanation for mid-ocean ridge depth
 discontinuities, Geology, 31, 399-402.
- Smith, D.K., Escartín, J., Schouten, H., Cann, J.R., 2008. Fault rotation and core complex formation:
 Significant processes in seafloor formation at slow-spreading mid-ocean ridges (Mid-Atlantic Ridge, 13_-15_N), Geochem. Geophys. Geosyst., 9(3), 10.1029/2007GC001699
- 979Smith, D.K., Escartín, J., Schouten, H., Cann, J.R., 2012. Active long-lived faults emerging along980slow-spreadingmid-oceanridges,Oceanography25(1),94–99,981http://dx.doi.org/10.5670/oceanog.2012.07.
- Tartarotti, P., Susini, S., Nimis, P., Ottolini, L., 2002. Melt migration in the upper mantle along the
 Romanche fracture zone (Equatorial Atlantic). Lithos 63, 125–149.
- Takahashi, N., (2001), Origin of Plagioclase Lherzolite from the Nikanbetsu Peridotite Complex,
 Hokkaido, Northern Japan: Implications for Incipient Melt Migration and Segregation in the
 Partially Molten Upper Mantle. Journal of Petrology 42, 39-54.
- 987 Thybo, H., 2006. The heterogeneous upper mantle low velocity zone. Tectonophysics 416, 53–79.
- Toomey D.R., Wilcock, W.S.D., Conder, J.A., Forsyth, D.W, Blundy, J.D., Parmentier, E.M.,
 Hammond, W.C., 2002. Asymmetric mantle dynamics in the MELT region of the East
 Pacific Rise. Earth and Planetary Science Letters 200, 287–295.
- 991 Toomey D.R., 2012. Piecing together rifts. Nature Geoscience 5, 235–236.
- Vine, F. J., Matthews, D. H., 1963. Magnetic Anomalies Over Oceanic Ridges, Nature, 199(4897),
 993 947–949. doi: 10.1038/199947a0.
- Warren, J.M., Shimizu, N., 2010. Cryptic variations in abyssal peridotite compositions: evidence for
 shallow-level melt infiltration in the oceanic lithosphere. Journal of Petrology 51, 395–423.
- Warren, J.M., 2016. Global variations in abyssal peridotite compositions. Lithos, 248-251, 193-219.
- Wegener, A., 1915. Die Entstehung der Kontinente und der Ozeae: Samml. Vieweg, Braunschweig,
 23, pp 1-94.

- White, S.C., Macdonald, K.C., Scheirer, D.S., Cormier, M.H., 1998. Distribution of isolated
 volcanoes on the flanks of the East Pacific Rise, 15.3°S-20°S. Journal of Geophysical
 Research 103, B12, 30,371-30,384.
- Whitney, D. L., Teyssier, C., Rey, P., Buck, W. R., 2013. Continental and oceanic core complexes,
 GSA Bulletin, 125, 3/4, 273–298, doi: 10.1130/B30754.1.

1005 Figure captions

1006 Figure 1. Uninterpreted (above) and interpreted (below). Shear wave velocity (Vs) sections along 1007 the tectonic equator (TE) for the Earth's first 300 km. The mantle is faster and the lithosphere is 1008 thicker in the western side with respect to the eastern side of the three major oceanic ridges, EPR— 1009 Eastern Pacific Ridge, MAR-Mid-Atlantic Ridge, IR-Indian Ridge. Upper asthenosphere 1010 superposes to low-velocity layer (LVZ), i.e., which is the main decoupling surface between 1011 lithosphere and mantle, allowing net rotation of lithosphere, i.e., a first-order eastward relative 1012 mantle flow, or westward drift of lithosphere. Secondary flow should be related to mantle obliquely 1013 upraised along oceanic ridges. Asymmetry between two sides of ridges is independent from age of 1014 oceanic lithosphere, shown at top in million years (modified after Panza et al., 2010).

1015



Figure 1



1018 Figure 2. Inferred kinematics of different particles of mantle rocks along a migrating mid-oceanic 1019 ridge, and related asymmetric upper mantle differentiation. Numbers from 1 to n indicate 1020 progressive age evolution of the rock markers within lithosphere and asthenosphere. Plates A and B 1021 move with velocity VA and VB, respectively. Vr is the velocity of the ridge; hS is the half-spreading 1022 rate. Since the ridge is moving westward, new sections of asthenospheric mantle migrate "eastward" 1023 relative to the lithosphere, permanently renewing mantle source of isostatically upwelling magma 1024 generated by depressurization below the ridge. During this mass transfer, the eastern part of upper 1025 asthenosphere becomes somewhat Fe-depleted, while a top to bottom mantle path decoupling occurs 1026 due to the relative "westward" drift of overlying lithosphere. The upper asthenosphere (LVZ) 1027 corresponds to the oblique ascent of partially molten mantle from lower asthenosphere. Two 1028 different paths of transfer exist: the main one (yellow paths) barely deviated in passing below the 1029 ridge, the other (red path) monitored by the suction effect from the strongly depressurized area 1030 below the spreading ridge and giving MORB (see Fig. 3 for details). Mantle lithosphere is created at 1031 the ridge through accretion of residues above the asthenospheric partial melting area, residues 1032 successively transferred upwards and laterally at hS rate within the mantle lithosphere on each side 1033 of the ridge. A top to bottom decoupling exists within the mantle lithosphere linked to the slow-1034 down by the eastward underlying flow of asthenosphere, and is responsible for detachment faults 1035 and mantle exhumation at the spreading ridge. Besides the mantle lithosphere composition, different 1036 on each side of the ridge, comes from a permanent refertilisation of the western plate whereas the 1037 eastern plate just preserves the barely refertilized feature of a harzburgitic residue generated during 1038 asthenosphere partial melting below axial ridge. See text for more details.

Figure 2



1041 Figure 3. P-T-H₂O diagrams, modified from Fig. 6 in Green et al. 2014. a) Experimentally 1042 determined solidi for hydrous silicate melt in fertile lherzolite for different water contents; blue 1043 dashed line: mantle adiabat; red star: MORB source mantle upwelling potential temperature at 230, 1044 150, 105 and 65 km; green dashed line: intraplate geotherm; orange dotted line: perturbation of 1045 oceanic intraplate geotherm by shear heating, adapted from Fig. 4 in Doglioni et al. 2005; orange 1046 dot: intraplate magma genesis at shallower depths (105 to 125 km vs. 110 to 160 km) taking into 1047 account shear heating within the lithosphere - asthenosphere decoupling zone. MORB are sourced 1048 from upwelling trace element depleted lherzolite from lower asthenosphere. Intraplate basalts, 1049 including OIB, are the products of upwelling of trace element enriched lherzolite from the middle 1050 and upper asthenosphere. See text for more details. b) Water storage capacity of lherzolitic mantle as 1051 a function of depth along the vapour-saturated solidus. Pargasite becomes unstable at >3 GPa, and 1052 water storage capacity drops to that (~200 ppm) which can be retained in NAMs (nominally 1053 anhydrous minerals) in lherzolite, inducing incipient melting of upper asthenosphere (LVZ). See text 1054 for more details.

Figure 3

