



HAL
open science

Phagocytic activity of human macrophages and *Drosophila* hemocytes after exposure to the neonicotinoid imidacloprid

Louise Walderdorff, Philippe Laval-Gilly, Laura Wechtler, Antoine Bonnefoy,
Jaïro Falla-Angel

► To cite this version:

Louise Walderdorff, Philippe Laval-Gilly, Laura Wechtler, Antoine Bonnefoy, Jaïro Falla-Angel. Phagocytic activity of human macrophages and *Drosophila* hemocytes after exposure to the neonicotinoid imidacloprid. *Pesticide Biochemistry and Physiology*, 2019, 160, pp.95-101. 10.1016/j.pestbp.2019.07.007 . hal-02960279

HAL Id: hal-02960279

<https://hal.univ-lorraine.fr/hal-02960279>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

1 **Phagocytic Activity of Human Macrophages and *Drosophila* Hemocytes after**
2 **Exposure to the Neonicotinoid Imidacloprid**

3

4 ***Louise WALDERDORFF (1)*, Philippe LAVAL-GILLY (1), Laura WECHTLER (1), Antoine***
5 ***BONNEFOY (2), Jairo FALLA-ANGEL (1)***

6 (1) Université de Lorraine, Inra, LSE, F-54000 Nancy, France

7 (2) Université de Lorraine, IUT de Thionville-Yutz, F-57970 Yutz, France

8

9 *corresponding author: louise.walderdorff@univ-lorraine.fr

10

11 **ABSTRACT**

12 Neonicotinoid insecticides are increasingly used in modern pest control and in conventional
13 agriculture. Their residues are frequently found in our environment and in our food leading to
14 chronic exposure of pollinating insects and humans. Indeed, evidence has become stronger that
15 chronic exposure to neonicotinoids might have a direct impact on the immune response of
16 invertebrates and vertebrates. Therefore, we compared the cellular immune response of human
17 macrophages (THP-1) and *Drosophila melanogaster* hemocytes (Schneider 2 cells) after exposure to
18 four different concentrations of the neonicotinoid imidacloprid. Cells were immune activated with
19 LPS (lipopolysaccharide) of *Escherichia coli* to compare the phagocytic activity of immune activated
20 and non-activated cells during pesticide exposure. *Drosophila* cells were more strongly affected by
21 the insecticide than human macrophages. Even though imidacloprid showed an adverse effect on
22 phagocytosis on both cells while immune activated, it decreased phagocytosis in *Drosophila* cells at
23 shorter exposure time and without immune activation.

24

25 Keywords

26 Imidacloprid, macrophage, phagocytosis, THP-1, *Drosophila*

27

28 1. INTRODUCTION

29 Having been launched on the market by Bayer Crop science in 1991, Imidacloprid (IMI) was the very
30 first insecticide of a new group of pesticides: the neonicotinoids (Mason et al., 2013). Neonicotinoid
31 insecticides are nicotinic acetylcholine receptor agonists in insects and are lethal through disruption
32 of the nervous system. All neonicotinoids are systematic insecticides with a high solubility in water
33 leading to the uptake of the pesticide by the plant's tissues. The pesticide's active molecule, which is
34 spread in the entire plant, thus acts directly on any sucking insect that would feed on the plant. Once
35 distributed in the whole plant it is also available for non-target insects such as pollinating insects as
36 they feed on contaminated nectar and pollen (Simon-Delso et al., 2015).

37 In the late 1980's the global insecticide market was dominated by carbamates, organophosphates
38 and pyrethroids, to which many pest insects have developed increasing resistance (Simon-Delso et
39 al., 2015). The neonicotinoids rapidly took over the global insecticide sales and by 2010 were
40 responsible for more than 25 percent of the market share with imidacloprid becoming the second
41 largest selling pesticide worldwide after Glyphosate (Jeschke et al., 2010). Despite the ban of
42 imidacloprid, clothianidin and thiamethoxam for outdoor application (they are still authorized in
43 greenhouses) in the European Union in May 2018 (Official Journal of the European Union, L 132/31,
44 2018), imidacloprid is still one of the most used insecticides in the world. In fact 75 percent of
45 neonicotinoids were used in Latin America, North America and Asia. Europe only accounted for 11
46 percent of the worldwide use (Bass et al., 2015).

47 The most common application and spread of neonicotinoids in agriculture is the use of seed coatings
48 or dressings which generates toxic dust during the process of sowing and can lead to contamination
49 of surrounding vegetation (Bonmatin et al., 2015). Further they are also used against sucking insects
50 in fruit and vegetable cultivations as well as in viticulture and hop growing. Residues of imidacloprid
51 and other neonicotinoids have been found in honeybees, honey and their pollen (Chauzat et al.,
52 2009; Codling et al., 2016; David et al., 2016; Hladik et al., 2016; Kasiotis et al., 2014) as well as fruits
53 and vegetables we consume (Fernández-Alba et al., 2000; Ko et al., 2014; Obana et al., 2002).

54 In the last decade neonicotinoids have been controversially discussed in literature as they have been
55 associated with a worldwide pollinator decline. While acute toxicity occurs rather seldom the cause
56 of this decline was associated with chronic pesticide exposure. There is a growing awareness that
57 neonicotinoids could weaken the immune response of pollinators. Indeed several studies have
58 shown a weakening effect of neonicotinoids and pathogens to pollinators (Abbo et al., 2017; Alaux et
59 al., 2010; Fauser-Misslin et al., 2014; Pettis et al., 2012; Vidau et al., 2011). The suppression of the
60 immune response to infectious diseases by neonicotinoids was not only shown for pollinators
61 (Brandt et al., 2016; Di Prisco et al., 2013; López et al., 2017) but also hypothesized for insectivorous
62 vertebrates like amphibians, birds, fish and mammals, when comparing disease outbreaks with the
63 use of neonicotinoids in the concerned regions and countries (Mason et al., 2013). A study of

64 Sánchez-Bayo and Goka (2005) supports this idea, claiming that medaka fish (*Oryzias latipes*)
65 exposed to imidacloprid were massively infested by a *Trichodina* ectoparasite when compared to
66 control groups without pesticides and groups exposed to zinc pyrethrin.

67 Also other reviews support this hypothesis and Gibbons et al. (2015) stated that imidacloprid and
68 clothianidin exert sublethal effects on vertebrates ranging from genotoxic and cytotoxic effects, and
69 impaired immune function, to reduced growth and reproductive success. Mostafalou and Abdollahi
70 (2013) mentioned human chronic diseases like cancer, Alzheimer, Parkinson or autoimmune diseases
71 that could be linked to sublethal pesticides exposure, even though they did not mention imidacloprid
72 in their work. Though Cimino et al. (2016) cited several studies of chronic exposure to imidacloprid
73 which showed adverse health effects on humans, still not much is known about the effect of
74 neonicotinoids on human health.

75 Innate immune responses in mammals and insects do show several functional similarities. Cells,
76 especially phagocytes, of these innate immune systems are comparable both in terms of morphology
77 and reactivity towards pathogen recognition (Browne et al., 2013). An important role in the primary
78 immune response is given to phagocytes. These cells are responsible among others for the
79 recognition and removal of foreign pathogens and the recruitment of other immune cells through
80 the release of cytokines (Chanput et al., 2014). Insect hemocyte's main function is phagocytosis of
81 pathogens and apoptotic material (Feldhaar and Gross, 2008) therefore being of major importance to
82 the organism. The most dominant hemocytes in *Drosophila* are plasmatocytes which are highly
83 motile and resemble monocytic macrophages in humans (Feldhaar and Gross, 2008). Insect
84 hemocytes are able to change their state from non-adherent to adherent upon pathogen recognition
85 and respond by "spreading" which enhances phagocytosis of small antigens (Negri et al., 2013). In
86 contrast to human monocytes, insect hemocytes are able to perform nodulation and encapsulation
87 of larger foreign objects like parasites as they adhere and spread over the surface of the foreign
88 object until they form overlapping layers (Negri et al., 2013). The encapsulation is often followed by
89 melanisation of the antigen (Schmid et al., 2008) and hemocytes are also able to secrete cytotoxic
90 compounds into the capsule (Negri et al., 2013). Imidacloprid could have a suppressive effect on this
91 cellular immune response of invertebrates as well as of vertebrates.

92 Hence in this study we investigate the impact of imidacloprid (IMI) on the phagocytic activity of
93 immune system cells of one invertebrate and one vertebrate species. We chose the cell lines
94 Schneider-2 from *Drosophila melanogaster* and THP-1, a human monocytic cell line that can be
95 differentiated into macrophages. To simulate an immune reaction we activated the cells with
96 lipopolysaccharide (LPS) of *Escherichia coli*.

97

98 2. MATERIALS AND METHODS

99 2.1. Cell culture and exposure

100 Both human THP-1 (Ref. ACC 16) and *D. melanogaster* Schneider-2 (Ref. ACC 130) cells were
101 purchased from Leibnitz – Institute DSMZ- Deutsche Sammlung von Mikroorganismen und
102 Zellkulturen GmbH. THP-1 cells were maintained in RPMI 1640 with 10 percent heat inactivated FBS
103 (fetal bovine serum), 2mM L-Glutamine, 50 U/mL penicillin and 50 µg/mL streptomycin at 37 °C and 5
104 percent CO₂. Schneider cells were kept in 44.75 percent Schneider’s insect medium, 44.75 percent
105 TC-100 insect medium, 10 percent heat inactivated FBS, 50 U/mL penicillin and 50 µg/mL
106 streptomycin at 20 °C.

107 Imidacloprid (37894 Sigma-Aldrich, powdery analytical standard) was dissolved in PBS (pH 7.4) at 300
108 µg/mL prior to experiments. Imidacloprid was further diluted in RPMI or Schneider medium in order
109 to obtain the following concentrations: 1; 25; 50 and 100 µg/mL IMI. During each experiment all cell
110 solutions have been tested with HPLC to determine the actual concentration of IMI that has been in
111 contact with cells to avoid measuring errors. An analytical standard of Imidacloprid solution (46341
112 Sigma-Aldrich) of 100 µg/mL in acetonitrile was used to obtain a standard curve for HPLC analyses.
113 Cells were incubated with imidacloprid at the cell particular conditions for 5 or 24 hours. Cell viability
114 was examined after 5 and 24 hours with a trypan blue solution (2 g/L in 0.9 percent NaCl solution).
115 Trypan blue is a diazo dye used for vital staining of cells (Tolnai, 1975). After the exposure to
116 imidacloprid cells were washed with PBS and incubated for 16 hours with LPS (lipopolysaccharide)
117 from *Escherichia coli* at concentrations of 0.1 µg/mL or 1 µg/mL for THP-1 and 1 µg/mL for Schneider
118 cells. LPS is an endotoxin and a component of the outer membrane of Gram-negative bacteria
119 resulting in a strong inflammatory response of the immune system.

120 2.2. Phagocytosis

121 Phagocytosis of THP-1 and Schneider cells was examined by using amine-modified polystyrene,
122 fluorescent yellow-green latex beads (1 µm mean particle size) after slight modifications of the
123 methodology of Bennisroune et al. (2012).

124 THP-1 cells were differentiated to their macrophage phenotype with 5 ng/mL of PMA (phorbol 12,
125 myristate 13-acetate) for 48 hours and washed once with PBS prior to pesticide exposure. Schneider
126 cells and differentiated THP-1 cells were grown on glass coverslips at a concentration of 7 x 10⁵
127 cells/mL in six-well or twelve-well plates. Cells were exposed to different concentrations of
128 imidacloprid from 1 µg/mL to 100 µg/mL, including a negative control without IMI, for either 5 or 24
129 hours at the particular cell conditions. 4 hours prior to the phagocytosis measure, 100 latex beads
130 per cell diluted in the cell particular medium were added to each well. After 4 hours of incubation
131 cells were washed with 3 percent BSA (Bovine serum albumin) dissolved in PBS in order to eliminate

132 latex beads which have not been absorbed by the cells (Komiyama et al., 2003). The glass coverslips
133 of differentiated THP-1 cells and Schneider cells were dipped three times in the BSA solution before
134 being dipped into PBS. Glass coverslips were then mounted on object plates and phagocytosis was
135 measured under a fluorescent microscope. A minimum of 250 cells were counted for each sample.
136 Phagocytosis was calculated as follows:

$$137 \quad \text{cells (phagocytosis)} = \frac{\text{number of cells ingesting at least 1 bead}}{\text{number of total cells}} * 100$$

138

139 2.3. Statistical analyses

140 All data was analysed using Addinsoft (2019) XLSTAT statistical and data analysis solution, Long
141 Island, NY, USA. For all tests a probability level of 0.05 was considered to be statistically significant.
142 After checking for normality and variance homogeneity using the Shapiro-Wilk's test and the
143 Bartlett's test, a one-way ANOVA was performed to identify differences between different IMI
144 groups of each LPS treatment, followed by a Dunnett's test for comparison of different IMI groups to
145 the control group or a Tukey's test of multiple comparison between different IMI groups of the same
146 LPS treatment. Data for viability of human THP-1 cells after 5 hours exposure time were log
147 transformed in order to fit parametric conditions. As parametric conditions were not given for the
148 viability of human THP-1 cells after 24 hours exposure time, even after log or arcsine transformation,
149 we used a non-parametric Kruskal-Wallis test.

150 Results were expressed as means of all replicates \pm SE (standard error).

151

152 3. RESULTS

153 3.1. *D. melanogaster* Schneider cells

154 Cell viability in Schneider cells of *D. melanogaster* was higher after 5 hours than after 24 hours
155 exposure to IMI. Cell viability varied between 88.4 ± 0.47 percent and 90.9 ± 0.96 percent after 5
156 hours and between 84.6 ± 1.73 percent and 80.4 ± 1.84 percent after 24 hours. No statistically
157 significant differences were found between the control without IMI and different IMI concentrations
158 neither after 5 hours (ANOVA $p=.233$ with $n=9$) nor after 24 hours (ANOVA $p=.480$ with $n=9$) (Figure
159 1).

160 Phagocytosis of *D. melanogaster* Schneider cells was tested after 5 and 24 hours of exposure to IMI
161 with or without immune activation with $1 \mu\text{g/mL}$ LPS of *Escherichia coli*.

162 Phagocytosis after 5 hours and no LPS activation showed a decline of phagocytosis with increasing
163 concentrations of IMI. 24.84 ± 1.71 percent of cells from the control group were able to phagocytize
164 a minimum of one latex bead. Percentages decreased continuously with increasing pesticide
165 concentrations until phagocytosis reached only 5.02 ± 0.86 percent at the highest IMI concentration,
166 which was statistically significant from the control (Dunnett $p=.035$, with $n=3$). When immune
167 activated with $1 \mu\text{g/mL}$ LPS, phagocytosis increased slightly up to 30.74 ± 3.26 percent in the control
168 group. IMI treated groups revealed decreasing percentages of phagocytosis compared to the control.
169 Statistically significant differences were determined between 25, 50 and $100 \mu\text{g/mL}$ IMI and the
170 control group (Tukey $p=.009$; $p=.025$; $p=.004$ respectively). Phagocytosis in these three treatment
171 groups varied from 21.37 ± 1.15 percent to 18.11 ± 0.95 percent.

172 After 24 hours of exposure to IMI and no immune activation phagocytosis ranged from 42.90 ± 4.99
173 percent in the control group to 27.85 ± 1.38 percent after IMI exposure, showing no statistical
174 significant difference between control and IMI (ANOVA $p=.222$ with $n=3$).

175 After immune activation with $1 \mu\text{g/mL}$ LPS for 16 hours following a 24 hours exposure to IMI we
176 observed a statistically significant decrease of phagocytosis with increasing pesticide concentrations
177 upon the lowest used IMI concentration (Tukey $p=.033$, $p=.0001$, $p=.0001$, $p<.0001$ for 1; 25; 50 and
178 $100 \mu\text{g/mL}$ IMI respectively). Values varied between 31.55 ± 1.92 percent in the control group and
179 12.92 ± 1.19 percent at the highest IMI concentration. At the lowest used IMI concentration of 1
180 $\mu\text{g/mL}$ only 24.05 ± 1.01 percent of cells were capable of phagocytosis, which is about a quarter less
181 compared to the control group. At $25 \mu\text{g/mL}$ the ability of cells for phagocytosis was reduced by one
182 third and at $50 \mu\text{g/mL}$ it was already reduced by more than half compared to the control (Figure 2, b).


183 3.2. Human THP-1 cells

184 Cell viability in THP-1 was measured after 5 and 24 hours exposure to IMI. Viability in human THP-1
185 cells was between 92.9 ± 1.06 and 90.3 ± 2.05 percent after 24 hours exposure while it ranged
186 between 84.9 ± 2.61 percent and 80.6 ± 2.27 percent after 5 hours. No statistical significant
187 differences in THP-1 viability were found between the control group and any tested IMI
188 concentration at 24 hours (Kruskal-Wallis $p=.857$ with $n=12$) or 5 hours (ANOVA $p=.88$ with $n=9$)
189 exposure to IMI (Figure 1).

190 Phagocytosis of differentiated THP-1 cells was stimulated by immune activation of either $0.1 \mu\text{g/mL}$
191 or $1 \mu\text{g/mL}$ LPS. Percentages of phagocytizing cells of the same population increased in average
192 about 16.4 percent while stimulated with LPS. Phagocytosis measured after an exposure of 5 hours
193 to IMI and no immune activation with LPS was ranging between 37.3 ± 5.02 percent and 55.3 ± 4.78
194 percent. Phagocytosis increased with immune activation and varied between 62.5 ± 4.31 percent to

195 72.2 ± 4.76 percent at 0.1 µg/mL LPS and between 52.1 ± 1.62 percent and 68.3 ± 3.21 percent at
196 1µg/mL LPS activation. No statistically significant difference between the control group and the
197 treatment groups was found for all tests after 5 hours exposure without, with 0.1 µg/mL or 1 µg/mL
198 LPS (ANOVA, p=.161, p=.624 and p=.070 respectively, with n=3).

199 No statistically significant differences in phagocytosis between different IMI concentrations and the
200 control were found after 24 hours of exposure and no immune activation with LPS (ANOVA p=.102
201 with n=3). Values varied between 84.07 ± 2.15 percent and 70.23 ± 2.65 percent. However,
202 differences between different IMI concentrations were found after immune activation with either 0.1
203 µg/mL or 1 µg/mL LPS. After activation with 0.1 µg/mL LPS phagocytosis decreased after 24 hours IMI
204 exposure in treated groups compared to the control group for up to 17.1 percent. The biggest
205 decreases in phagocytosis were found at 25 µg/mL and 100 µg/mL IMI which was also confirmed by
206 statistical analyses (Tukey p=.038 and p=.044 respectively). 81.33 ± 1.43 percent of cells of the
207 control group were capable of phagocytizing at least one latex bead while this percentage decreased
208 with increasing IMI concentrations except for cells at 50 µg/ml IMI which were able to phagocytize
209 more latex beads than cells at 25 µg/mL IMI. Phagocytosis after 24 hours and later activation with 1
210 µg/mL LPS (Figure 2) ranged from 81.21 ± 4.69 percent at the control group to 67.2 ± 4.99 percent at
211 50µg/mL IMI, which was statistically different from the control group (Tukey p=.034).



212


213 **Figure 1: Viability of *D. melanogaster* Schneider cells (a,b) and human THP-1 cells (c,d) after 5 and 24**
 214 **hours exposure to IMI. No statistical differences were found between the control group without IMI and**
 215 **the IMI concentrations neither for Schneider after 5 and 24 hours nor THP-1 after 5 and 24 hours (ANOVA:**
 216 **p=.233; p=.480; p=.880; Kruskal-Wallis p=.857 and n=9; n=9; n=9; n=12 respectively).**

217

218 4. DISCUSSION

219 In the last decade neonicotinoid insecticides have been associated with a worldwide pollinator
 220 decline as they might act as an immune suppressor in insects. But evidence has become stronger
 221 that they might not only weaken the immune response in insects but also in insectivorous
 222 vertebrates (Mason et al., 2013). Further residues of neonicotinoids have also been detected in
 223 human urine as we are exposed to contaminated food (Nomura et al., 2013; Taira et al., 2013). A few
 224 studies exist on human chronic exposure to neonicotinoids but still not much is known on their
 225 impact on human immune cells (Han et al., 2018). To investigate this effect, macrophage-like
 226 immune system cells of two different species, *Drosophila melanogaster* and human, were tested *in*
 227 *vitro* on their cellular reactivity when confronted with four different concentrations of IMI (1, 25, 50
 228 and 100 µg/mL). The cells of the two organisms had similar characteristics and were able to

229 phagocytize particles. Thus we measured the phagocytic capacity of the two cell lines after pesticide
 230 exposure. Further cells were exposed to LPS of *Escherichia coli*, which acts as a pathogen associated
 231 molecular pattern (PAMP). Hence cells confronted with LPS were immune activated leading to an
 232 immune reaction or an inflammatory response. We were then able to compare the effect of IMI on
 233 immune activated and non-activated cells.




234

235 **Figure 2: Phagocytosis of *D. melanogaster* Schneider cells (a,b) and human THP-1 cells (c,d) after 5 and 24 hours exposure**
 236 **to IMI and immune activation with LPS. Stars indicate statistically significant differences from the control group without**
 237 **IMI (ANOVA $p < .05$ with $n = 3$).**

238 Without immune activation IMI significantly decreased phagocytosis of Schneider cells after 5 hours
 239 of exposure at the highest tested concentration. Immune activation with LPS increased this inhibiting
 240 effect leading to a significant decrease of phagocytosis upon 25 μg/mL IMI. These results match with
 241 results of Di Prisco et al. (2013) who found clothianidin, another neonicotinoid insecticide, to
 242 negatively affect the activation of the Toll-pathway of *Drosophila melanogaster*. The Toll-pathway in
 243 turn plays an important role on immune responses including the activation of phagocytosis (Valanne
 244 et al., 2011). Similar results on phagocytosis were also obtained after an exposure of 24 hours. IMI
 245 significantly decreased phagocytosis in Schneider cells after immune activation upon our lowest
 246 tested concentration of 1 μg/mL. It thus seems that the immune activation with LPS enforces the
 247 toxic effect of IMI in *Drosophila* cells as immune stress increases.

248 As for human macrophages IMI significantly decreased phagocytosis only after 24 hours exposure at
249 higher IMI concentrations when cells were immune activated with either 0.1 µg/mL or 1 µg/mL LPS.
250 No significant alteration of phagocytosis was found in macrophages after the shorter exposure time
251 of 5 hours with or without immune activation. Several studies have shown a dose-dependent
252 suppression of phagocytosis of phagocytic vertebrate cells alongside other pollutants (Bennasroune
253 et al., 2012; Couleau et al., 2012; Dunier et al., 1991; Roszell and Anderson, 1994) suggesting that
254 concentrations used were not high enough to induce effects on phagocytosis of THP-1 cells after
255 short time exposure. In THP-1 cells LPS results in an activation of the NF-κB transcription factor
256 (Chanput et al., 2014). Di Prisco et al. (2017) tested clothianidin on the THP-1 cell line showing that
257 NF-κB was down regulated by the neonicotinoid insecticide. However, it is to mention that we were
258 only differentiating between cells that were able to phagocytise at least one fluorescent bead and
259 those who did not phagocytize any beads at all. We did not consider the number of beads that were
260 ingested by the cells. As shown in Figure 3 cells were able to engulf many beads, making it impossible
261 to count their number. We did though see cells that clearly ingested less beads than others when
262 observed under the microscope. Therefore, it is possible that with our method we might not have
263 been able to detect an effect of IMI on the efficiency of phagocytosis. IMI could be influencing
264 different aspects of the phagocytic process. With our method we cannot clearly say if IMI can
265 influence the rate of the uptake of bacteria or the degradation of bacteria once engulfed.



266

267 *Figure 3 : Phagocytized fluorescent latex beads by Schneider 2 cells of D. melanogaster. White arrows show cells that*
268 *engulfed a high number of beads, grey arrows show cells that engulfed a smaller number of beads and black arrows show*
269 *cells that did not phagocytise any fluorescent latex bead. The scale bar is 10 µm.*

270

271 IMI already showed an effect on Drosophila Schneider cells even when cells were not stimulated with
272 LPS whereas effects in human THP-1 cells only occurred if cells were confronted with LPS and thus
273 initiating an immune reaction. Having in mind that IMI is an insecticide, with insects being the

274 desired target, it is not surprising that Schneider cells are reacting more sensitive to it than human
275 THP-1 cells. Cresswell *et al.* (2012) interpreted in their results that temperature might play a role
276 during IMI intoxication in honey and bumblebees. Schneider cells as well as bee hemocytes were
277 incubated at 22 °C while THP-1 cells were incubated at 37 °C. These temperature differences are due
278 to different cell characteristics but it is not to neglect that this may play a role in the intoxication
279 processes in the two cell lines. Camp and Buchwalter (2016) and Song *et al.* (1997) found higher
280 chronic toxicity of IMI on aquatic invertebrates at lower temperatures while Van den Brink *et al.*
281 (2016) found that summer populations of aquatic insects were stronger effected by IMI than winter
282 populations. Apart from intoxication processes temperature also plays a role during the immune
283 response. Indeed Linder *et al.* (2008) suggested that colder temperatures had rather beneficial
284 effects on the immune response of *Drosophila* as heat shock proteins may improve immune function.
285 According to our results IMI did not reduce viability of human THP-1 nor *Drosophila*'s Schneider cells
286 at any tested concentration from 1 µg/mL to 100 µg/mL IMI. We were thus working with sublethal
287 doses in our experimental conditions. Hence EC₅₀ values are dependent on cell and tissue type as it
288 was already assumed by Benzidane *et al.* (2011). Concentrations used during experiments were
289 ranging from 1 µg/mL to 100 µg/mL, which were quite high when compared to environmental
290 concentrations. Chauzat *et al.* (2009) and Kasiotis *et al.*(2014) detected 5.7 to 73.9 ng/g imidacloprid
291 in pollen while LD₅₀ values for honey bees varied from 4.5 ng – 81 ng/bee for oral ingestion (Nauen *et*
292 *al.*, 2001; Cresswell, 2011) and from 18 ng – 102 ng/bee during contact (Iwasa *et al.*, 2004; Nauen *et*
293 *al.*, 2001). In *Drosophila* the LD₅₀ at oral ingestion was estimated at 0.41 - 1.39 µg/g (Arain *et al.*,
294 2014). It is, though, to mention that IMI has a very short half-life in organisms. IMI showed a similar
295 metabolism in honey bees as in *Drosophila melanogaster*. In bees the half-life of IMI accounts of 4.5
296 hours (Chauzat *et al.*, 2009) and is reduced to several metabolites like 6-chloronicotinic acid and urea
297 derivative (Suchail *et al.*, 2004). Consequently concentrations of IMI in honeybees could be way
298 higher than the actual ones found, depending on when the bee has ingested contaminated nectar for
299 the last time. Further these metabolites might have other toxicological effects on the host organisms.
300 Indeed Suchail *et al.* (2004) claims that the mortality of bees is more due to the action of the
301 metabolites than the actual parent molecule. The EC₅₀ value of honey bee Kenyon cells was
302 estimated at around 6.3 µg/mL IMI (Déglise, *et al.*, 2002). As IMI is an agonist on the nicotinic
303 acetylcholine receptors effects on neuronal cells are expected to be higher than on immune system
304 cells. It is therefore not implausible that effects on the two cell types used appeared at higher
305 concentrations. These indications should be kept in mind when analysing *in vitro* toxicity test as we
306 are missing metabolic processes.

307 To conclude we can summarize that the concentrations from 1 µg/mL to 100 µg/mL of IMI did reduce
308 phagocytic activity in both human macrophages and *Drosophila* hemocytes after immune activation

309 with LPS of *Escherichia coli*. LPS clearly intensified the inhibiting effects of IMI on cell phagocytic
310 activity. Phagocytosis in *Drosophila* cells was more strongly inhibited than in human macrophages.
311 While we observed no effect of IMI on phagocytosis of human macrophages without immune
312 activation, IMI did reduce phagocytosis in *Drosophila* cells after even short exposure and no immune
313 activation.

314

315 Acknowledgements

316 This work was funded by « Communauté d'agglomérations Portes de France – Thionville ».

317 Conflict of interest

318 The authors report no conflicts of interest to be declared.

319 References

- 320 Abbo, P.M., Kawasaki, J.K., Hamilton, M., Cook, S.C., DeGrandi-Hoffman, G., Li, W.F., Liu, J., Chen,
321 Y.P., 2017. Effects of Imidacloprid and *Varroa destructor* on survival and health of European
322 honey bees, *Apis mellifera*. *Insect Sci.* 24, 467–477.
- 323 Alaux, C., Brunet, J.-L., Dussaubat, C., Mondet, F., Tchamitchan, S., Cousin, M., Brillard, J., Baldy, A.,
324 Belzunces, L.P., Le Conte, Y., 2010. Interactions between *Nosema* microspores and a
325 neonicotinoid weaken honeybees (*Apis mellifera*). *Environ. Microbiol.* 12, 774–782.
- 326 Arain, M., Hu, X.-X., Li, G.-Q., 2014. Assessment of Toxicity and Potential Risk of Butene-fipronil Using
327 *Drosophila melanogaster*, in Comparison to Nine Conventional Insecticides. *Bull. Environ.*
328 *Contam. Toxicol.* 92, 190–195. <https://doi.org/10.1007/s00128-013-1155-8>
- 329 Bass, C., Denholm, I., Williamson, M.S., Nauen, R., 2015. The global status of insect resistance to
330 neonicotinoid insecticides. *Pestic. Biochem. Physiol.* 121, 78–87.
- 331 Bennisroune, A., Rojas, L., Foucaud, L., Goulaouic, S., Laval-Gilly, P., Fickova, M., Couleau, N.,
332 Durandet, C., Henry, S., Falla, J., 2012. Effects of 4-nonylphenol and/or diisononylphthalate
333 on THP-1 cells: impact of endocrine disruptors on human immune system parameters. *Int. J.*
334 *Immunopathol. Pharmacol.* 25, 365–376.
- 335 Benzidane, Y., Lapied, B., Thany, S.H., 2011. Neonicotinoid insecticides imidacloprid and clothianidin
336 affect differently neural Kenyon cell death in the cockroach *Periplaneta americana*. *Pestic.*
337 *Biochem. Physiol.* 101, 191–197. <https://doi.org/10.1016/j.pestbp.2011.09.005>

338 Bonmatin, J.-M., Giorio, C., Girolami, V., Goulson, D., Kreuzweiser, D.P., Krupke, C., Liess, M., Long,
339 E., Marzaro, M., Mitchell, E.A.D., 2015. Environmental fate and exposure; neonicotinoids and
340 fipronil. *Environ. Sci. Pollut. Res.* 22, 35–67.

341 Brandt, A., Gorenflo, A., Siede, R., Meixner, M., Büchler, R., 2016. The neonicotinoids thiacloprid,
342 imidacloprid, and clothianidin affect the immunocompetence of honey bees (*Apis mellifera*
343 L.). *J. Insect Physiol.* 86, 40–47. <https://doi.org/10.1016/j.jinsphys.2016.01.001>

344 Browne, N., Heelan, M., Kavanagh, K., 2013. An analysis of the structural and functional similarities of
345 insect hemocytes and mammalian phagocytes. *Virulence* 4, 597–603.

346 Camp, A., Buchwalter, D., 2016. Can't take the heat: Temperature-enhanced toxicity in the mayfly
347 *Isonychia bicolor* exposed to the neonicotinoid insecticide imidacloprid. *Aquat. Toxicol.* 178,
348 49–57.

349 Chanput, W., Mes, J.J., Wichers, H.J., 2014. THP-1 cell line: an in vitro cell model for immune
350 modulation approach. *Int. Immunopharmacol.* 23, 37–45.

351 Chauzat, M.-P., Carpentier, P., Martel, A.-C., Bougeard, S., Cougoule, N., Porta, P., Lachaize, J.,
352 Madec, F., Aubert, M., Faucon, J.-P., 2009. Influence of pesticide residues on honey bee
353 (Hymenoptera: Apidae) colony health in France. *Environ. Entomol.* 38, 514–523.

354 Cimino, A.M., Boyles, A.L., Thayer, K.A., Perry, M.J., 2016. Effects of neonicotinoid pesticide exposure
355 on human health: a systematic review. *Environ. Health Perspect.* 125, 155–162.

356 Codling, G., Al Nagggar, Y., Giesy, J.P., Robertson, A.J., 2016. Concentrations of neonicotinoid
357 insecticides in honey, pollen and honey bees (*Apis mellifera* L.) in central Saskatchewan,
358 Canada. *Chemosphere* 144, 2321–2328. <https://doi.org/10.1016/j.chemosphere.2015.10.135>

359 Commission implementing regulation (EU) 2018/783 of 29 May 2018 amending Implementing
360 Regulation (EU) No 540/2011 as regards the conditions of approval of the active substance
361 imidacloprid; Official Journal of the European Union, L 132/31, 2018

362 Couleau, N., Techer, D., Pagnout, C., Jomini, S., Foucaud, L., Laval-Gilly, P., Falla, J., Bennisroune, A.,
363 2012. Hemocyte responses of *Dreissena polymorpha* following a short-term in vivo exposure
364 to titanium dioxide nanoparticles: Preliminary investigations. *Sci. Total Environ.* 438, 490–
365 497.

366 Cresswell, J.E., 2011. A meta-analysis of experiments testing the effects of a neonicotinoid insecticide
367 (imidacloprid) on honey bees. *Ecotoxicology* 20, 149–157.

368 Cresswell, J.E., Page, C.J., Uygun, M.B., Holmbergh, M., Li, Y., Wheeler, J.G., Laycock, I., Pook, C.J., de
369 Ibarra, N.H., Smirnoff, N., Tyler, C.R., 2012. Differential sensitivity of honey bees and bumble
370 bees to a dietary insecticide (imidacloprid). *Zoology* 115, 365–371.
371 <https://doi.org/10.1016/j.zool.2012.05.003>

372 David, A., Botías, C., Abdul-Sada, A., Nicholls, E., Rotheray, E.L., Hill, E.M., Goulson, D., 2016.
373 Widespread contamination of wildflower and bee-collected pollen with complex mixtures of
374 neonicotinoids and fungicides commonly applied to crops. *Environ. Int.* 88, 169–178.

375 Déglise, P., Grünewald, B., Gauthier, M., 2002. The insecticide imidacloprid is a partial agonist of the
376 nicotinic receptor of honeybee Kenyon cells. *Neurosci. Lett.* 321, 13–16.
377 [https://doi.org/10.1016/S0304-3940\(01\)02400-4](https://doi.org/10.1016/S0304-3940(01)02400-4)

378 Di Prisco, G., Cavaliere, V., Annoscia, D., Varricchio, P., Caprio, E., Nazzi, F., Gargiulo, G., Pennacchio,
379 F., 2013. Neonicotinoid clothianidin adversely affects insect immunity and promotes
380 replication of a viral pathogen in honey bees. *Proc. Natl. Acad. Sci.* 110, 18466–18471.

381 Di Prisco, G., Iannaccone, M., Ianniello, F., Ferrara, R., Caprio, E., Pennacchio, F., Capparelli, R., 2017.
382 The neonicotinoid insecticide Clothianidin adversely affects immune signaling in a human cell
383 line. *Sci. Rep.* 7, 13446.

384 Dunier, M., Siwicki, A.K., Demaël, A., 1991. Effects of organophosphorus insecticides: Effects of
385 trichlorfon and dichlorvos on the immune response of carp (*Cyprinus carpio*): III. In Vitro
386 effects on lymphocyte proliferation and phagocytosis and in Vivo effects on humoral
387 response. *Ecotoxicol. Environ. Saf.* 22, 79–87. [https://doi.org/10.1016/0147-6513\(91\)90049-](https://doi.org/10.1016/0147-6513(91)90049-U)
388 U

389 Fauser-Misslin, A., Sadd, B.M., Neumann, P., Sandrock, C., 2014. Influence of combined pesticide and
390 parasite exposure on bumblebee colony traits in the laboratory. *J. Appl. Ecol.* 51, 450–459.

391 Feldhaar, H., Gross, R., 2008. Immune reactions of insects on bacterial pathogens and mutualists.
392 *Microbes Infect.* 10, 1082–1088.

393 Fernández-Alba, A.R., Tejedor, A., Agüera, A., Contreras, M., Garrido, J., 2000. Determination of
394 imidacloprid and benzimidazole residues in fruits and vegetables by liquid chromatography–
395 mass spectrometry after ethyl acetate multiresidue extraction. *J. AOAC Int.* 83, 748–755.

396 Gibbons, D., Morrissey, C., Mineau, P., 2015. A review of the direct and indirect effects of
397 neonicotinoids and fipronil on vertebrate wildlife. *Environ. Sci. Pollut. Res.* 22, 103–118.

398 Han, W., Tian, Y., Shen, X., 2018. Human exposure to neonicotinoid insecticides and the evaluation of
399 their potential toxicity: An overview. *Chemosphere* 192, 59–65.

400 Hladik, M.L., Vandever, M., Smalling, K.L., 2016. Exposure of native bees foraging in an agricultural
401 landscape to current-use pesticides. *Sci. Total Environ.* 542, Part A, 469–477.
402 <https://doi.org/10.1016/j.scitotenv.2015.10.077>

403 Iwasa, T., Motoyama, N., Ambrose, J.T., Roe, R.M., 2004. Mechanism for the differential toxicity of
404 neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Prot.* 23, 371–378.
405 <https://doi.org/10.1016/j.cropro.2003.08.018>

406 Jeschke, P., Nauen, R., Schindler, M., Elbert, A., 2010. Overview of the status and global strategy for
407 neonicotinoids. *J. Agric. Food Chem.* 59, 2897–2908.

408 Kasiotis, K.M., Anagnostopoulos, C., Anastasiadou, P., Machera, K., 2014. Pesticide residues in
409 honeybees, honey and bee pollen by LC–MS/MS screening: Reported death incidents in
410 honeybees. *Sci. Total Environ.* 485–486, 633–642.
411 <https://doi.org/10.1016/j.scitotenv.2014.03.042>

412 Ko, A.-Y., Rahman, M.M., Abd El-Aty, A.M., Jang, J., Park, J.-H., Cho, S.-K., Shim, J.-H., 2014.
413 Development of a simple extraction and oxidation procedure for the residue analysis of
414 imidacloprid and its metabolites in lettuce using gas chromatography. *Food Chem.* 148, 402–
415 409. <https://doi.org/10.1016/j.foodchem.2013.10.055>

416 Komiyama, K., Okaue, M., Miki, Y., Ohkubo, M., Moro, I., Cooper, E.L., 2003. Non-specific cellular
417 function of *Eisenia fetida* regulated by polycyclic aromatic hydrocarbons: The 7th
418 international symposium on earthworm ecology· Cardiff· Wales· 2002. *Pedobiologia* 47, 717–
419 723.

420 Linder, J.E., Owers, K.A., Promislow, D.E., 2008. The effects of temperature on host–pathogen
421 interactions in *D. melanogaster*: Who benefits? *J. Insect Physiol.* 54, 297–308.

422 López, J.H., Krainer, S., Engert, A., Schuehly, W., Riessberger-Gallé, U., Crailsheim, K., 2017. Sublethal
423 pesticide doses negatively affect survival and the cellular responses in American foulbrood-
424 infected honeybee larvae. *Sci. Rep.* 7, 40853.

425 Mason, R., Tennekes, H., Sánchez-Bayo, F., Jepsen, P.U., 2013. Immune suppression by neonicotinoid
426 insecticides at the root of global wildlife declines. *J. Env. Immunol Toxicol* 1, 3–12.

427 Mostafalou, S., Abdollahi, M., 2013. Pesticides and human chronic diseases: Evidences, mechanisms,
428 and perspectives. *Toxicol. Appl. Pharmacol.* 268, 157–177.
429 <https://doi.org/10.1016/j.taap.2013.01.025>

430 Nauen, R., Ebbinghaus-Kintscher, U., Schmuck, R., 2001. Toxicity and nicotinic acetylcholine receptor
431 interaction of imidacloprid and its metabolites in *Apis mellifera* (Hymenoptera: Apidae). *Pest*
432 *Manag. Sci.* 57, 577–586.

433 Negri, P., Maggi, M., Correa-Aragunde, N., Brasesco, C., Eguaras, M., Lamattina, L., 2013. Nitric oxide
434 participates at the first steps of *Apis mellifera* cellular immune activation in response to non-
435 self recognition. *Apidologie* 44, 575–585.

436 Nomura, H., Ueyama, J., Kondo, T., Saito, I., Murata, K., Iwata, T., Wakusawa, S., Kamijima, M., 2013.
437 Quantitation of neonicotinoid metabolites in human urine using GC-MS. *J. Chromatogr. B*
438 941, 109–115. <https://doi.org/10.1016/j.jchromb.2013.10.012>

439 Obana, H., Okihashi, M., Akutsu, K., Kitagawa, Y., Hori, S., 2002. Determination of acetamiprid,
440 imidacloprid, and nitenpyram residues in vegetables and fruits by high-performance liquid
441 chromatography with diode-array detection. *J. Agric. Food Chem.* 50, 4464–4467.

442 Pettis, J.S., Johnson, J., Dively, G., 2012. Pesticide exposure in honey bees results in increased levels
443 of the gut pathogen *Nosema*. *Naturwissenschaften* 99, 153–158.

444 Roszell, L., Anderson, R., 1994. Inhibition of phagocytosis and superoxide production by
445 pentachlorophenol in two leukocyte subpopulations from *Fundulus heteroclitus*. *Mar.*
446 *Environ. Res.* 38, 195–206. [https://doi.org/10.1016/0141-1136\(94\)90007-8](https://doi.org/10.1016/0141-1136(94)90007-8)

447 Sánchez-Bayo, F., Goka, K., 2005. Unexpected effects of zinc pyrethrin and imidacloprid on Japanese
448 medaka fish (*Oryzias latipes*). *Aquat. Toxicol.* 74, 285–293.

449 Schmid, M.R., Brockmann, A., Pirk, C.W., Stanley, D.W., Tautz, J., 2008. Adult honeybees (*Apis*
450 *mellifera* L.) abandon hemocytic, but not phenoloxidase-based immunity. *J. Insect Physiol.*
451 54, 439–444.

452 Simon-Delso, N., Amaral-Rogers, V., Belzunces, L.P., Bonmatin, J.M., Chagnon, M., Downs, C., Furlan,
453 L., Gibbons, D.W., Giorio, C., Girolami, V., Goulson, D., Kreuzweiser, D.P., Krupke, C.H., Liess,
454 M., Long, E., McField, M., Mineau, P., Mitchell, E.A.D., Morrissey, C.A., Noome, D.A., Pisa, L.,
455 Settele, J., Stark, J.D., Tapparo, A., Van Dyck, H., Van Praagh, J., Van der Sluijs, J.P.,
456 Whitehorn, P.R., Wiemers, M., 2015. Systemic insecticides (neonicotinoids and fipronil):
457 trends, uses, mode of action and metabolites. *Environ. Sci. Pollut. Res.* 22, 5–34.
458 <https://doi.org/10.1007/s11356-014-3470-y>

459 Song, M.Y., Stark, J.D., Brown, J.J., 1997. Comparative toxicity of four insecticides, including
460 imidacloprid and tebufenozide, to four aquatic arthropods. *Environ. Toxicol. Chem.* 16, 2494–
461 2500.

462 Suchail, S., Debrauwer, L., Belzunces, L.P., 2004. Metabolism of imidacloprid in *Apis mellifera*. *Pest*
463 *Manag. Sci.* 60, 291–296.



464 Taira, K., Fujioka, K., Aoyama, Y., 2013. Qualitative profiling and quantification of neonicotinoid
465 metabolites in human urine by liquid chromatography coupled with mass spectrometry.

466 Tolnai, S., 1975. A method for viable cell count. *TCA Man. Tissue Cult. Assoc.* 1, 37–38.
467 <https://doi.org/10.1007/BF00914435>

468 Valanne, S., Wang, J.-H., Rämét, M., 2011. The *Drosophila* toll signaling pathway. *J. Immunol.* 186,
469 649–656.

470 Van den Brink, P.J., Van Smeden, J.M., Bekele, R.S., Dierick, W., De Gelder, D.M., Noteboom, M.,
471 Roessink, I., 2016. Acute and chronic toxicity of neonicotinoids to nymphs of a mayfly species
472 and some notes on seasonal differences. *Environ. Toxicol. Chem.* 35, 128–133.

473 Vidau, C., Diogon, M., Aufauvre, J., Fontbonne, R., Viguès, B., Brunet, J.-L., Texier, C., Biron, D.G., Blot,
474 N., El Alaoui, H., 2011. Exposure to sublethal doses of fipronil and thiacloprid highly increases
475 mortality of honeybees previously infected by *Nosema ceranae*. PLoS One 6, e21550.
476



Phagocytosis of latex beads after immune activation with LPS

Phagocytic cell

Fluorescent latex bead

Detailed description: This legend block is enclosed in a rounded rectangle. It contains a title 'Phagocytosis of latex beads after immune activation with LPS'. Below the title, there are two items: a purple circular cell with a blue outline, labeled 'Phagocytic cell', and a small yellow dot, labeled 'Fluorescent latex bead'.