

Bioavailability and transfer of elevated Sm concentration to alfalfa in spiked soils

Ruoyu Hu, Thierry Beguiristain, Alexis A. de Junet, Corinne Leyval

► To cite this version:

Ruoyu Hu, Thierry Beguiristain, Alexis A. de Junet, Corinne Leyval. Bioavailability and transfer of elevated Sm concentration to alfalfa in spiked soils. Environmental Science and Pollution Research, 2020, 10.1007/s11356-020-09223-z. hal-02961811

HAL Id: hal-02961811 https://hal.univ-lorraine.fr/hal-02961811

Submitted on 8 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bioavailability and transfer of elevated Sm concentration to alfalfa in spiked soils

Ruoyu HU, Thierry BEGUIRISTAIN, Alexis De JUNET, Corinne LEYVAL*

Université de Lorraine, CNRS, LIEC, F-54000 Nancy, France * Corresponding author: corinne.leyval@univ-lorraine.fr, tel. 33(0)372745204

Abstract Rare earth elements (REEs) have been widely used in recent decades and their exploitation has led to industrial REE emission and to contaminated soils especially in former mining areas. This raised people's concerns on the accumulation and toxicity of REEs in soils and plants, and consequences on plant health. Although many studies dealt with REE in soils and plants, there is still a need to precise their toxicity, bioavailability and transfer to plants in contaminated sites in order to restore such ecosystems. We studied the bioavailability and transfer of a REE to *Medicago sativa* grown on two contaminated soils differing in their chemical characteristics. A pot experiment was set up in a growth chamber where two natural soils were spiked or not with samarium (Sm) as a model REE. Two chemical extractants were tested to estimate the bioavailability of Sm in the soil, its decrease with time and its transfer to the plants. Results showed that DTPA extractable Sm was well correlated with Sm uptake in alfalfa shoots. The experiment pointed out a significantly lower in the less acidic soil than in the other. The uptake of Sm from soil to alfalfa shoots depended on the soil pH and on the spiking concentration. The soil to plant transfer factor was low (<0.08) but a 30% reduction of alfalfa biomass was observed when the soils were spiked with 100 to 200 mg kg⁻¹ of Sm.

Keywords:

Rare earth element, Transfer, Soil, Bioavailability, Plant, Ageing

1. Introduction

Rare earth elements (REEs) are a group of elements in the periodic table consisting of 15 lanthanide group elements from lanthanum (La) to lutetium (Lu), along with the pseudo lanthanide (Tyler 2004). According mainly to the geochemical behaviour of the REEs, they were divided into light (LREEs; La to Sm) and heavy REEs (HREEs; europium to Lu) (Brioschi et al. 2013). Because of their magnetic, fluorescent and catalytic properties, REEs, especially LREEs are nowadays widely used in many industries especially high-technology industries like electronics and medicine (Loell et al. 2011a; Pagano 2016). In natural soils, REE concentrations range from 40 to 700 mg·kg⁻¹ all over the world and are not considered as toxic for the environment. However, the higher demand and mining of REEs in the past decades have led to a notable increase in REEs in the environment (MacMillan et al. 2019) and high soil REE concentrations in former mining areas (Chao et al. 2016; Khan et al. 2017). For example, REE concentration in soil of Bayan Obo rare earth mining area in China is above 2.7×10^4 mg·kg⁻¹ due to the mining activity (Liang et al. 2014). Landfills with improperly discarded materials containing REEs might also be a source of contamination.

REE concentrations in plants are very low in natural soils (Tyler 2004), but their phytotoxicity may be a concern in contaminated areas (Carpenter et al. 2015). REEs are not considered as essential elements for plants, but positive effects of REEs at low concentrations were reported on plant growth (Diatloff et al. 1995a; Xu et al.

2002; Hong et al. 2003; Xiangsheng et al. 2006; Cheng et al. 2015). In China, REEs have even been used to promote plant growth and have been widely applied in agriculture since the 1990s. However, when a high dose of REE fertiliser has been applied, plant growth was limited and even a toxic effect was found (Hong et al. 2003). Diatloff et al. (1995a, b) studied the toxicity of lanthanum (La) and cerium (Ce) on corn and mungbean seedlings in nutrient solutions and showed a strong inhibition of mungbean root elongation and plant growth with Ce (even at 0.2 μ M). Thomas et al. (2014) studied the phytotoxicity of La, Ce and Y (yttrium) on germination and growth of native plants and crops in two artificial soils with different pH. However, there is little information on the phytotoxicity of REEs in contaminated soils. The excessive amounts of REEs are not only harmful to plants but are also able to be taken up by plants and enter the food chain, resulting in their accumulation in organisms and finally in human beings (Loell et al. 2011a; Turra et al. 2013). Therefore, studies regarding the effects of REEs in contaminated soils on plants, and their transfer from soil to plants should be carried out (Gonzalez et al. 2014). The LREE uptake by plants involves calcium (Ca²⁺ ion) channels and HREE uptake uses aluminium (Al³⁺ ion) channels (Gonzalez et al. 2014; Yuan et al. 2017). The analysis of REEs in different plants on mining and control areas was performed for health risk assessment and showed an enrichment of LREEs in the former (Zhuang et al. 2017) or to identify potential hyperaccumulating plants (Khan et al. 2017).

The bioavailability of REEs has been discussed in many studies but many of them considered natural soils where their concentrations are low, or agricultural soils with fertilizer inputs. The mobility of REEs in soils is affected by REE sources and forms, soil particle size, pH and organic matter content (Tyler 2004; Babula et al. 2008; MacMillan et al. 2019). The bioavailability of REEs and transfer to plants in soils contaminated due to industrial or mining activities has been less considered. Different methods or development of methods have been studied to estimate/predict the uptake of metals in different plants. Among these methods, simple extractants have not been widely investigated for REEs. Using sequential chemical extractions, Cao et al. (2000b) showed that bioavailability of REEs differed with the REEs. Li et al. (2001) compared different extractants and found that 0.1 M HCl extractable fraction of REEs was well correlated with their concentration in wheat roots and shoots in cultivated soils in China. Li et al. (2018) used CaCl₂ to correlate lanthanide toxicity to soil invertebrates with bioavailability.

This work aimed to study the bioavailability and transfer of a REE to a crop plant *Medicago sativa* grown on two contaminated soils differing in their chemical characteristics. A pot experiment was set up in a growth chamber where two natural soils were spiked or not with elevated concentrations of Samarium (Sm) as a model REE. The bioavailability of Sm was estimated using two chemical extractants in the spiked and control soils just after spiking, after two weeks of ageing and at plant harvest. Sm phytoavailability was also estimated by plant accumulation at harvest. Plant growth, P and Ca plant uptake were also estimated.

2. Material and methods

2.1. Experiment design

The soils used for the experiments were collected from the surface horizon (5-25 cm) of two sampling sites in north-eastern France: Maron (48°38'45.75"N, 6°4'48.44"E) and Bouzule (48°44'7.95"N, 6°19'9.64"E). They both belong to Luvisol with different pH values. The physicochemical parameters of the two sampled soils are shown in Table 1. Soil samples were air-dried, ground, sieved to 2 mm, homogenised and stored in a dry and

dark room ready to be used in the experiments. The Bouzule soil was partially amended with pre-sieved (2 mm), washed and dried sand (30w/70w) to avoid compaction and allow seedling growth. A ten times concentrated solution of SmCl₃•6H₂O in deionized water was added to 1/10 of both soils. The concentrated spiked soils were dried at 50 °C, homogenised, and added to the rest of the unspiked soils and homogenised to obtain 100 and 200 mg·kg⁻¹ Sm. 1 kg of soils were placed in 1kg pots with eight replicates and three Sm conditions (0, 100, 200 mg·kg⁻¹) according to LREE concentrations in abandoned mining sites (Chao et al. 2016). Deionized water was added to reach 70% of water holding capacity (WHC) and pots were placed in a dark room at 4°C (T0) for 2 weeks for the ageing of the Sm contamination.

After soil ageing, 4-5-day-old alfalfa (*Medicago sativa*) seedlings germinated on sterilised vermiculite were transplanted to the pots (T1) with 15 plants in each pot. Pots with plants were transferred to a growth chamber with 16 h photoperiod, 22°C/18°C day/night, 250 µmol photons m⁻² s⁻¹, 80% relative humidity for 6 weeks. During the 6-week growing period, the pots were watered twice a week and a nutrient solution (Redon et al. 2009) was added once per week to maintain 70% WHC.

2.2. Sampling and analysis

Soil samples were collected at T0 and T1 in centrifuge tubes and dried at 80 °C for 72h. Alfalfa plants and soil samples were harvested (T2) 28 days (Maron) or 42 days (Bouzule) after transplantation. Pots with Maron soil were harvested earlier due to the poor growth of plants on this soil. Shoots and roots were harvested, gently rinsed with deionized water then dried at 60 °C for 72 h, and the dry weight of shoots and roots of each pot was recorded. Soil samples were collected at T2 in centrifuge tubes and dried at 80 °C for 72 h. All soil and shoot samples were ground in a mortar with a pestle before the analysis.

To analyse the total concentration of Sm in soil, 200 mg of ground soil sample was transferred into an acid-cleaned Teflon tube designed for CEM Mars 5 Digestion Microwave System® and 10 mL of *Aqua regia* (nitric acid: hydrochloric acid =1:3 v/v) was added. Digestion was processed under the adapted programme (15 min digestion at 25°C and 10 min digestion at 170°C). After the digestion and cooling down procedure, the digested samples were filtered (DigiFILTER, SCP Science® 010-500-070), and the filtered solutions were transferred into 15 mL centrifuge tubes, adjusted to 10 mL with ultra-pure deionized water 18 MW and preserved at 4 °C till the analysis.

The bioavailable concentrations of Sm in soils were extracted by two different extractants. The DTPA extracting solution consists of diethylenetriaminepentaacetic acid (DTPA) (0.005M), CaCl₂ (0.01M) and triethanolamine (0.1M) (Carter and Gregorich 2007; Kim et al. 2015). One gram of ground soil sample was transferred into a 15 mL centrifuge tube, with 10 mL of DTPA solution and homogenised on a rotary shaker with 99 rpm for 2 hours. Samples were centrifuged at 3227 g for 15 min (20 °C) and the supernatant was filtered in 15 mL centrifuge tubes, adjusted to 10 mL with ultra-pure deionized water 18 M W and preserved at 4 °C till the analysis. The second extractant is CaCl₂. One gram of ground sample was transferred into a 15 mL centrifuge tubes, adjusted to 10 mL with the same homogenization and centrifugation method as above. The supernatant was filtered in 15 mL centrifuge tubes, adjusted to 10 mL with the same homogenization and centrifugation method as above. The supernatant was filtered in 15 mL centrifuge tubes, adjusted to 10 mL with the same homogenization and centrifugation method as above. We and preserved at 4 °C till the analysis.

Dried shoot samples were digested in nitric acid (HNO₃) and hydrogen peroxide (H₂O₂) solution. Six mL of 69% HNO₃ (CAS: 7697-37-2) and 3 mL of 30 % H₂O₂ (w/w H₂O) (CAS: 7722-84-1) were added into DigiTUBEs containing 200 mg of ground shoot sample at 20 °C for a night. Then the DigiTUBEs with the sample solution was placed on DigiPREP MS 48 & TS (SCP Science® 010-500-205 & 010-500-275) hot plate system covered with DigiPROBE Watch Glass (SCP Science® 010-500-081) processing the digestion programme adapted to plant tissue with 5 min solubilization at 45 °C, 10 min pre-digestion at 65 °C and 60 min digestion at 95 °C. After cooling down, the solution was filtered as previously described and kept at 4 °C until the analysis.

The total and the bioavailable Sm concentrations in soil samples and the concentration of Sm in alfalfa shoots were analysed using coupled plasma-optical emission spectroscopy (ICP-OES) analytical technique using a Varian Inc.® (Part A) - 720/730-ES ICP-OES. Calcium (Ca), sodium (Na), potassium (K) and magnesium (Mg) concentrations in alfalfa shoots were also analysed using the same technique and are presented as supplementary data (Online Resource 1). All the analysis used NCS Certified Reference Material NCS DC 73030 soil (Sm recovery rate = 85.1%) and Labmix24 oriental basma tabacco leaves (track elements) - IC-INCT-OBTL-5 (Sm recovery rate = 70%) as references. The soil-plant transfer factor (SPTF) (Papastefanou et al. 1999; Cao et al. 2000a; Chyi 2003; Šmuc et al. 2012) was calculated as the ratio between Sm concentration in alfalfa shoots and the total concentration of Sm in soils.

Statistical analysis was performed with RStudio software (Version 1.1.463 – \bigcirc 2009-2018 RStudio, Inc.). One-way and two-way analysis of variance (ANOVA 1 and 2) and the Tukey honestly-significant-difference (HSD) post-hoc test were performed for the significance of the data with a 95% significant level (p-value ≤ 0.05).

3. Results

3.1. Total and extractable Sm in both soils

Sm concentration in unspiked Maron and Bouzule soils was $14.83 \pm 0.37 \text{ mg} \cdot \text{kg}^{-1}$ and $4.99 \pm 0.46 \text{ mg} \cdot \text{kg}^{-1}$, respectively. After spiking with 100 and 200 mg \cdot \text{kg}^{-1} Sm, the concentrations were $82.76 \pm 12.25 \text{ mg} \cdot \text{kg}^{-1}$ and $169.48 \pm 47.11 \text{ mg} \cdot \text{kg}^{-1}$ for Maron soil, and $86.44 \pm 32.62 \text{ mg} \cdot \text{kg}^{-1}$ and $151.49 \pm 63.09 \text{ mg} \cdot \text{kg}^{-1}$ for Bouzule soil, respectively.

Both CaCl₂ and DTPA extractants were used for Maron and Bouzule soils at 3 sampling dates: at spiking (T0), after 2 ageing weeks (T1) and at plant harvest (T2), and in unspiked soils. For Maron soil CaCl₂ extractable Sm was below the detection limit (0.1 mg·kg⁻¹) in unspiked soil. In spiked soils, CaCl₂ extractable Sm was 0.2 ± 0.02 mg·kg⁻¹ at T0, 0.1 ± 0.01 mg·kg⁻¹ at T2 in 100 mg·kg⁻¹ spiked soil, and 0.39 ± 0.03 mg·kg⁻¹ at T0 and 0.18 ± 0.02 mg·kg⁻¹ at T2 in 200 mg·kg⁻¹ spiked soil. For Bouzule soil, CaCl₂ extractable Sm was below the detection limit in all samples.

The DTPA extractable Sm in unspiked Maron and Bouzule soils was also low: $4.16 \pm 0.58 \text{ mg} \cdot \text{kg}^{-1}$ at T0, $4.13 \pm 0.35 \text{ mg} \cdot \text{kg}^{-1}$ at T1 and $3.27 \pm 0.45 \text{ mg} \cdot \text{kg}^{-1}$ at T2 in Maron soil, without significant difference between the 3 sampling dates. Much lower DTPA extractable Sm was found in Bouzule soil with 0.36 ± 0.05

mg·kg⁻¹ at T0, 0.30 \pm 0.03 mg·kg⁻¹ at T1 and 0.20 \pm 0.005 mg·kg⁻¹ at T2, with a significant difference between T0 and T2.

The DTPA extractable concentration of Sm in the spiked soils decreased as a function of time (Figure 1). It decreased by 16% and 11% respectively in 100 and 200 mg·kg⁻¹ spiked Maron soil during the 2-week ageing period from T0 to T1. For Bouzule soil, there was a significant 24% and 29% decrease of DTPA extractable Sm in 100 and 200 mg·kg⁻¹ spiked soil respectively during the 2-week ageing (T0 to T1). A significant decrease was also observed between T1 and T2, during the 4 weeks' alfalfa growing time, with 24% and 26% decrease respectively in 100 and 200 mg·kg⁻¹ spiked Maron soil. During the 6 weeks' alfalfa growing time (between T1 and T2), the available Sm concentration in Bouzule spiked soil also significantly decreased with a 48% and 40% decrease at 100 and 200 mg·kg⁻¹ Sm spiked concentration.

3.2. Alfalfa growth in Sm spiked and unspiked soils

In Maron soil, the shoot biomass showed a significant decrease in the 200 mg kg⁻¹ Sm spiked soil compared to the unspiked control but the root biomass showed no significant difference between control and spiked soils. Shoot and root biomass in Bouzule soil was significantly lower in Sm spiked than in the unspiked soil (Figure 2). It was significant already with 100 mg·kg⁻¹ Sm spiked concentration (p-value ≤ 0.05) and no further decrease was observed at 200 mg kg⁻¹. For Maron soil, the shoot biomass showed a significant decrease in the 200 mg kg⁻¹ Sm spiked soil compared to the unspiked control but the root biomass showed no significant difference between control and spiked soils.

3.3. Sm transfer to plants

The concentration of Sm in alfalfa shoots was significantly higher in the spiked soils than in the controls and was twice higher with 200 than with 100 mg kg⁻¹ Sm spiked soils (p-value = 0.05) (Figure 3). The soil-plant transfer factor (SPTF) was around 0.05-0.08 in Maron soil, and it was not significantly different between spiked and unspiked soils. However, in Bouzule soil, the SPTF was around 0.02 to 0.035 and was significantly lower in spiked than in unspiked soil (Figure 3). The Sm uptake by plants as a function of DTPA-extractable concentration of Sm in soil was also calculated (Figure 4) and showed a positive linear correlation ($R^2 = 0.9334$, linear regression model).

We analysed P and Ca concentrations in plants. There were no significant differences in P and Ca concentrations in alfalfa shoots between unspiked and spiked soils (data not shown). P and Ca content in shoots was also not significantly different between spiked and unspiked Maron soil. However, in Bouzule spiked soil, the content of P and Ca in shoots was significantly lower than in unspiked soil (Figure 5).

4. Discussion

4.1. Assessment of bioavailability of Sm using different extractants

Different extractants and extraction methods have been applied to study the bioavailability of REEs in the soil or their phytoavailability and potential uptake by plants. In our study, we performed a single extraction method using two neutral extractants: DTPA and CaCl₂. Our results showed that CaCl₂ extractant had a low extracting ability, while the phytoavailable Sm in the soil was well predicted by DTPA extraction as the Sm

plant uptake accounted for 22.9% of DTPA extractable Sm (with R²=0.93). According to Rao et al. (2010), the bioavailability of REEs can be overestimated by strong acid reagents such as CH₃COOH and HCl. Meanwhile, the complexing extractants such as DTPA and EDTA showed moderate extraction ability and have been applied in different studies. EDTA was used to indicate that 10% to 16% of REEs were phytoavailable in natural soils in central Germany (Loell et al. 2011a, b). EDTA showed a stronger extracting ability than DTPA as it can extract both carbonate and organically bound fractions of metals in soils with low carbonate content (Sahuquillo et al. 2003). For Censi et al. (2017), DTPA gave a good indication of the plant-bioavailable fraction of REEs and other trace elements in rural, urban and industrial soils in Sicily. The neutral salt 0.01M CaCl₂ was recommended as a good extractant of phytoavailability (Menzies et al. 2007; Kim et al. 2015). However, in our study, CaCl₂ could not be used to represent alfalfa available Sm in both soils due to its weak extraction ability, while DTPA showed a good indication of alfalfa available Sm in the soil.

4.2. Bioavailability of high concentration of Sm in soils and influence of ageing

In our study, the bioavailable Sm, estimated by DTPA extraction, accounted for 40% of the total Sm concentration (extracted by *aqua regia*) in freshly spiked Maron soil and 25% in freshly spiked Bouzule soil. Despite the same spiking concentrations in both soils, it was significantly higher in the acidic Maron soil than the more neutral Bouzule soil. In a study of four types of Chinese soils, with both control and spiked (200 mg·kg⁻¹ REEs) soils, the red earth with lowest pH had the highest exchangeable REEs (Liang et al. 2005). A higher REE availability in lower pH soil was also reported in a study comparing total REE contents and the potentially bioavailable fraction using EDTA in three different soils in Germany (Diatloff et al. 1995b). Diatloff et al. (1996) found that the Ca(NO₃)₂ extractable REEs decreased with the soil pH increase when adding CaCO₃ and increased when soil pH decreased by adding CaSO₄•2H₂O. Yufeng et al. (2003) applied artificial acid rainwater to Chinese red soil and showed an increase in the REE concentrations in the soil leaching solution after the acid rainwater application. Our result of higher bioavailable Sm in Maron soil further confirmed pH as an important parameter influencing the bioavailability of REEs.

Despite soil pH and soil composition, the organic matter in soil with a large amount of negatively charged groups is able to adsorb or chelate cations as REEs and thereby decrease their availability and transfer (Tyler 2004; Loell et al. 2011b; Fiket et al. 2017). In our study, the bioavailability of Sm is higher in Maron soil even though the organic matter was higher than in Bouzule soil, indicating that the pH had a major influence on REE bioavailability. Ca^{2+} was reported to have an influence on the bioavailability of LREE to plant, as LREE are reported to use the same ion channels as Ca^{2+} when entering plant root (Yuan et al. 2017). In our study, there was no indication that Ca concentration influenced the bioavailability of the LREE Sm, but calcium concentration in both soils was similar.

Other than the above-mentioned factors, the ageing significantly affected the bioavailability of Sm in the spiked soils. In our study, the bioavailable fraction of Sm significantly decreased within the two weeks of ageing. Cao et al. (2000b) studied the bioavailability of REEs in natural soils using the same model plant alfalfa (*Medicago sativa*) and sequential extractions. They showed that exchangeable REE fraction contributed to the bioavailable REEs, which explained the decreased bioavailability of REEs, including Sm, with time as REEs were adsorbed to soil particles. In the unspiked soils, the bioavailable fraction of Sm was much lower than in the spiked soils (26 and 6 % for Maron and Bouzule, respectively), which was also observed for natural and spiked

soils by Liang et al. (2005). The bioavailable fraction of REEs accounted for only 10.1 % of total REEs and 13.8% of Sm in natural soil (Loell et al. 2011b). We also observed a further decrease of the bioavailable Sm from T1 to T2 down to 25 % in Maron and 9 % in Bouzule soil which could be due to the ageing effect and the presence of alfalfa plants as well. Therefore, REE toxicity studies with spiked soils (Thomas et al., 2014) should include an ageing period of the soil before to introduce the plants.

4.3. Impact of high concentration of Sm on plant growth

Although some studies have mentioned a positive influence of REEs on plant growth at low concentration in soil (Hong et al. 2003; Cheng et al. 2015), other studies reported the inhibition of plant growth and toxic effects of high concentrations REEs to plants (Carpenter et al. 2015). However, most of the studies on the effects of REEs on plant growth were performed in liquid media (Asher and Smith 1995; Diatloff et al. 1995a, b; d'Aquino et al. 2009) on two or three LREEs (yttrium, lanthanum and cerium) and not in soil. Few data are focusing on the effect of single elements (Pagano 2016). In our study, we focused on the effect of a single REE on alfalfa. REE toxicity and transfer to plants may differ with plant species and REEs. This was observed with elevated La, Y and Ce concentrations in artificial spiked soil where lower IC25s values for Ce and some plants were observed (Thomas et al. 2014). However, we showed a strong decrease, around 30%, of alfalfa shoot biomass when the Sm concentration in soil reached 100 to 200 mg·kg⁻¹. Carpenter et al. (2015) showed that the IC25s of native plants in artificial soils spiked with 6 REEs was between 100 and 300 mg·kg⁻¹ REEs, but the ageing of the contamination was not considered neither in the experiment of Liang et al. (2005) where ryegrass weight was inhibited by spiking REEs on natural soils. Since the spiking was performed with SmCl₃, a potential effect of Cl addition might be considered. Salinity, for example addition of NaCl, can be harmful to alfalfa growth and lead to decreased concentration of shoot K, Ca and P (Li et al. 2010; Ashrafi et al. 2018). However, compared to these experiments, the added Cl concentration in our study was much lower. Further, the concentration of Ca, and P, and Na and K in plants was not significantly different between SmCl₃ spiked and non-spiked soil (Online Resource 1), suggesting no side effect of the addition of Cl.

4.4. Transfer of Sm to plants

The SPTF of Sm was rather low: 0.05 and about 0.08 in the control and spiked Maron soil, respectively, and 0.035 and about 0.02 in the control and spiked Bouzule soil. The difference between control and spiked soils can be explained by the source and ageing time of Sm. In studies focussing on natural soils, the transfer of REEs to plants, especially to wheat seeds (Liang et al. 2005) or fruits (Cheng et al. 2015) was also low (around 0.03). According to Tyler (2004), the concentrations of REEs in the plant are generally low due to the low SPTF, ranging between 0.04 to 0.09 and even lower in grass (Tyler and Olsson 2001) and vegetables (Bibak et al. 1999). In an experiment using REE-containing coal fly ash amendment He et al. (2019) showed a transfer factor from soil to alfalfa shoots of the light REEs La and Ce, and of the heavy REE Y ranging from 0.01 to 0.10. The transfer factor depended on the element and was higher for Ce than the two other REEs. Our results for Maron soil fit with this range of data and the SPTF is lower in Bouzule soil which was mainly influenced by the availability of Sm in the soil. Moreover, the preferable neutral soil for alfalfa lead to better growth in Bouzule soil and a dilution effect occurred. Other plants reported to be hyperaccumulators, such as *Phytolacca Americana* that can accumulate up to 623 mg·kg⁻¹REEs or the Chinese fern *Dicranopteris dichotoma*, showed

higher SPTF values, up to 75, depending on the soil properties and REE concentrations (Chen et al. 2016; van der Ent et al. 2018). Our study focused on REE transfer from a contaminated soil to a widespread crop plant with rapid growth, where, although the SPTF value is lower, the plant can be strongly impacted by the presence of REEs.

5. Conclusions

Although the SPTF of Sm to alfalfa represented a low percentage, the presence of Sm at high concentration inhibited the growth of alfalfa. In areas with a high natural abundance of REEs, in post-mining sites, or with the increasing risk of uncontrolled landfills, non-adapted plants might then be strongly impacted. pH was a key parameter of the bioavailability of Sm since higher Sm availability and higher SPTF were found in the acid soil. The bioavailable Sm decreased during the ageing process in the soils, suggesting interactions between Sm and soil constituents. A decrease with time of the bio- and phytoavailability of REE in contaminated soils can be expected and help native or crop plants to cope with the presence of REEs. However, in order to further reduce the bioavailability and the transfer to plants of REEs in contaminated soils, and restore such ecosystems, other soil parameters could be considered such as Ca content and organic matter. Rhizospheric processes, such as root exudation or the contribution of root-associated microorganisms, which are known to affect the mobility and availability of metallic trace elements, should also be investigated for REEs.

Acknowledgements

This work was supported by the French National Research Agency through the national program "Investissements d'avenir" with the reference ANR-10-LABX-21- RESSOURCES21. The authors would like to give thanks to the technical staff of LIEC especially David BILLET, Christine FRIRY and Hervé MARMIER, and to the Experimental phytotronic platform of Lorraine (PEPLor, Université de Lorraine).

References

- Asher CJ, Smith FW (1995) Rare earth elements and plant growth: II. responses of corn and mungbean to low concentrations of lanthanum in dilute, continuously flowing nutrient solutions. J Plant Nutr 18:1977–1989
- Ashrafi E, Razmjoo J, Zahedi M (2018) Effect of salt stress on Growth and Ion accumulation of alfalfa (Medicago sativa L.) cultivars. J Plant Nutr 41:818–831
- Babula P, Adam V, Opatrilova R, Zehnalek J, Havel L, Kizek R (2008) Uncommon heavy metals, metalloids and their plant toxicity: A review. Environ Chem Lett 6:189–213
- Bibak A, Stümp S, Knudsen L, Gundersen V (1999) Concentrations of 63 elements in cabbage and sprouts in Denmark. Commun Soil Sci Plant Anal 30:2409–2418
- Brioschi L, Steinmann M, Lucot E, Pierret MC, Stille P, Prunier J, Badot PM (2013) Transfer of rare earth elements (REE) from natural soil to plant systems: Implications for the environmental availability of anthropogenic REE. Plant Soil 366:143–163
- Cao X, Chen Y, Gu Z, Wang X (2000a) Determination of trace rare earth elements in plant and soil samples by inductively coupled plasma-mass spectrometry. Int J Environ Anal Chem 76:295–309
- Cao X, Wang X, Zhao G (2000b) Assessment of the bioavailability of rare earth elements in soils by chemical

fractionation and multiple regression analysis. Chemosphere 40:23-28

- Carpenter D, Boutin C, Allison JE, Parsons JL, Ellis DM (2015) Uptake and Effects of Six Rare Earth Elements (REEs) on Selected Native and Crop Species Growing in Contaminated Soils. PLoS One 10:e0129936
- Carter MR, Gregorich EG (2007) Soil Sampling and Methods of Analysis, Second Edition, 2nd Editio. CRC Press, Boca Raton, USA
- Censi P, Cibella F, Falcone EE, Cuttitta G, Saiano F, Inguaggiato C, Latteo V (2017) Rare earths and trace elements contents in leaves: A new indicator of the composition of atmospheric dust. Chemosphere 169:342–350
- Chao Y, Liu W, Chen Y, Chen W, Zhao L, Ding Q, Wang S, Tang Y-T, Zhang T, Qiu R-L (2016) Structure, Variation, and Co-occurrence of Soil Microbial Communities in Abandoned Sites of a Rare Earth Elements Mine. Environ Sci Technol 50:11481–11490
- Chen Z, Chen Z, Bai L (2016) Rare earth element migration in gullies with different Dicranopteris dichotoma covers in the Huangnikeng gully group, Changting County, Southeast China. Chemosphere 164:443–450
- Cheng J, Ding C, Li X, Zhang T, Wang X (2015) Rare earth element transfer from soil to navel orange pulp (Citrus sinensis Osbeck cv. Newhall) and the effects on internal fruit quality. PLoS One 10:1–15
- Chyi LL (2003) Trace elements in soils and plants, 4th Editio. CRC Press, Boca Raton, USA
- d'Aquino L, de Pinto MC, Nardi L, Morgana M, Tommasi F (2009) Effect of some light rare earth elements on seed germination, seedling growth and antioxidant metabolism in Triticum durum. Chemosphere 75:900–905
- Diatloff E, Asher CJ, Smith FW (1996) Concentrations of rare earth elements in some Australian soils. Aust J Soil Res 34:735–747
- Diatloff E, Smith FW, Asher CJ (1995a) Rare earth elements and plant growth: III. Responses of corn and mungbean to low concentrations of cerium in dilute, continuously flowing nutrient solutions. J Plant Nutr 18:1991–2003
- Diatloff E, Smith FW, Asher CJ (1995b) Rare earth elements and plant growth: I. Effects of lanthanum and cerium on root elongation of corn and mungbean. J Plant Nutr 18:1963–1976
- Fiket Ž, Medunić G, Furdek Turk M, Ivanić M, Kniewald G (2017) Influence of soil characteristics on rare earth fingerprints in mosses and mushrooms: Example of a pristine temperate rainforest (Slavonia, Croatia). Chemosphere 179:92–100
- Gonzalez V, Vignati DAL, Leyval C, Giamberini L (2014) Environmental fate and ecotoxicity of lanthanides: Are they a uniform group beyond chemistry? Environ Int 71:148–157
- He H, Fan C, Peng Q, Wu M, Zheng J, Wu G (2019) Bioaccumulation and translocation of rare earth elements in two forage legumes grown in soils treated with coal fly ash. Chem Geol 528:119284
- Hong F, Wang L, Liu C (2003) Study of lanthanum on seed germination and growth of rice. Biol Trace Elem Res 94:273–286
- Khan AM, Behkami S, Yusoff I, Md Zain S Bin, Bakar NKA, Bakar AFA, Alias Y (2017) Geochemical characteristics of rare earth elements in different types of soil: A chemometric approach. Chemosphere 184:673–678
- Kim RY, Yoon JK, Kim TS, Yang JE, Owens G, Kim KR (2015) Bioavailability of heavy metals in soils: definitions and practical implementation—a critical review. Environ Geochem Health 37:1041–1061

- Li F, Shan X, Zhang S (2001) Evaluation of single extractants for assessing plant availability of rare earth elements in soils. Commun Soil Sci Plant Anal 32:2577–2587
- Li J, Verweij RA, van Gestel CAM (2018) Lanthanum toxicity to five different species of soil invertebrates in relation to availability in soil. Chemosphere 193:412–420
- Li R, Shi F, Fukuda K, Yang Y (2010) Effects of salt and alkali stresses on germination, growth, photosynthesis and ion accumulation in alfalfa (Medicago sativa L.). Soil Sci Plant Nutr 56:725–733
- Liang T, Li K, Wang L (2014) State of rare earth elements in different environmental components in mining areas of China. Environ Monit Assess 186:1499–1513
- Liang T, Zhang S, Wang L, Kung H Te, Wang Y, Hu A, Ding S (2005) Environmental biogeochemical behaviors of rare earth elements in soil-plant systems. Environ Geochem Health 27:301–311
- Loell M, Albrecht C, Felix-Henningsen P (2011a) Rare earth elements and relation between their potential bioavailability and soil properties, Nidda catchment (Central Germany). Plant Soil 349:303–317
- Loell M, Reiher W, Felix-Henningsen P (2011b) Contents and bioavailability of rare earth elements in agricultural soils in Hesse (Germany). J Plant Nutr Soil Sci 174:644–654
- MacMillan GA, Clayden MG, Chételat J, Richardson MC, Ponton DE, Perron T, Amyot M (2019) Environmental Drivers of Rare Earth Element Bioaccumulation in Freshwater Zooplankton. Environ Sci Technol 53:1650–1660
- Menzies NW, Donn MJ, Kopittke PM (2007) Evaluation of extractants for estimation of the phytoavailable trace metals in soils. Environ Pollut 145:121–130
- Pagano G (2016) Rare Earth Elements in Human and Environmental Health, 1st edn. Jenny Stanford, Singapore
- Papastefanou C, Manolopoulou M, Stoulos S, Ioannidou A, Gerasopoulos E (1999) Soil-to-plant transfer of 137Cs, 40K and 7Be. J Environ Radioact 45:59–65
- Rao CRM, Sahuquillo A, Lopez-Sanchez JF (2010) Comparison of single and sequential extraction procedures for the study of rare earth elements remobilisation in different types of soils. Ananlytica Chim Acta 662:128–136
- Redon PO, Béguiristain T, Leyval C (2009) Differential effects of AM fungal isolates on Medicago truncatula growth and metal uptake in a multimetallic (Cd, Zn, Pb) contaminated agricultural soil. Mycorrhiza 19:187–195
- Sahuquillo A, Rigol A, Rauret G (2003) Overview of the use of leaching/extraction tests for risk assessment of trace metals in contaminated soils and sediments. TrAC Trends Anal Chem 22:152–159
- Šmuc NR, Dolenec T, Serafimovski T, Dolenec M, Vrhovnik P (2012) Geochemical characteristics of rare earth elements (REEs) in the paddy soil and rice (Oryza sativa L.) system of Kočani Field, Republic of Macedonia. Geoderma 183–184:1–11
- Thomas PJ, Carpenter D, Boutin C, Allison JE (2014) Rare earth elements (REEs): Effects on germination and growth of selected crop and native plant species. Chemosphere 96:57–66
- Turra C, De Nadai Fernandes EA, Bacchi MA, Sarriés GA, Júnior FB, Reyes AEL (2013) Rare earth elements in citrus production systems. J Plant Nutr 36:762–771
- Tyler G (2004) Rare earth elements in soil and plant systems A review. Plant Soil 267:191-206
- Tyler G, Olsson T (2001) Plant uptake of major and minor mineral elements as influenced by soil acidity and liming. Plant Soil 230:307–321

- van der Ent A, Echevarria G, Baker AJM, Morel JL (2018) Agromining: Farming for Metals. Springer International Publishing, Cham
- Xiangsheng L, Jiachen W, Jun Y, Yubin F, Yanping W, He Z (2006) Application of Rare Earth Phosphate Fertilizer in Western Area of China. J Rare Earths 24:423–426
- Xu X, Zhu W, Wang Z, Witkamp GJ (2002) Distributions of rare earths and heavy metals in field-grown maize after application of rare earth-containing fertilizer. Sci Total Environ 293:97–105
- Yuan M, Guo M-N, Liu W-S, Liu C, van der Ent A, Morel JL, Huot H, Zhao W-Y, Wei X-G, Qiu R-L, et al (2017) The accumulation and fractionation of Rare Earth Elements in hydroponically grown Phytolacca americana L. Plant Soil 421:67–82
- Yufeng Z, Zhenghua W, Xiaorong W, Lemei D, Yijun C (2003) Mobility of the Rare Earth Elements with Acid Rainwater Leaching in the Soil Column. Bull Environ Contam Toxicol 67:399–407
- Zhuang M, Zhao J, Li S, Liu D, Wang K, Xiao P, Yu L, Jiang Y, Song J, Zhou J, et al (2017) Concentrations and health risk assessment of rare earth elements in vegetables from mining area in Shandong, China. Chemosphere 168:578–582

	Maron	Bouzule
Soil pH	5.34	6.68
Sand: silt: clay (%)	81:362:557	167:513:320
$CaCO_3 (g \cdot kg^{-1})$	0	0
Organic carbon (g·kg ⁻¹)	40.1	15.8
Total Nitrogen (g·kg ⁻¹)	3.1	1.7
Organic matter (g·kg ⁻¹)	69.4	27.4
C/N	13.0	9.6
$P(mg\cdot kg^{-1})$	8	91
CEC cobalthexamine $(\text{cmol}^+ \cdot \text{kg}^{-1})$	23.2	16.1
Saturation rate (%)	85	100
$Ca (cmol^+ kg^{-1})$	18.1	13.1
Mg (cmol ⁺ ·kg ⁻¹)	1.07	1.79
Na (cmol ⁺ ·kg ⁻¹)	0.07	0.11
K (cmol ⁺ ·kg ⁻¹)	0.50	1.14
Fe (cmol ⁺ ·kg ⁻¹)	0.03	0.01
$Mn (cmol^+ kg^{-1})$	0.73	0.30
Al (cmol ⁺ ·kg ⁻¹)	1.45	0.05

Table 1 Physico-chemical characteristics of the soils used in the experiment



Figure1 DTPA extractable Sm concentration (mg kg⁻¹) in (a) Maron and (b) Bouzule spiked soils as a function of time (T0=0, T1=14 days, T2 = 42 days (Maron) and 56 days (Bouzule)) and spiked Sm concentration (100 and 200 mg kg⁻¹). Different letters within a soil indicate significant differences (two-way ANOVA, p<0.05).



Figure 2 Shoot and root dry weight (mg) of alfalfa plants in (a) Maron and (b) Bouzule soil as a function of spiked Sm concentration (0, 100 and 200 mg kg⁻¹). Mean \pm SD (n= 8). Different letters within a soil indicate significant differences (two-way ANOVA, p<0.05).



Figure 3 Sm concentration (mg kg⁻¹) in alfalfa shoots growing in (a) Maron and (b) Bouzule soil as a function of Sm spiked concentration (0, 100 mg kg⁻¹ and 200 mg kg⁻¹) and soil-plant transfer factor. Mean \pm SD (n= 8). Different letters within a soil indicate significant differences (two-way ANOVA, p<0.05).



Figure 4 Sm concentration (mg kg⁻¹) in plants as a function of the DTPA extractable Sm in spiked and unspiked soils.



Figure 5 Content of (a) Sm (μ g), (b) Ca (mg) and (c) P (mg) in alfalfa shoots growing in two soils as a function of Sm concentration (0, 100 ppm and 200 ppm). Mean ± SD (n=8). Different letters within a soil indicate significant differences (two-way ANOVA, p<0.05).

Online supplementary material



Figure caption: Concentration of (a) Na, (b) Ca, (c) K, (d) Mg and (e) P (mg kg⁻¹) in alfalfa shoots growing in two soils as a function of Sm concentration (0, 100 ppm and 200 ppm). Mean ± SD (n=8). Different letters within a soil indicate significant differences (two-way ANOVA, p<0.05)