



HAL
open science

How does a domestication process modulate oogenesis and reproduction performance in Eurasian perch?

Amine Khendek, Maud Alix, Sandrine Viot, Yannick Ledoré, Camille Rousseau, Robert Mandiki, Patrick Kestemont, Tomas Policar, Pascal Fontaine, Sylvain Milla

► To cite this version:

Amine Khendek, Maud Alix, Sandrine Viot, Yannick Ledoré, Camille Rousseau, et al.. How does a domestication process modulate oogenesis and reproduction performance in Eurasian perch?. *Aquaculture*, 2017, 473, pp.206-214. 10.1016/j.aquaculture.2017.02.003 . hal-02994633

HAL Id: hal-02994633

<https://hal.univ-lorraine.fr/hal-02994633>

Submitted on 8 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 **How does a domestication process modulate oogenesis and reproduction**
2 **performance in Eurasian perch?**

3

4 **Amine Khendek^a, Maud Alix^a, Sandrine Viot^a, Yannick Ledoré^a, Camille**
5 **Rousseau^a, Robert Mandiki^b, Patrick Kestemont^b, Tomas Policar^c, Pascal**
6 **Fontaine^a, Sylvain Milla^{a*}**

7 ^a *Unit Research Animal and Functionality of Animal Products (URAFPA), University of Lorraine. Boulevard des*
8 *Aiguillettes, BP 236, 54506 Vandoeuvre-Lés-Nancy, France.*

9 ^b *Research Unit in Environmental and Evolutionary Biology (URBE), NARILIS, University of Namur (FUNDP),*
10 *Rue de Bruxelles, 61, B-5000 Namur, Belgium.*

11 ^c *University of South Bohemia in Ceske Budejovice, Faculty of Fisheries and Protection of Waters, South*
12 *Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Zatisi 728/II., 389 25 Vodnany,*
13 *Czech Republic.*

14 * *Corresponding author. Tel.: +33 3 83 68 55 95 fax: +33 3 83 68 40 01;*

15 *e-mail address: Sylvain.Milla@univ-lorraine.fr*

16

17

18

19

20

21

22

23

24

25

26

27

28 **Abstract:**

29 The domestication process is accompanied with adaptation of the animals to captive
30 conditions. It induces changes at different levels thereby affecting a variety of biological
31 functions. While there is abundant literature on the domestication effects on growth and stress
32 response in teleosts, the effects on reproduction have received limited attention. In this work,
33 we investigated the domestication effect on reproductive ability of Eurasian perch (*Perca*
34 *fluviatilis* L.), a promising candidate for the development of European aquaculture and whose
35 reproductive physiological processes through the domestication history has not been
36 questioned yet.

37 To address this question, two populations of F1 and “domesticated” females were submitted
38 to a photothermal program allowing the control of gonadogenesis advancement. Gonado-
39 somatic index, 17 β -estradiol levels and oocyte diameter were significantly higher in
40 domesticated population than in F1 one. In contrast, testosterone, 11-ketotestosterone, and
41 vitellogenin levels were found to be higher in F1 females than in their domesticated
42 counterparts. Lower reproductive performance were observed in the domesticated population
43 compared to the F1 one in terms of survival rate, hatching rate, percentage of eyed stage
44 larvae and percentage of malformations. In conclusion, this study shows that, in this example
45 of domestication process, despite a positive effect on the precocious gonadogenesis
46 advancement and vitellogenesis progress this domestication route negatively affected the
47 reproductive performance.

48 **Keywords:** *Perca fluviatilis*, Recirculating Aquatic System, GnRH, Gonadotropins, Sexual
49 steroids.

50

51

52

53

54

55 **1. Introduction**

56 Price (1984) defined the domestication as “the process whereby a population of animals
57 becomes adapted to humans and captivity conditions through one or several combinations of
58 genetic changes throughout the generations”. Digard (2003) proposed to define domestication
59 as “the action that men exercise over animals they own, even by raising them”. Thus the list
60 of domesticated fish species goes from the range of 25-30 in the first definition to more than
61 200 according to the definition of Digard (2003). This shows that the concept of
62 domestication still divides opinions within the scientific community. However, everyone
63 agrees that fish populations may undergo, through the domestication process, changes in their
64 genome due to adaptation to environmental stressors, genetic selections (deliberate or not) or
65 inbreeding phenomena (Balon 2004 ; Price 1984). It induces changes at anatomical,
66 physiological and molecular levels thereby affecting a variety of biological functions (e.g.
67 Brummet et al. 2004). There is abundant literature on the domestication effects on growth
68 (Withler and Beacham 1994; Hassin et al. 1997; Tymchuck et al. 2006) and stress response in
69 teleosts (Pickering and Pottinger 1989, 1997; Vandeputte and Prunet 2002; Douxfils et al.
70 2011). On the one side, it has been shown that globally domestication reduces fish stress
71 response, by rendering them less fearful as showed in salmonids (Pickering and Pottinger
72 1989,1997; Vandeputte and Prunet 2002), or less sensitive to acute handling stressors in
73 Eurasian perch (Douxfils et al. 2011). But in European sea bass (*Dicentrarchus labrax*) it has
74 been shown that heritability of cortisol response to stress was low (Volckaert et al. 2012). On
75 the other side, the domestication seems to boost the growth performance (Withler and
76 Beacham 1994; Hassin et al. 1997; Tymchuck et al. 2006).

77 Contrary to growth and stress response, the effects of domestication process on reproduction
78 have received limited attention. Moreover, it appears that this field of research has conducted
79 to conflicting results. In the ide (*Leuciscus idus*), Krejszeff et al. 2009 showed that
80 domestication positively influenced the ovulatory responses following the Ovopel™
81 hormonal treatment. In contrast, they showed that domestication exhibited a negative effect
82 on embryo survival. More recently, a study on the same species showed different results by
83 pointing out a lower ovulatory response to Ovopel™ in the pond-cultured group relatively to
84 wild fish (Ciesla et al. 2013) and no domestication effect was observed on the embryo
85 survival. Moreover, the effect of domestication can be different from one species to another.
86 In Atlantic salmon (*Salmo salar*) domestication led to an 83% decrease of the hatching rates
87 (Jonsson and Jonsson 2006). In cod (*Gadus morhua*), domestication negatively affected
88 fertilization rates, hatching and larvae survival (Salze et al. 2006). In brown trout (*Salmo*
89 *trutta*) domestication negatively affected sperm quality (Dannewitz et al. 2004) but had a
90 positive effect on spawning and egg weight (Randak et al. 2006). Consequently, notable
91 differences also exist between the fish families, the fish species and the fish populations.
92 These differences led us to investigate the domestication effects on reproductive ability of
93 Eurasian perch (*Perca fluviatilis*), a species whose domestication history is still recent, and
94 belonging to a fish family poorly investigated.

95 Eurasian perch is a freshwater fish species highly valuable in Europe (Fontaine, 2009). The
96 perch market is local and the current production of market-size perch in Europe is thus unable
97 to satisfy the demand, which is mainly centered in the Alpine region (Setälä et al. 2008;
98 Watson, 2008). Consequently to a relative decline of wild stocks due to intensive fisheries,
99 and to an increasing demand for small sized perch especially in Switzerland, intensive perch
100 culture has been developed in the past two decades, and Eurasian perch has become a relevant
101 species in European aquaculture (Fontaine 2004; Watson 2008). Up to now, only one study

102 has examined the effect of domestication on Eurasian perch reproduction (Kristan et al. 2012).
103 Results showed a positive effect of domestication on fertilization rates, but a negative effect
104 on ovulation and hatching. Actually, there is no detailed study aiming at understanding the
105 influence of domestication on both the mating performance and the reproductive mechanisms
106 underlying these effects. Our hypothesis is that disruption in the level of the endocrine actors
107 that control the reproductive cycle explains the changes of the reproductive performance
108 along the domestication process.

109 The aim of this study was to compare two perch populations (F1 and “domesticated”),
110 exposed to a special photothermal program allowing the control of gonadogenesis
111 advancement, in terms of morphological parameters (gonado-somatic index, hepato-somatic
112 index), levels of GnRH and gonadotropins gene expression, sexual steroids and vitellogenin,
113 as well as the oocyte developmental stages and reproductive performance.

114

115 **2. Materials and methods**

116

117 *2.1 Fish*

118 Two populations of F1 and “domesticated” perch were studied in this experiment, both
119 originally came from the Geneva Lake (French-Swiss border). F1 breeders came from wild
120 broodstock (F0) collected as eggs directly taken from Geneva Lake. According to the local
121 producer (Lucas Perches, Hampont, France), “domesticated” breeders are generated from
122 seven generations fish raised in captivity, even if the introduction of wild individuals during
123 reproduction seasons cannot be totally excluded. The domestication history of these fish is
124 thus a "black box". The F1 and domesticated breeders were issued from an artificial
125 reproduction completed on November 1st, 2011 and were then raised on the producer
126 facilities, before coming to the URAFPA-MAN facilities located in Nancy (France) on

127 February 6th, 2012, where they were placed in two 6000 L. tanks for growth. At their arrival
128 to our facilities, fish of both populations were 3-months-old. From their hatching, the two
129 studied fish populations (F1 and “domesticated”) have thus been raised in equal and
130 standardized husbandry conditions, at the Lucas perches facilities, fish were raised at a
131 temperature range of 18°C-20°C and a photoperiod of 13L/11D with a blue light or a red
132 light. At the URAFPA-MAN facilities fish were kept at 20°C and a photoperiod of 16L/8D
133 during the growth period. Fish were daily fed to apparent satiation during the growth period
134 and during the experiment (Le Gouessant pellets n°5, proteins 47%, lipids 13%, ash 6.90%
135 and fiber 2.2%). Fish were handled in accordance with national and international guidelines
136 for protection of animal welfare.

137

138 *2.2 Experimentation and water quality management*

139 Experiment started on May 17th, 2013, fish were 18 months old and never spawned before.

140 Breeders of both populations were transferred to 500 L. independent experimental tanks (3
141 tanks per population with 48 fish per each tank), each tank had his own filtration system and
142 25% of the water was changed every month. At the beginning of the experiment, fish weight
143 ranged between 67-137.8g for the “domesticated” fish and in the range of 70.4-129.8g for the
144 F1 ones. Both populations were exposed to a photothermal program enable to induce and
145 drive the gonadogenesis (Fontaine et al. 2015) (Fig. 1).

146 In each tank, water quality was measured three times per week. pH was maintained between
147 7.0 and 7.5 by NaCO₃ additions. The dissolved oxygen was maintained over of 6 mg/L. The
148 total ammonia (0,36 ± 0,53 mg/L) and nitrite (0,08 ± 0,12 mg/L) nitrogen were measured
149 using a CARY I spectrophotometer and remained always below 1 mg/L.

150 Fish were sampled at days T0, T35 (decrease of temperature), T157 (vernalization period) and
151 295 (increasing of temperature) after the start of the photothermal program (Fig. 1). Six

152 females per tank were caught at each sampling period. The first sampling occurred just before
153 the initial photoperiod decrease. At each sampling date, fish were anesthetized into a Tricaine
154 methanesulfonate (MS-222) bath (120 mg/L, Sigma). Blood was sampled from the caudal
155 vein using a syringe, then stored on ice in heparinized microtubes until centrifuged at 3000g
156 for 10 min (Centrifuge Jouan C-412). Aliquots of plasma were stored at -80°C until ELISA
157 analysis was performed. Then, each fish was euthanatized by overanesthetizing in Tricaine
158 methanesulfonate (MS-222) bath (240 mg/L), weighted and dissected for gonad and liver
159 weighting and calculation of the gonadosomatic index ($GSI=100\times \text{gonad weight}/\text{total fish}$
160 weight) and hepatosomatic index ($HSI= 100\times \text{liver weight}/\text{total fish weight}$). For gene
161 expression analysis, the whole brain and pituitary were snap frozen (separately) in liquid
162 nitrogen and stored at -80°C.

163

164 *2.3 Gonadal histology*

165 Samples of ovaries were stored in a Bouin-Holland solution for 1 week, washed once with
166 water, twice with 70 % ethanol and stored in absolute ethanol (Abdulfatah et al. 2010). Then,
167 fragments of gonads were cut into 6-mm-thick slices, dehydrated with OTTIX solution
168 (DIAPATH SpA, Italy/MM France) and embedded in a paraffin substitute dubbed Diawax
169 (DIAPATH SpA, Italy/MM France) (adapted from Langeron 1942). Sections of 6 µm were
170 cut from the Diawax block and stained with a Masson's trichrome-haematoxylin Gill III
171 (Merck, Darmstadt, Germany), 0.5 % phloxine B (VWR, California, USA) and 0.5 % light
172 green (Sigma, Saint Quentin Fallavier, France) according to Rinchar and Kestemont (1996).
173 The stage of oocyte development was determined according to Wallace and Selman (1981).
174 Observations have been completed with a light upright optical microscope (Nikon Eclipse Ni-
175 U), Nikon BR software (Nikon France, Champigny sur Marne, France).

176

177 *2.4 Total RNA extraction and reverse transcription*

178 Total RNA was extracted from the brain and pituitary using TRIzol reagent (VWR,
179 CTRIZ02) according to the manufacturer's protocol. Concentration of total RNA was
180 measured by UV absorption spectrometry using a Jasco V-630 Spectrophotometer. DNase
181 treatment (DNase I Thermo Fischer Scientific, EN0521) was performed according to
182 manufacturer's recommendations. Reverse transcription was performed with M-MLV reverse
183 transcriptase (Sigma, M1302) according to the manufacturer's recommendations. For each
184 sample 1µg of total RNA was used. Reaction mixes have been done following manufacturer's
185 recommendations. Each reverse transcription mix was incubated at 37°C for 50 min and then
186 heated at 90°C for 10 min, c-DNA samples were finally kept at -80°C.

187

188 *2.5 Gene expression analysis*

189 Primers were designed based on a multiple alignment of sequences available in the NCBI data
190 base, targeting conserved regions of the FSHb and LHb-subunits, the GnRH 3 as well as the
191 β-Actin as housekeeping gene.

192 In order to isolate and to sequence *P. fluviatilis* FSHb, LHb-subunits and the GnRH 3 cDNAs
193 from the brain, a PCR reaction was performed. PCR conditions were as follows: denaturing
194 cycle of 95°C for 2 min, followed by 35 cycles of 95°C for 1 min, 58°C for 1 min, 72°C for 1
195 min, followed by a final elongation step of 72°C for 5 min. PCR products were checked on
196 1% agarose gel electrophoresis. The single obtained band corresponding to the expected
197 product size was excised from the gel and the amplicon was purified with E.Z.N.A. Pure Kit
198 (VWR, D6493-01). Purified products were cloned employing the InsTAclone PCR Cloning
199 KIT (Thermo Fischer Scientific, K1214), and plasmids of 2 days cultures of bacterial colonies
200 (*Escherichia coli*). Plasmids were analyzed by PCR using M7 universal primers and agarose
201 gel to confirm the size of the inserted cDNA. Automated sequencing was performed

202 (ABI3730XL DNA sequencer, Macrogen, Korea). Upon gene identification, primers for
203 quantification were designed on two different exons with the FastPCR software V6.5. (Table
204 1).

205 Real-time PCR was carried out with a StepOne Plus q-PCR Cyclor (Applied Technologies)
206 using QUANTA SYBR green reagent (VWR, 733-1382) in a 20 μ L reaction volume (5 μ L
207 diluted cDNA, 2.5 μ L (600 nM) of each primer, 10 μ L of 2-fold diluted SYBR-Green
208 solution) under the following conditions: 95°C initial denaturation for 10 min, followed by 40
209 cycles of 95°C denaturation for 30s, 60°C primer annealing and extension for 1 min. All
210 PCRs were run in duplicate. Expression was determined by the comparative CT method
211 (DDCT) considering a calibrator sample (pooled brain or pituitary cDNA).

212 A melting curve analysis was carried out to ensure assay specificity. The absence of amplified
213 genomic DNA was validated by comparing the CT obtained in samples with and without
214 DNase treatments and by designing the primers on two different exons. Samples were
215 normalized to β -Actin mRNA.

216

217 *2.6 Hormone assays*

218 17 β -Estradiol (E2, ng/mL) was assayed on 50 μ L of plasma using the DIAsource E2-
219 ELISA kit (DIAsource, KAP0621). When necessary, dilution from 1/20 to 1/70 of the plasma
220 samples was performed. Sensitivity was 5 pg/ml. CV intra-assay varied between 3.1 and 2.6%
221 and CV inter-assays varied between 4.7 and 2.4% for low and high levels, respectively.

222 Testosterone (T, ng/mL) was assayed on 25 μ L of plasma using the DIAsource Testosterone
223 ELISA Kit (DIAsource, KAPD1559). When necessary, a dilution 1/2 of the plasma samples
224 was performed. Sensitivity was 83 pg/mL, CV intra-assay varied between 1.5 and 9.5 % and
225 CV inter-assays varied between 7.6 and 8.7% for low and high levels, respectively. The
226 androgen 11-ketotestosterone (11KT, ng/mL) was assayed on 50 μ L of plasma diluted 1/50 in

227 ELISA buffer using the Cayman 11-keto Testosterone EIA Kit (Cayman, 582751) and
228 sensitivity was 1.3 pg/mL. CV intra-assay varied between 2.6 and 8.4 % and CV inter-assays
229 varied between 9.4 and 12.1% for low and high levels, respectively. Plasma
230 17 α ,20 β -dihydroxy-4-pregnen-3-one (DHP, ng/mL) was assayed using
231 radioimmunoassay according to Fostier and Jalabert (1986) adapted by Mandiki et al. (2005),
232 following two extractions with cyclohexane/ethyl acetate (v/v). The DHP antibody was
233 provided by Dr A. Fostier (INRA, Rennes, France).
234 Cortisol was assayed on 20 μ L of plasma using the DIAsource Cortisol ELISA Kit
235 (DIAsource, KAPDB 270). The intra-assay coefficient of variation was 5.6%, the inter-assay
236 coefficient of variation was 6.9%. The detection limit was 2.5 ng/mL.

237

238 *2.7 Vitellogenin assay*

239 Vitellogenin was indirectly estimated with Alkali-labile phosphate levels (ALP, μ g/mL). ALP
240 levels were assayed on 80 μ L of plasma. Plasma samples were first centrifuged for
241 precipitation of high molecular weight proteins with trichloroacetic acid (20%). The pellet
242 was then washed with absolute ethanol, chloroform solution - di-ethyl ether - absolute ethanol
243 (1: 2: 2), acetone and finally the di-ethyl ether. The dried pellet was then dissolved in a NaOH
244 solution (2N) brought to 100°C for 15 min, and then the phosphate release was neutralized by
245 adding cold HCl (2N). Phosphates are then assayed by the method of Stanton (1968) using a
246 phosphorus standard solution for calibration, addition of molybdcic acid and reducer solutions
247 and finally by reading the absorbance at 660 nm with a spectrophotometer.

248

249 *2.8 Aromatase assay*

250 Aromatase activity was measured as the specific release of tritiated water produced by
251 conversion of [1 β -H³]-androstenedione to estrone, as previously described by Noakson et al.

252 (2001) with some modifications (Mandiki et al. 2005). Fifteen milligrams of frozen tissue
253 (gonads) were homogenized in a buffer solution (50 mM potassium phosphate, 250 mM
254 sucrose and 10 mM dithiothreitol pH 7,4; 1:10). A tenth of the homogenate were added of a
255 NADPH-substrate system consisting of 1 mM NADPH, 25 nM Androst-4-ene-3, 17-dione,
256 [1β - $^3\text{H}(\text{N})$] 275 nM Androst-4-ene-3, 17-dione in the buffer solution and incubated with
257 gentle shaking for 60 min at 30°C. The reaction was stopped by adding trichloroacetic acid.
258 Samples were then centrifuged and the supernatant extracted with chloroform, the bottom
259 contained the protein. The aqueous phase was washed with dextran (5%)-activated charcoal
260 (0.5%), centrifuged, and an aliquot of the supernatant was added to a liquid scintillating to
261 determine the radioactivity due to the presence of [^3H] H_2O . For this purpose a liquid
262 scintillation counter was used. The Lowry method was used to determine protein
263 concentration after denaturing the protein with NaOH. The aromatase activity was expressed
264 in $\text{fmol min}^{-1} \text{mg}^{-1}$ protein.

265

266 *2.9 Spawning season and reproductive performance*

267 Spawning season started on March 25th, 2014 and finished on April 3th, 2014. 77% of the
268 females spawned during the spawning season in the F1 population and 78% of the females
269 spawned in the “domesticated” one. Spawning was induced by temperature increase at the end
270 of the photothermal programme and no hormonal treatment was used during the experiment.
271 Males and females were kept together in the same tanks. Females were stripped every day
272 during the spawning season at 5:00 AM. When a female was ovulating, ribbons were
273 collected in a dry basin and fertilized with 1mL of sperm originated from three different
274 males, in order to optimize the fertilization success. Ribbons were then transferred into
275 incubators and maintained at constant temperature ($13 \pm 0,5^\circ\text{C}$). After 24h, 48h, 72h and 120h
276 post fertilization, samples of approximately 200 eggs from each ribbon were observed to

277 calculate survival rates. Hatching started at 8 days post fertilization. Larvae were measured
278 and analyzed for malformations (sclerosis, lordosis, yolk bag malformations) with a binocular
279 microscope, on fixed embryos (n=6 spawns).

280

281 *2.10 Statistical analysis*

282 Data are expressed as mean \pm standard deviation with individual fish as the experimental unit.
283 Normality of data and homogeneity of variances were tested using Shapiro–Wilk and Levene
284 test, and data were either log-transformed or arcsin square-root transformed (for data
285 expressed in percentage) when needed. When data respected the assumptions of normality, a
286 two-way analysis of variance (ANOVA) with two independent factors (population and time)
287 was performed followed by LSD post hoc tests to identify significant differences. The level of
288 significance used in all tests was $p < 0.05$. When data were not normally distributed, a
289 Kruskal–Wallis test was performed followed by the nonparametric Mann–Whitney test. All
290 statistical analyses were performed using the STATISTICA software (StatSoft, Tulsa, USA).

291

292 **3. Results**

293

294 *3.1. GSI and gonadal development stages*

295 During the first month of the experiment (T0 to T35), females of both populations had a low
296 GSI (below 1%) (Fig. 2). Significant increase of the GSI was observed after five months
297 (T157) ($P < 0.05$), no significant difference was observed between the two populations until
298 T157. At T295, the “domesticated” females showed a significantly higher GSI ($p < 0.05$) in
299 comparison with the F1 ones (Fig. 2, $P < 0.05$). Histological analysis revealed that during the
300 first five weeks of the photothermal program, both female populations showed immature
301 gonads with more than 90% of protoplasmic oocytes. Since T157 both populations showed

302 more advanced gonad maturation and no significant difference between the two populations
303 was pointed out (Fig. 3). However, the protoplasmic oocyte diameter was significantly more
304 elevated for the “domesticated” females ($109\pm 19\mu\text{m}$) than for the F1 ones ($104\pm 11\mu\text{m}$) at T35
305 ($p<0.05$). The same observation was noticed at T295 with ($152\pm 13\mu\text{m}$) for “domesticated”
306 females and ($138\pm 12\mu\text{m}$) for the F1 ones ($p<0.05$). At T295, “domesticated” females showed
307 a more advanced gonadal development than the F1 ones with respectively 54% and 39% of
308 late vitellogenesis oocytes (Fig. 3), even if there was no significant difference in terms of the
309 oocyte diameter for vitellogenic oocytes.

310

311 *3.2. Sexual steroids, vitellogenin and aromatase*

312 Values for plasma 17β -estradiol (E2) were the lowest levels at T0 and T35 for both female
313 populations (less than 0.5 ng/mL) (Fig. 4A), with a significantly higher level ($p<0.05$) for
314 domesticated females at T35 in comparison with F1 ones. At T157, plasma E2 levels
315 significantly increased for both populations in comparison with T35 levels. At T295, a drop of
316 E2 levels was noticed in the F1 population, so E2 values were significantly higher ($p<0.05$) in
317 domesticated females than in F1 ones.

318 Plasma Testosterone (T) levels started to increase since T157 until the end of the experiment
319 for both populations (Fig. 4B). Significant difference was observed at T157 when plasma T
320 levels were significantly higher for F1 females in comparison with the “domesticated” fish
321 ($p<0.05$). Plasma vitellogenin levels followed a similar pattern of change as those observed
322 for T (Fig 4C), with a significant difference ($p<0.05$) at T157 in favor of F1 females. 11-
323 Ketotestosterone (11-KT) levels remained low for both female populations until T295, when
324 11-KT levels significantly increased but no significant difference between the two groups was
325 observed (Fig. 4D).

326 Plasma DHP levels showed an increasing pattern along the experimental time-course for both
327 populations (Fig. 5A) and no significant difference was observed between the two groups. In
328 the same way, the aromatase showed an increasing activity, but no significant difference was
329 registered between the two groups (Fig. 5B).

330

331 *3.3. GnRH and GtHs gene expressions*

332 GnRH mRNA levels did not show any significant difference between the two populations
333 (Fig. 6A, B), but showed a two-fold increase at T35 and remained at this level until T157.

334 β FSH mRNA levels showed a quick increase between T0 and T35 for both populations (Fig.
335 6C, D), and were significantly higher in the F1 perch brains in comparison with their
336 “domesticated” counterparts at T157 and decreased at the last sampling time. In both brain
337 and pituitary tissues (Fig. 6E, F), β LH mRNA levels increased coincidentally with advancing
338 ovarian development. Significant differences in favor of the F1 population were observed at
339 T35 and T295 in the brain ($p < 0.05$) (Fig. 6E) and pituitary ($p < 0.005$) (Fig. 6F).

340

341 *3.4. Plasma cortisol*

342 Plasma cortisol levels showed a significant difference between the two groups with higher
343 values found in the F1 population ($p < 0.05$) at T35 than the “domesticated” fish (Fig. 7). But,
344 the difference between the two groups was not significant at the other sampling times.

345

346 *3.5. Reproductive performance*

347 The development rates at 24h, 48h, 72h and 120h post-fertilization were evaluated for the two
348 populations. The F1 population showed a better survival rate than the “domesticated” one at
349 the four observation times (Table 4) ($p < 0.05$). The F1 population showed a higher eyed stage
350 larvae percentage and a higher hatching rate than the “domesticated” population. F1

351 population larvae were of better quality than their “domesticated” counterparts in regard of
352 percentage of deformities ($p < 0.05$).

353

354 **4. Discussion**

355

356 In the present study, changes in gonadosomatic index in the population of “domesticated” fish
357 corresponds to the profile described by Migaud et al. (2002), and the maximum level of GSI
358 reached at the last sampling time ($18.5 \pm 3.2\%$), is consistent with the level obtained in
359 Eurasian perch females by Kristan et al. (2012). These GSI values were significantly higher
360 than those registered in F1 fish ($15.8 \pm 3.2\%$), being themselves lower than those recorded by
361 Migaud et al. (2003) and Kristan et al. (2012) in wild Eurasian perch populations (F0). In
362 agreement with the above GSI results, “domesticated” females showed a slightly more
363 advanced gonadal development than the F1 ones with respectively 54% and 39% of late
364 vitellogenesis oocytes at the end of the photothermal program, and a more elevated
365 protoplasmic oocyte diameter than the F1 ones. This further supports the more advanced
366 gonadogenesis in the “domesticated” fish by comparison to their F1 counterparts. These
367 results do not corroborate previous observations made by Kristan et al. (2012) in Eurasian
368 perch females. Differences between farmed and wild perch have been suggested to be caused
369 mainly by unsuitable feed for the farmed fish (Bell et al. 1997; Kestemont et al. 1999;
370 Izquierdo et al. 2001; Kestemont et al. 2008; Henrotte et al. 2010). These authors stated that,
371 to address lower reproduction in farmed perch, it is necessary to improve the diet by adding
372 prey fish and a larger proportion of natural ingredients to the feed. In the experiment of
373 Kristan et al. (2012) wild fish were fed with prey fish and farmed ones with both industrial
374 food and prey fish, while in our study, both populations were fed only by industrial food. This
375 could be one element explaining the discrepancy between the two studies.

376

377 Moreover, we cannot rule out the implication of other external factors (e.g. type of
378 photothermal program) and internal factors (e.g fish age) in the monitoring of divergent data.
379 Our wild population is a F1 generation issued from F0 broodstock caught as eggs from the
380 wild (Geneva Lake) and reared in artificial conditions during their whole life. The wild
381 population studied by Kristan (2012) is a wild population reared in ponds and transferred in
382 artificial conditions for the experiment, which implies acclimation phenomena due to the
383 transfer to artificial conditions, which may in turns alter gonad maturation and reproduction.
384 In addition, we cannot exclude the influence of internal factors like fish age that has been
385 shown to influence reproductive capacity in teleosts (Abi-Ayad, 1998). Our fish were
386 eighteen months old for both populations, whereas fish studied by Kristan (2012) were three
387 years old. The geographical origins, and by consequence, potentially the genetic patrimony, is
388 another internal factor that can impact fish reproduction and justify divergent results.

389

390 While the gonadogenesis was more advanced in the “domesticated” population, the final
391 reproductive outcomes were markedly better in the F1 fish. The fertilization rate monitored at
392 24h was almost 3-fold higher in the F1 fish ($78\pm 35\%$) in comparison with the “domesticated”
393 group ($28\pm 29\%$). The fecundity rates observed by Kristan (2012) were more elevated (90%)
394 in both populations. Kouril and Hamackova (1999) reported 60-95% fertilization of eggs in
395 wild perch after artificial stripping which is in agreement with our values for the F1
396 population. This difference of fertilization rates is an additional indication that the egg quality
397 in Eurasian perch is highly variable between the studies and also within the same population.
398 The highest contrast between farmed and F1 perch in the parameters of egg quality was
399 recorded after the evaluation of hatching rates. A really poor percentage of hatching rate was
400 obtained for the farmed fish ($2\pm 2\%$), slightly lower than the data provided by Kristan et al.

401 (2012) in other farmed population. This poor ova quality could be explained by the fact that
402 “domesticated” fish exhibited a faster gametogenesis than the F1 ones and they did not
403 accumulate enough nutrients for embryo development. This better egg quality in F1 fish was
404 also confirmed by comparing the percentages of malformations between both populations.
405 Anyway these results and those of Kristan et al. (2012) suggest that “domesticated”
406 broodstock in Eurasian perch exhibit lower reproductive performance than fish at an earlier
407 stage of domestication.

408

409 In the present study, the better reproductive performance in F1 are *a priori* surprising
410 regarding the gonadogenesis and oocyte development results. To understand the physiological
411 mechanisms that are behind these differences, we investigated some physiological actors
412 driving the control of oogenesis control. The estrogen 17- β estradiol (E2) levels were higher
413 in the “domesticated” fish than in the F1 fish, which is in agreement with the GSI results.
414 Contrarily, testosterone levels were higher in F1 females than in “domesticated” ones. We
415 tried to understand these opposite results by measuring aromatase activity (enzyme which
416 converts testosterone to 17-beta estradiol). No significant difference in aromatase activity was
417 found between the two populations while we expected a decline in aromatase activity for the
418 F1 population. Our observation cannot explain this difference in sexual steroids levels
419 between the two populations. However, E2 being the principal mediator of vitellogenin
420 production and incorporation in the oocytes (Lubzens et al. 2010), these more elevated E2
421 levels in domesticated fish should have been accompanied with higher levels of vitellogenin
422 in the plasma. However opposite results were found. These conflicting observations might be
423 explained by three simultaneous phenomena: firstly Eurasian perch displays strongly elevated
424 testosterone levels by comparison to other fish species and these testosterone levels are 20-50-
425 fold higher than the E2 level (Milla et al. 2009a). Secondly, like E2, testosterone is a

426 stimulator of vitellogenin production in fish (Ramezani-Fard et al. 2013). Thirdly, more
427 elevated testosterone levels were measured in the F1 perch than in the domesticated ones.
428 All together, we speculate that the higher vitellogenin level measured in F1 fish would be
429 more explained by the differential testosterone level between both groups rather than the
430 classically described E2 involvement. Concerning 11 keto-testosterone, given the low level
431 observed in comparison with the testosterone level and the absence of contrasted
432 concentrations between both groups, we consider that this sexual steroid was not a major actor
433 explaining this divergence of oogenesis progress between F1 and domesticated fish.
434 Moreover, to explain the divergence in oogenesis and reproductive performance between the
435 two populations, we investigated the plasma $17\alpha,20\beta$ -dihydroxy-4-pregnen-3-one (DHP),
436 which is the maturation inducing steroid (MIS) in percids (Nagahama et al. 1987). Our
437 hypothesis was that, a difference in kinetics of the final oocyte maturation between the two
438 populations would be at the origin of the reproductive performance divergence. But plasma
439 DHP levels which shows a classical kinetics of change did not display different levels
440 between the two female groups. However it has been reported that another MIS is associated
441 with the final oocyte maturation control in some perciform species: it is the $17\alpha,20\beta,21$ -
442 trihydroxy-4-pregnen-3-one (20β -S) (Thomas 1988; Asahina et al. 1991 ; Modesto and
443 Canario 1993; King et al. 1994), we can imagine that in perch, final oocyte maturation could
444 be controlled by the combined action of DHP and 20β -S, but we cannot exclude the
445 hypothesis that the DHP could be the principal MIS in perch but it has to be sampled more
446 frequently during the final oocyte maturation phase. All combined, our hypothesis is that this
447 domestication process displayed distinct effects according to the time-course of oogenesis.
448 Indeed, farmed fish exhibited an earlier gonadogenesis and vitellogenesis than F1 ones, then
449 turning to depreciative effects on final oogenesis, oocyte quality, finally traduced by poor
450 reproductive performance.

451

452 Several studies have reported the effect of domestication on reproduction efficiency in many
453 species (Dannewitz et al. 2004; Salze et al. 2005; Randak et al. 2006; Jonsson and Jonsson
454 2006; Krejszeff et al. 2009; Kristan et al. 2012; Ciesla et al. 2013). These data suggest that
455 physiological changes occur during the domestication process due to farming conditions.
456 However, these studies do not provide information about the endocrine mechanisms in
457 broodstock explaining these changes in reproductive capacity. As the level of sex steroids is
458 modified in our study, we hypothesized that a disruption of the gonadotropic axis occurred
459 during the domestication process. Significant differences between the two populations were
460 monitored in the level of gonadotropin gene expressions in the brain and pituitary, while
461 GnRH gene expression was not affected. However, we only investigated the GnRH3 form in
462 our experiment, but there is another form involved during the reproduction in fish which is the
463 GnRH1 (Yaron et al. 2003) and which could be affected as well during the domestication
464 process. These results support the hypothesis of a gonadotropic axis disruption, starting at the
465 gonadotropic level affecting in turns the secretion of sexual steroids in the gonad. As the level
466 of FSH and LH transcripts were higher in F1 fish, in particular at the two last sampling times,
467 we think that the more markedly rise in testosterone level in this population is due to LH rise,
468 then leading to a better reproductive performance in F1 fish. Indeed, blood plasma
469 testosterone is regulated after exposure to the LH agonist hCG in perch females at the end of
470 the reproductive cycle (Milla et al. 2009a). However, we cannot rule out the fact that the
471 measures of FSH and LH transcript levels might not reflect the circulating levels of GTHs.
472 Moreover, the hormonal differences observed between the two populations might reflect a
473 differential threshold of action of these hormones between the two populations.
474 Kristan (2012) proposed that the stress state of fish would explain the differences in the
475 reproductive performance between the two populations. Even if there was not any problem in

476 terms of water quality, infection or mortality, the level of cortisol were not low during this
477 experiment and the level was higher in F1 fish before the alveoli cortical stage of the cycle.
478 Consequently, in accordance with the proposal of Kristan, we cannot rule out the fact that the
479 differential cortisol levels or sensitivity to stressful conditions led to differential advancement
480 of early gonadogenesis between the two groups of fish, stress and cortisol being major
481 modulators of fish reproduction (Barton 2002; Milla et al. 2009b; Schreck 2010). But, this
482 precocious difference alone would not be sufficient to explain the major differences in the
483 reproductive performance, as fish were probably in the same level of stress state since the
484 vitellogenesis stage.

485 In conclusion, this study showed that this domestication route negatively affected the
486 reproductive performance. However, some conflicting results were observed about the
487 reproductive physiological parameters: on the one hand, this domestication process advanced
488 the start of gonadogenesis until the start of the vitellogenesis period, probably caused by more
489 precocious E2 rise. Then, the lower level of gonadotropins transcripts and testosterone
490 potentially led to a drop in vitellogenin in comparison to F1 fish, further affecting the
491 reproductive success. This study showed that this domestication process negatively affected
492 reproduction, so more attention should be paid by the farmers to the choice of renewing
493 broodstock in particular when they maintain fish coming from several generations of
494 domestication in their facilities,

495

496 **Funding.** This study was supported by the FranceAgrimer establishment (French
497 establishment of national agricultural and marine products) contract number: 2013-1566, and
498 partly supported by projects CENAKVA (No. CZ.1.05/2.1.00/01.0024) and CENAKVA II
499 (No. LO1205 under the NPU I program).

500

501 **Authors contributions.** Amine Khendek was responsible for the carrying out data analyses
502 and wrote the main part of the manuscript, and performed the gene expression analysis.
503 Sylvain Milla, Pascal Fontaine and Tomas Policar initiated the research and/or managed the
504 work and gave final approval for submission. Sandrine Viot and Camille Rousseau
505 performed the plasma steroid assays and histological analysis, Maud Alix managed the
506 spawning season and compared the reproductive performance of both populations, Yannick
507 Ledoré took care of fish daily and managed the samplings; SNM Mandiki and Patrick
508 Kestemont performed the enzymatic activity assay and RIA assay for aromatase activity and
509 DHP. All authors read and approved the manuscript.

510

511 **Acknowledgments.** The authors are grateful to the INRA institute of Champenoux, France for
512 access to histological equipment and their help, and to PB2P team (University of Lorraine,
513 France) for access to the bacterial culture equipment for cloning GnRH and GtHs genes. We
514 would like to thank Vincent Joussemet for his help during the spawning season (University of
515 Lorraine, France) and all the members of team DAC (University of Lorraine, France) who
516 helped during fish sampling.

517

518

519

520

521 **References**

- 522 Abdulfatah, A., Fontaine, P., Kestemont, P., Gardeur, J.N., Marie, M., 2010. Effects of photothermal kinetics
523 and amplitude of photoperiod decrease on the induction of the reproduction cycle in female Eurasian
524 perch (*Perca fluviatilis*). *Aquacult.* 322, 169 - 176.
- 525 Abi-Ayad, S.M., 1998. Etude expérimentale de la biologie de la reproduction de la perche fluviatile (*Perca*
526 *fluviatilis*). Effet de la composition en acides gras de la série (n-3) de l'alimentation des géniteurs sur la
527 qualité des oeufs et des larves. PhD thesis Liège, Belgium.
- 528 Asahina, K., Zhu, Y., Aida, K., Higashi, T., 1991. Synthesis of 17 α ,21-dihydroxy-4-pregnen-3,20-dione,
529 17 α ,20,21-dihydroxy-4-pregnen-3-one, and 17 α ,20,21-trihydroxy-4-pregnen-3-one in the ovaries of

530 tobi-numeri dragonet, *Repomucenus beniteguri*, *Callionymidae Teleostei*. In: Scott A.P., Sumpter J.P.,
531 Kime D.E. and Rolfe M.S. Reproductive Physiology of Fish, Sheffield, pp. 80-82.

532 Balon, E.K., 2004. About the oldest domesticates among fishes. *J. Fish Biol.* 65, 1 - 27.

533 Barton, B.A., 2002. Stress in fishes: a diversity of responses with particular reference to changes in circulating
534 corticosteroids. *Integ. And Comp. Biol.* 42, 517 - 525.

535 Bell, J.G., Farndale, B.M., Bruce, M.P., Navas, J.M., Carillo, M., 1997. Effect of broodstock dietary lipid on
536 fatty acid compositions of eggs from sea bass (*Dicentrarchus labrax*). *Aquacult.* 149, 107 - 119.

537 Brummett, R., Angoni, D., Pouomogne, V., 2004. On-farm and on-station comparison of wild and domesticated
538 Cameroonian populations of *Oreochromis niloticus*. *Aquacult.* 242, 157 - 164.

539 Ciesla, M., Jonczyk, R., Gozdowski, D., Sliwinski, J., Rechulicz, J., Andrzejewski, W., 2013. Changes in ide
540 (*Leuciscus idus*) female's reproductive parameters after stimulation with carp pituitary homogenate
541 (CPH) and Ovopel: the effect of domestication?. *Aquacult. Int.* 22, 77 - 88.

542 Dannewitz, J., Petersson, E., Dahl, J., Prestegaard, T., Lof, A., Jarvi, T., 2004. Reproductive success of hatchery-
543 produces and wild-born brown trout in an experimental stream. *J. Appl. Ecol.* 41, 355 - 364.

544 Digard, J.P., 2003. La domestication animale revisitée par l'anthropologie. *Bulletin de la Soci. Zool. de France.*
545 128, 283 - 296.

546 Douxfils, J., Mathieu, C., Mandiki, S.N.M., Milla, S., Henrotte, E., Wang, N., Vandecan, M., Dieu, M., Dauchot,
547 N., Pigneur, L.M., Li, X., Rougeot, C., Melard, C., Silvestre, F., Van Doninck, K., Raes, M., Kestemont,
548 P., 2011. Physiological and proteomic evidences that domestication process differentially modulates the
549 immune status of juvenile Eurasian perch (*Perca fluviatilis*) under chronic confinement stress. *Fish*
550 *Shellfish Immun.* 31, 1113 - 1121.

551 Fontaine, P., 2004. Eurasian perch culture, a way of diversification for freshwater aquaculture. *J. Anim. Prod.*
552 17, 189 - 193.

553 Fontaine, P., 2009. Development of European inland fish culture and domestication of new species. *Cah. Agric.*
554 18, 144 - 147.

555 Fontaine, P., Wang, N., Hermelink, B., 2015. Broodstock management and control of the reproductive cycle. In:
556 Kestemont P., Dabrowski K., Summerfelt F.C. (Eds.), *Biology and culture of percid fishes*. Springer,
557 Dordrecht, pp. 103-122.

558 Fostier, A., Jalabert, B., 1986. Stereoidogenesis in rainbow trout (*Salmo gairdneri*) at various preovulatory
559 stages: changes in plasma hormone level and in vivo responses of the ovary to salmon gonadotropin. *Fish*
560 *Physiol. Biochem.*, 2, 87 - 99.

561 Hassin S., de Monbrison, D., Hanin Y., Elizur A., Zohar Y., Popper D.M., 1997. Domestication of the white
562 grouper *Epinephelus aeneus*, 1. Growth and reproduction. *Aquacult.* 156, 305 - 316.

563 Henrotte, E., Mandiki, R.S.N.M., Prudencio, A.T., Vandecan, M., Méléard, C., Kestemont, P., 2010. Egg and
564 larval quality, and egg fatty acid composition of Eurasian perch breeders (*Perca fluviatilis*) fed different
565 dietary DHA/EPA/AA ratios. *Aquacult. Res.* 41, 53 - 61.

566 Izquierdo, M.S., Fernández-Palacios, H., Tacon, A.G.J., 2001. Effect of broodstock nutrition on reproductive
567 performance of fish. *Aquacult.* 197, 25 - 42.

568 Jonsson, B., Jonsson, N., 2006. Cultured Atlantic salmon in nature: a review of their ecology and interaction
569 with wild fish. *ICES J. Mar. Sci.* 63, 1162 - 1181.

570 Kestemont, P., Cooremans, J., Abi-Ayad, S.M., Mélard, C., 1999. Cathepsin L in eggs and larvae of perch *Perca*
571 *fluviatilis*: variations with developmental stage and spawning period. *Fish Physiol. Biochem.* 21, 59 - 64.

572 Kestemont, P., Henrotte, E., Wang, N., Hamza, N., Paulsen, H., Overton, J., 2008. Feeding and nutrition of
573 European percid broodstock and early life stages. In: Fontaine P., Kestemont P., Teletchea F. and Wang
574 N. (Eds.), *Proceeding of abstracts and short communications of the workshop Percid Fish Culture From*
575 *Research to Production*. Namur, pp. 28-34.

576 King, V.W., Thomas, P., Sullivan, C.V., 1994. Hormonal regulation of final maturation of striped bass oocytes
577 *in vitro*. *Gen. Comp. Endocrinol.* 96, 223 - 233.

578 Kouril, J., and Hamackova, J., 1999. Artificial of perch propagation of European perch (*Perca fluviatilis*) by
579 means of a GnRH analogue. *Czech J. Anim. Sci.* 44, 309 - 316.

580 Krejszef, S., Targońska, K., Źarski, D., Kucharczyk, D. 2009. Domestication affects spawning of the ide
581 (*Leuciscus idus*) preliminary study. *Aquacult.* 295, 145 - 147.

582 Kristan, J., Stejskal, V., Policar, T., 2012. Comparison of reproduction characteristics and broodstock mortality
583 in farmed and wild Eurasian Perch (*Perca fluviatilis* L.) females during spawning season under controlled
584 conditions. *Turkish J. Fish. Aquat. Sci.* 12, 191 - 197.

585 Langeron, M., 1942. *Précis de Microscopie: Technique-Expérimentation-Diagnostic*. Masson et Cie, Paris. 1339
586 pp (in French).

587 Lubzens, E., Cerda, J., Young, G., Bobe, J., 2010. Oogenesis in teleost fish: how fish eggs are formed. *Gen.*
588 *Comp. Endocrinol.* 165, 367 - 389.

589 Mandiki, S.N.M., Babiak, I., Bopopi, J.M., Leprieur, F., Kestemont, P., 2005. Effects of sex steroids and their
590 inhibitors on endocrine parameters and gender growth differences in Eurasian perch (*Perca fluviatilis*)
591 juveniles. *Steroids.* 70, 85 - 94.

592 Migaud, H., Fontaine, P., Sulisty, I., Kestemont, P., Gardeur, J.N., 2002. Induction of out-of-season spawning
593 in Eurasian perch *Perca fluviatilis*: effects of rates of cooling and cooling durations on female
594 gametogenesis and spawning. *Aquacult.* 205, 253 - 267.

595 Migaud, H., Mandiki, R., Gardeur, J.N., Kestemont, P., Fontaine, P., 2003. Synthesis of sex steroids in final
596 oocyte maturation and induced ovulation in female Eurasian perch, (*Perca fluviatilis*). *Aquat. Living*
597 *Resour.* 16, 380 - 388.

598 Milla, S., Mandiki, R., Hubermont, P., Rougeot, C., Melard, C., Kestemont, P., 2009a. Ovarian steroidogenesis
599 inhibition by constant photothermal conditions is caused by a lack of gonadotropin stimulation in
600 Eurasian perch. *Gen. Comp. Endocrinol.* 163, 242 - 250.

601 Milla, S., Wang, N., Mandiki, R., Kestemont, P., 2009b. Corticosteroids: friends or foes of teleost fish
602 reproduction? *Comp. Biochem. Physiol. A.* 153, 242 - 251.

603 Modesto, T., Canario, A.V.M., 1993. The sex steroids of toadfish, (*Halobatrachus didactylus*). In *Abstr. 12th*
604 *Int. Congr. Comp. Endocrinol.* Toronto.

605 Nagahama, Y., 1987. Endocrine control of oocyte maturation. In *Hormones and Reproduction in Fishes,*
606 *Amphibians, and Reptiles*. Edited by D.O. Norris and R.E. Jones. pp. 171- 201. Plenum, New York, USA.

607 Noaksson, E., Tjarnlund, U., Bosveld, A.T.C., Balk, L., 2001. Evidence for endocrine disruption in Perch (*Perca*
608 *fluviatilis*) and Roach (*Rutilus rutilus*) in a remote Swedish lake in the vicinity of a public refuse dump
609 *Toxicol. And Appl. Pharmacol.* 174, 160 - 176.

610 Pickering, A.D., Pottinger, T.G., 1989. Stress responses and disease resistance in salmonid fish: effects of
611 chronic elevation of plasma cortisol. *Fish Physiol. Biochem.* 7, 253 - 258.

612 Pickering, A.D., Pottinger, T.G., 1997. Genetic basis to the stress response: selective breeding for stress-tolerant
613 fish. In: *Fish stress and health in aquaculture*, G.K. Iwama, A.D. Pickering, J.P. Sumpter, C.B. Schreck
614 (eds), 171-193. Cambridge University Press, Cambridge, UK.

615 Price, E.O. 1984. Behavioral aspects of animal domestication. *Q. Rev. Biol.* 59, 1 - 32.

616 Ramezani-Fard, E., Kamarudin, M.S., Harmin, S.A., 2013. Endocrine control of oogenesis in teleosts. *Asian J.*
617 *Anim. Vet. Adv.* 8, 205 - 215.

618 Randák, T., Kocour, M., Žlábek, V., Policar, T., Jarkovský, J., 2005. Effect of culture conditions on reproductive
619 traits of brown trout *Salmo trutta* l. *Bull. Fr. Pêche Piscic.* 383, 1-12.

620 Rinchar, J., Kestemont, P., 1996. Comparative study of reproductive biology in single and multiple spawner
621 cyprinid fish: I. Morphological and histological features. *J. Fish Biol.* 49, 883 - 894.

622 Salze, G., Tocher, D.R., Roy, W.J., Robertson, D.A., 2005. Egg quality determinants in cod (*Gadus morhua* L.):
623 egg performance and lipids in eggs from farmed and wild broodstock. *Aquacul. Res.* 46, 1488 - 1499.

624 Schreck, C.B., 2010. Stress and fish reproduction: The roles of allostasis and hormesis. *Gen. Comp. Endocrinol.*
625 165, 549 - 556.

626 Setälä, J., Laitinen, J., Virtanen, J., Saarni, K., Nielsen, M., Honkanen, A., 2008. Spatial integration of
627 freshwater fish markets in the Northern Baltic Sea area. *Fish. Res.* 92, 196 - 206.

628 Stanton, M.G., 1968. Colorimetric determination of inorganic phosphate in the presence of biological material
629 and adenosine triphosphate. *J. Anal. Biochem.* 22, 27 - 34.

630 Thomas, P., 1988. Changes in the plasma levels of maturation-inducing steroids in several perciform fishes
631 during induced ovulation. *Am. Zool.* 28, 53 - 63.

632 Tymchuk, W.E., Biagi, C., Withler, R., Devlin, R.H., 2006. Growth and behavioral consequences of
633 introgression of a domesticated aquaculture genotype into a native strain of Coho Salmon. *T. Am. Fish.*
634 *Soc.* 153, 442 - 455.

635 Vandeputte, M., Prunet, P., 2002. Génétique et adaptation chez les poissons : domestication, résistance au stress
636 et adaptation aux conditions de milieu. *INRA Prod. Anim.* 15, 365 - 371.

637 Volckaert, F.A.M., Hellems, B., Batargias, C., Louro, B., Massault, C., Van Houdt, J. K.J, Haley, C., de
638 Koning, D.J. Canario, A.V.M., 2012. Heritability of cortisol response to confinement stress in
639 European sea bass *dicentrarchus labrax*. *Genet. Select. Evol.* 44, 15.

640 Wallace, R.A., Selman, K., 1981. Cellular dynamic aspects of oocyte growth in teleosts. *Am. Zool.* 21, 325 -
641 343.

642 Watson, L., 2008. The European market for perch (*Perca fluviatilis*). In: P. Fontaine, P. Kestemont, F. Teletchea
643 and N. Wang (Eds.), *Proceeding of abstracts and short communications of the workshop. Percid Fish*
644 *Culture from Research to Production*, Namur, Belgium, 10 - 14.

645 Withler, R.E., Beacham, T.D., 1994. Genetic variation in body weight and flesh colour of coho salmon
646 (*Oncorhynchus kisutch*) in British Columbia. *Aquacult.* 119, 135 - 148.

647 Yaron, Z., Gur, G., Melamed, P., Rosenfeld, H., Elizur, A., Levavi-Sivan, B., 2003. Regulation of fish
648 gonadotropins. *Int. Rev. Cytol.* 225, 131 - 185.

649

650 **Legends**

651

652 **Fig. 1. The photothermal program (adapted from Fontaine et al. 2015).** The photothermal
653 program was used to induce perch reproduction. The dark curve corresponds to the
654 photoperiod variations. The light curve corresponds to the temperature variations. T0, T35,
655 T157, T295 are the four sampling times which represent days 0, 35, 157 and 295 after the
656 start of the experiment.

657

658 **Fig. 2. GSI changes (%) in F1 and domesticated females.** (9 females per sampling period +
659 Standard deviation). Sampling times T0, T35, T157 and T295 correspond respectively to days
660 0, 35, 157 and 295 after the beginning of the photothermal program. The asterisk represents a
661 significant difference between the 2 fish populations ($p < 0.05$).

662

663 **Fig. 3. Different developmental stages observed in domesticated and F1 perch female**
664 **ovaries.** (9 females per sampling time + Standard deviation). D and F1 refer to Domesticated
665 and F1 respectively. Sampling periods T0, T35, T157 and T295 correspond respectively to
666 days 0, 35, 157 and 295 after the beginning of the photothermal program.

667

668 **Fig. 4. Plasma sexual steroids and vitellogenin levels variation. (A) E2 levels variation**
669 **(ng/mL) in F1 and domesticated females. (B) Testosterone levels variation (ng/mL) in**
670 **F1 and domesticated females. (C) Vitellogenin levels variation ($\mu\text{g ALP/mL}$) in F1 and**
671 **domesticated females. (D) 11KT variations (pg/mL) in F1 and domesticated females.** (9
672 females per time per population + standard deviation). Sampling periods T0, T35, T157 and
673 T295 correspond respectively to days 0, 35, 157 and 295 after the beginning of the
674 photothermal program. The asterisk (*) represents a significant difference ($p < 0.05$) and (**)
675 ($p < 0.001$).

676

677 **Fig.5. Plasma DHP and aromatase activity. (A) Plasma DHP levels (ng/mL) variation in**
678 **domesticated and F1 perch females. (B) Aromatase activity variation (fmol/min.mg) in**
679 **domesticated and F1 perch females.** (9 females per time per population + standard
680 deviation). Sampling periods T0, T35, T157 and T295 correspond respectively to days 0, 35,
681 157 and 295 after the beginning of the photothermal program.

682

683 **Fig. 6. GnRH 3 and gonadotropins gene expression quantification in the brain and**
684 **pituitary. (A) GnRH 3 mRNA levels in F1 and domesticated female brains. (B) GnRH 3**
685 **mRNA levels in F1 and domesticated female pituitaries. (C) βFSH mRNA levels in F1**
686 **and domesticated female brains. (D) βFSH mRNA levels in F1 and domesticated female**
687 **pituitaries. (E) βLH mRNA levels in F1 and domesticated female brains.. (F) βLH**
688 **mRNA levels in F1 and domesticated female pituitaries.** (9 females per time per population
689 + standard deviation). Sampling periods T0, T35, T157 and T295 correspond respectively to
690 days 0, 35, 157 and 295 after the beginning of the photothermal program. mRNA levels have
691 been quantified by q-PCR. The asterisk (*) represents a significant difference ($p < 0.05$) and
692 (**) ($p < 0.001$).

693

694 **Fig. 7. Cortisol levels variation (ng/mL) in F1 and domesticated females.** (9 females per
695 time per population + standard deviation). Sampling times T0, T35, T157 and T295
696 correspond respectively to days 0, 35, 157 and 295 after the beginning of the photothermal
697 program. The asterisk (*) represents a significant difference ($p < 0.05$).

698

699