



HAL
open science

Epidemiological surveillance of SARS-CoV-2 by genome quantification in wastewater applied to a city in the northeast of France: Comparison of ultrafiltration- and protein precipitation-based methods

Isabelle Bertrand, Julie Challant, Helene Jeulin, Cédric Hartard, Laurence Mathieu, Severine Lopez, Evelyne Schvoerer, Sophie Courtois, Christophe Gantzer

► To cite this version:

Isabelle Bertrand, Julie Challant, Helene Jeulin, Cédric Hartard, Laurence Mathieu, et al.. Epidemiological surveillance of SARS-CoV-2 by genome quantification in wastewater applied to a city in the northeast of France: Comparison of ultrafiltration- and protein precipitation-based methods. *International Journal of Hygiene and Environmental Health*, 2021, 233, pp.113692. 10.1016/j.ijheh.2021.113692 . hal-03167302

HAL Id: hal-03167302

<https://hal.univ-lorraine.fr/hal-03167302>

Submitted on 11 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Epidemiological surveillance of SARS-CoV-2 by genome quantification in wastewater applied to a**
2 **city in the northeast of France: comparison of ultrafiltration- and protein precipitation-based**
3 **methods**

4
5 Isabelle Bertrand, ^{a*} Julie Challant, ^a Hélène Jeulin, ^{a,b} Cédric Hartard, ^{a,b} Laurence Mathieu, ^{a,c}
6 Séverine Lopez, ^a Scientific Interest Group Obépine, ^d Evelyne Schvoerer, ^{a,b} Sophie Courtois, ^e
7 Christophe Gantzer^a

8
9 ^a Université de Lorraine, CNRS, LCPME, F-54000 Nancy, France

10 ^b Laboratoire de Virologie, CHRU de Nancy Brabois, F-54500 Vandœuvre-lès-Nancy, France

11 ^c EPHE, PSL, UMR CNRS 7564, LCPME, F-54000 Nancy, France

12 ^d The steering committee of the scientific interest group Obépine is composed of Jean-Luc Bailly
13 (Laboratoire Microorganismes, Génome et Environnement, LMGE, UMR 6023 CNRS-Université
14 Clermont Auvergne, Clermont-Ferrand, France), Christophe Gantzer (Laboratoire de Chimie
15 Physique et Microbiologie pour les Matériaux et l'Environnement, LCPME, UMR 7564 CNRS-
16 Université de Lorraine, Nancy, France), Soizick Le Guyader (Ifremer Nantes, France), Yvon Maday
17 (Institut Carnot Smiles, Sorbonne Université, Laboratoire Jacques-Louis Lions, UMR 7598 and
18 Institut Universitaire de France), Vincent Maréchal (Sorbonne Université, INSERM, Centre de
19 Recherche Saint-Antoine, 75012, Paris, France), Jean-Marie Mouchel (Sorbonne Université, CNRS,
20 EPHE, UMR 7619 Metis, Paris, France, e-LTER Zone Atelier Seine), Laurent Moulin (Laboratoire R&D
21 Eau de Paris, France), Remy Teyssou (Val de Grâce, Unité de Virologie de l'IRBA. Institut de
22 Recherche Biomédicale des Armées, Brétigny, France), Sébastien Wurtzer (Laboratoire R&D Eau de
23 Paris, France).

24 ^e SUEZ, CIRSEE, 38 rue du Président Wilson, F-78230 Le Pecq, France

25

26 *Corresponding author: LCPME, Campus Brabois Santé, Bâtiment AB, 9 Avenue de la Forêt de Haye,
27 BP 20199, F-54505 Vandoeuvre-lès-Nancy, France
28 Email: isabelle.bertrand@univ-lorraine.fr

29

30 **Abstract**

31 The aim of the present study was to develop a simple, sensitive, and specific approach to quantifying
32 the SARS-CoV-2 genome in wastewater and to evaluate this approach as a means of epidemiological
33 surveillance. Twelve wastewater samples were collected from a metropolitan area in north-eastern
34 France during April and May 2020. In addition to the quantification of the SARS-CoV-2 genome, F-
35 specific RNA phages of genogroup II (FRNAPH GGII), naturally present in wastewater, were used as
36 an internal process control for the viral concentration and processing of RT-PCR inhibitors. A
37 concentration method was required to allow the quantification of the SARS-CoV-2 genome over the
38 longest possible period. A procedure combining ultrafiltration, phenol-chloroform-isoamyl alcohol
39 purification, and the additional purification of the RNA extracts was chosen for the quantification of
40 the SARS-CoV-2 genome in 100-mL wastewater samples. At the same time, the COVID-19 outbreak
41 was evaluated through patients from the neighbouring University Hospital of Nancy, France. A
42 regular decrease in the concentration of the SARS-CoV-2 genome from $\sim 10^4$ gc/L to $\sim 10^2$ gc/L of
43 wastewater was observed over the eight weeks of the study, during which the population was
44 placed under lockdown. The SARS-CoV-2 genome was even undetectable during one week in the
45 second half of May and present but non-quantifiable in the last sample (28 May). A concordant
46 circulation in the human community was highlighted by virological diagnosis using respiratory
47 samples, which showed a decrease in the number of COVID-19 cases from 677 to 52 per week over
48 the same period. The environmental surveillance of COVID-19 using a reliable viral quantification
49 procedure to test wastewater is a key approach. The real-time detection of viral genomes can allow
50 us to predict and monitor the circulation of SARS-CoV-2 in clinical settings and survey the entire
51 urban human population.

52

53 **Keywords:** SARS-CoV-2; Wastewater; Concentration Methods; Prevalence

54 **Highlights**

- 55 • Wastewater-based epidemiology is an interesting approach for SARS-CoV-2.
- 56 • Ultrafiltration is a reliable method to concentrate SARS-CoV-2 from wastewater.
- 57 • FRNAPH naturally present in wastewater can be used as internal process control.

58

59 **1. Introduction**

60 In December 2019, the first outbreak of a new coronavirus disease (COVID-19) was reported in
61 Wuhan, China. A new coronavirus linked to Severe Acute Respiratory Syndrome (SARS-CoV-2) was
62 identified. It displayed more than 80% sequence homology with the previous SARS-CoV, identified
63 in 2003 (Cheung et al., 2020). By March 2020, 114 countries reported outbreaks and the WHO
64 declared a pandemic linked to SARS-CoV-2 (WHO, 2020). France was severely affected, with more
65 than 2,550,000 confirmed COVID-19 infections and more than 62,000 deaths by the end of 2020.

66 Even though many cases of COVID-19 are asymptomatic, when symptomatic it is characterized
67 mainly by fever and respiratory symptoms, but there are also a variety of other symptoms
68 frequently associated with the disease. These include gastrointestinal complaints, including
69 anorexia, diarrhoea, and vomiting/nausea at frequencies of up to 27%, 12% and 10%, respectively
70 (Cheung et al., 2020). SARS-CoV-2 is able to productively replicate in human gut enterocytes (Lamers
71 et al., 2020) and the SARS-CoV-2 genome has been detected in the stools of both symptomatic and
72 asymptomatic patients (Cai et al., 2020; Gao et al., 2020; Holshue et al., 2020; Tang et al., 2020;
73 Wölfel et al., 2020; Zhang et al., 2020). Viral RNA may be detected in stools up to 10 days after viral
74 clearance from the respiratory tract, regardless of the severity of the disease (Cheng et al., 2020).
75 The viral concentration can be high, potentially reaching 10^7 genome copies (gc)/g (Wölfel et al.,
76 2020).

77 As expected, very high concentrations of the SARS-CoV-2 genome have also been detected in
78 wastewater (Ahmed et al., 2020; Hindson, 2020; Medema et al., 2020; Nemudryi et al., 2020;
79 Randazzo et al., 2020; Wu et al., 2020a; Wurtzer et al., 2020). Concentrations may reach close to
80 10^6 gc/L (Wu et al., 2020a; Wurtzer et al., 2020). The presence of SARS-CoV-2 in wastewater
81 indicates a potential health risk, but also a data source that can be used for epidemiological
82 purposes (Lodder and de Roda Husman, 2020; Thompson et al., 2020). Wastewater-based
83 epidemiology is not a new idea. It has been used to detect chemicals (Choi et al., 2020) and other

84 viruses, such as norovirus, hepatitis A virus, poliovirus, and Aichivirus (Asghar et al., 2014; Hellmer
85 et al., 2014; Lodder et al., 2012, 2013). Many countries (including The Netherlands, Spain, France,
86 Australia, and Israel) now support the idea of using this type of epidemiological approach in the
87 surveillance of human populations and to possibly launch an early warning system to predict future
88 outbreaks (Ahmed et al., 2020; Bar Or et al., 2020; Medema et al., 2020; Wu et al., 2020; Wurtzer
89 et al., 2020).

90 Whatever its purpose — whether epidemiological or to assess public health risk — the first step is
91 to define a method of quantifying SARS-CoV-2 in wastewater. The methodologies that have been
92 used to date are highly diverse. The concentration methods were based on various principles,
93 including ultrafiltration (Ahmed et al., 2020; Medema et al., 2020; Nemudryi et al., 2020; Trottier et
94 al., 2020); protein precipitation (Kumar et al., 2020; La Rosa et al., 2020b; Wu et al., 2020a);
95 electronegative membranes (Ahmed et al., 2020) and ultracentrifugation (Wurtzer et al., 2020).
96 Recovery rates are currently unknown or extrapolated from those of similar viruses such as porcine
97 epidemic diarrhoea virus (PEDV) and mengovirus (MgV) (Randazzo et al., 2020).

98 The aim of this study was to define the dynamics of viral concentration in wastewater during the
99 first lockdown of the French population. This objective required a simple, specific, and sensitive
100 approach to quantifying the SARS-CoV-2 genome in wastewater. In the present study, we took
101 advantage of the wide circulation of SARS-CoV-2 in north-eastern France to compare two methods
102 of virus concentration. We assessed the recovery rate of SARS-CoV-2 by virus quantification directly
103 in four unconcentrated wastewater samples. We also included an internal process control, using F-
104 specific RNA phages of genogroup II (FRNAPH GGII). We applied the most reliable method to
105 investigate the temporal variations of SARS-CoV-2 genome concentrations in wastewater over 12
106 weeks in an area of France with 250,000 inhabitants, which has been significantly impacted by
107 COVID-19.

108

109 **2. Material and methods**

110 **2.1. Sewage samples**

111 Samples were collected between 2 April and 28 May 2020 in the French Grand Est region. In that
112 region, the population was under lockdown from 17 March to 10 May 2020. A total of 12 water
113 samples (400 mL) were collected from a wastewater treatment plant (WWTP) weekly or twice
114 weekly. The average volume of influent treated in this WWTP was $\sim 118,000 \text{ m}^3/\text{day}$ ($\sim 250,000$
115 inhabitants). The wastewater samples were captured at regular intervals, just after decantation, but
116 before activated sludge treatment. The samples were stored at -20°C until analysis.

117 **2.2. Clinical samples**

118 At the same time, 19,850 samples were examined by the laboratory of the local University Hospital
119 to ascertain the presence of the SARS-CoV-2 genome in nasopharyngeal swabs. These samples were
120 collected between week 10 and week 22 from suspected COVID-19 patients hospitalized in the local
121 university hospital (60%) and from retirement home residents (40%). More than 95% of the people
122 sampled during this period were resident in the urban area of the WWTP. The procedure used for
123 the detection of the SARS-CoV-2 genome was based on primers and probes designed to target two
124 RdRp (RNA-dependent RNA polymerase) gene segments (RdRp_IP2 and RdRp_IP4), in accordance
125 with a procedure developed by the French National Reference Centre for Respiratory Viruses
126 (Institut Pasteur, Paris, France).

127

128 **2.3. Concentration procedures**

129 Two concentration procedures, based on ultrafiltration and on PEG 6000 precipitation, respectively,
130 were compared in the present study.

131 For the ultrafiltration procedure, a Centricon[®] Plus-70 centrifugal ultrafilter with a cut-off of 100 kD
132 (Merck Millipore) was used. Before processing the samples, the ultrafilter was washed with 60 mL
133 deionised water by centrifugation at $1,500 \times g$ for 15 min to remove the trace amounts of glycerine,

134 in accordance with the manufacturer's recommendations. Two volumes of 50 mL wastewater were
135 both filtered by centrifugation at $1,500 \times g$ for 15 min. After each centrifugation step, the
136 concentrate was recovered by inverting the system and applying centrifugation ($1,000 \times g$ for 2 min).
137 The resulting concentrate's volume was around 1.5 mL. The ultrafilter was then washed with 3.5 mL
138 of deionised water. The washing solution was added to the 1.5 mL concentrate to produce the final
139 concentrate sample (5 mL). In order to recover the maximum amount of virus genome from the
140 ultrafilter, two further washing steps were undertaken, each using 5 mL NucliSENS® lysis buffer
141 (bioMérieux) for an incubation time of 5 min. The entire volume (15 mL) was then used for nucleic
142 acid extraction. Twelve water samples were subjected to this procedure.

143 PEG 6000 precipitation was performed in a 250-mL centrifuge bottle containing 3 g beef extract
144 powder, 3 g NaCl, and 0.37 g glycine for 100 mL of wastewater. After the dissolution of the beef
145 extract powder, 20 g of PEG 6000 were added. The sample was gently stirred at 4°C for 2 h and then
146 maintained at 4°C overnight. The pellet obtained after centrifugation at $4,500 \times g$ and 4°C for 45
147 min was resuspended in deionised water to obtain a concentrated 5 mL sample. Ten millilitres of
148 NucliSENS® lysis buffer were then added to the concentrate. After incubation for 10 min at room
149 temperature, the entire volume (15 mL) was used for nucleic acid extraction. This concentration
150 method was tested on the first four water samples of our study.

151 In tandem with each concentration procedure, 5 mL samples of unconcentrated water were used
152 for viral genome extraction. They were submitted to the same procedure as the concentrated
153 samples. Ten millilitres of NucliSENS® lysis buffer was added to the water, which was incubated at
154 room temperature for 10 min, prior to the nucleic acid extraction.

155 **2.4. Nucleic acid extraction**

156 Both concentrated and unconcentrated wastewater samples reached the same volume of 15 mL
157 after the lysis step. Phenol-chloroform-isoamyl alcohol purification was then undertaken with both
158 concentrated and unconcentrated samples. This was realized in a 50 mL conical tube, containing 4

159 g of a mixture of high-vacuum silicon grease (Dow Corning®) and silicon dioxide (Sigma) (90:10 w/w),
160 which was labelled the separation tube. The sample in lysis buffer (15 mL) was transferred to the
161 separation tube and 15 mL phenol-chloroform-isoamyl alcohol (25:24:1 pH 7.8-8.2, Acros
162 Organics™) were added. After 15 s of vigorous stirring by hand and centrifugation (3,500 × g for 5
163 min), the hydrophile supernatant (15 mL) was recovered. The nucleic acid extraction was continued
164 using 70 µL of magnetic silica beads and the NucliSENS® easyMAG™ platform (bioMérieux). The
165 extracted nucleic acids were eluted in 100 µL of elution buffer.

166 A complementary step was applied to some of the samples to remove residual environmental
167 inhibitors using OneStep PCR Inhibitor Removal kit (ZymoResearch). The RNA samples were stored
168 at -80°C until viral genome quantification. A negative control made of sterile phosphate buffered
169 saline was included at the beginning of each nucleic acid extraction procedure.

170 **2.5. Viral genome quantification**

171 Viral RNA quantification was performed using real-time RT-PCR and RT-digital droplet PCR (RT-
172 ddPCR).

173 For the FRNAPH GGII genome, the VTB4-Fph GII set published by Wolf et al. (2010) was used. For
174 the SARS-CoV-2 genome, two primer sets were selected: the “RdRp_IP4” set developed by the
175 Pasteur Institute (Paris, France: [https://www.who.int/docs/default-source/coronaviruse/real-time-
176 rt-pcr-assays-for-the-detection-of-sars-cov-2-institut-pasteur-paris.pdf?sfvrsn=3662fcb6_2](https://www.who.int/docs/default-source/coronaviruse/real-time-rt-pcr-assays-for-the-detection-of-sars-cov-2-institut-pasteur-paris.pdf?sfvrsn=3662fcb6_2)) for the
177 RdRp gene and the “E” set developed by Corman et al. (2020) for the envelope protein (E) gene. The
178 specificity of these sets for other respiratory viruses, including human coronaviruses, has been
179 previously described (Corman et al., 2020; Etievent et al., 2020). The SARS-CoV-2 and FRNAPH GGII
180 genomes were quantified in the same RNA extracts. For the RdRp_IP4, E, and VTB4-Fph GGII sets,
181 quantification was performed using an RNA UltraSens™ One-Step Quantitative RT-PCR system
182 (Applied Biosystems™). The RdRp_IP4 and E sets were applied to 5 µL of RNA in a 25-µL reaction
183 volume with final concentrations of 0.4 µM for each primer and 0.2 µM for the probe. The VTB4-

184 Fph GGII set was used on 2 μ L of RNA in a 20- μ L reaction volume with final concentrations of 1 μ M
185 for each primer and 0.3 μ M for the probe. A StepOnePlus Real-Time PCR System (Applied
186 Biosystems™) was used for these three real-time RT-PCR assays. For the RdRp_IP4 and E sets, the
187 RT step was performed at 50 °C for 20 min and PCR amplification was performed at 95°C for 2 min,
188 followed by 50 cycles of 15 s at 95°C and 30 s at 58 °C. For the VTB4-Fph GGII set the RT step was
189 performed at 50 °C for 30 min and PCR amplification was performed at 95°C for 2 min, followed by
190 45 cycles of 15 s at 95°C and 40 s at 58 °C. Negative and positive controls were included in each
191 experiment. Quantification was carried out using standard curve ranges. RNA extracted from
192 patients who had tested positive for SARS-CoV-2 was quantified using ddRT-PCR with the E set, as
193 described below. Quantified RNA was then used to obtain the standard curve for both RdRp_IP4
194 and E genes. The nCoV-ALL-Control plasmid (Eurofins genomic) was also used for the standard curve
195 of the E gene. This plasmid containing ampicillin resistance gene was maintained in TOP10
196 chemically competent *E. coli* (Invitrogen) and quantified using Qubit 4 fluorometer (Invitrogen). At
197 last, RNA extracted from GA phage suspension was used to obtain the standard curve for FRNAPH
198 GGII. RNA extracted from GA phage suspension was quantified by ddRT-PCR using VTB4-Fph the
199 GGII set as described below. The standard curves ranged from 1×10^{-1} to 1×10^4 gc/reaction for the
200 RdRp_IP4 and E genes of SARS-CoV-2 and from 2.8×10^{-1} to 2.8×10^4 gc/reaction for the GA phage.
201 The limit of detection (LoD) was 1 gc/RT-qPCR reaction for the RdRp and E genes. The limit of
202 quantification (LoQ) ranged from 1 to 10 gc/RT-qPCR reaction for the RdRp gene and reached 1
203 gc/RT-qPCR reaction for the E gene. When we take the analytical volumes of the wastewater sample
204 concentrations subjected to RT-qPCR analysis into consideration, the LoQ ranged from 2×10^2 to
205 2×10^3 gc/L of wastewater for the RdRp gene and reached 2×10^2 gc/L of wastewater for the E gene.
206 The ddRT-PCR assays were performed using the E set of SARS-CoV-2 and the VTB4-Fph GGII set for
207 GA phage. Amplifications were carried out in a 20- μ L reaction mixture containing 5 μ L of RNA and
208 15 μ L of One-Step RT-ddPCR™ Kit for Probes (Bio-Rad). The reaction mix contained 0.9 μ M of each

209 primer and 0.3 μ M of the probe. The samples were placed in the droplet generator using 70 μ L of
210 generator oil to generate up to 20,000 droplets per sample. The resulting picolitre droplet emulsions
211 (40 μ L) were transferred to a Veriti 96-Well Thermal Cycler (Applied Biosystems). After
212 amplification, the plate was transferred to the QX100™ Droplet Reader (Bio-Rad) and QuantaSoft™
213 Software (Bio-Rad) was used to measure the number of positive droplets per well. Droplets were
214 designated positive or negative based on their fluorescence amplitude, using thresholding. The
215 starting concentration of each target RNA molecule was then calculated, by modelling a Poisson
216 distribution.

217 For each sample and each targeted virus (SARS-CoV-2 and FRNAPH GGII) the recovery rate was
218 calculated as follows: recovery rate = genome copies (gc) in 5 mL unconcentrated sample
219 $\times 20 \times 100 /$ gc in 100 mL of the concentrate. The high concentration of the FRNAPGH GGII genome
220 allowed for the determination of the recovery rate in all wastewater samples. It was also possible
221 to determine the recovery rate for SARS-CoV-2 in 5 of the 12 samples.

222 **2.6. Estimation of RT-qPCR inhibition**

223 The estimation of RT-qPCR inhibition in the wastewater samples was based on the concentrations
224 of FRNAPH GGII detected. Logarithmic dilutions (1/10 and 1/100) were performed in PCR-grade
225 water following viral RNA extraction. The RT-qPCR assay was then carried out on both undiluted and
226 diluted RNA extracts. The percentage of inhibition was estimated for undiluted and 1/10 samples
227 by taking the concentration obtained from the 1/100 samples as a reference, due to the high dilution
228 of potential inhibitors.

229 **2.7. Statistical analysis**

230 The statistical analyses were performed using XLSTAT 2020.3.1.2 software. As normal distribution
231 of the data and the homogeneity of variances could not be met, two non-parametric tests were
232 used. A Kruskal-Wallis test was performed on k-independent samples ($k > 3$) to compare the viral
233 genome copy values obtained using the different concentration and purification methods. A

234 Wilcoxon signed-rank test was performed on paired samples to compare the recovery rates
235 following the two methods and the genome concentrations of the two targeted genes. *P* values <
236 0.05 were considered statistically significant.

237

238 **3. Results and discussion**

239 **3.1. Concentration method for SARS-CoV-2 quantification in wastewater**

240 The epidemiological surveillance of SARS-CoV-2 by quantifying its genome in wastewater requires
241 reliable methods of concentration and detection. Beginning our study in a region that was highly
242 impacted by COVID-19 allowed us to estimate recovery rates for SARS-CoV-2.

243 FRNAPH are usually present in wastewater at concentrations that are relatively stable over time and
244 around the world (Lucena et al., 2003). We obtained just under 2.5×10^7 gc/L which was high enough
245 to allow us to evaluate both recovery rates and PCR inhibition problems. Using these
246 bacteriophages, the presence of PCR inhibitors was detected at even a low volume (5 mL) of
247 unconcentrated wastewater. Indeed, the PCR inhibition varied between 88% and 100% in undiluted
248 RNA extracts (n=5) and between 14% and 42% in 1/10 diluted RNA extracts (n=5). Following phenol-
249 chloroform purification, the PCR inhibitors had been completely removed from 5 mL of wastewater.
250 In these five wastewater samples, the level of contamination by SARS-CoV-2 also enabled the
251 quantification of the virus in 5 mL unconcentrated samples and showed that, following phenol-
252 chloroform purification, the concentrations of SARS-CoV-2 genome detected in the undiluted
253 samples had multiplied by between a factor of 2 and a factor of 10. The positive impact of this
254 method on viral RNA extraction from environmental samples such as wastewater, sediments and
255 animal stool samples has already been shown by previous studies (Miura et al., 2011; Hartard et al.,
256 2015). The phenol-chloroform purification method was systematically applied in the following
257 experiments. A complementary method for the removal of PCR inhibitors in RNA extracts was tested
258 during the comparison of the concentration methods (as described below).

259 Two concentration procedures, based on protein precipitation using PEG 6 000 and on ultrafiltration
260 using the Centricon® 70-Plus 100 kD device, respectively, were compared. Recovery rates for both
261 FRNAPH GGII and SARS-CoV-2 could be determined (Table 1) for both methods in four distinct
262 wastewater samples collected in April 2020, since these samples tested positive for the SARS-CoV-
263 2 genome even when only 5 mL of unconcentrated wastewater was analysed. The mean recovery
264 rate of FRNAPH GGII reached $47.7 \pm 18.5\%$ and $39.2 \pm 23.9\%$ using ultrafiltration and protein
265 precipitation, respectively. Because some of the results obtained for FRNAPH GGII showed residual
266 PCR inhibition, the RNA extracts previously obtained by both methods were then purified using a
267 PCR inhibitor removal kit. This treatment led to mean recovery rates of $47.1 \pm 10.4\%$ and
268 $43.3 \pm 20.3\%$ for ultrafiltration and protein precipitation, respectively. This supplementary
269 treatment therefore led to a slight increase in the mean recovery rate of protein precipitation. It
270 also led to a decrease in the standard deviation in both concentration methods. The concentration
271 values obtained for FRNAPH GGII using the two concentration methods, with or without the use of
272 the PCR inhibitor removal kit, were not significantly different (p value = 0.098, Kruskal-Wallis test).
273 For SARS-CoV-2, the mean recovery rates of the RdRp_IP4 gene reached $64.1 \pm 50.2\%$ and
274 $32.4\% \pm 20.2\%$ using ultrafiltration and protein precipitation, respectively. Additional purification of
275 the RNA extracts led to mean recovery rates of $55.8 \pm 46.9\%$ and $23.5 \pm 15.0\%$ for ultrafiltration and
276 protein precipitation, respectively. Thus, the mean recovery rate obtained for SARS-CoV-2
277 (RdRp_IP4 gene) using ultrafiltration were twice as high as those obtained using protein
278 precipitation. Moreover, the highest recovery rate was obtained using ultrafiltration in the four
279 wastewater samples. As observed in the case of the phage, additional purification reduced the
280 standard deviation values. The genome concentration values were not significantly different
281 between the two concentration methods (p value = 0.088, Kruskal-Wallis test). Amplification of the
282 E gene was then performed on the purified RNA extracts, resulting in mean recovery rates of
283 $64.0 \pm 41.6\%$ and $45.0 \pm 44.6\%$ for ultrafiltration and protein precipitation, respectively (Table 1).

284 The statistical analysis of the data obtained for both the RdRp_IP4 and E genes showed significantly
285 higher recovery rates for ultrafiltration than for protein precipitation (p value = 0.009, Wilcoxon
286 signed-rank test).

287 Recovery rates and the inhibition of molecular detection methods applied for SARS-CoV-2 in
288 wastewater have been poorly described. Nevertheless, our SARS-CoV-2 recovery rate was higher
289 than those described in a recent review paper (La Rosa et al., 2020a) reporting five studies that
290 described recovery rates for coronavirus in water matrices. But these recovery rates were all defined
291 using artificial spiking with different types of coronaviruses (bovine enteric coronavirus,
292 transmissible gastroenteritis virus [TGEV], murine hepatitis virus, and SARS-CoV). The recovery rate
293 did not exceed 28% for bovine enteric coronavirus seeded in wastewater (Collomb et al., 1986) and
294 it varied from 0 to 21.4% for SARS-CoV seeded in wastewater ($n=4$) (Wang et al., 2005). Our higher
295 recovery rates for the SARS-CoV-2 naturally present in the wastewater samples could be primarily
296 explained by the choice of concentration method. Increasing PEG concentration might lead to a
297 decrease in the standard deviation values as suggested by the recovery rate of $41.9 \pm 6.5\%$ for TGEV
298 seeded in concentrated surface water (Blanco et al., 2019). Two of the studies reviewed by La Rosa
299 and colleagues (2020a) used adsorption/elution on glass wool with the objective of analysing large
300 volumes (5–50 L) of surface water (Blanco et al., 2019) and dechlorinated tap water (Abd-Elmaksoud
301 et al., 2014). Such methods are commonly used as primary concentration methods and are followed
302 by secondary concentration methods — PEG precipitation in the case of these two studies.
303 However, the combination of concentration methods used on large volumes of water commonly
304 leads to a decrease in recovery rates. By working on municipal wastewater spiked with different
305 enveloped and non-enveloped viruses, Ye et al. (2016) have shown that ultrafiltration is a more
306 suitable method for wastewater than ultracentrifugation or PEG precipitation because the recovery
307 rate of ultrafiltration is 25% compared with 5% by ultracentrifugation or PEG precipitation. Since Ye
308 and colleagues (2016) used culture as their detection method, the low recovery rates may be

309 explained by the possible inactivation of coronavirus by the greater ultracentrifugal forces and the
310 lower precipitation capacity of enveloped viruses compared with non-enveloped viruses in PEG
311 precipitation. Wurtzer and colleagues (2020) showed the ability of ultracentrifugation to
312 concentrate SARS-CoV-2 from wastewater, but the recovery rate was not given, and detection was
313 done using RT-qPCR instead of culture methods. It is important to note that the ultrafiltration
314 method could be used for the detection of infectious SARS-CoV-2 in wastewater, as shown with
315 FRNAPH by Medema et al. (2020). The environmental conditions may have a negative impact on the
316 integrity of the SARS-CoV-2 particles, with implications for detection methods and viral risk
317 assessment; this requires further research.

318 The procedure combining ultrafiltration, phenol-chloroform purification, and additional purification
319 of the RNA extracts was chosen to monitor the concentrations of the SARS-CoV-2 genome in
320 wastewater during longitudinal wastewater sampling. Our approach based on ultrafiltration may be
321 performed with different devices such as Centricon® (Ahmed et al., 2020; Medema et al., 2020),
322 Corning® Spin-X® UF (Nemudryi et al., 2020) or Vivaspin® (Trottier et al., 2020) concentrators.

323

324 **3.2. Quantification of the SARS-CoV-2 genome in wastewater**

325 Our longitudinal study was performed in a French geographical area with one of the highest
326 prevalence rates of COVID-19. The population was placed under lockdown from 17 March until 11
327 May 2020, but the number of cases detected at the University Hospital increased continuously until
328 the end of March, when the wastewater sampling period started. Twelve wastewater samples,
329 collected between 2 April and 28 May 2020, were analysed in one replicate using both the
330 ultrafiltration procedure on 100 mL and the direct analysis of 5 mL to quantify the SARS-CoV-2 and
331 FRNAPH GGII genomes.

332 The FRNAPH GGII genome could be quantified in both 5 mL and 100 mL of wastewater in all the
333 samples (Figure 1). The mean concentrations reached $2.1 \times 10^7 \pm 1.1 \times 10^7$ gc/L and

334 $1.6 \times 10^7 \pm 1.4 \times 10^7$ gc/L in unconcentrated and concentrated samples, respectively. From these
335 concentration values, the recovery rates of FRNAPH GGII ranged from 14.1% to 133.8%.

336 We propose a recovery rate of over 10% as a quality control, to validate the results. This goes beyond
337 the current ISO standards for the molecular detection of viruses in water and food, which require
338 recovery efficiencies of over 1% (International Organization for Standardization - ISO 2019; Lowther
339 et al. 2019). Quantifying FRNAPH at the same time as SARS-CoV-2 also allowed for the normalisation
340 of faecal material content, which may vary daily, depending on rainfall or variations in faecal
341 content. In our samples, FRNAPH genomes ranged from 1.1×10^7 to 6.7×10^7 gc/L, showing that
342 the faecal pollution was relatively stable during the period in question. No significant rainfall events
343 were reported during the sampling period. To compare the variations in SARS-CoV-2 genome
344 concentrations in different wastewater samples may require the normalisation of genome content.
345 Other variables, such as temperature, pH, turbidity, total suspended solids, 5-day biological oxygen
346 demand (BOD5), antecedent dry days, and conductivity may explain the variability of faecal
347 indicator bacteria (FIB) (Mohammed et al., 2018; Paule-Mercado et al., 2016) and probably of other
348 faecal pathogens.

349 The SARS-CoV-2 genome could be quantified in 5 mL of wastewater, without using any
350 concentration procedures, between 2 April and 30 April (Figure 1). By analysing 100 mL of
351 wastewater, a decrease in the concentration of the SARS-CoV-2 genome was observed over the
352 course of 8 weeks (Figure 1). The SARS-CoV-2 genome was even undetectable during one week (22
353 and 25 May) and present but below the LoQ in the last sample (28 May). The concentrations of RdRp
354 and E genes in the twelve samples appeared to be similar (p value = 0.496, Wilcoxon signed-rank
355 test). Moreover, the concentration values obtained using real-time RT-qPCR and ddRT-PCR for the
356 E gene were similar (p value = 0.734, Wilcoxon signed-rank test). This suggests that RT-PCR inhibitors
357 were effectively removed by our protocol. Indeed, ddRT-PCR is far less influenced by inhibitors than
358 RT-qPCR (Sun et al., 2019).

359 The decrease in the genome concentrations detected in wastewater may be related to the decrease
360 in the number of COVID-19 cases observed at the Nancy University Hospital (Figure 1). Between 2
361 April and 23 April, the number of cases decreased from 677/week to 286/week and the
362 concentration values of the SARS-CoV-2 genome detected in 100 mL of wastewater decreased from
363 1.2×10^4 gc/L to 3.0×10^3 gc/L. During this period, the virus genome could be detected in only 5 mL
364 of wastewater and decreased only from 1×10^4 gc/L to 7×10^3 gc/L. The analysis of only 5 mL of
365 wastewater could be less representative of the variation in viral genome concentration. After a
366 single data point showing an increase in genome concentration in wastewater in both 5 mL and 100
367 mL (30 April), the SARS-CoV-2 genome was no longer detected in 5 mL of wastewater and the
368 concentrations dropped below 10^3 gc/L in the 100 mL analysis. For three of the samples during this
369 period, either no genes were detected or only one gene was detected, and the number of cases
370 varied between 27/week and 52/week. Nevertheless, SARS-CoV-2 was still detected in wastewater
371 in the last sample (28 May) collected 17 days after the end of the lockdown. We observed a parallel
372 decrease in cases in patients and genome concentration in wastewater, confirming the link between
373 the circulation of the virus in the human population and its presence in wastewater. This clearly
374 confirms the findings of other studies showing such a relationship (Wurtzer et al., 2020). One of the
375 first studies on the detection of the SARS-CoV-2 genome in wastewater (Medema et al., 2020) even
376 showed that it was possible to detect the virus in wastewater during the early stages of the COVID-
377 19 epidemic.

378

379 **4. Conclusion**

380 We developed a method of concentrating SARS-CoV-2 from wastewater with one of the highest
381 recovery rates described in the literature. The quantification of the SARS-CoV-2 genome was carried
382 out in both concentrated and unconcentrated wastewater which was naturally contaminated; this
383 gives our results a high degree of reliability. Moreover, this study provides additional data to validate

384 proof-of-concept for a link between the outbreak in the human community and the concentrations
385 of faecally excreted viruses in wastewater. This method can contribute to the monitoring of the
386 epidemic and improve the management of potential viral recirculation.

387

388 **Declaration of interest**

389 None

390

391 **Funding**

392 This work has benefited from the financial support of the French Government's Ministry of Higher
393 Education and Research and Innovation (Ministère de l'enseignement supérieur, de la recherche et
394 de l'innovation [MESRI]). This study was supported by the French scientific interest group Obépine
395 for the surveillance of SARS-CoV-2 in wastewater. All the authors' salaries were funded by either
396 Lorraine University (France), the University Hospital of Nancy (France) or SUEZ (France).

397

398 **Acknowledgements**

399 The authors would like to thank Sibel Berger for her contribution to the acquisition of the RT-qPCR
400 assays for SARS-CoV-2 at LCPME. The authors would also like to thank Christophe Merlin and Hélène
401 Guilloteau for their contributions to the plasmid replication.

402

403 **References**

404 **XX**

405 Ahmed, W., Angel, N., Edson, J., Bibby, K., Bivins, A., O'Brien, J.W., Choi, P.M., Kitajima, M., Simpson,
406 S.L., Li, J., Tschärke, B., Verhagen, R., Smith, W.J.M., Zaugg, J., Dierens, L., Hugenholtz, P.,
407 Thomas, K.V., Jochen, F. Mueller, J.F., 2020. First confirmed detection of SARS-CoV-2 in

408 untreated wastewater in Australia: A proof of concept for the wastewater surveillance of
409 COVID-19 in the community. *Sci Total Environ.* 728, 138764.
410 <https://doi.org/10.1016/j.scitotenv.2020.138764>

411 Asghar, H., Diop, O.M., Weldegebriel, G., Malik, F., Shetty, S., Bassioni, L. El, Akande, A.O.,
412 Maamoun, E. Al, Zaidi, S., Adeniji, A.J., Burns, C.C., Deshpande, J., Oberste, M.S., Lowther, S.A.,
413 2014. Environmental surveillance for polioviruses in the global polio eradication initiative. *J.*
414 *Infect. Dis.* 210 (Issue suppl_1), S294–S303. <https://doi.org/10.1093/infdis/jiu022>

415 Bar Or, I., Yaniv, K., Shagan, M., Ozer, E., Erster, O., Mendelson, E., 2020. Regressing SARS-CoV-2
416 sewage measurements onto COVID-19 burden in the population: a proof-of-concept for
417 quantitative environmental surveillance. medRxiv preprint.
418 <https://doi.org/10.1101/2020.04.26.20073569>

419 Cai, J., Xu, J., Lin, D., Yang, Z., Xu, L., Qu, Z., Zhang, Y., Zhang, H., Jia, R., Liu, P., Wang, X., Ge, Y., Xia,
420 A., Tian, H., Chang, H., Wang, C., Li, J., Wang, J., Zeng, M., 2020. A case series of children with
421 2019 novel coronavirus infection: clinical and epidemiological features. *Clin. Infect. Dis.*
422 *ciaa198*. <https://doi.org/10.1093/cid/ciaa198>

423 Cheung, K.S., Hung, I.F., Chan, P.P., Lung, K., Tso, E., Liu, R., Ng, Y., Chu, M.Y., Chung, T.W., Tam, A.R.,
424 Yip, C.C., Leung, K.-H., Yim-Fong Fung, A., Zhang, R.R., Lin, Y., Cheng, H.M., Zhang, A.J., To, K.K.,
425 Chan, K.-H., Yuen, K.-Y., Leung, W.K., 2020. Gastrointestinal manifestations of SARS-CoV-2
426 infection and virus load in fecal samples from the Hong Kong cohort and systematic review
427 and meta-analysis. *Gastroenterol.* 159 (1), 81–95.
428 <https://doi.org/10.1053/j.gastro.2020.03.065>

429 Choi, P.M., Thomas, K.V., O'Brien, J.W., Mueller, J.F., 2020. Mining population exposure and
430 community health via wastewater-based epidemiology. In: Jiang G., Li X. (eds) *A New Paradigm*
431 *for Environmental Chemistry and Toxicology*. Springer, Singapore.

432 Corman, V.M., Landt, O., Kaiser, M., Molenkamp, R., Meijer, A., Chu, D.K.W., Bleicker, T., Brünink, S.,
433 Schneider, J., Schmidt, M.L., Mulders, D.G.J.C., Haagmans, B.L., van der Veer, B., van den Brink, S.,
434 Wijsman, L., Goderski, G., Romette, J.L., Ellis, J., Zambon, M., Peiris, M., Goossens, H., Reusken, C.,
435 Koopmans, M.P.G., Drosten, C., 2020. Detection of 2019 novel coronavirus (2019-nCoV) by real-
436 time RT-PCR. Euro Surveill.25 (3), 2000045. [https://doi.org/10.2807/1560-](https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045)
437 [7917.ES.2020.25.3.2000045](https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045)

438 Etievant, S., Bal, A., Escuret, V., Brengel-Pesce, K., Bouscambert, M., Cheynet, V., Generenaz, L., Oriol,
439 G., Destras, G., Billaud, G., Josset, L., Frobert, E., Morfin, M., Gaymard, A., 2020. Performance
440 Assessment of SARS-CoV-2 PCR Assays Developed by WHO Referral Laboratories. J. Clin. Med. 9,
441 E1871. <https://doi.org/10.3390/jcm9061871>

442 Gao, Q.Y., Chen, Y.X., Fang, J.Y., 2020. 2019 Novel coronavirus infection and gastrointestinal tract.
443 J. Dig. Dis. 21 (3), 125–126. <https://doi.org/10.1111/1751-2980.12851>

444 Hartard, C., Rivet, R., Banas, S., Gantzer, C., 2015. Occurrence of and sequence variation among F-
445 specific RNA bacteriophage subgroups in feces and wastewater of urban and animal origins.
446 Appl Environ Microbiol. 81 (8) , 6505–6515. <https://doi.org/10.1128/AEM.01905-15>

447 Hellmér, M., Paxéus, N., Magnus, L., Enache, L., Arnholm, B., Johansson, A., Bergström, T., Norder,
448 H., 2014. Detection of pathogenic viruses in sewage provided early warnings of hepatitis A
449 virus and norovirus outbreaks. Appl. Environ. Microbiol. 80 (21), 6771–6781.
450 <https://doi.org/10.1128/AEM.01981-14>

451 Hindson, J., 2020. COVID-19: faecal–oral transmission? Nat. Rev. Gastroenterol. Hepatol. 17 (5),
452 259. <https://doi.org/10.1038/s41575-020-0295-7>

453 Holshue, M., DeBolt, C., Lindquist, S., Lofy, K., Wiesman, J., Bruce, H., Spitters, C., Ericson, K.,
454 Wilkerson, S., Tural, A., Diaz, G., Cohn, A., Fox, L., Patel, A., Gerber, S., Kim, L., Tong, S., Lu, X.,
455 Lindstrom, S., Pallansch, M., Weldon, W., Biggs, H., Uyeki, T., Pillai, S., 2020. First case of 2019

456 novel coronavirus in the United States. *N. Engl. J. Med.* 382 (10), 929–936.
457 <https://doi.org/10.1056/NEJMoa2001191>

458 Institut Pasteur, Paris. Protocol: Real-Time RT-PCR Assays for the Detection of SARS-CoV-2. Available
459 online: [https://www.who.int/docs/default-source/coronaviruse/real-time-rt-pcr-assays-for-the-](https://www.who.int/docs/default-source/coronaviruse/real-time-rt-pcr-assays-for-the-detection-of-sars-cov-2-institut-pasteur-paris.pdf?sfvrsn=3662fcb6_2)
460 [detection-of-sars-cov-2-institut-pasteur-paris.pdf?sfvrsn=3662fcb6_2](https://www.who.int/docs/default-source/coronaviruse/real-time-rt-pcr-assays-for-the-detection-of-sars-cov-2-institut-pasteur-paris.pdf?sfvrsn=3662fcb6_2) (accessed on 30 March
461 2020).

462 Kumar, M., Kumar Patel, A., Shah, A.V., Raval, J., Rajpara, N., Joshi, M., Joshid, C.G., 2020. First proof
463 of the capability of wastewater surveillance for COVID-19 in India through detection of genetic
464 material of SARS-CoV-2. *Sci. Total Environ.* 746, 141326.
465 <https://doi.org/10.1016/j.scitotenv.2020.141326>

466 Lamers, M. M., Beumer, J., van der Vaart, J., Knoops, K., Puschhof, J., Breugem, T. I., Ravelli, R. B. G.
467 J., van Schayck, P., Mykytyn, A. Z., Duimel, H. Q., van Donselaar, E., Riesebosch, S., Kuijpers, H.
468 J. H., Schippers, D., van de Wetering, W. J., de Graaf, M., Koopmans, M., Cuppen, E., Peters, P.
469 J., Haagmans, B. L., Clevers, H., 2020. *Science*. SARS-CoV-2 productively infects human gut
470 enterocytes. 369(6499):50-54. <https://doi.org/10.1126/science.abc1669>

471 La Rosa, G., Bonadonna, L., Lucentini, L., Kenmoe, S., Suffredini, E., 2020a. Coronavirus in water
472 environments: Occurrence, persistence and concentration methods - A scoping review. *Water*
473 *Res.* 179, 115899. <https://doi.org/10.1016/j.watres.2020.115899>

474 La Rosa, G., Iaconelli, M., Mancini, P., Bonanno Ferraro, G., Veneri, C., Bonadonna, L., Lucentini, L.,
475 Suffredini, E., 2020b. First detection of SARS-CoV-2 in untreated wastewaters in Italy. *Sci Total*
476 *Environ.* 736, 139652. <https://doi.org/10.1016/j.scitotenv.2020.139652>

477 Lodder, W. J., Buisman, A. M., Rutjes, S. A., Heijne, J. C., Teunis, P. F., de Roda Husman, A. M., 2012.
478 Feasibility of quantitative environmental surveillance in poliovirus eradication strategies. *Appl*
479 *Environ Microbiol.* 78 (11), 3800–3805. <https://doi.org/10.1128/AEM.07972-11>

480 Lodder, W.J., Rutjes, S.A., Takumi, K., de Roda Husman, A.M., 2013. Aichi virus in sewage and surface
481 water, the Netherlands. *Emerg. Infect. Dis.* 19 (8), 1222–1230.
482 <https://doi.org/10.3201/eid1908.130312>

483 Lodder, W., de Roda Husman, A.M., 2020. SARS-CoV-2 in wastewater: potential health risk, but also
484 data source. *Lancet Gastroenterol Hepatol.* 5 (6), 533–534. [https://doi.org/10.1016/S2468-
485 1253\(20\)30087-X](https://doi.org/10.1016/S2468-1253(20)30087-X)

486 Lowther, J.A., Bosch, A., Butot, S., Ollivier, J., Mäde, D., Rutjes, S.A., Hardouin, G., Lombard, B., In't
487 Veld, P., Leclercq, A., 2019. Validation of EN ISO method 15216 - Part 1 - Quantification of
488 hepatitis A virus and norovirus in food matrices. *Int. J. Food Microbiol.* 288:82–90.
489 <https://doi.org/10.1016/j.ijfoodmicro.2017.11.014>.

490 Lucena, F., Méndez, X., Morón, A., Calderón, E., Campos, C., Guerrero, A., Cárdenas, M., Gantzer, C.,
491 Schwartzbrod, L., Skrabber, S., Jofre, J., 2003. Occurrence and densities of bacteriophages
492 proposed as indicators and bacterial indicators in river waters from Europe and South
493 America. *J Appl Microbiol.* 94 (5), 808–15. <https://doi.org/10.1046/j.1365-2672.2003.01812.x>

494 Medema, G., Heijnen, L., Elsinga, G., Italiaander, R., Brouwer, A., 2020b. Presence of SARS-
495 Coronavirus-2 RNA in sewage and correlation with reported COVID-19 prevalence in the early
496 stage of the epidemic in The Netherlands. *Environ. Sci. Technol. Lett.* 7, 7, 511–516.
497 <https://dx.doi.org/10.1021/acs.estlett.0c00357>

498 Miura, T., Masago, Y., Sano, D., Omura, T., 2011. Development of an effective method for recovery
499 of viral genomic RNA from environmental silty sediments for quantitative molecular detection.
500 *Appl Environ Microbiol.* 77 (12), 3975–81. <https://doi.org/10.1128/AEM.02692-10>

501 Mohammed, H., Hameed, I.A., Razak Seidu, R., 2018. Comparative predictive modelling of the
502 occurrence of faecal indicator bacteria in a drinking water source in Norway. *Sci Total Environ.*
503 628-629, 1178–1190. <https://doi.org/10.1016/j.scitotenv.2018.02.140>

504 Nemudryi, A., Nemudraia, A., Wiegand, T., Surya, K., Buyukyoruk, M., Wilkinson, R., Cicha C.,
505 Vanderwood, K.K., Wilkinson, R., Wiedenheft, B., 2020. Temporal detection and phylogenetic
506 assessment of SARS-CoV-2 in municipal wastewater. *Cell Rep Med.* 1(6), 100098.
507 <https://doi.org/10.1016/j.xcrm.2020.100098>

508 Paule-Mercado, M.A., Ventura, J.S., Memon, S.A., Jahng, D., Kang, J.-H., Lee; C.-H., 2016. Monitoring
509 and predicting the fecal indicator bacteria concentrations from agricultural, mixed land use
510 and urban stormwater runoff. *Sci Total Environ.* 550, 1171–1181.
511 <http://dx.doi.org/10.1016/j.scitotenv.2016.01.026>

512 Randazzo, W., Truchado, P., Cuevas-Ferrando, E., Simón, P., Allende, A., Sánchez, G., 2020b. SARS-
513 CoV-2 RNA titers in wastewater anticipated COVID-19 occurrence in a low prevalence area.
514 *Water Res.* 181, 115942. <https://doi.org/10.1016/j.watres.2020.115942>

515 Sun, B., Bosch, A., Myrmel, M., 2019. Extended direct lysis method for virus detection on berries
516 including droplet digital RT-PCR or real time RT-PCR with reduced influence from inhibitors. *J*
517 *Virol Methods* 271, 113638. <https://doi.org/10.1016/j.jviromet.2019.04.004>

518 Tang, A., Tong, Z., Wang, H., Dai, Y., Li, K., Liu, J., Wu, W., Yuan, C., Yu, M., Li, P., Yan, J., 2020.
519 Detection of novel coronavirus by RT-PCR in stool specimen from asymptomatic child, China.
520 *Emerg. Infect. Dis. J.* 26 (6), 1337–1339. <http://dx.doi.org/10.3201/eid2606.200301>

521 Thompson, J. R., Nancharaiah, Y. V., Gu, X., Lee W. L., Rajal, V. B., Haines, M. B., Girones, R., Ng L. C.,
522 Alm, E.J., Wuertz, S., 2020. Making waves: Wastewater surveillance of SARS-CoV-2 for
523 population-based health management. *Water Res.* 184, 116181.
524 <https://doi.org/10.1016/j.watres.2020.116181>

525 Trottier, J., Darques, R., Ait Mouheb, N., Partiot, E., Bakhache, W., Deffieu, M.S., Gaudin, R., 2020.
526 Post-lockdown detection of SARS-CoV-2 RNA in the wastewater of Montpellier, France. *One*
527 *Health* 10, 100157. <https://doi.org/10.1016/j.onehlt.2020.100157>

528 Wang, X.W., Li, J.S., Guo, T.K., Zhen, B., Kong, Q.X., Yi, B., Li, Z., Song, N., Jin, M., Xiao, W.J., Zhu, X.M.,
529 Gu, C.Q., Yin, J., Wei, W., Yao, W., Liu, C., Li, J.F., Ou, G.R., Wang, M.N., Fang, T.Y., Wang, G.J., Qiu,
530 Y.H., Wu, H.H., Chao, F.H., Li, J.W., 2005. Concentration and detection of SARS coronavirus in
531 sewage from Xiao tang Shan hospital and the 309th hospital. *J. Virol. Methods* 128 (1–2), 156–
532 161.

533 WHO (World Health Organization) [https://www.who.int/dg/speeches/detail/who-director-general-s-](https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020)
534 [opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020](https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020) Last accessed 23 March
535 2020.

536 Wolf, S., Hewitt, J., Greening, G.E., 2010. Viral multiplex quantitative PCR assays for tracking sources
537 of fecal contamination. *Appl. Environ. Microbiol.* 76 (5), 1388–1394.
538 <https://doi.org/10.1128/AEM.02249-09>

539 Wölfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Mueller, M.A., Niemeyer, D.,
540 Vollmar, P., Rothe, C., Hoelscher, M., Bleicker, T., Bruenink, S., Schneider, J., Ehmann, R.,
541 Zwirgmaier, K., Drosten, C., Wendtner, C., 2020. Virological assessment of hospitalized cases
542 of coronavirus disease 2019. *Nature* 581 (7809), 465–469. [https://doi.org/10.1038/s41586-](https://doi.org/10.1038/s41586-020-2196-x)
543 [020-2196-x](https://doi.org/10.1038/s41586-020-2196-x)

544 Wu, F., Zhang, J., Xiao, A., Gu, X., Lee, L.W., Armas, F., Kauffman, K., Hanage, W., Matus, M., Ghaeli,
545 N., Endo, N., Duvallet, C., Poyet, M., Moniz, K., Washburne, A.D., Erickson, T.B., Chai, P.R.,
546 Thompson, J., Alm, E.J., 2020a. SARS-CoV-2 Titers in Wastewater Are Higher than Expected
547 from Clinically Confirmed Cases. *mSystems*5(4), e00614-20.
548 <https://doi.org/10.1128/mSystems.00614-20>

549 Wu, F., Xiao, A., Zhang, J., Moniz, K., Endo, N., Armas, F., Bonneau, R., Brown, M.A., Bushman, M.,
550 Chai, P.R., Duvallet, C., Erickson, T.B., Foppe, K., Ghaeli, N., Gu, X., Hanage, W.P., Huang, K.H.,
551 Lee, W.L., Matus, M., McElroy, K.A., Nagler, J., Rhode, S.F., Santillana, M., Tucker, J.A., Wuertz,
552 S., Zhao, S., Thompson, J., Alm, E.J. 2020b. SARS-CoV-2 titers in wastewater foreshadow

553 dynamics and clinical presentation of new COVID-19 cases. medRxiv preprint.
554 <https://doi.org/10.1101/2020.06.15.20117747>

555 Wurtzer, S., Marechal, V., Mouchel, J.M., Maday, Y., Teyssou, R., Richard, E., Almayrac, J.L., Moulin,
556 L., 2020. Evaluation of lockdown effect on SARS-CoV-2 dynamics through viral genome
557 quantification in waste water, Greater Paris, France, 5 March to 23 April 2020. EuroSurveill.
558 25(50), pii=200077. <https://doi.org/10.2807/1560-7917.ES.2020.25.50.2000776>

559 Ye, Y., Ellenberg, R.M., Graham, K.E., Wigginton, K.R., 2016. Survivability, partitioning, and recovery
560 of enveloped viruses in untreated municipal wastewater. Environ. Sci. Technol. 50, 5077–
561 5085. <https://doi.org/10.1021/acs.est.6b00876>

562

563 **Table 1: Recovery rates (%) of SARS-CoV-2 and FRNAPH GGII genomes in wastewater samples using two concentration methods (protein**
 564 **precipitation and ultrafiltration) both alone and combined with the use of a PCR inhibitor removal kit.**

565

Virus / Primers set	SARS-CoV-2 / RdRp_IP4		SARS-CoV-2 / E		FRNAPH GGII / VTB4-Fph GGII	
Concentration method	PP	UF	PP	UF	PP	UF
Without PCR Inhibitor removal kit						
Sample 1	56.0	131.0	ND	ND	58.2	49.1
Sample 2	12.0	56.0	ND	ND	4.5	26.0
Sample 3	41.8	60.2	ND	ND	43.9	71.1
Sample 4	19.8	9.2	ND	ND	50.0	44.4
Mean ± SD	32.4 ± 20.2	64.1 ± 50.2	ND	ND	39.2 ± 23.9	47.7 ± 18.5
With PCR Inhibitor removal kit						
Sample 1	40.5	119.0	108.0	119.0	68.2	52.7
Sample 2	27.0	61.0	23.0	45.0	19.1	37.5
Sample 3	22.4	32.6	43.0	70.0	37.2	58.9
Sample 4	4.3	10.6	6.0	22.0	50.0	39.4
Mean ± SD	23.5 ± 15.0	55.8 ± 46.9	45.0 ± 44.6	64.0 ± 41.6	43.3 ± 20.3	47.1 ± 10.4

566

567 PP: protein precipitation, UF: ultrafiltration, ND: not done

568 **Figure 1. Quantitative time-course monitoring of SARS-CoV-2 and FRNAPH-GGII genomes in**
569 **wastewater samples in the Nancy metropolitan area and evolution of the number of cases per**
570 **week at the local University Hospital.** Quantification in 100 mL of wastewater (ultrafiltration) of the
571 SARS-CoV-2 genome targeting RdRp (full triangles) and E (full squares) genes and of the FRNAPH
572 GGII genome (full diamonds). Quantification in 5 mL of wastewater: open forms. Negative samples
573 of the SARS-CoV-2 genome are marked with an asterisk. The sample positive for the RdRp gene but
574 below the LoQ is marked with a plus sign. The number of cases of COVID-19 detected at the local
575 University Hospital is shown per week (bars). The limit of detection (LoD) for RdRp and E genes
576 (2×10^2 gc/L) is represented by a white arrow on the left y-axis. The lines given for genome
577 concentration values are only designed to provide a visual guide.

578

