



HAL
open science

Fetal biometry in ultrasound: A new approach to assess the long-term impact of simulation on learning patterns

G. Ambroise Ambroise Grandjean, C. Bertholdt, S. Zuily, M. Fauvel, G. Hossu, P. Berveiller, O. Morel

► To cite this version:

G. Ambroise Ambroise Grandjean, C. Bertholdt, S. Zuily, M. Fauvel, G. Hossu, et al.. Fetal biometry in ultrasound: A new approach to assess the long-term impact of simulation on learning patterns. *Journal of Gynecology Obstetrics and Human Reproduction*, 2021, 50 (8), pp.102135. 10.1016/j.jogoh.2021.102135 . hal-03231243

HAL Id: hal-03231243

<https://hal.univ-lorraine.fr/hal-03231243>

Submitted on 24 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Fetal biometry in ultrasound: a new approach to assess the long-term impact of simulation on learning patterns

G. Ambroise Grandjean^{1,2,3}, C. Bertholdt^{1,2}, S. Zuily⁴, M. Fauvel⁵, G. Hossu⁵, P. Berveiller^{6,7}, O. Morel^{1,2}

¹ Université de Lorraine, IADI - INSERM, F-54000, Nancy, France

² CHRU Nancy, Department of Obstetrics and Gynecology, F-54000 Nancy, France

³ Université de Lorraine, Midwifery Department, Nancy F-54000, France

⁴ Université de Lorraine, Hôpital virtuel de Lorraine, Nancy F-54000, France

⁵ CHRU Nancy, Université de Lorraine, CIC-IT, F-54000 Nancy, France

⁶ CHI Poissy Saint-Germain-en-Laye, Department of Obstetrics and Gynecology, F-78300, Poissy, France

⁷ Université Versailles Saint-Quentin, EA 7404 - Gamètes Implantation Gestation, F-78180, Montigny le Bretonneux, France

ABSTRACT

Context: Simulation-based education (SBE) has demonstrated its acceptability and effectiveness in improving ultrasound training. Because of the high cost of its implementation (investment in equipment and supervision), a pragmatic assessment of the transfer of skills learned in SBE to clinical practice and the identification of its optimal scheduling conditions have been requested to optimize its input.

Objectives: To quantify the long-term impact of simulation-based education (SBE) on the adequate performance of ultrasound fetal biometry measurements (I). The secondary objective was to identify the temporal patterns that enhanced SBE input in learning (II).

Methods: Trainees were arbitrarily assigned to a 6-month course in obstetric ultrasound with or without an SBE workshop. In the SBE group, the workshop was implemented 'before' or at an 'early' or a 'late-stage' of the course. Those who did not receive SBE were the control group. The ultrasound skills of all trainees were prospectively collected, evaluated by calculating the delta between OSAUS (Objective Structured Assessment of Ultrasound Skills) scores before and after the course (I). Concomitantly, the accuracy of trainees' measurements was assessed throughout the course by verifying their correlation with the corresponding measurements by their supervisors. The percentage of trainees able to perform five consecutive sets of correct measurements in the control group and in each SBE subgroup were compared (II).

Results: The study included 61 trainees (39 SBE and 22 controls). Comparisons between groups showed no significant difference in the quantitative assessment of skill enhancement (difference in the pre- and post-internship OSAUS score: 1.09 ± 0.87 in the SBE group and 0.72 ± 0.98 in the control group) (I). Conversely, the predefined acceptable skill level was reached by a significantly higher proportion of trainees in the 'early' SBE subgroup (74%, compared with 30% in the control group, $P < 0.01$) (II).

Conclusions: The quantitative assessment does not support the existence of long-term benefits from SBE training, although the qualitative assessment confirmed SBE helped to raise the minimal level within a group when embedded in an 'early' stage of a practical course.

1
2
3 **Context:** Simulation-based education (SBE) has demonstrated its acceptability and effectiveness in improving
4 ultrasound training. Because of the high cost of its implementation (investment in equipment and supervision), a
5 pragmatic assessment of the transfer of skills learned in SBE to clinical practice and the identification of its
6 optimal scheduling conditions have been requested to optimize its input.

7 **Objectives:** To quantify the long-term impact of simulation-based education (SBE) on the adequate performance
8 of ultrasound fetal biometry measurements (I). The secondary objective was to identify the temporal patterns
9 that enhanced SBE input in learning (II).

10 **Methods:** Trainees were arbitrarily assigned to a 6-month course in obstetric ultrasound with or without an SBE
11 workshop. In the SBE group, the training was implemented 'before' or at an 'early' or a 'late-stage' of the course.
12 Those who did not receive SBE were the control group. The ultrasound skills of all trainees were prospectively
13 collected, evaluated by calculating the delta between OSAUS (Objective Structured Assessment of Ultrasound
14 Skills) scores before and after the course (I). Concomitantly, the accuracy of trainees' measurements was
15 assessed throughout the course by verifying their correlation with the corresponding measurements by their
16 supervisors. The percentage of trainees able to perform five consecutive sets of correct measurements in the
17 control group and in each SBE subgroup were compared (II).

18 **Results:** The study included 61 trainees (39 SBE and 22 controls) were recruited. Comparisons between groups
19 showed no significant difference in the quantitative assessment of skill enhancement (difference in the pre- and
20 post-internship OSAUS score: 1.09 ± 0.87 in the SBE group and 0.72 ± 0.98 in the control group) (I). Conversely,
21 the predefined acceptable skill level was reached by a significantly higher proportion of trainees in the 'early' SBE
22 subgroup (74%, compared with 30% in the control group, $P=0.01$)(II).

23 **Conclusions:** The quantitative assessment does not support the existence of long-term benefits from SBE
24 training, although the qualitative assessment confirmed SBE helped to raise the minimal level within a group
25 when embedded in an 'early' stage of a practical course.

26 27 28 **INTRODUCTION**

29
30 Ultrasound is used in many specialties because of its accessibility (1,2) and the ability to study items moving in
31 real time at the patient's bedside (3). These advantages give it a key role in the monitoring and management of
32 pregnancies, especially by fetal biometric measurements (4–9). It is nonetheless operator-dependent (10,11), a
33 characteristic that causes errors and inaccuracy that can account for more than 10% of the variability of some
34 parameter measurements (11). Improving intra- and interobserver reproducibility is therefore a major objective
35 in obstetrics.

36 As we await the automated measurement techniques currently under development (12–14), it remains
37 essential to improve the teaching of fetal biometry to limit the variability and optimize the quality and quality
38 control of these measurements (15). High-fidelity simulation-based education (SBE) in ultrasound appears to be a
39 promising solution (16). Its acceptability and its short-term utility for improving performance have now been
40 established (17–21), although its benefits for the quality and cost of obstetric care have not been clearly
41 confirmed (22). Currently, only Tolsgaard et al. have used an experimental approach to explore these aspects,
42 offering pathways for assessing the medico economical impact of SBE (23). Their approach relies especially on
43 the demonstration of a potential synergistic effect of SBE (16) that could make it possible to accelerate progress
44 for the same quantity of teaching.

45 SBE remains expensive (both the equipment and logistics). It is therefore important to verify how the
46 positive impacts of SBE on progression modify clinical practice in the long term and to assess the importance of
47 the potential synergistic effect as a function of the timing of its use during the learning process.

48 Accordingly, the principal objective of our study was to compare the clinical progress over the long term
49 between students who were or were not exposed to simulation training. The secondary objective was to study

50 the effect of the timing of the exposure (by comparing the different timing of the SBE workshops relative to the
51 course) on the capacity to acquire and maintain the level of competence required.

52

53 **METHODS**

54

55 A systematic, prospective, and continuous collection of data about the practice of ultrasound for measuring fetal
56 biometry during the initial phase of learning these skills took place at the University Hospital of Nancy in the
57 maternity ward between November 1, 2017, and October 31, 2018.

58

59 **Participants**

60 The participants were student sonographers recruited during their initial training in the departments of
61 obstetrics and gynecology, medical gynecology, and midwifery of the University of Lorraine. The inclusion criteria
62 were enrollment in the specialized diploma program in obstetrics and gynecology or medical gynecology (first
63 through eighth semester of their internship-residency, or enrollment in the second cycle of their midwifery
64 studies). The exclusion criteria were: the failure to undergo an initial assessment or absence from the course for
65 a duration exceeding one month during the study period.

66

67 **Description of the METHOD project**

68 This study is part of a university project aimed at optimizing the teaching and learning of ultrasound
69 measurements of fetal biometry (METHOD project: MEasurement TeachHing in Obstetrics Design). French
70 students currently learn ultrasound as apprentices in practical clinical training courses. For the METHOD
71 pedagogical project, workshops with ultrasound simulators were progressively included into this curriculum.
72 None of the participants included in the project had previously received this type of instruction. Three teachers
73 (OM, CB and GA) coordinated this project and ran the clinical course. During the courses, students were
74 supervised by experienced clinicians (obstetricians or midwives). The simulation workshop was developed and
75 led by an instructor trained in teaching by simulation (GA).

76

77

78

79 - Theoretical instruction

80 Before the clinical training course, all participants attended in person a two-hour theoretical presentation, the
81 same for all, about the methodology and practice of ultrasound fetal biometry.

82

83

84 - Pedagogical program of the simulation workshops

85 As recommended by Tolgaard et al., the workshops were conducted in groups of two participants. During a
86 single 3-hour workshop with a simulator (Simbionix US Mentor®), the participants practiced a series of
87 increasingly difficult exercises under the constant supervision of the teacher (GA). The program included in
88 particular exercises of hand-eye coordination and of knobology (24), as well as the performance of fetal
89 biometry.

90

91 - Pre- and post-course quantitative evaluations

92 All participants included in the study were assessed by an OSAUS score before and after the course (Fig 1). The
93 conditions of a stand-alone examination were reproduced with pregnant women (15–41 weeks of pregnancy)
94 who volunteered, and each participant was assessed individually by one of the instructors (OM, CB or GA) as they
95 took ultrasound fetal biometry measurements. The METHOD OSAUS score (customized for fetal biometry) was
96 used to rate their skill level from 0 through 5 (25). The examination could not exceed 10 minutes, and no
97 supervision or suggestions were offered during it. The pre-workshop assessments were scheduled early during or
98 before the course and always before the simulator workshop (double-blinded assessment, both instructor and
99 participant blinded about exposure to the simulation). The post-course assessments were scheduled after the

100 last day of the course. To limit the potential evaluation bias associated with knowledge of the student's exposed-
101 unexposed status for the post-course OSAUS scores, these were assessed by four clinicians not involved in the
102 study (single-blinding).

103
104 - Continuous qualitative assessment

105 An online platform (the METHOD Logbook) was available to participants to enable the continuous collection of
106 data about the instruction. It enabled participants to report in real time the results of the fetal biometry
107 measurements they performed themselves during their training course (head circumference - HC, abdominal
108 circumference - AC, femur length – FL, and estimated fetal weight – EFW). These examinations were
109 systematically verified by an experienced clinician. For each examination performed, the students recorded both
110 their set of measurements and that of their supervising clinician in the METHOD logbook. The METHOD Logbook
111 platform determined the validity of each parameter measured, based on the difference between the student's
112 measurements and those of the supervisor, that is, on the agreement of the student's and supervisor's
113 measurements, with the latter considered as the reference measurements. Validation depended on an
114 interoperator variability less than 0.8 of the Z-score for the EFW and less than 0.8, 0.7 and 0.6 of the Z-scores for
115 FL, AC and HC, respectively. These cutoffs were defined by taking into account the variability inherent to each
116 biometric parameter. These differences in variability are explained by their different levels of technical difficulty
117 and the different distributions for each type of measurement (11).

118 Each time students completed a set of measurements and entered it into the Logbook, they received
119 feedback (for each parameter measurement and for EFW), presented as an LC–CUSUM (Learning Curve Using
120 Cumulative Sum control charts) (26–28). The display setting for the METHOD Logbook showed a rising curve for
121 each validated examination (strong agreement with the reference measurement) and a falling curve for each
122 non-validated measurement (low agreement with the reference measurement) (Figure 2).

123
124
125 **Study plan and procedures**

126
127 - Timing of exposure

128 The participants were randomly allocated to the control (no simulation workshop exposure during the
129 observation period) and simulation (exposure to SBE) groups. The SBE workshops were planned randomly at
130 different times on the course schedule, with the participants distributed within both the control and simulation
131 (SBE exposure) groups. Within the simulation group, participants were distributed into three subgroups, the
132 'before' group (exposure before the course), the 'early 'course group (interval between the beginning of the
133 course and exposure < 2 months) and the 'late' course group (interval between exposure and the end of the
134 course < 2 months); these groups defined different exposure timing patterns (Figure 1).

135
136 - Endpoints

137 The primary endpoint was the percent delta (that is, percent change) in progression during the clinical training
138 course, defined by the difference between the pre- and post-course OSAUS scores. We hypothesized that the
139 mean change in progression differed between the simulation and control groups. The secondary endpoint was
140 defined by the proportion of participants exposed to SBE able to acquire a threshold skill level according to their
141 exposure timing pattern. This threshold level was defined by the validation of five consecutive EFW in the
142 METHOD Logbook.

143
144
145 **Statistical analysis:**

146
147 - Calculation of study size

148 In the usual learning conditions of initial training, the student's skill level is expected to progress. When this
149 progression is assessed by a score, a delta/change between the scores obtained before and after the training
150 expresses this progression (corresponding to the progression delta described above).

151 For the METHOD project, the expected progression was estimated from data from preliminary studies
152 assessing the validity and predictive value of the METHOD OSAUS score (25). Based on these studies, the skill
153 level in fetal biometry of a beginner in obstetric ultrasound corresponded to OSAUS scores ranging between 1.5
154 ± 0.4 and 1.8 ± 0.7 . After practice experience for longer than 6 months or > 20 supervised examinations, the
155 operators reached an intermediate skill level corresponding to scores of 3.3 ± 0.6 and 3.3 ± 0.8 respectively
156 (25,29).

157 Given these data, we estimated the expected progression of participants without SBE to be at least 1.25
158 points on the OSAUS. Our hypothesis was that SBE would enable greater progress for the exposed participants
159 (delta 1.75 points on the OSAUS). We calculated that we required a minimum of 20 participants in the exposed
160 and unexposed groups to validate this hypothesis. In the SBE group, at least 8 participants were necessary in
161 each subgroup to enable a comparative qualitative analysis between the different timing patterns. This required
162 that we increase the size of the SBE group. [The modest headcount within subgroups did not allow to lead a](#)
163 [relevant quantitative assessment for each timing pattern.](#)

164
165 - Statistical tests
166 The quantitative data are expressed as their means and standard deviations, with their range (minimum and
167 maximum). The normality of the data was tested by the Shapiro Wilk test. Intergroup differences were studied
168 by Student's t test (with equal variance). For the qualitative variables, Chi-square or Fisher tests were used
169 according to the distribution of the data. A P value less than 0.05 was considered significant.

170 The statistical analysis was performed with R software, version 3.6.2 (2019-12-12).

171
172 - Ethics and reporting of the results
173 This study was reported to the CNIL (no. A2017-3). It is presented in accordance with the guidelines for Health
174 Care Simulation Research and CONSORT (extension for pragmatic trials).

175
176

177 RESULTS

178
179 **Participants:**
180 The study included 61 participants (36 residents and 25 midwifery students) (Figure 3). [Details of the cohort](#)
181 [demographic characteristics are available in Table 1.](#) The mean observation period was 5 months and 15 days \pm
182 62 days.

183 After the allocation of the minimum number of individuals within each group and subgroup, there remained
184 18 additional participants. These participants were distributed between the control group, and the early and late
185 course subgroups (3, 14 and 1 additional participant, respectively). The distribution of the additional participants
186 favored, for pedagogical reasons, early access to SBE.

187 At the beginning of the study, the pre-course OSAUS scores were similar in the exposed and control groups
188 (2.61 vs 2.7 respectively, $P=0.83$) (Table 2).

189
190
191 Quantitative assessment
192 The final post-course OSAUS score and the global delta for progression did not differ significantly between the
193 SBE and control groups (Table 2). In comparison with the estimates from preliminary studies, the initial skill level
194 of the participants was better than expected (2.63 ± 1.01 versus 1.25) and the global progression lower ($0.95 \pm$
195 0.92 versus 1.75).

196
197

198
199 Qualitative assessment
200 The METHOD Logbook platform was used by 51 (82%) participants who generated LC-CUSUM, a percentage that
201 did not differ between the SBE (79%) and control (87%) groups (Table 3). The proportion of participants able to
202 reach the expected skill level (five consecutive validated examinations) was significantly higher in the 'early'
203 group (74% vs 30% in the control group, $P < 0.05$).

204 The mean number of examinations necessary to reach this skill level ranged from 8.2 ± 3.5 to 13.5 ± 4.8
205 and did not differ significantly between the groups (Table 3).
206
207
208
209
210
211

212 **DISCUSSION**

213 **Main findings and interpretation of results**

214 At the end of a long observation period, the participants exposed to the simulation workshop did not progress
215 significantly more than the control group. Nonetheless, the effect of the simulation might have been masked,
216 especially because the participants' initial level was higher than expected, which left less margin to observe
217 progression.
218

219 Moreover, the proportion of participants able to reach the skill level expected was significantly higher in
220 the subgroup of participants exposed to SBE near the beginning of the course. This result is crucial for
221 understanding the impact of SBE on clinical practice. Standardizing the skill level guarantees a minimum
222 threshold level of practice and is thus a major step toward improving the global quality of care. The number of
223 practitioners able to reach this threshold skill level thus constitutes a more useful outcome measure than the
224 mean or individual progression for assessing the potential clinical impact of a training activity.

225 These differences observed between the different timing patterns suggest the existence of different
226 learning processes. This is probably explained by the optimized use of SBE by the participants who faced these
227 clinical situations around the same time. This observation raises the question of the relevance of the "never the
228 first time on the patient" adage, which was initially an argument for the development of simulation in the
229 training of health professionals (HAS report 2012) (30). If this approach remains ethically justified for invasive
230 procedures, it fails to incorporate the benefit of learning from mistakes, as described by L. Dyre et al (31).
231 Learning from errors probably helps to optimize SBE, with students using the simulator to resolve problems
232 encountered in clinical practice. This mechanism appears to optimize on the one hand training on the simulator,
233 and on the other hand, increase the ability to transfer the skills acquired on the simulator to clinical practice.
234 Among the skills used for ultrasound, hand-eye coordination and visual-cognitive skills (ability to mentally
235 represent an ultrasound diagnosis) may comprise the major elements that explain this synergistic effect (16). A
236 lower effect of learning from mistakes at the end of the course and a short interval between SBE and the final
237 assessment may explain the less good performance observed in the late-course group.
238

239 **Strengths and limitations**

240 The substantial number of participants combined with the duration of the observation in the METHOD study is
241 one of its strengths. Only a single other study had a similar sample size (32), and for all of the observations
242 concerning abdominal ultrasound (33), the mean observation periods were less than a month, compared with
243 more than five months in ours.

244 This study also represents the first assessment of the effect of the timing of the exposure (by comparing the
245 different timing of the SBE workshops relative to the course) on the capacity to acquire and maintain the level of
246 competence required. The simultaneous use of quantitative outcome measures (score) and qualitative measures
247 (LC-CUSUM), all relevant to clinical practice and not to performance on the simulator, is the second major

248 strength of this study. This methodological choice enabled us to avoid the known bias that results from the
249 extrapolation to clinical settings of performance observed with a simulator (21).

250 The improvement of the performance of participants exposed to SBE as well as the clinical training
251 course is due partly to the supplementary effect induced by the addition of training support for the SBE group.
252 This inequality bias in principle is a limitation attributable to the fact that all studies up to now have simply
253 assessed the interest of SBE, with no experiments proposing randomization between SBE alone and clinical
254 practice alone (34). Because learning ultrasound is indissociable from access to clinical practice, these conditions
255 cannot ethically be reproduced outside of an experimental context; this bias is thus inevitable. It should
256 nonetheless be noted that the experimental and short-term comparison between SBE alone (i.e. without clinical
257 practice) and practical training alone confirmed that SBE augments the students' progress (35).

258 The methods and the pedagogical process used in the METHOD study can be implemented easily in other
259 medical schools and institutions. But the single-center nature of the data collection limits the possibility of
260 generalizing the results, especially because of possible biases associated with local particularities in education
261 methods and in the practice of obstetric ultrasound.

262

263 **Practical applications**

264 The experience acquired with the METHOD project also provides valuable information about the practical use of
265 the OSAUS score. At the conclusion of the study, the investigators identified in particular pitfalls related to the
266 difficulty in quantifying the participants' progress with a delta between two scores: the magnitude of this delta
267 depends principally on the operator's initial level rather than on his or her weekly progress. This effect is
268 explained by the nonlinear nature of the progress (more rapid and marked in the early learning phase). Given the
269 intrinsic difficulty of obstetric ultrasound, progression later on appears slower and more difficult to quantify, and
270 it is characterized by a lower rise in the score. This characteristic explains the difficulty in showing a difference in
271 progression in the METHOD study due to the participants' initial skill levels, which were higher than anticipated.
272 In the future, this experience will lead us to limit the use of changes in OSAUS scores to the observation of
273 beginning operators and the comparison of groups with comparable initial levels and similar observation periods.

274 The continuous massive expansion of simulation in training health professionals calls for the
275 development of scientific tools making it possible to reinforce the practical applicability of experimental results
276 and the relevance of the studies. The publication of guidelines for Health Care Simulation Research (36)
277 constitutes a first step, but the question of the cost-benefit relation for SBE must be explored in greater depth
278 (37,38). In the light of the results of the METHOD study, this investigation must consider the timing of the SBE. At
279 the same time, the systematization of the outcome measures used to assess the impact of simulation should
280 make it possible to optimize comparisons between the studies as well as the work of meta-analysis. This
281 procedure must be based on a Delphi procedure (39) to define a set of homogeneous outcomes enabling the
282 assessment and comparison of procedures involving SBE. The possibility of concrete application of these results
283 to our educational practices depends on these final two stages.

284

285

286

287

288

289

290

291

292

293

294

Conclusion:

The assessment of the learning phase by the METHOD OSAUS score does not allow us to conclude that
there is a difference in the mean progression between participants exposed to SBE and the control group. On the
other hand, the association of SBE with clinical practice at the onset of the learning phase enabled a significantly
higher number of participants to reach the required skill level, compared with the control group.


ACKNOWLEDGMENT


The authors would like to thank the investigators, coordinators, participants and patients for their involvement in
the study. We also thank Jo Ann Cahn for ensuring the manuscript final translation and proofreading.

- 295
296
297
298 1. Jenssen C, Gilja OH, Serra AL, Piscaglia F, Dietrich CF, Rudd L, Sidhu P. European Federation of Societies
299 for Ultrasound in Medicine and Biology (EFSUMB) Policy Document Development Strategy - Clinical Practice
300 Guidelines, Position Statements and Technological Reviews. *Ultrasound Int Open* 2019; **5**: E2-10.
- 301 2. American Institute of Ultrasound in Medicine. AIUM Practice Guideline for the Performance of Obstetric
302 Ultrasound Examinations. *Journal of Ultrasound in Medicine* 2013; **32**: 1083-101.
- 303 3. Moore CL, Copel JA. Point-of-Care Ultrasonography. *New England Journal of Medicine* 2011; **364**:
304 749-57.
- 305 4. Massoud M, Duyme M, Fontanges M, Combourieu D. Courbe d'estimation de poids foetal 2014 par le
306 Collège français d'échographie foetale (CFEF). *J Gynecol Obstet Biol Reprod* 2016; **45**: 80-5.
- 307 5. Papageorgiou AT, Ohuma EO, Altman DG, Todros T, Ismail LC, Lambert A. International standards for
308 fetal growth based on serial ultrasound measurements: the Fetal Growth Longitudinal Study of the intergrowth-
309 21st Project. *Lancet* 2014; **384**: 869-79.
- 310 6. Salomon LJ, Alfirevic Z, Berghella V, Bilardo C, Hernandez-Andrade E, Johnsen SL, et al. Practice
311 guidelines for performance of the routine mid-trimester fetal ultrasound scan. *Ultrasound Obstet Gynecol* 2011;
312 **37**: 116-26.
- 313 7. Henriksen T. The macrosomic fetus: a challenge in current obstetrics. *Acta Obstet Gynecol Scand* 2008;
314 **87**: 134-45.
- 315 8. Boulvain M, Senat M-V, Perrotin F, Winer N, Beucher G, Subtil D, et al. Induction of labour versus
316 expectant management for large-for-date fetuses: a randomised controlled trial. *Lancet* 2015; **385**: 2600-5.
- 317 9. Chien LY, Whyte R, Aziz K, Thiessen P, Matthew D, Lee SK, et al. Improved outcome of preterm infants
318 when delivered in tertiary care centers. *Obstet Gynecol* 2001; **98**: 247-52.
- 319 10. Benacerraf BR, Minton KK, Benson CB, Bromley BS, Coley BD, Doubilet PM, et al. Proceedings: Beyond
320 Ultrasound First Forum on Improving the Quality of Ultrasound Imaging in Obstetrics and Gynecology. *J*
321 *Ultrasound Med* 2018; **37**: 7-18.
- 322 11. Sarris I, Ioannou C, Chamberlain P, Ohuma E, Roseman F, Hoch L, et al. Intra- and interobserver
323 variability in fetal ultrasound measurements. *Ultrasound Obstet Gynecol* 2012; **39**: 266-73.
- 324 12. Hur H, Kim YH, Cho HY, Park YW, Won H-S, Lee M-Y, et al. Feasibility of three-dimensional
325 reconstruction and automated measurement of fetal long bones using 5D Long Bone. *Obstet Gynecol Sci* 2015;
326 **58**: 268-76.
- 327 13. Ambroise Grandjean G, Hossu G, Banasiak C, Ciofolo-Veit C, Raynaud C, Rouet L, et al. Optimization of
328 Fetal Biometry With 3D Ultrasound and Image Recognition (EPICEA): protocol for a prospective cross-sectional
329 study. *BMJ Open* 2019; **9**: e031777.
- 330 14. Ambroise Grandjean G, Hossu G, Bertholdt C, Noble P, Morel O, Grangé G. Artificial intelligence
331 assistance for fetal head biometry: Assessment of automated measurement software. *Diagn Interv Imaging*
332 2018; **99**: 709-16.
- 333 15. Sarris I, Ioannou C, Dighe M, Mitidieri A, Oberto M, Qingqing W, et al. Standardization of fetal
334 ultrasound biometry measurements: improving the quality and consistency of measurements. *Ultrasound Obstet*
335 *Gynecol* 2011; **38**: 681-7.
- 336 16. Tolsgaard MG. A multiple-perspective approach for the assessment and learning of ultrasound skills.
337 *Perspect Med Educ* 2018; **7**: 211-3.
- 338 17. Chalouhi GE, Bernardi V, Ville Y. Ultrasound simulators in obstetrics and gynecology: state of the art.
339 *Ultrasound Obstet Gynecol* 2015; **46**: 255-63.
- 340 18. Le Lous M, Tsatsaris V, Tesnière A, Grangé G. Improving students' ability to perform a standardized fetal
341 biometry plane using ultrasound simulators. *J Gynecol Obstet Hum Reprod* 2017; **46**: 439-443

- 342 19. Gueneuc A, De Garnier J, Dommergues M, Rivière M, Ville Y, Chalouhi GE. Impact de l'intégration de la
343 simulation en échographie obstétricale dans le cursus des étudiants sages-femmes. *Gynécologie Obstétrique*
344 *Fertilité & Sénologie* 2019; **47**: 776-782
- 345 20. Madsen ME, Konge L, Nørgaard LN, Tabor A, Ringsted C, Klemmensen AK, et al. Assessment of
346 performance measures and learning curves for use of a virtual-reality ultrasound simulator in transvaginal
347 ultrasound examination. *Ultrasound Obstet Gynecol* 2014; **44**: 693-9.
- 348 21. Tolsgaard MG, Madsen ME, Ringsted C, Oxlund BS, Oldenburg A, Sorensen JL, et al. The effect of dyad
349 versus individual simulation-based ultrasound training on skills transfer. *Med Educ* 2015; **49**: 286-95.
- 350 22. Kirkpatrick D. Great Ideas Revisited. Techniques for Evaluating Training Programs. Revisiting Kirkpatrick's
351 Four-Level Model. *Training and Development*. 1996; **50**: 54-9.
- 352 23. Tolsgaard MG, Tabor A, Madsen ME, Wulff CB, Dyre L, Ringsted C, et al. Linking quality of care and
353 training costs: cost-effectiveness in health professions education. *Med Educ* 2015; **49**: 1263-71.
- 354 24. Enriquez JL, Wu TS. An introduction to ultrasound equipment and knobology. *Crit Care Clin*. janv 2014;
355 **30**:25-45, v.
- 356 25. G. Ambroise Grandjean, P.Gabriel, G. Hossu, S. Zuily, O. Morel, P. Berveiller. Training in fetal ultrasound
357 biometry : prospective assesment of Objective Structured Assessment of Ultrasound Skills (OSAUS) efficiency.
358 *submitted to Journal of Gynecology Obstetrics and Human Reproduction* 2020.
- 359 26. Biau DJ, Williams SM, Schlup MM, Nizard RS, Porcher R. Quantitative and individualized assessment of
360 the learning curve using LC-CUSUM. *Br J Surg* 2008; **95**: 925-9.
- 361 27. Biau DJ, Porcher R, Salomon LJ. CUSUM: a tool for ongoing assessment of performance. *Ultrasound*
362 *Obstet Gynecol* 2008; **31**: 252-5.
- 363 28. Weerasinghe S, Mirghani H, Revel A, Abu-Zidan FM. Cumulative sum (CUSUM) analysis in the
364 assessment of trainee competence in fetal biometry measurement. *Ultrasound Obstet Gynecol* 2006; **28**:
365 199-203.
- 366 39. Tolsgaard MG, Ringsted C, Dreisler E, Klemmensen A, Loft A, Sorensen JL, et al. Reliable and valid
367 assesment of ultrasound operator competence in obstetrics and gynecology. *Ultrasound Obstet Gynecol* 2014;
368 **43**: 437-43.
- 369 30. Haute Autorité de Santé. Simulation en santé 2012. [Disponible en ligne: [https://www.has-
370 sante.fr/jcms/c_930641/fr/simulation-en-sante](https://www.has-sante.fr/jcms/c_930641/fr/simulation-en-sante)]. Cité le 1 juin 2020.
- 371 31. Dyre L, Tabor A, Ringsted C, Tolsgaard MG. Imperfect practice makes perfect: error management
372 training improves transfer of learning. *Medical Education* 2017; **51**: 196-206.
- 373 32. Andreasen LA, Tabor A, Nørgaard LN, Ringsted C, Sandager P, Rosthøj S, et al. Multicenter randomized
374 trial exploring effects of simulation-based ultrasound training on obstetricians' diagnostic accuracy: value for
375 experienced operators. *Ultrasound Obstet Gynecol* 2020; **55**: 523-9. 34.
- 376 33. Le Lous M, Tsatsaris V, Tesnière A, Grangé G. Improving students' ability to perform a standardized
377 foetal biometry plane using ultrasound simulators. *J Gynecol Obstet Hum Reprod* 2017; **46**:439-443.
- 378 34. Chalouhi GE. Re: Sustained effect of simulation-based ultrasound training on clinical performance: a
379 randomized trial. *Ultrasound Obstet Gynecol* 2015; **46**: 264-5.
- 380 35. Chao C, Chalouhi GE, Bouhanna P, Ville Y, Dommergues M. Randomized Clinical Trial of Virtual Reality
381 Simulation Training for Transvaginal Gynecologic Ultrasound Skills. *J Ultrasound Med* 2015; **34**: 1663-7.
- 382 36. Cheng A, Kessler D, Mackinnon R, Chang TP, Nadkarni VM, Hunt EA, et al. Reporting Guidelines for
383 Health Care Simulation Research: Extensions to the CONSORT and STROBE Statements. *Simul Health* 2016; **11**:
384 238-48.
- 385 37. Tolsgaard MG, Tabor A, Madsen ME, Wulff CB, Dyre L, Ringsted C, et al. Linking quality of care and
386 training costs: cost-effectiveness in health professions education. *Med Educ*. 2015; 49: 1263-71.
- 387 38. Osborne B, Parange N, Thoirs K. The effectiveness of the use of high fidelity simulators in obstetric
388 ultrasound training: A systematic review. *Australas J Ultrasound Med* 2015; **18**: 107-11.
- 389 39. de Villiers MR, de Villiers PJT, Kent AP. The Delphi technique in health sciences education research. *Med*
390 *Teach* 2005; **27**: 639-43.
- 391

392	
393	Tables & Figures :
394	
395	Table 1 : Demographic characteristics of the trainees cohort.
396	
397	Table 2: Allocation and results of pre- and post-course assessment in the different groups
398	
399	Table 3: Logbook utilization rates and LC-CUSUM (learning curve cumulative sum) analysis
400	
401	Figure 1: Presentation of the different timing of the METHOD cohort participants' simulation exposure and of
402	their assessments.
403	
404	Figure 2: Example of four LC-CUSUM (Learning Curve Using Cumulative Sum control charts) obtained at the end
405	of the course for the measurement of the estimated fetal weight (each curve represents the individual
406	progression of one participant).
407	
408	Figure 3: CONSORT3 flow diagram of enrollments, allocations, follow-up and analysis for METHOD study.
409	





Trainees assessed for eligibility (n=63)

Excluded (n=1)

- Did not attend pre-course lecture and pre-OSAUS¹ assessment

Trainees eligible for random allocation (n=62)

Allocated to "Simulation" group (n=39)

Allocated to "Control" group (n=23)

Excluded (n=1)
• Leave of absence

Lost to follow-up :

- quantitative assessement² (n=0)
- qualitative assessement³ (n=8)

Lost to follow-up :

- quantitative assessement² (n=1)
- qualitative assessement³ (n=4)

Analyzed :

- quantitative assessment (n=39)
- qualitative assessment (n=31)
Excluded from analysis (n=0)

Analyzed :

- quantitative assessment (n=22)
- qualitative assessment (n=19)
Excluded from analysis (n=1)

Table 1 : Demographic characteristics of the trainee cohort

Training curriculum	Mean age ¹	Mean year of medical education ¹	Gender ²	
			Female	Male
Midwife trainees (n=25)	26.2 ± 0,6	4 ± 0	22	1
Residents (n=36)	29.9 ± 2.1	8.8 ± 1.9	29	7

¹ Results are presented as mean ± standard deviation

² Results are presented as absolute numbers

Table 2: Allocation and results of pre- and post-course assessment in the different groups

	Total (n=61)	Control group (n=22)	SBE* group (n=39)	P-value
Pre-OSAUS1	2.63 ± 1.01 [0.4; 4.8]	2.7 ± 1.1 [0.4; 4.2]	2.61 ± 0.97 [0.4; 4.8]	0.83
Post-OSAUS2	3.58 ± 0.72 [1.8; 4.8]	3.38 ± 0.69 [1.8; 4.8]	3.7 ± 0.72 [1.8; 4.8]	0.1
Delta3	0.95 ± 0.92 [-1; 3.2]	0.71 ± 0.98 [-1; 2.6]	1.09 ± 0.72 [-1; 3.2]	0.41

Results are reported as mean ±SD (range).

¹OSAUS, pre-course assessment via OSAUS METHOD score (Objective Assessment of Ultrasound Skills – Measurement Teaching in Obstetrics Design)

²Post-OSAUS, post-course assessment via OSAUS METHOD score (Objective Assessment of Ultrasound Skills – Measurement Teaching in Obstetrics Design)

³ Difference between post-course and pre-course OSAUS scores

*Simulation based education (in ultrasound)

Table 3: Logbook utilization rates and LC-CUSUM (learning curve cumulative sum) analysis

	Control group (n=21)	SBE* group (n=39)	'Before' (n=8)	'Early' stage (n=22)	'Late' stage (n=9)
Logbook users ¹	20 (95)	31 (79)	4 (50)	19 (86)	8 (89)
Effective skill level reached ²	6 (30)	17 (55)	2 (50)	14 (74)	1 (13)
Numbers of exams required to reach the skill ³	10.7 ± 4.5 [5; 16]	9.5 ± 4.3 [5; 18]	11 ± 2.8 [9; 13]	8.6 ± 4 [5; 18]	18 /

*Simulation based education

¹ Results are presented as absolute numbers (percentages)

² Effective skill level is considered as reached when 5 consecutive correct exams were performed (correct exam corresponding to a difference less than 0.8 Z-score between trainee and supervisor for estimated fetal weight). Results are presented in absolute numbers (percentage)

³ Results are presented as means ±SD [range].