

An investigation of the fate of petroleum contaminants (LNAPL) under fluctuating groundwater table levels induced by climate change

Amélie Cavelan, Fabrice Golfier, Catherine Lorgeoux, Stéfan Colombano, Jacques Deparis, Hossein Davarzani, Anne-Julie Tinet, Constantin Oltean, Noële Enjelvin, Pierre Faure

▶ To cite this version:

Amélie Cavelan, Fabrice Golfier, Catherine Lorgeoux, Stéfan Colombano, Jacques Deparis, et al.. An investigation of the fate of petroleum contaminants (LNAPL) under fluctuating groundwater table levels induced by climate change. DEEPSURF International Conference 2021, Oct 2021, Nancy, France. , 2021. hal-03445662

HAL Id: hal-03445662 https://hal.univ-lorraine.fr/hal-03445662

Submitted on 24 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



DEEPSURF 2021



TOWARD SOLUTIONS FOR ENERGY AND ECOLOGICAL TRANSITION

AN INVESTIGATION OF THE FATE OF PETROLEUM CONTAMINANTS (LNAPL) UNDER FLUCTUATING GROUNDWATER TABLE LEVELS INDUCED BY CLIMATE CHANGE

Amélie Cavelan 1*, Fabrice Golfier 2, Catherine Lorgeoux 2, Stéfan Colombano 3, Jacques Deparis 3, Hossein Davarzani 3, Anne-Julie Tinet 2, Constantin Oltean², Noële Enjelvin⁴, Pierre Faure¹



¹Université de Lorraine, CNRS, LIEC, Nancy, France, ²Université de Lorraine, CNRS, GeoRessources, Nancy, France, ³BRGM, Orléans, France, ⁴Université de Lorraine, INRAE, LSE, GISFI, Vandoeuvre-lès-Nancy, France. *Corresponding author: amelie.cavelan@univ-lorraine.fr

1. LIGHT-NON-AQUEOUS-PHASE-LIQUIDS (LNAPLs)?

- o Derived from petroleum hydrocarbons (gasoline, diesel fuel...)
- Viscous liquids with a density < 1 g/cm3 and a lifetime > 10 yrs
- Composition: mainly aliphatic hydrocarbons and BTEX
- o 40% of soil and groundwater contaminations

VOC, CO, CH How does the intensity of GWL fluctuations affect LNAPLs? Biodegradation Rates Biodegradation Dissolution LNAPL Dissolution **LNAPL** longevity **Dissolved LNAPL** *Tuctuations* Groundwater flow intensity Future LNAPL behavior? Gas emissions +++ USZ **USZ** VOC, CO₂, CH₄ Gas pushed to the surface Residual LNAPL trients re-Biodegradation \ Spreading of Spreading of LNAPLS LNAPLS

Seasonal/pump-induced GWL variations favor the LNAPLs mobilization and release

Trapped LNAPL+ air + water

Groundwater flow

Fig.1. Inlfuence of the groundwater table level (GWL) fluctuations on LNAPL mobilization and release processes. (i) initial, (ii) low (dry season), (iii) high water table level (wet season). VOC: volatil organic carbon. SZ: saturated zone. USZ: unsaturated zone.

2. CLIMATE CHANGE, GROUNDWATER AND LNAPLS

THE PRESENCE OF GROUNDWATER TABLE FLUCTUATIONS:

- o favors the vertical spreading of the LNAPLs across the soil (Fig.1)
- o favors the release of the LNAPL compounds the in the air and groundwater
- o favors the natural attenuation of the LNAPL source/ reduces the LNAPL longevity




Fig. 2. Measured and projected increase of the surface temperature, extreme events frequency, groundwater withdrawal and level variations intensity from 1900 to 2100. Data from the IPCC (2021), the European Environment Agency, and Wada (2016).

CLIMATE CHANGE MAY:

3. OUR WORK

Gas analysis: fluxes

and composition

(IPCC predictions)

- o change precipitation and groundwater recharge intensity and frequency (Fig.2)
- o increase the intensity and frequency of the extreme climatic events.

2 columns = 2 scenarios of


groundwater table fluctuations

- o increase groundwater use/decrease mean groundwater table levesl.
- o increase the amplitude of seasonal and pump-induced groundwater table level variations.

WHAT WE NEED TO KNOW:

The effect of increased groundwater level fluctuations on the LNAPLs' mobilization, release and natural attenuation rates, and the consequences for short/long-term groundwater and air quality at contaminated sites.

Lysimetric scale (2m³) 150 -"Current" scenario 100 100 120 "Extreme" scenario 100 100 120 20 Climatic scenarios 1yr = 2 months









Fig. 3. Lysimetric columns (2m³) used to simulate the effect on LNAPL of two scenarios of precipitations and groundwater level variations over 14 months.

Laboratory scale



2D tanks



1D centimetric columns

THE GOALS:

Dissolved

LNAPL

Groundwater flow

- o Improve the monitoring techniques of LNAPLs contaminated sites in combining different in-situ and ex-situ monitorging techniques
- o Improve the understanding of LNAPLs behavior under various climatic conditions (current or future)

WHAT WE MONITOR?

Lysimetric columns

- o Gaseous surface emissions (flux, composition, µGC)
- o The LNAPL saturation distribution in soil (electrical resistivity and permittivity) o Other soil physical-chemical properties (pH, RedOx, water tension, temperature)
- o The concentration and composition of the dissolved LNAPL phase (suction cups,
- SPE, GC/MS, TPH, DOC, FTIR)

IN THE LABORATORY:

- o Calibrations of in-situ signals (tensiometers, permittivity, resistivity) under controlled saturations of LNAPL/water/air (2 tanks/1D colunms)
- o Numerical modeling of the LNAPL transport and mobilization.

effects on the fate of LNAPL contaminated sites.

(past regional records)

o Optimization of analytical protocols for contaminated water samples.

IN THE COMING MONTHS:

o Application in field for the real-time monitoring of a contaminated site. o Upscaling of modelisation at the field scale for prediction of climate change













