



HAL
open science

Cross-talk between non-alcoholic fatty liver disease and cardiovascular disease: Implications for future trial design

Pierpaolo Pellicori, Muthiah Vaduganathan, João Pedro Ferreira, Faiez Zannad, Arun J Sanyal

► To cite this version:

Pierpaolo Pellicori, Muthiah Vaduganathan, João Pedro Ferreira, Faiez Zannad, Arun J Sanyal. Cross-talk between non-alcoholic fatty liver disease and cardiovascular disease: Implications for future trial design. *Endocrinology, Diabetes & Metabolism*, 2021, pp.101281. 10.1016/j.diabet.2021.101281 . hal-03469195

HAL Id: hal-03469195

<https://hal.univ-lorraine.fr/hal-03469195v1>

Submitted on 7 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cross-Talk Between Non-Alcoholic Fatty Liver Disease and Cardiovascular Disease: Implications for Future Trial Design

Short Title: NAFLD and Cardiovascular Disease.

¹Pierpaolo Pellicori MD, ²Muthiah Vaduganathan, MD MPH ^{3,4} João Pedro Ferreira, MD PhD
³Faiez Zannad, MD PhD ⁵Arun J. Sanyal, MD

¹Robertson Institute of Biostatistics and Clinical Trials Unit, University of Glasgow, University Avenue, Glasgow, G12 8QQ.

²Division of Cardiovascular Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts.

³Centre d'Investigation Clinique Plurithématique Pierre Drouin-INSERM CHU de Nancy, Nancy, France & Université de Lorraine, FCRIN INI-CRCT (Cardiovascular and Renal Clinical Trialists) Network, France

⁴Cardiovascular R&D Centre (UnIC), Faculty of Medicine, University of Porto, Porto, Portugal.

⁵Dept. of Internal medicine, Virginia Commonwealth University School of Medicine, Richmond, VA 23298

Word count: 6,723

Corresponding author: Dr Pierpaolo Pellicori

Robertson Institute of Biostatistics and Clinical Trials Unit, University of Glasgow, University Avenue, Glasgow, G12 8QQ, UK. Tel: +44 0 141 330 4744; Fax: +44 0 141 330 5094

Email: pierpaolo.pellicori@glasgow.ac.uk

Conflict of interest: Arun J. Sanyal: President of Sanyal Bio. He has stock options in Genfit, Akarna, Tiziana, Indalo, Durect Inversago and Galmed. He has served as a consultant to Astra Zeneca, Nitto Denko, Conatus, Nimbus, Salix, Tobira, Takeda, Janssen, Gilead, Terns, Birdrock, Merck, Valeant, Boehringer-Ingelheim, Bristol Myers Squibb, Lilly, Hemoshear, Zafgen, Novartis, Novo Nordisk, Pfizer, Exhalenz and Genfit. He has been an unpaid consultant to Intercept, Echosens, Immuron, Galectin, Fractyl, Syntlogic, Affimune, Chemomab, Zydus, Nordic Bioscience, Albireo, Prosciento, Surrozen. His institution has received grant support from Gilead, Salix, Tobira, Bristol Myers, Shire, Intercept, Merck, Astra Zeneca, Malinckrodt, Cumberland and Novartis. He receives royalties from Elsevier and UptoDate. Virginia Commonwealth University, where he is a full time employee, also has ownership interests in Sanyal Biotechnology. Muthiah Vaduganathan has received research grants and/or serves on advisory boards for American Regent, Amgen, AstraZeneca, Baxter Healthcare, Bayer AG, Boehringer Ingelheim, Cytokinetics, and Relypsa, and has participated on clinical endpoint committees for studies sponsored by Galmed, Novartis, and the NIH. Pierpaolo Pellicori has received travel support from Boehringer Ingelheim and consultation fees from Novartis. JPF does not report any conflict of interest. Faiez Zannad is the founder of CardioRenal and the Global Cardiovascular Clinical Trialist Forum.

Abstract

The natural history of non-alcoholic fatty liver disease (NAFLD) is still not fully elucidated. Patients with NAFLD have a low risk of liver complications, unless substantial liver fibrosis has developed. On the other hand, NAFLD has been linked with excess metabolic and cardiovascular complications. Therapies targeting common pathways may benefit both NAFLD and underlying cardiometabolic risk. Therefore, there is a rationale for considering cardiovascular endpoints in the context of NAFLD trials and, vice-versa, to consider the concomitant presence of NAFLD in drug development for cardiometabolic disorders. This manuscript provides a framework for consideration for future trials examining the inter-relationship between cardiovascular disease and NAFLD.

Key words: NAFLD, NASH, trials, cardiovascular, endpoints

Introduction

Excess fat in the liver, or steatosis, in the absence of a history of pathological alcohol consumption, characterizes a heterogeneous metabolic condition referred to as non-alcoholic fatty liver disease (NAFLD)¹. NAFLD has two principal phenotypes: (a) a nonalcoholic fatty liver (NAFL) which is characterized by excess macrovesicular steatosis alone or in combination with minimal inflammation, and (b) nonalcoholic steatohepatitis (NASH) which is defined by the presence of macrovesicular steatosis, lobular inflammation and hepatocellular ballooning typically in a centrilobular distribution². The principal risk factors for NAFLD include obesity, type 2 diabetes mellitus (T2DM), hypertension, and hypertriglyceridemia³, the same risk factors that are also associated with cardio-metabolic disease and chronic kidney disease. The development of NAFLD, especially NASH, further leads to increased LDL-cholesterol due to continued synthesis of cholesterol in the liver despite increased hepatic free cholesterol concentrations⁴. It is therefore unsurprising that NAFLD and cardiometabolic disease frequently co-exist and the epidemiology of NAFLD closely tracks with that of obesity and T2DM⁵⁻⁷. Furthermore, cardio-metabolic events are the most common cause of mortality in patients with NAFLD especially before the development of advanced fibrosis^{8,9}.

While there is substantial literature on the importance of cardio-metabolic outcomes in patients with NAFLD^{10,11} and serious but silent liver disease in the T2DM population, the care of these patients is often siloed and suboptimal, ignoring the totality of the disease burden on the patient (Table 1). This is in part due to the lack of suitable, easy-to-deploy diagnostics and approved therapies for NAFLD. There is also an unmet need to increase awareness of NAFLD within

populations at risk for cardio-metabolic outcomes. NAFLD may independently drive cardiometabolic risk, especially when significant liver fibrosis has developed¹². Indeed, the development of progressive NAFLD includes metabolic perturbation, lipid overload, cell stress, inflammation and fibrosis, which are also seen in cardio-metabolic disorders¹³. It is plausible that therapies targeting common pathways may yield benefits for both NAFLD and underlying cardio-metabolic disease providing strong rationale for consideration of NAFLD-related endpoints in the context of clinical trials for cardiovascular disease.

Many drugs are also metabolized in the liver and the pharmacokinetic profile of such agents could be substantially altered by the presence of NAFLD with advanced fibrosis^{14,15}. The possibility of unexpected toxicity due to altered drug exposure to the liver leading to drug-induced liver injury is another reason why it is important to consider the concomitant presence of NAFLD in drug development for cardiometabolic disorders.

In light of these intersecting issues, we provide a framework for consideration for future trials examining the inter-relationship between cardiovascular disease and NAFLD.

Identifying NAFLD in Clinical Practice

NAFLD is widely prevalent and approximately 25-30% of the adult population have hepatic steatosis, worldwide⁹. Of these, about 20-25% might have NASH¹⁶. The clinical outcomes in those with either NAFL or NASH with early stage disease are mainly cardio-metabolic and cancer related, whereas liver related outcomes become more prominent only when stage 2

fibrosis or higher stages of disease develop^{12,17}. It has also become clear that disease activity drives fibrosis and that, even without therapeutic intervention, both disease activity and fibrosis can wax and wane¹⁸⁻²⁰. From a hepatological perspective, it is therefore important to identify those with NASH, especially those with high disease activity and fibrosis stage 2 or higher. The identification of those with cirrhosis, who can be clinically silent until a complication such as variceal hemorrhage or hepatocellular cancer occur, is also important.

The diagnostic approach for NAFLD is rapidly evolving although there is not one standard globally accepted algorithm. In routine clinical settings, an elevated aspartate transaminase (AST) and alanine aminotransferase (ALT) often is the first trigger for evaluation of NAFLD. The AST and ALT are classical measures of tissue injury and are widely available and used. However, they are severely limited by low sensitivity and specificity for NASH, the phenotype that is most likely to lead to cirrhosis²¹. The full spectrum of NAFLD can be seen with normal AST and ALT and these are useful when positive but not very informative when normal. An elevated AST and ALT are also not necessarily reflective of NAFLD and may be due to other common conditions such as alcohol-or drug- induced liver disease or injury, hepatitis or iron overload. A mild elevation in liver enzymes is also commonly observed in patients with heart failure, frequently due to elevated central venous pressure²². Prior to entering patients into cardiovascular clinical trials, which have often allowed patients with AST and ALT up to 1.5-2 times the upper limit of normal, it is both good clinical practice and prudent to evaluate such patients for the presence and cause of underlying liver disease.

The key questions to be addressed in the diagnostic evaluation for NAFLD include the presence of excess fat in the liver, assessment of disease activity and fibrosis stage.

How to diagnose excess liver fat and fibrosis?

There are now several well-validated tools for the measurement of excess hepatic lipid content. In routine clinical practice, their positive predictive values are high because the pre-test probability of a positive test is substantially enhanced in obese individuals with T2DM or other cardio-metabolic risk factors²³. A commonly used tool is the transient elastography which provides a measure of hepatic steatosis from the attenuation characteristics of sound waves in the liver; the continuous attenuation parameter (CAP) has both high sensitivity and specificity in detecting NAFLD but is not as good at differentiating grades of excess steatosis²⁴. Sonography can reveal an echogenic liver as a feature of hepatic steatosis but lacks relative sensitivity and specificity²⁵. MRI-based tools provide a more accurate and direct measurement of hepatic triglyceride content and their changes; the magnetic resonance imaging-derived proton density fat fraction (MRI-PDFF) is a widely used tool in the context of clinical trials^{26,27}. It is however limited by its cost and availability, and might not be suitable for every patient, for instance those unable to fit in the MRI machine or who have claustrophobia. In many clinical trials where fat content is a key outcome measure, a transient elastography is often used as a screening tool to reduce screen-fail rates inherent to the use of MRI-PDFF²⁸.

Measures of steatosis however are insufficient to provide information regarding underlying liver tissue injury, inflammation and fibrosis. There are currently no established non-invasive tools that outperform an AST and ALT for the assessment of disease activity. An ALT and AST over

40 IU/l has been associated with presence of steatohepatitis and fibrosis, respectively²⁹. Several simple laboratory aids such as the AST:platelet ratio (APRI) and the FIB-4 index (based on age, AST, ALT, platelets) provide a high negative predictive value. Specifically, a FIB-4 index <1.3 can effectively exclude over 90% of individuals with stage 2 fibrosis or higher³⁰. Transient elastography or MRI-based tool such as two-dimensional magnetic resonance elastography (2D-MRE) are often used to gain specificity in the identification of those with fibrosis stage 2 or higher, but this approach is limited by their poor positive predictive value. Both transient elastography and MRI measure liver stiffness which can be affected by hepatic congestion, inflammation, fibrosis, cholestasis, and post-prandial blood flow³¹. Ultrasound-based methods to measure liver stiffness are in development but have not been validated to the extent that transient elastography and 2D-MRE have.

Based on these considerations, while there remain areas of uncertainty, most patients can be identified to have NAFLD with low, intermediate, or high risk of liver-related outcomes and the use of a liver biopsy to evaluate the disease is increasingly being relegated to indeterminate cases and in clinical trials for NASH³².

Clinical profile and natural history of patients with NAFLD

NAFLD has been reported at even young ages, but is usually first diagnosed in middle-aged individuals, frequently in the background of several comorbidities that substantially increase CV risk⁹. Observational studies report that up to 50% of patients with NAFLD are obese, and rate of prevalent T2DM, hypertension, and hyperlipidemia might be as high as 70%. Smoking habits and coronary artery disease are less frequent³³; genetic predisposition has been implicated in up

to 20% of patients with NAFLD³⁴. Over half of patients diagnosed to have NAFLD report to continue to drink, regularly or occasionally, a substantial amount of alcohol, which might confound the exact diagnosis and have important consequences on disease progression^{35,36}. NAFLD is usually asymptomatic, but reduced exercise tolerance and fatigue might limit daily living activities in some, and lead to a poorer quality of life compared with healthier controls³⁷. The use of CV therapies that are commonly used for primary or secondary prevention, such as statins, is rarely reported in observational studies conducted in patients with NAFLD (*Table 1*). When reported, therapeutic use appears suboptimal with respect to CV risks faced by these patients.

The natural history of NAFLD and NASH is not fully elucidated. Current knowledge about the natural course of the disease is based on placebo arms of clinical trials and longitudinal cohorts with multiple liver biopsies performed. It has been previously suggested that disease progression follows a linear trajectory with one-stage progression every seven years for those with NASH³⁸. This is now known to be imprecise; it is more likely that the course of the disease is bi-directional with some patients showing regression whilst others progress¹⁸. Disease activity, measured by the severity of lobular inflammation and hepatocellular ballooning, is a key determinant of fibrosis change³⁹. Weight gain⁴⁰ and increasing AST and ALT have been associated with worsening disease activity¹⁸. Conversely, weight loss has been linked to decreased disease activity⁴¹. While a 10% weight loss threshold has been suggested to identify those who experience fibrosis improvement⁴², recent data indicate that fibrosis can change without changes in weight¹⁸. The rate of fibrosis progression to cirrhosis depends on the baseline

stage of fibrosis and ranges from less than 10% in 10 years in those with stage 1 disease to 5-10% annually for those with bridging fibrosis^{18,43}.

From a clinical point of view, liver-related outcomes are almost entirely related to the development of cirrhosis; by then, many patients with NAFLD might have already suffered the clinical consequences of an overlapping cardio-metabolic disease (*Figure 1*).

Liver-related endpoints in cardiovascular trials

Patient populations.

From a hepatological perspective, the population to be selected should be linked to the mechanism of action of the intervention to be tested and the outcome to be measured. Thus, for metabolic agents that are primarily focused on reducing atherogenic risk, assessment of hepatic steatosis is often used in phase 2A trials to obtain proof of concept that the drug is potentially active. On the other hand, agents which may impact fibrogenic pathways or have complex effects on cardio-metabolic risk, may also improve NASH and hepatic fibrosis. In such cases, the entry criteria for the trial should include a measure of hepatic steatosis and liver stiffness. A transient elastography (CAP >300 db/m and liver stiffness measurement (LSM) >8 kp) or MRI-PDFF (>5% steatosis)/2D-MRE (>2.8 kp) are the best choices for this purpose⁴⁴. Depending on the objectives of a given study, these criteria could be used as inclusion or for stratification.

In trials specifically targeting those with established heart failure, there is a high probability of having underlying NASH with advanced fibrosis^{45,46}. In such studies, the presence of previously

undiagnosed cirrhosis, but also the concomitant presence of renal and cardiac dysfunction, may alter the pharmacokinetics of drugs administered and it is prudent to obtain a measure of liver stiffness at baseline^{14,15}. If LSM is greater than 12.6 or 16 kp, the threshold values for 90% sensitivity and specificity respectively for the diagnosis of cirrhosis, study sponsors will need to decide whether to include or exclude such patients. However, in HF, the use of transient elastography or MRI may be confounded by the effect of hepatic congestion on liver stiffness which can result in over-diagnosis of cirrhosis. A FIB-4 >3.2 also can be used to diagnose cirrhosis with high specificity but the sensitivity of this cut-off is poor²⁸. For drugs that target underlying lipotoxicity or with direct antifibrotic activity, patients with both significant cardiac dysfunction and NASH with advanced fibrosis (stages 3 and 4) could be treated with such agents to improve both cardiac and liver clinical outcomes.

New onset liver dysfunction in a trial including such patients could reflect either decompensation of underlying cirrhosis, drug induced liver injury, or might be secondary to development or worsening of heart failure. It might be recommended to measure liver enzymes and bilirubin about 4 weeks apart during the screening period of the trial, to evaluate their possible fluctuations prior to initiation of therapy. This might help to interpret their changes during the course of the trial.

Endpoints.

Hepatic steatosis: changes in hepatic steatosis alone are not very meaningful clinically.

Decrease in steatosis has not been shown to impact how a patient feels, functions or

survives and thus does not meet criteria for a reasonably accepted surrogate primary endpoint. .

Normalization of liver enzymes: In phase 2A trials of NASH, a reduction or normalization of ALT is the most reliable indicator for drug benefit and the likelihood that larger histology-based trials will be positive. However, by themselves, normalization of liver enzymes has not been linked to clinical outcome improvement and also does not meet criteria for therapeutic efficacy in advanced phase trials.

Fibrosis regression: The clinical relevance of fibrosis regression depends on the clinical context in which it occurs. If a drug improves hepatic fibrosis by treating the root cause of NASH (i.e. the underlying metabolic disturbance causing steatosis in the first place), it is likely to reflect improvement in the overall disease state. On the other hand, regression of fibrosis with a pure anti-fibrotic agent which does not address hepatic steatosis, tissue injury and inflammation may or may not result in overall improvement in the health status of the patient. Reduced fibrosis, measured by transient elastography, may be considered as a secondary endpoint in cardiovascular trials. MR elastography can also be used for this purpose especially if cardiac MRI is being performed as part of the trial.

Reduced progression to cirrhosis: The onset of cirrhosis is associated with an annual 3-4% risk of clinical decompensation and an increase in all-cause mortality and liver related mortality in those with NAFLD^{47,48}. Reduced progression to cirrhosis is expected to translate into reduced liver related outcomes and mortality, and it is generally accepted

as a surrogate endpoint in clinical trials for NASH^{49,50}. For cardiovascular trials targeting underlying metabolic/inflammatory/fibrotic pathways, reduced progression to cirrhosis (diagnosed by a high specificity test such as MRE or transient elastography) would provide evidence of improvement in liver status above and beyond the cardiovascular benefits of the agent.

Reduced progression to MELD score of 15: The model for end-stage liver disease (MELD) score is widely accepted and validated as a marker for three-month mortality risk in those with cirrhosis regardless of etiology. It is currently used to determine organ allocation for liver transplantation. A MELD score of 15 or higher identifies those where long term outcomes with liver transplant are superior to medical therapy⁵¹. In trials which might include patients with cirrhosis (i.e. heart failure trials), this may be included in a composite endpoint for long-term outcomes. An increase in MELD score from 12 or less to 15 or higher is considered an acceptable surrogate endpoint in trials of NASH with cirrhosis^{49,50}.

Reduced liver outcomes: It is unlikely that cardiovascular trials will include patients with advanced liver disease at risk of liver outcomes within a 1-2-year time frame.

Cardio-metabolic endpoints in trials for NAFLD

There are a growing number of clinical trials for NAFLD or NASH (*Table 2*). While cardiovascular outcomes are considered to be mainly safety measures in such trials, it is reasonable to consider these as part of the efficacy of treatments specifically for those agents that address

underlying root causes for both the underlying cardiometabolic risk and liver disease. The baseline cardio-metabolic measures that should be considered are noted below.

Atherogenic risk profile: Multiple studies have shown that the development of NAFLD and particularly NASH is associated with increasing small dense LDL-C, which are particularly atherogenic.^{52,53} There is also increased hyperinsulinemia and elevated apolipoprotein-B in patients with NASH⁵². It is also noteworthy that insulin resistance is both a cause and effect of NAFLD and NASH and the association of NAFLD with metabolic syndrome is mutual and bi-directional^{54,55}. With progression to cirrhosis, both the hypertriglyceridemia and hypercholesterolemia improve reflecting a decrease in de novo lipogenesis⁵². In pre-cirrhotic populations in clinical trials, it is recommended that detailed lipid profiling be performed in phase 2A or 2B trials to determine if these are reasonable surrogates to be included in phase 3 trials.

Myocardial function: left ventricular hypertrophy and diastolic dysfunction are commonly described in those with NAFLD and are associated with the severity of hepatic fibrosis^{45,46}. When the mechanism of action of a drug suggests that there could be beneficial effects on both the heart and the liver, MRI or echocardiographic assessment of ventricular structure and function might be performed. Electrocardiography studies should be included in all NASH trials to ensure that any agent is not causing QTc prolongation or rhythm abnormalities.

Serum markers: Elevated natriuretic peptides levels reflect increased myocardial wall stress, and are powerful predictors of future CV events in cardiovascular medicine. However, an inverse association exists between natriuretic peptides levels with obesity, liver enzymes and the amount of liver fat in patients with NAFLD^{56,57}. Therefore, the potential of natriuretic peptides as diagnostic tool for identifying early, subclinical cardiac dysfunction in patients with NAFLD might be questionable, as well as their use as surrogate endpoint in this setting. Elevated troponin levels, reflecting myocardial injury, are directly associated with liver enzymes, and could improve prognostic stratification in patients with NAFLD⁵⁸. Markers of fibrosis, which may be involved in the transition to both liver and heart failure, could be also considered but perhaps lack specificity.

Renal outcomes: Currently, based on regulatory advice, most trials for NAFLD exclude individuals with chronic kidney disease (CKD). However, the prevalence of CKD increases with NAFLD progression especially in those with cirrhosis⁵⁹. In trials targeting NAFLD subpopulations with a high prevalence of CKD, renal impairment studies to establish the pharmacokinetic-pharmacodynamics of novel first-in-class agents should be performed early in the course of development and appropriate renal safety strategies integrated in to trial design to mitigate risk and still perform the trial in the relevant “intended use” population.

MACE events: These events are routinely measured in current phase 2B and 3 clinical trials for NASH and outcomes adjudicated by independent adjudication committees using current best practices for this purpose.

Hospitalization from heart failure: The reported incidence of hospitalization from heart failure is low to non-existent in populations currently being enrolled in trials for NASH. This is surprising, considering the high number of CV risk factors and comorbidities, therefore the high risk of developing heart failure, of patients with NAFLD. Overall, these findings suggest a low awareness of heart failure amongst hepatologists, and lack of standardised and robust criteria to diagnose heart failure, particularly in the presence of a normal left ventricular systolic function at imaging.

Candidate cardiovascular therapies for NAFLD

It might be possible that treatments targeting an overt cardio-metabolic dysfunction could resolve some of the pathophysiological mechanisms that promote a progression from NAFLD to fibrosis and development of liver-related complications, and perhaps improve outcomes.

Elevated levels of aldosterone are common amongst patients with NAFLD and cirrhosis, as well as in those with CV risk factors and CVD, and promote endothelial dysfunction, fibrosis and inflammation. Hyperaldosteronism is a valid target of treatment in those with cardiac dysfunction and perhaps in those with NAFLD. In a single centre, open label, randomised trial conducted in 31 adults with biopsy-proven NAFLD, treatment for 52 weeks with spironolactone

25 mg/day combined to vitamin E vs vitamin E monotherapy, led to a decreased NAFLD liver fat score, an index of steatosis⁶⁰.

Expert review of current evidence suggests that statins might prevent development of hepatic complications⁶¹. For those in whom statins cannot be used or are poorly tolerated, ezetimibe or PCSK9-inhibitors might be considered to improve hepatic steatosis and reduce liver enzymes^{62,63}.

Sodium-glucose cotransporter type-2 (SGLT2) inhibitors improve glycemic control, decrease blood pressure and body weight, and lead to important beneficial effects on CV morbidity and mortality in patients with T2DM and heart failure⁶⁴. A series of small studies suggest that SGLT2-inhibitors are a promising class of drugs even for liver protection in NAFLD⁶⁵.

Glucagon-like peptide-1 receptor agonists (GLP-1RA) promote satiety and reduce food intake, leading to weight loss, and prevent atherosclerotic and renal events in patients with T2DM, which makes them attractive therapies for patients with NAFLD⁶⁶. In a multi-center, double-blind, randomized, placebo-controlled phase-2 trial in NASH, treatment with liraglutide for 48 weeks led to a greater rate of histological resolution of disease, and slower progression to fibrosis, compared with placebo⁶⁷. A recently completed 72-week multi-center, randomized, double-blind placebo-controlled trial suggested that another GLP-1RA, semaglutide, also reduced histological features of NASH, compared with placebo⁶⁸. To date, no mechanistic link is established between liver protection and reduced cardiovascular events associated with use of SGLT2i or GLP-1RA.

The unmet need for CV risk reduction in NAFLD: future perspectives

Placing too much emphasis only on liver-related surrogate endpoints might delay discovery, or repurposing of potentially effective treatments, lead to waste of significant amount of resources and, more importantly, increase patients' risk of developing side effects associated with use of therapies that are subsequently found to be clinically meaningless. Some of these concerns are also contemplated in the guidance for companies developing therapies for patients with NASH recently issued by U.S. Food and Drug Administration, in which it is recommended that sponsors should consider trial endpoints that reflect robust clinical benefits (ie: histological resolution of NASH, reduced progression to, or reversal of fibrosis, need for a liver transplant), including improved survival, to support an accelerated approval process.

Targeting comorbidities and CV risk factors might be important for preventing CV complications and cancer in patients with NAFLD. However, the annual rate of these events might be low, and trials would require large sample size and long follow-up to produce convincing evidence. It is also unknown whether spontaneous resolution of less severe forms of NAFLD or NASH, as observed in >1/3 of patients randomised to placebo in clinical trials, might decrease future risk, further limiting trial feasibility and power⁶⁹. To increase event rates, many CV outcome trials use composite endpoints: a similar strategy might be applied to trials conducted in patients with NAFLD. Development of CV, but also cancer, kidney- or liver – related complications are all clinically relevant endpoints that have a significant impact on people health and well-being in people with NAFLD, and will ultimately lead to disability and death. Considering recurrent non-fatal and fatal events would further increase power. Alternative

statistical methods that prioritize the more important clinical event (for instance, death over changes in liver function tests) between matched pairs of patients exposed to treatment and controls might be useful in these scenarios⁷⁰.

Testing multiple doses of the same treatment, or various treatments simultaneously within this very heterogeneous population⁷¹ using innovative trial designs would perhaps speed up discovery and avoid exposure of patients with NAFLD to futile therapies for long periods.

Flexible, adaptive trial designs would initially identify safety issues and signs of efficacy on key metabolic targets (ie: reduction in liver fat content by imaging or normalization of liver enzymes) to screen treatments and doses that should progress further, allowing a direct transition between a phase 2 to a phase 3 trial, maximizing time and resources. These trials might identify which medicines work best for which patients and whether some patients require higher doses, or several medicines, to reduce risk to target levels. Important biological and behavioral differences exist between men and women that might be relevant to explain the epidemiology of many diseases, including NAFLD, and influence response to treatment⁷². Considering these often neglected issues in future trial design, analysis, and interpretation will potentially advance personalized approaches to the management of NAFLD. Matching a particular patient profile to a specific treatment could help tailor treatment to the individual patient's needs and fulfil aims of precision medicine.

Conclusions

There is a strong rationale for considering cardiovascular endpoints in the context of NAFLD trials. A detailed characterization of the baseline cardio-metabolic profile and a careful collection

and adjudication of adverse events will improve understanding of safety and efficacy of interventions in patients with NAFLD, and clarify its natural history. To achieve all of this, a closer collaboration between cardiologists, endocrinologists and hepatologists is desirable. .

Acknowledgments: The current manuscript summarizes the content of discussions started by this group in December 2018 at CardioVascular Clinical Trialists forum. This meeting is a unique platform to bring together academia, industry representatives, regulatory and health technology experts, payers, patients, and patient advocacy group experts to facilitate multidisciplinary discussions to advance the next phase in evidence generation for this high-risk entity.

References

- 1) Ludwig J, Viggiano TR, McGill DB, Oh BJ. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc.* 1980;55(7):434-438.
- 2) Kleiner DE, Brunt EM, Van Natta M, Behling C, Contos MJ, Cummings OW, Ferrell LD, Liu YC, Torbenson MS, Unalp-Arida A, Yeh M, McCullough AJ, Sanyal AJ. Design and validation of a histological scoring system for nonalcoholic fatty liver disease. *Hepatology.* 2005;41(6):1313-1321.
- 3) Rinella ME. Nonalcoholic fatty liver disease: a systematic review. *Jama.* 2015;313(22):2263-2273.
- 4) Min HK, Kapoor A, Fuchs M, Mirshahi F, Zhou H, Maher J, Kellum J, Warnick R, Contos MJ, Sanyal AJ. Increased hepatic synthesis and dysregulation of cholesterol metabolism is associated with the severity of nonalcoholic fatty liver disease. *Cell Metab.* 2012;15(5):665-674.

- 5) Estes C, Razavi H, Loomba R, Younossi Z, Sanyal AJ. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology*. 2018;67(1):123-133.
- 6) Estes C, Anstee QM, Arias-Loste MT, Bantel H, Bellentani S, Caballeria J, Colombo M, Craxi A, Crespo J, Day CP, Eguchi Y, Geier A, Kondili LA, Kroy DC, Lazarus JV, Loomba R, Manns MP, Marchesini G, Nakajima A, Negro F, Petta S, Ratziu V, Romero-Gomez M, Sanyal A, Schattenberg JM, Tacke F, Tanaka J, Trautwein C, Wei L, Zeuzem S, Razavi H. Modeling NAFLD disease burden in China, France, Germany, Italy, Japan, Spain, United Kingdom, and United States for the period 2016-2030. *J Hepatol*. 2018;69(4):896-904.
- 7) Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1789-1858.
- 8) Adams LA, Lymp JF, St Sauver J, Sanderson SO, Lindor KD, Feldstein A, Angulo P. The natural history of nonalcoholic fatty liver disease: a population-based cohort study. *Gastroenterology*. 2005;129(1):113-121.
- 9) Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology*. 2016;64(1):73-84.
- 10) Targher G, Byrne CD, Lonardo A, Zoppini G, Barbui C. Non-alcoholic fatty liver disease and risk of incident cardiovascular disease: A meta-analysis. *J Hepatol*. 2016;65(3):589-600.

- 11) Mantovani A, Petracca G, Beatrice G, Tilg H, Byrne CD, Targher G. Non-alcoholic fatty liver disease and risk of incident diabetes mellitus: an updated meta-analysis of 501 022 adult individuals. *Gut*. 2021;70(5):962-969.
- 12) Angulo P, Kleiner DE, Dam-Larsen S, Adams LA, Bjornsson ES, Charatcharoenwitthaya P, Mills PR, Keach JC, Lafferty HD, Stahler A, Haflidadottir S, Bendtsen F. Liver Fibrosis, but No Other Histologic Features, Is Associated With Long-term Outcomes of Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology*. 2015;149(2):389-397 e310.
- 13) Friedman SL, Neuschwander-Tetri BA, Rinella M, Sanyal AJ. Mechanisms of NAFLD development and therapeutic strategies. *Nat Med*. 2018;24(7):908-922.
- 14) Cobbina E, Akhlaghi F. Non-alcoholic fatty liver disease (NAFLD) - pathogenesis, classification, and effect on drug metabolizing enzymes and transporters. *Drug Metab Rev*. 2017;49(2):197-211.
- 15) Merrell MD, Cherrington NJ. Drug metabolism alterations in nonalcoholic fatty liver disease. *Drug Metab Rev*. 2011;43(3):317-34.
- 16) Spengler EK, Loomba R. Recommendations for Diagnosis, Referral for Liver Biopsy, and Treatment of Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis. *Mayo Clin Proc*. 2015;90(9):1233-1246.
- 17) Hagström H, Nasr P, Ekstedt M, Hammar U, Stål P, Hultcrantz R, Kechagias S. Fibrosis stage but not NASH predicts mortality and time to development of severe liver disease in biopsy-proven NAFLD. *J Hepatol*. 2017;67(6):1265-1273.
- 18) Kleiner DE, Brunt EM, Wilson LA, Behling C, Guy C, Contos M, Cummings O, Yeh M, Gill R, Chalasani N, Neuschwander-Tetri BA, Diehl AM, Dasarathy S, Terrault N,

- Kowdley K, Loomba R, Belt P, Tonascia J, Lavine JE, Sanyal AJ. Association of Histologic Disease Activity With Progression of Nonalcoholic Fatty Liver Disease. *JAMA Netw Open*. 2019;2(10):e1912565.
- 19) McPherson S, Hardy T, Henderson E, Burt AD, Day CP, Anstee QM. Evidence of NAFLD progression from steatosis to fibrosing-steatohepatitis using paired biopsies: implications for prognosis and clinical management. *Journal of hepatology*. 2015;62(5):1148-1155.
- 20) Pais R, Charlotte F, Fedchuk L, Bedossa P, Lebray P, Poynard T, Ratziu V; LIDO Study Group. A systematic review of follow-up biopsies reveals disease progression in patients with non-alcoholic fatty liver. *J Hepatol*. 2013;59(3):550-556.
- 21) Feldstein AE, Wieckowska A, Lopez AR, Liu YC, Zein NN, McCullough AJ. Cytokeratin-18 fragment levels as noninvasive biomarkers for nonalcoholic steatohepatitis: a multicenter validation study. *Hepatology*. 2009;50(4):1072-1078.
- 22) van Deursen VM, Damman K, Hillege HL, van Beek AP, van Veldhuisen DJ, Voors AA. Abnormal liver function in relation to hemodynamic profile in heart failure patients. *J Card Fail*. 2010;16(1):84-90.
- 23) Younossi ZM. Non-alcoholic fatty liver disease - A global public health perspective. *Journal of hepatology*. 2019;70(3):531-544.
- 24) Pu K, Wang Y, Bai S, Wei H, Zhou Y, Fan J, Qiao L. Diagnostic accuracy of controlled attenuation parameter (CAP) as a non-invasive test for steatosis in suspected non-alcoholic fatty liver disease: a systematic review and meta-analysis. *BMC Gastroenterol*. 2019;19(1):51.

- 25) Saadeh S, Younossi ZM, Remer EM, Gramlich T, Ong JP, Hurley M, Mullen KD, Cooper JN, Sheridan MJ. The utility of radiological imaging in nonalcoholic fatty liver disease. *Gastroenterology*. 2002;123(3):745-750.
- 26) Caussy C, Reeder SB, Sirlin CB, Loomba R. Noninvasive, Quantitative Assessment of Liver Fat by MRI-PDFF as an Endpoint in NASH Trials. *Hepatology*. 2018;68(2):763-772.
- 27) Noureddin M, Lam J, Peterson MR, Middleton M, Hamilton G, Le TA, Bettencourt R, Changchien C, Brenner DA, Sirlin C, Loomba R. Utility of magnetic resonance imaging versus histology for quantifying changes in liver fat in nonalcoholic fatty liver disease trials. *Hepatology*. 2013;58(6):1930-1940.
- 28) Zhang X, Wong GL, Wong VW. Application of transient elastography in nonalcoholic fatty liver disease. *Clin Mol Hepatol*. 2020;26(2):128-141.
- 29) Bril F, McPhaul MJ, Caulfield MP, Clark VC, Soldevilla-Pico C, Firpi-Morell RJ, Lai J, Shiffman D, Rowland CM, Cusi K. Performance of Plasma Biomarkers and Diagnostic Panels for Nonalcoholic Steatohepatitis and Advanced Fibrosis in Patients With Type 2 Diabetes. *Diabetes Care*. 2020;43(2):290-297.
- 30) Shah AG, Lydecker A, Murray K, Tetri BN, Contos MJ, Sanyal AJ; Nash Clinical Research Network. Comparison of noninvasive markers of fibrosis in patients with nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol*. 2009;7(10):1104-1112.
- 31) ASL-ALEH Clinical Practice Guidelines: Non-invasive tests for evaluation of liver disease severity and prognosis. *J Hepatol*. 2015;63(1):237-264.

- 32) Leoni S, Tovoli F, Napoli L, Serio I, Ferri S, Bolondi L. Current guidelines for the management of non-alcoholic fatty liver disease: A systematic review with comparative analysis. *World J Gastroenterol*. 2018;24(30):3361-3373.
- 33) Chang Y, Ryu S, Sung KC, Cho YK, Sung E, Kim HN, Jung HS, Yun KE, Ahn J, Shin H, Wild SH, Byrne CD. Alcoholic and non-alcoholic fatty liver disease and associations with coronary artery calcification: evidence from the Kangbuk Samsung Health Study. *Gut*. 2019;68(9):1667-1675.
- 34) Kozlitina J. Genetic Risk Factors and Disease Modifiers of Nonalcoholic Steatohepatitis. *Gastroenterol Clin North Am*. 2020;49(1):25-44.
- 35) VanWagner LB, Ning H, Allen NB, Ajmera V, Lewis CE, Carr JJ, Lloyd-Jones DM, Terrault NA, Siddique J. Alcohol Use and Cardiovascular Disease Risk in Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology*. 2017;153(5):1260-1272.e1263.
- 36) Ekstedt M, Franzén LE, Holmqvist M, Bendtsen P, Mathiesen UL, Bodemar G, Kechagias S. Alcohol consumption is associated with progression of hepatic fibrosis in non-alcoholic fatty liver disease. *Scand J Gastroenterol*. 2009;44(3):366-374.
- 37) Sayiner M, Stepanova M, Pham H, Noor B, Walters M, Younossi ZM. Assessment of health utilities and quality of life in patients with non-alcoholic fatty liver disease. *BMJ Open Gastroenterol*. 2016;3(1):e000106.
- 38) Singh S, Allen AM, Wang Z, Prokop LJ, Murad MH, Loomba R. Fibrosis progression in nonalcoholic fatty liver vs nonalcoholic steatohepatitis: a systematic review and meta-analysis of paired-biopsy studies. *Clin Gastroenterol Hepatol*. 2015;13(4):643-654 e641-649; quiz e639-640.

- 39) Schuppan D, Surabattula R, Wang XY. Determinants of fibrosis progression and regression in NASH. *Journal of hepatology*. Feb 2018;68(2):238-250.
- 40) Kim Y, Chang Y, Cho YK, Ahn J, Shin H, Ryu S. Obesity and Weight Gain Are Associated With Progression of Fibrosis in Patients With Nonalcoholic Fatty Liver Disease. *Clin Gastroenterol Hepatol*. 2019;17(3):543-550.e542.
- 41) Koutoukidis DA, Astbury NM, Tudor KE, Morris E, Henry JA, Noreik M, Jebb SA, Aveyard P. Association of Weight Loss Interventions With Changes in Biomarkers of Nonalcoholic Fatty Liver Disease: A Systematic Review and Meta-analysis. *JAMA Intern Med* 2019;179(9):1262-1271..
- 42) Vilar-Gomez E, Martinez-Perez Y, Calzadilla-Bertot L, Torres-Gonzalez A, Gra-Oramas B, Gonzalez-Fabian L, Friedman SL, Diago M, Romero-Gomez M. Weight Loss Through Lifestyle Modification Significantly Reduces Features of Nonalcoholic Steatohepatitis. *Gastroenterology*. 2015;149(2):367-378 e365; quiz e314-365.
- 43) Sanyal AJ, Harrison SA, Ratziu V, Abdelmalek MF, Diehl AM, Caldwell S, Shiffman ML, Aguilar Schall R, Jia C, McColgan B, Djedjos CS, McHutchison JG, Subramanian GM, Myers RP, Younossi Z, Muir AJ, Afdhal NH, Bosch J, Goodman Z. The Natural History of Advanced Fibrosis Due to Nonalcoholic Steatohepatitis: Data From the Simtuzumab Trials. *Hepatology*. 2019;70(6):1913-1927.
- 44) Siddiqui MS, Harrison SA, Abdelmalek MF, Anstee QM, Bedossa P, Castera L, Dimick-Santos L, Friedman SL, Greene K, Kleiner DE, Megnien S, Neuschwander-Tetri BA, Ratziu V, Schabel E, Miller V, Sanyal AJ; Liver Forum Case Definitions Working Group. Case definitions for inclusion and analysis of endpoints in clinical trials for

nonalcoholic steatohepatitis through the lens of regulatory science.

Hepatology.2018;67(5):2001-2012.

45) Simon TG, Bamira DG, Chung RT, Weiner RB, Corey KE. Nonalcoholic Steatohepatitis is Associated with Cardiac Remodeling and Dysfunction. *Obesity (Silver Spring)*. 2017;25(8):1313-1316.

46) Canada JM, Abbate A, Collen R, Billingsley H, Buckley LF, Carbone S, Trankle CR, Idowu MO, Kadariya D, Van Tassell B, Sanyal AJ, Siddiqui MS. Relation of Hepatic Fibrosis in Nonalcoholic Fatty Liver Disease to Left Ventricular Diastolic Function and Exercise Tolerance. *Am J Cardiol*. 2019;123(3):466-473.

47) Sanyal AJ, Banas C, Sargeant C, Luketic VA, Sterling RK, Stravitz RT, Shiffman ML, Heuman D, Coterrell A, Fisher RA, Contos MJ, Mills AS. Similarities and differences in outcomes of cirrhosis due to nonalcoholic steatohepatitis and hepatitis C. *Hepatology*.2006;43(4):682-689.

48) Vilar-Gomez E, Calzadilla-Bertot L, Wai-Sun Wong V, Castellanos M, Aller-de la Fuente R, Metwally M, Eslam M, Gonzalez-Fabian L, Alvarez-Quiñones Sanz M, Conde-Martin AF, De Boer B, McLeod D, Hung Chan AW, Chalasani N, George J, Adams LA, Romero-Gomez M. Fibrosis Severity as a Determinant of Cause-Specific Mortality in Patients With Advanced Nonalcoholic Fatty Liver Disease: A Multi-National Cohort Study. *Gastroenterology*.2018;155(2):443-457.e417.

49) Sanyal AJ, Friedman SL, McCullough AJ, Dimick-Santos L. Challenges and opportunities in drug and biomarker development for nonalcoholic steatohepatitis: findings and recommendations from an American Association for the Study of Liver

- Diseases-U.S. Food and Drug Administration Joint Workshop. *Hepatology*. 2015;61(4):1392-1405.
- 50) U.S. Food and Drug Administration. *Noncirrhotic Nonalcoholic Steatohepatitis With Liver Fibrosis: Developing Drugs for Treatment*. 2018.
- 51) Merion RM, Schaubel DE, Dykstra DM, Freeman RB, Port FK, Wolfe RA. The survival benefit of liver transplantation. *Am J Transplant*. 2005;5(2):307-313.
- 52) Siddiqui MS, Fuchs M, Idowu MO, Luketic VA, Boyett S, Sargeant C, Stravitz RT, Puri P, Matherly S, Sterling RK, Contos M, Sanyal AJ. Severity of nonalcoholic fatty liver disease and progression to cirrhosis are associated with atherogenic lipoprotein profile. *Clin Gastroenterol Hepatol*. 2015;13(5):1000-1008.e1003.
- 53) Bril F, Sninsky JJ, Baca AM, Superko HR, Portillo Sanchez P, Biernacki D, Maximos M, Lomonaco R, Orsak B, Suman A, Weber MH, McPhaul MJ, Cusi K. Hepatic Steatosis and Insulin Resistance, But Not Steatohepatitis, Promote Atherogenic Dyslipidemia in NAFLD. *J Clin Endocrinol Metab*. 2016;101(2):644-652.
- 54) Loomba R, Friedman SL, Shulman GI. Mechanisms and disease consequences of nonalcoholic fatty liver disease. *Cell*. 2021;184(10):2537-2564.
- 55) Lonardo A, Leoni S, Alswat KA, Fouad Y. History of Nonalcoholic Fatty Liver Disease. *Int J Mol Sci*. 2020;21(16):5888.
- 56) Wang TJ, Larson MG, Levy D, Benjamin EJ, Leip EP, Wilson PW, Vasani RS. Impact of obesity on plasma natriuretic peptide levels. *Circulation*. 2004;109(5):594-600.
- 57) Johansen ML, Schou M, Rasmussen J, Rossignol P, Holm MR, Chabanova E, Dela F, Faber J, Kistorp C. Low N-terminal pro-brain natriuretic peptide levels are associated

with non-alcoholic fatty liver disease in patients with type 2 diabetes. *Diabetes Metab.* 2019;45(5):429-435.


- 58) Lazo M, Rubin J, Clark JM, Coresh J, Schneider AL, Ndumele C, Hoogeveen RC, Ballantyne CM, Selvin E. The association of liver enzymes with biomarkers of subclinical myocardial damage and structural heart disease. *J Hepatol.* 2015;62(4):841-847.
- 59) Byrne CD, Targher G. NAFLD as a driver of chronic kidney disease. *J Hepatol.* 2020;72(4):785-801.
- 60) Polyzos SA, Kountouras J, Mantzoros CS, Polymerou V, Katsinelos P. Effects of combined low-dose spironolactone plus vitamin E vs vitamin E monotherapy on insulin resistance, non-invasive indices of steatosis and fibrosis, and adipokine levels in non-alcoholic fatty liver disease: a randomized controlled trial. *Diabetes Obes Metab.* 2017;19(12):1805-1809.
- 61) Athyros VG, Alexandrides TK, Bilianou H, Cholongitas E, Doumas M, Ganotakis ES, Goudevenos J, Elisaf MS, Germanidis G, Giouleme O, Karagiannis A, Karvounis C, Katsiki N, Kotsis V, Kountouras J, Liberopoulos E, Pitsavos C, Polyzos S, Rallidis LS, Richter D, Tsapas AG, Tselepis AD, Tsioufis K, Tziomalos K, Tzotzas T, Vasiliadis TG, Vlachopoulos C, Mikhailidis DP, Mantzoros C. The use of statins alone, or in combination with pioglitazone and other drugs, for the treatment of non-alcoholic fatty liver disease/non-alcoholic steatohepatitis and related cardiovascular risk. An Expert Panel Statement. *Metabolism.* 2017;71:17-32.

- 62) Nakade Y, Murotani K, Inoue T, Kobayashi Y, Yamamoto T, Ishii N, Ohashi T, Ito K, Fukuzawa Y, Yoneda M. Ezetimibe for the treatment of non-alcoholic fatty liver disease: A meta-analysis. *Hepatol Res.* 2017;47(13):1417-1428.
- 63) Theocharidou E, Papademetriou M, Reklou A, Sachinidis A, Boutari C, Giouleme O. The Role of PCSK9 in the Pathogenesis of Non-alcoholic Fatty Liver Disease and the Effect of PCSK9 Inhibitors. *Curr Pharm Des.* 2018;24(31):3654- 365
- 64) Fontes-Carvalho R, Santos-Ferreira D, Raz I, Marx N, Ruschitzka F, Cosentino F. Protective effects of SGLT-2 inhibitors across the cardiorenal continuum: two faces of the same coin. *Eur J Prev Cardiol.* 2021 Feb 28;zwab034. doi: 10.1093/eurjpc/zwab034. Epub ahead of print. PMID: 33659986.
- 65) Xing B, Zhao Y, Dong B, Zhou Y, Lv W, Zhao W. Effects of sodium-glucose cotransporter 2 inhibitors on non-alcoholic fatty liver disease in patients with type 2 diabetes: A meta-analysis of randomized controlled trials. *J Diabetes Investig.* 2020;11:1238-1247.
- 66) Kristensen SL, Rørth R, Jhund PS, Docherty KF, Sattar N, Preiss D, Køber L, Petrie MC, McMurray JJV. Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials [published correction appears in *Lancet Diabetes Endocrinol.* 2020 Mar;8(3):e2]. *Lancet Diabetes Endocrinol.* 2019;7(10):776-785.
- 67) Armstrong MJ, Gaunt P, Aithal GP, Barton D, Hull D, Parker R, Hazlehurst JM, Guo K; LEAN trial team, Abouda G, Aldersley MA, Stocken D, Gough SC, Tomlinson JW, Brown RM, Hübscher SG, Newsome PN. Liraglutide safety and efficacy in patients with non-

- alcoholic steatohepatitis (LEAN): a multicentre, double-blind, randomised, placebo-controlled phase 2 study. *Lancet*. 2016;387(10019):679-690.
- 68) Newsome PN, Buchholtz K, Cusi K, Linder M, Okanou T, Ratziu V, Sanyal AJ, Sejling AS, Harrison SA; NN9931-4296 Investigators. A Placebo-Controlled Trial of Subcutaneous Semaglutide in Nonalcoholic Steatohepatitis. *N Engl J Med*. 2021;384(12):1113-1124.
- 69) Han MAT, Altayar O, Hamdeh S, Takyar V, Rotman Y, Etzion O, Lefebvre E, Safadi R, Ratziu V, Prokop LJ, Murad MH, Nouredin M. Rates of and Factors Associated With Placebo Response in Trials of Pharmacotherapies for Nonalcoholic Steatohepatitis: Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol*. 2019;17(4):616-629.e626
- 70) Ferreira JP, Jhund PS, Duarte K, Claggett BL, Solomon SD, Pocock S, Petrie MC, Zannad F, McMurray JJV. Use of the Win Ratio in Cardiovascular Trials. *JACC Heart Fail*. 2020;8(6):441-450.
- 71) Lonardo A, Arab JP, Arrese M. Perspectives on Precision Medicine Approaches to NAFLD Diagnosis and Management. *Adv Ther*. 2021;38(5):2130-2158.
- 72) Lonardo A, Nascimbeni F, Ballestri S, Fairweather D, Win S, Than TA, Abdelmalek MF, Suzuki A. Sex Differences in Nonalcoholic Fatty Liver Disease: State of the Art and Identification of Research Gaps. *Hepatology*. 2019;70(4):1457-1469.
- 73) Younossi ZM, Stepanova M, Rafiq N, Makhlof H, Younoszai Z, Agrawal R, Goodman Z. Pathologic criteria for nonalcoholic steatohepatitis: interprotocol agreement and ability to predict liver-related mortality. *Hepatology*. 2011;53(6):1874- 1882

- 74) Sebastiani G, Alshaalan R, Wong P, Rubino M, Salman A, Metrakos P, Deschenes M, Ghali P. Prognostic Value of Non-Invasive Fibrosis and Steatosis Tools, Hepatic Venous Pressure Gradient (HVPG) and Histology in Nonalcoholic Steatohepatitis. *PLoS One*. 2015;10(6):e0128774.
- 75) Leung JC, Loong TC, Wei JL, Wong GL, Chan AW, Choi PC, Shu SS, Chim AM, Chan HL, Wong VW. Histological severity and clinical outcomes of nonalcoholic fatty liver disease in nonobese patients. *Hepatology*. 2017;65(1):54- 64.
- 76) Ito T, Ishigami M, Ishizu Y, Kuzuya T, Honda T, Hayashi K, Nishimura D, Toyoda H, Kumada T, Goto H, Hirooka Y. Utility and limitations of noninvasive fibrosis markers for predicting prognosis in biopsy-proven Japanese non-alcoholic fatty liver disease patients. *J Gastroenterol Hepatol*. 2019;34(1):207- 214.
- 77) Rafiq N, Bai C, Fang Y, Srishord M, McCullough A, Gramlich T, Younossi ZM. Long-term follow-up of patients with nonalcoholic fatty liver. *Clin Gastroenterol Hepatol*. 2009;7(2):234- 238

Central Illustration/Figure 1. Unmet Need for Cardiovascular Risk Reduction in Non-alcoholic Fatty Liver Disease (NAFLD). NAFLD is a very heterogeneous condition, often asymptomatic. While patients with NAFLD develop liver complications only in the presence of significant liver fibrosis, NAFLD has been linked with excess metabolic and cardiovascular complications, and non-liver related cancers. Therefore, there is a rationale for considering CV endpoints in the context of NAFLD trials, and vice versa, to consider the concomitant presence of NAFLD in drug development for cardiometabolic disorders.



Study	Recruitment (year)	FU, Years	Location	Participants, n	BMI kg/m ²	Women, (%)	Age, Years	Comorbidities	Therapy	Outcome
Vilar-Gomez ⁴⁸	1995-2013	5.5	Europe, Asia, Australia, Cuba	458; 159: bridging fibrosis 299: cirrhosis	33 (9)	52%	56 (11)	Smokers 17% HTN 61% CVD 9% Cancer 5% CKD 13% Diabetes 67% Alcohol excess 14%	Statin 24% Metformin 39% Insulin 24%	N. Deaths: 37 (85% LR) Other outcomes: 37 LT 41 HCC 10 CAD 4 Stroke 30 non-LR Ca Cirrhosis associated with greater mortality of LRE. Patients with bridging fibrosis have higher rates of vascular events and non-LR Ca.
Hagström ¹⁷	1971-2009	20	Sweden	646; NASH: 383 Fibrosis 0:25% Fibrosis 1:24% Fibrosis 2:23% Fibrosis 3: 9% Fibrosis 4: 3%	28 (4)	38%	48 (14)	Smokers 24% HTN 30% Diabetes 14% Hyperlipidaemia 8%	NR	N. Deaths: 214 Causes of death: 37% CV 26% non-LR Ca 8% LR; rest: Other causes. NAFLD: ↑ Risk of liver disease NAFLD: ↑ Risk of CVD Advanced Fibrosis, but not NASH, ↑ Risk of mortality
Younossi ⁷³	NR	12	USA	209; NASH: 131 Non-NASH: 78	37 (10) 35 (11)	69% 50%	49 (14) 48 (15)	NASH/non-NASH: Diabetes 24%/15%	NR	NASH/non-NASH: ACM: 31%/31% LRM: 13%/1% NASH: ↑ Risk of LRM Advanced Fibrosis: ↑ Risk of LRM
Sebastiani ⁷⁴	2004-2013	5	Canada	148 (NASH); Fibrosis 0:15% Fibrosis 1:36% Fibrosis 2:15% Fibrosis 3: 19% Fibrosis 4: 15%	31 (5)	30%	50 (11)	HTN 39% Diabetes 33%	NR	Deaths: 9 (6%). Clinical outcomes (ACM/hepatic complications): 24 (16%). Non-invasive assessment of fibrosis predicted outcome.
Angulo ¹²	1975-2005	13	6 Countries	619 (97 no FU); Fibrosis 0:52% Fibrosis 1:23% Fibrosis 2:14% Fibrosis 3: 9%	31 (26-37)	62%	49 (38-60)	Smokers 9% HTN 31% Diabetes 38%	Statin 10%	ACM/Liver Transplant:193 (37%) Liver death/transplant: 18 (3%) ↑ Fibrosis, but no other histologic features of steatohepatitis, ↑

				Fibrosis 4: 3%							Risk of Death/LRE
Leung ⁷⁵	2006-2015	~4	Hong Kong	307; 27% advanced fibrosis.	28 (25-31)	44	51 (11)	HTN 55% Diabetes 55%	NR		Deaths: 6 (1 LR) CV events: 22 (including 8 strokes and 8 MIs) 5 LRE (including 2 HCC) 6 Not-LR Ca Patients with BMI<25 Kg/m ² (n=72) were less likely to have fibrosis and had better outcomes.
Ito ⁷⁶	1999-2014	7	Japan	246; (NASH 63%) Fibrosis 0:25% Fibrosis 1:39% Fibrosis 2:15% Fibrosis 3: 17% Fibrosis 4: 3%	26 (24-29)	48	55 (41-63)	HTN 42% Diabetes 45% Dyslipidaemia 71%	NR		Deaths: 8 (LR 75%) Other events: 16 Ca (not-LR),9 HCC, 12 CV. Advanced Fibrosis (≥3) ↑ Risk of LR-Ca, not-LR Ca, and mortality.
Rafiq ⁷⁷	NR	~11 (up to 28)	USA	173; NASH: 72 Non-NASH: 101	34 (10) 34 (10)	69 54	51 (13) 49 (15)	Diabetes ~70%	NR		Deaths: 78, due to CAD (n=22), not-LR Ca (n=14) or LR (includes HCC; n=12). No difference in ACM between NASH and non-NASH; patients with NASH had greater LR mortality than Non-NASH.

Table 1. Selected studies summarizing characteristics and outcome of patients with biopsy proven NAFLD. Abbreviations used: FU – follow-up; BMI – body mass index; NA - not applicable; HTN – Hypertension; CVD – cardiovascular disease; LT – Liver transplant; LR – liver related; LRE – liver related events; LRM – liver related mortality; CAD – coronary artery disease; ACM – all cause mortality; Ca – Cancer; HCC – Hepatocellular carcinoma; MI – myocardial infarction.

ClinicalTrials.gov	Drug	Patients	N of patients	Trial status	Key Outcomes
NCT02548351	Obeticholic Acid	NASH	2,480	A, not R	Liver outcomes & ACM
NCT02704403	Elafibranor	NASH	2,157	T	Safety; Liver outcomes & ACM; Cardio-metabolic markers and CV events
NCT03900429	Resmetirom	NASH	2,000	R	Liver outcomes & ACM
NCT04849728	Lanifibranor	NASH	2,000	R	Liver outcomes & ACM
NCT04104321	Aramchol	NASH	2,000	R	Liver outcomes & ACM
NCT03028740	Cenicriviroc	NASH	1,779	T	Liver outcomes & ACM
NCT04822181	Semaglutide	NASH	1,200	R	Liver outcomes & ACM
NCT04365868	Belapectin	NASH	1,010	R	Liver outcomes & ACM
NCT04951219	Resmetirom	NAFLD	1,000	R	Safety; Liver & metabolic markers
NCT03439254	Obeticholic acid	NASH & cirrhosis	919	A, not R	Liver markers
NCT04197479	Resmetirom	NAFLD	700	R	Safety; Liver & metabolic markers
NCT04321031	PF-06865571/ PF-05221304	NASH	450	R	Safety (includes abnormal ECGs); Liver & metabolic markers
NCT01617772	Atorvastatin or L-Carnitine	NASH	440	U	Liver markers
NCT03449446	Selonsertib, Firsocostat, Cilofexor, and Combinations	NASH	395	C	Safety; Liver markers
NCT04065841	Tropifexor and licogliflozin	NASH	380	R	Safety; Liver & metabolic markers
NCT02855164	Tropifexor	NASH	351	T	Safety; Liver & metabolic markers
NCT04173065	VK2809	NASH	337	R	Safety; Liver markers
NCT04378010	EDP-305	NASH	336	R	Safety; Liver markers
NCT04906421	TVB-2640	NASH	330	R	Safety; Liver markers
NCT04583423	MK-3655	NASH	328	R	Safety; Liver markers
NCT02970942	Semaglutide	NASH	320	C	Safety; Liver, inflammatory & metabolic markers
NCT04880187	AXA1125	NASH	273	R	Safety; Liver & metabolic markers
NCT04267393	BMS-986263	NASH	270	R	Safety (includes abnormal ECGs); Liver & metabolic markers
NCT04052516	Icosabutate	NASH	264	R	Liver markers
NCT04171765	BFKB8488A	NASH	260	R	Safety; Liver markers
NCT04193982	Saroglitazar/Vitamin E	NAFLD	250	R	Safety; Liver & metabolic markers

NCT04147195	LYS006 +/- Tropifexor	NAFLD & NASH	250	R	Safety; Liver & metabolic markers
NCT03008070	Lanifibranor	NASH	247	C	Liver, inflammatory & metabolic markers
NCT04771273	BI 456906	NASH	240	R	Safety; Liver & metabolic markers
NCT05011305	Saroglitazar	NASH	240	R	Safety; Liver & metabolic markers
NCT02781584	Selonsertib, Firsocostat, Cilofexor, Fenofibrate and/or Vascepa®	NASH	220	C	Safety; Liver & metabolic markers
NCT02443116	NGM282	NASH	220	C	Safety; Liver markers
NCT04929483	BIO89-100	NASH	216	R	Safety; Liver & metabolic markers
NCT04198805	Vitamin E/Omega-3 fatty acid	NAFLD	200	R	Safety; Liver, inflammatory & metabolic markers
NCT04166773	Tirzepatide	NASH	196	R	Safety; Liver & metabolic markers
NCT04048876	CC-90001	NASH	195	R	Safety; Liver & metabolic markers
NCT03517540	Tropifexor & cenicriviroc	NASH	193	C	Safety; Liver markers
NCT03551522	Seladelpar	NASH	181	C	Safety (includes abnormal ECGs); Liver, inflammatory & metabolic markers
NCT04399538	PF-06865571/PF-05221304	NAFLD	180	R	Safety (includes abnormal ECGs); Liver & metabolic markers
NCT04773964	MET642	NASH	180	R	Safety (includes abnormal ECGs); Liver & metabolic markers
NCT03969719	PF-06835919	NASH	164	C	Safety; Liver, inflammatory & metabolic markers
NCT03486899	BMS-986036	NASH	160	A, not R	Safety; Liver markers
NCT04767529	EFX	NASH	160	R	Safety; Liver & metabolic markers
NCT03812029	EYP001a	NASH	160	R	Safety; Liver markers
NCT03486912	BMS-986036	NASH	155	A, not R	Safety (includes abnormal ECGs); Liver & metabolic markers
NCT03912532	NGM282	NASH	152	A, not R	Safety; Liver markers
NCT04210245	Aldafermin	NASH	150	R	Safety; Liver markers
NCT04932512	ION224	NASH	150	R	Safety; Liver & metabolic markers
NCT04142749	Oltipraz	NAFLD	144	R	Safety; Liver, inflammatory & metabolic markers
NCT03938246	TVB-2640	NASH	141	R	Safety; Liver, inflammatory & metabolic markers
NCT04501406	Pioglitazone	NASH	138	R	Safety; Liver & metabolic markers

NCT04944992	Efinopegdutide	NAFLD	130	R	Safety; Liver & metabolic markers
NCT03763877	PXL770	NAFLD	120	C	Safety; Liver, inflammatory & metabolic markers
NCT03925701	Vildagliptin	NAFLD & T2DM	120	R	Liver & metabolic markers
NCT04761848	Cilostazol	NAFLD	120	R	Safety; Liver, inflammatory & metabolic markers
NCT04702490	MET409 +/- empagliflozin	T2DM and NASH	120	A, not R	Safety (includes abnormal ECGs) and tolerability
NCT04321343	PXL065	NASH	120	R	Safety; Liver, inflammatory & metabolic markers
NCT03350165	Pemafibrate	NAFLD	118	C	Safety; Liver markers
NCT04505436	HM15211	NASH	112	R	Safety; Liver markers
NCT03976401	EFX	NASH	110	A, not R	Safety; Liver markers
NCT03987074	Semaglutide, Firsocostat, Cilofexor	NASH	109	C	Safety; Liver markers
NCT03061721	Saroglitazar magnesium	NAFLD	106	C	Safety; Liver & metabolic markers
NCT03371355	ISIS 703802	NAFLD & T2D	105	C	Safety; Liver, inflammatory & metabolic markers
NCT04048135	BIO89-100	NAFLD & NASH	101	A, not R	Safety; Liver & metabolic markers
NCT04328077	TERN-101	NASH	101	C	Safety; Liver markers
NCT03723252	Dapagliflozin	NASH	100	R	Liver, inflammatory & metabolic markers
NCT04235205	Elobixibat + cholestyramine	NAFLD	100	A, not R	Safety; Liver & metabolic markers
NCT03656744	HTD1801	NASH & T2DM	100	C	Safety; Liver & metabolic markers
NCT01930123	IV Fructose infusion	NAFLD	100	R	Liver & metabolic markers

Table 2. Phase II and III trials registered on clinicaltrials.gov as for 25th of August 2021, currently recruiting (R), active and not recruiting (A, not R), recently completed (C), terminated (T) or with unknown status (U) conducted in adults with NASH or NAFLD, enrolling at least 100 participants. Abbreviations used: ECG – electrocardiogram; IV – intravenous.