



**HAL**  
open science

## Physicochemical degradation of phycocyanin and means to improve its stability: A short review

Aida Adjali, Igor Clarot, Zilin Chen, Eric Marchioni, Ariane Boudier

### ► To cite this version:

Aida Adjali, Igor Clarot, Zilin Chen, Eric Marchioni, Ariane Boudier. Physicochemical degradation of phycocyanin and means to improve its stability: A short review. *Journal of Pharmaceutical Analysis*, 2021, 10.1016/j.jpha.2021.12.005 . hal-03511791

**HAL Id: hal-03511791**

**<https://hal.univ-lorraine.fr/hal-03511791v1>**

Submitted on 10 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Journal Pre-proof



Physicochemical degradation of phycocyanin and means to improve its stability: A short review

Aida Adjali, Igor Clarot, Zilin Chen, Eric Marchioni, Ariane Boudier

PII: S2095-1779(21)00148-9

DOI: <https://doi.org/10.1016/j.jpha.2021.12.005>

Reference: JPHA 701

To appear in: *Journal of Pharmaceutical Analysis*

Received Date: 8 October 2021


Revised Date: 22 December 2021

Accepted Date: 26 December 2021

Please cite this article as: A. Adjali, I. Clarot, Z. Chen, E. Marchioni, A. Boudier, Physicochemical degradation of phycocyanin and means to improve its stability: A short review, *Journal of Pharmaceutical Analysis* (2022), doi: <https://doi.org/10.1016/j.jpha.2021.12.005>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Xi'an Jiaotong University. Production and hosting by Elsevier B.V. All rights reserved.



## **Physicochemical degradation of phycocyanin and means to improve its stability: A short review**

Aida Adjali <sup>a</sup>, Igor Clarot <sup>a</sup>, Zilin Chen <sup>b,c</sup>, Eric Marchioni <sup>d</sup>, Ariane Boudier <sup>a,\*</sup>

<sup>a</sup> *Université de Lorraine, CITHEFOR, F-54000 Nancy, France;*

<sup>b</sup> *Key Laboratory of Combinatorial Biosynthesis and Drug Discovery, Ministry of Education, Hubei Province Engineering and Technology Research Center for Fluorinated Pharmaceuticals, and Wuhan University School of Pharmaceutical Sciences, Wuhan, 430071, China*

<sup>c</sup> *State Key Laboratory of Transducer Technology, Chinese Academy of Sciences, Beijing 100080, China*

<sup>d</sup> *Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France*

\* Corresponding author:

Ariane Boudier

Tel: +33 372 747 228

E-mail: [Ariane.boudier@univ-lorraine.fr](mailto:Ariane.boudier@univ-lorraine.fr) (A. Boudier)

**Abstract**

The cyanobacterium *Arthrospira platensis*, spirulina, is a source of pigments such as phycobiliprotein, phycocyanin. Phycocyanin is used in the food, cosmetic, and pharmaceutical industries because of its antioxidant, anti-inflammatory, and anticancer properties. The different steps involved in extraction and purification of this protein can alter the final properties. In this review, the stability of phycocyanin (pH, temperature, and light) is discussed, considering the physicochemical parameters of kinetic modeling. The optimal working pH range for phycocyanin is between 5.5 and 6.0 and it remains stable up to 45 °C; however, exposure to relatively high temperatures or acidic pH decreases its half-life and increases the degradation kinetic constant. Phycobiliproteins are sensitive to light and preservatives such as mono- and di-saccharides, citric acid, or sodium chloride appear to be effective stabilizing agents. Encapsulation within nano- or micro- structured materials such as nanofibers, microparticles, or nanoparticles, can also preserve or enhance its stability.

**Keywords:** *Arthrospira platensis*; spirulina; phycocyanin stability; preservatives; encapsulation.

## 1. Introduction

Spirulina or *Arthrospira platensis* is a microscopic cyanobacterium that is considered to be an environmental friendly microalga. This organism is characterized by its rapid growth and consumption of little energy and water per kilogram [1]. This cyanobacterium has received significant attention because of its high protein content, accounting for up to 70% of its dry mass. Among the various nutrients (vitamins A, B12, B6, C, D, beta-carotene, and minerals) that constitute spirulina, phycocyanin, a phycobiliprotein responsible for the blue-green color of spirulina represents up to 20% of the dry mass of total cellular proteins [2,3]. The different applications of phycocyanin are shown in Fig. 1.

In terms of nutrition, this cyanobacterium globally represents a source of interest because of its high protein and antioxidant contents, which possess reactive oxygen scavenging and metal chelating activity [4]. Metal chelating activity eliminates toxins such as heavy metals (iron, copper, and cadmium) by ion exchange and chelation in 5-membered chelated rings [5]. Antioxidant and detoxification properties are also exploited for cosmetic applications in masks, creams, and gels.

Phycobiliproteins, including phycocyanin, are used as a natural blue dye [6] in food (chewing gum, dairy products, ice, and jellies), cosmetics (lipsticks), and in medicine as biochemical tracers in immunoassays because of their fluorescent properties [7].

Phycocyanin has the potential for drug development. It exerts anti-tumor effects and exhibits apoptotic characteristics such as DNA fragmentation, nuclear condensation, membrane bubble formation, and cell shrinkage [8]. In addition, its use in combination with anti-inflammatory (piroxicam [9]), anticancer (topotecan [10], doxorubicin [11]) drugs appears to be promising, even in multi-drug resistant cells [11]. The antioxidant activity of phycocyanin is also relevant in the alleviation of many diseases, including inflammation, cancer, and other disorders caused by oxidative stress [12]. The bioactivity of phycocyanin and other compounds extracted from algae has been extensively reviewed in literature [8,13,14], which is beyond the scope of this review.

Phycocyanin extraction from *A. platensis* as a protein-pigment complex appears to be easy because of its high solubility in water. Various production, separation, and purification methods have been carried out and extensively covered by other reviews [7,14–19] and are therefore beyond the scope of this paper. However, these methods are considered to present a major obstacle with respect to the stability of phycocyanin during the extraction and purification processes. This molecule appears to be highly sensitive to environmental stress, particularly temperature, pH, and light [6,20]. This instability also limits its use in food and cosmetic domains for the development of products because it causes precipitation and partial discoloration by changing from blue to green or by total discoloration. This may also result in unwanted effects, such as a loss of antioxidant

capacity [21]. Therefore, numerous stability studies have been performed to determine the optimal conditions for use by evaluating kinetic parameters such as the order of the reaction, the kinetic constant, or the half-life as a function of degradation conditions. These studies were performed to gather information on the change in food quality, as previously described for other antioxidants [22]. To avoid phycocyanin degradation and to improve its shelf-life, the use of stabilizing agents or direct encapsulation in different types of particles was explored. These methods are widely used in food products and cosmetic formulations. This review discusses the experimental methods and physicochemical parameters used to study the *in vitro* degradation of phycocyanin. This will provide practical keys for experimentations to define the conditions for its proper use or to test new stabilizers. The different stabilization techniques (*via* preservatives or formulation processes) will also be presented which is original from other review articles [23].

## 2. Structural, physical, and chemical characterizations of phycocyanin

Phycocyanin (220 kDa) is composed of a polypeptide chain and chromophores called phycocyanobilin (Figs. 2A and B). The polypeptide chain comprises  $\alpha$ - and  $\beta$ -subunits with molecular masses between 18 and 20 kDa [24]. The two subunits form monomers ( $\alpha\beta$ ), which organize into trimer ( $(\alpha\beta)_3$ ) and further into hexamers ( $(\alpha\beta)_6$ ). Each monomer contains three chromophores, which are positioned at the Cys-84 residues of subunit  $\alpha$ , and Cys-84 and Cys-155 of the  $\beta$  subunit.

UV-vis absorbance is conventionally used to evaluate the purity and degradation of phycocyanin because the protein is characterized by a maximum absorbance at 620 nm with two weaker bands at 360 nm and 650 nm [25]. The ratio Absorbance at 620 nm/Absorbance at 280 nm ( $A_{620}/A_{280}$ ) is commonly used to evaluate phycocyanin purity after extraction and purification. This highlights the purity of the protein compared to the contaminating proteins (non-specific absorbance at 280 nm). In general, a purity ratio between 0.7 and 3.9 indicates that phycocyanin is provided as food grade and a higher purity ratio indicates reagent/analytical grade [16]. High purity can be obtained when the ratio is  $> 4$  [26]. The  $A_{620}/A_{280}$  ratio is also used to monitor protein denaturation [25,27]. This is explained by a change in conformation of the tetrapyrrole chromophores due to the de-folding of the protein. This imposes a particular conformation of the chromophore inducing a modification in the absorbance ratio [27].

### 3. Degradation studies

#### 3.1 Experimental conditions

Table 1 presents an overview of the literature for the experimental conditions used in the degradation studies of phycocyanin.

These features have many similarities. Food grade phycocyanin is generally used for studying its degradation. The degradation is commonly evaluated by UV-visible spectrophotometry based on the spectroscopic properties of the protein. The study variables were the temperature (25 to 100 °C), pH of the medium (3.0 to 8.0), and in a few studies, the presence of light. The initial concentration of phycocyanin did not vary in the studies and remained between 0.3 and 3.7 mg/mL. The results were modeled using different mathematical models, and the degradation of phycocyanin generally follows first-order kinetics characterized by a linear decrease in a semi-logarithmic plot of decreasing variables during the reaction time. Authors commonly determine the kinetic constant ( $k$ ), half-life ( $t_{1/2}$ ), and activation energy ( $E_a$ ). However, some studies monitored the results using a reaction order of  $n=6$  or a Weibull model [29,31,39]. Böcker et al. [29] showed low correlation coefficients with first-order kinetics. Consequently, their degradation profile was modeled using a progressive approach with a non-linear decrease and a reaction order of  $n=6$ . This deviation may be related to different treatment parameters and differences in the source of phycocyanin, as suggested by the authors. This study examined a food coloring formula containing 15.7% (*m/m*) phycocyanin. According to the authors, this formulation is relevant for applications in the food industry compared to conditions reported in other publications. The high reaction order ( $n = 6$ ) was explained as an empirical reaction order that describes the decrease in coloring activity better than other tested reaction orders. A high reaction order indicates that other parallel reactions, consecutive reactions, or intermediate products are formed, influencing the course of the reaction [29]. In another study, the Weibull equation showed a better fit in the description of the decrease in the absorbance of food-grade phycocyanin and reagent [31] compared to the first-order kinetic model used in other studies [6,28,30]. The authors suggested that differences in the origin of the samples and the methodologies of the heat treatment may be the reason for the best fit made by the Weibull model with respect to the first-order model. Table 2 [6,28,30,36] presents some detailed results after fitting the data to the first-order kinetic model obtained in the studies.

#### 3.2. Effect of phycocyanin purity

The degradation kinetics of reagent grade and food-grade phycocyanin were compared [31]. The absorbance values of both showed a decrease with time and temperature, similar to other reports [28]. However, experiments using reactive grade phycocyanin showed significantly lower thermal stability than food grade phycocyanin ( $k = 1.24 \times 10^{13} \pm 2.2 \times 10^{13}$  vs.  $1.1 \times 10^{14} \pm 2.7 \times 10^{14} \text{ h}^{-1}$ , respectively). The explanation



of such a phenomenon may rely on the processing of the protein: gentle treatment to best preserve the color when used as a food grade, whereas a more extensive purification and stabilization process, including lyophilization when used as a reactive grade. This may affect the native structure of the protein, which is sensitive to environmental stress [31].

### 3.3. Effect of temperature

Various studies have investigated the effect of temperature on the breakdown of phycocyanin. At values close to room temperature (25 to 47 °C), phycocyanin solution degrades very slowly ( $t_{1/2} = 309.4 \pm 12.0$  min at 47 °C and at pH=6) [6]. Some authors have reported the absence of degradation up to 45 °C [36]. Between 47 and 69 °C, the degradation rate was higher in relation to the reported half-lives ( $14.5 \pm 4.2$  min and at pH=6) (Table 2) [6,28,30,36]. These profiles have also been reported in various studies [28–30,37]. At temperatures above 70 °C, the denaturation of the protein was accelerated ( $t_{1/2} = 9.7 \pm 1.6$  min at 74 °C and pH = 6) [6], consistent with other studies [29,36]. The values of activation energies reported in Table 2 were in the range of several hundred kJ/mol, showing a strong temperature dependence of the degradation of phycocyanin. From these reports, a temperature less than 45 °C is optimal for preserving phycocyanin stability.

### 3.4. Effect of pH

Acidity of a medium is an important element that destabilizes phycocyanin, leading to its degradation. pH of the medium also modifies the spectral properties and color of the protein. Phycocyanin solution at neutral pH is perceived as blue and at acidic pH as green. Exploiting this sensitivity, pH indicator polylactic acid/polyethylene oxide (PLA/PEO) ultrafine fibers containing phycocyanin was developed [41]. This is important for food industries. Phycocyanin solutions under acidic pH (3.0 and 4.0 at room temperature) exhibited lower absorbance at 620 nm and stronger absorbance at 280 nm compared to pH 5, 6, and 7 at temperatures of 55 and 65 °C for the same protein concentration. This is explained by the precipitation of proteins [36] and probably by a change in the conformation of the protein. For pH values greater than 4 and up to 6, the chromophore retains its extended geometry; however, when the pH is lower, it folds into a cyclic conformation, modifying its spectral properties [25,42]. The stability can also be dependent on the cyanobacterium strain from which it is extracted [37]. When *Spirulina* was used, the best conditions for protein stability were pH 5.5 and 6.0.

Various studies have demonstrated phycocyanin stability as a function of both pH and temperature (Table 2). At temperatures between 50 and 55 °C and pH 6.0, phycocyanin remained stable [28]. It was also stable between 57 and 65 °C and at pH 5.0; however, between 50 and 65 °C and at pH 7.0, phycocyanin denatured and became unstable. These results are consistent with the results from other studies [6,30,36]. Although several researchers have performed experiments between pH 5.0 and 8.0 to

simulate the conditions of some tests (for example, antioxidant assay, using 1,1-diphenyl-2-picrylhydrazyl or 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid, or in cell culture). The results confirmed that the optimal range for maintaining the phycocyanin stability is a pH value between 5.5 and 6.0.

### 3.5. Effect of light

Few studies have explored the impact of light on the degradation of phycocyanin in solution at a constant temperature (25 °C) [36–39]. A methodology to study the relationship between light and temperature on phycocyanin degradation using correlation models was described by Escalante et al. [39]. After exposure to lamps emitting an intensity of 50 and 100  $\mu\text{mol}/\text{m}^2\cdot\text{s}$ , the phycocyanin concentration decreased in a dose-dependent manner as a function of time. At the same light intensity, phycocyanin solution adjusted to pH 6.0 showed less degradation than those adjusted to pH 5.0 and pH 7.0. The final protein concentration decreased by approximately 20% after continuous exposure to 100  $\mu\text{mol}/\text{m}^2\cdot\text{s}$  for 36 h, regardless of the pH of the medium. The authors observed a slightly higher level of degradation after exposure to 100  $\mu\text{mol}/\text{m}^2\cdot\text{s}^1$  total intensity than after exposure to an intensity of 50  $\mu\text{mol}/\text{m}^2\cdot\text{s}^1$ . Similar results were reported by other authors with accelerated degradation when the protein was stored at 40 °C vs. 4 °C [38]. Therefore, the best storage conditions are in the dark.

### 3.6. Recommendations on stability studies

As discussed in Section 2 and Table 1, UV-vis absorbance is conventionally used to evaluate the purity and degradation of phycocyanin. For a wider valorization of this protein, particularly in the pharmaceutical context, stability studies under predefined stressful conditions (e.g., ICH Q1B, CPMP/ICH/279/95 [43] for photostability) will be necessary and separation methods must be used. To further describe the stability and degradation products formed (e.g., peptide sequence composition), high-resolution analytical methods are justified. There are very few studies in the literature that describe systems devoted to phycocyanin stability analysis, such as high-performance liquid chromatography (HPLC) coupled with diode array detection (DAD) [44] or mass spectrometry (MS) [45], which can provide essential information on the structure (characterization) and content of the various impurities formed during the degradation phase. Electrophoretic methods (capillary [46] or gel [47]) have also been described for phycocyanin analysis, and many developments are expected in the future for stability testing.

## 4. Methods to improve phycocyanin stability

Preservatives are necessary to ensure that the manufactured food remains safe and intact for a long time (shelf life). Salts, sugars, and vinegar have been used for centuries as food preservatives. As described above, phycocyanin is sensitive to environmental

stress. The degradation of the protein fraction significantly impacts the maintenance of color and bioactivity of phycocyanin [48], which are major elements in industries. This explains the use of preservatives and the encapsulation in particulate forms to preserve its color and prevent its denaturation. In this section, only studies describing results on protein stability were reviewed.

#### 4.1 Use of preservatives

Preservatives are used to enhance the stability of phycocyanin. (Table 3 [6,28,31-32,34-36,40,49-51])

The chemical structure and the concentrations of the preservatives used are important because the resulting mixture must remain safe for humans. They must not denature the protein or alter its optical and antioxidant properties. Therefore, some substances are excluded because of safety reasons (e.g., sodium azide and dithiothreitol [6]) or because it is too important to induce protein precipitation (e.g., NaCl 5% [6]). The selected compounds were mono- or disaccharides (glucose, fructose, saccharose, trehalose, lactose, maltose, and sorbitol), inorganic salts (sodium chloride and calcium chloride), and organic acids (citric acid, ascorbic acid, and benzoic acid). In some studies, the compounds are associated [35]; in others, natural matrices have been tested (conventional honey and honey from Manuka (*Leptospermum scoparium*) [50]), and polymers were also used [32]. A pigment, biopterin- $\alpha$ -glucoside, was also selected to improve the photostability of phycocyanin [40].

Although all the protocols are not completely comparable (in terms of concentration of phycocyanin, composition, and pH of the medium), some conditions appeared to be interesting for improving the stability of phycocyanin, particularly towards thermodegradability. Concerning the mono- and disaccharides, several studies showed that these compounds can improve protein stability [28,31,35,49,50]. Some compounds were better than the others. For example, glucose 20% (*m/m*) or sorbitol 50% (*m/m*) induced a two-fold increase in the half-life of the protein compared to the control, and each control was better than that of saccharose 20% (*m/m*) (half-life nearly similar to the control) [49]. Others stated the importance of the concentrations of mono- or disaccharides rather than the compound itself [6]. Therefore, they selected glucose or saccharose, both at 20% (*m/m*), and both allowed in preserving more than 62% of the protein after 15 min at 60 °C vs. only 47% without any preservatives [6]. The same study emphasized the use of NaCl at high concentrations (2.5% to 20%), which was also tested by Wu et al. [36]. However, the solution became turbid at concentrations of 5% (*m/m*) of salt. Citric acid preserved 67% phycocyanin after 45 days vs. less than 3% for the control protein without any preservatives [35]. This was explained by its capacity to decrease the pH of the medium from 7 to 6 (optimal pH for the protein) and by its chelating capacity. The use of citric acid has also been described as a stabilizer for other natural proteins

(whey protein) [35]. Concerning the photodegradation of phycocyanin, the presence of the pigment biopterin- $\alpha$ -glucoside prevented its degradation and decoloration [40].

#### 4.2. Particulate forms

The development of particulate forms is common in the food, cosmetics, and pharmaceutical industries. Studies dealing with the stability of phycocyanin in particulate forms are detailed in Table 4 [25,47,49,42-65].

In addition to the enhancement of the stability of the protein [58], the purpose is to develop active packaging to preserve food because of its antioxidant properties to prolong food half-life and avoid its deterioration [56] and the development of a stabilized phycocyanin-based blue food [25].

The formulation of phycocyanin is a challenge because it is a protein (specific folding that can be modified or even degraded during the process) that is highly soluble in water (difficulty in remaining compartmentalized or encapsulated with low or no uncontrolled release from the particle), and sensitive to pH and temperature (which may limit the use of some formulation protocols). The results from these studies showed that encapsulation efficiency (encapsulated and/or at the surface of the particle) of the protein is high that is it by electrospraying up to 75% encapsulation was achieved [61], or approximately 100% by microencapsulation [60]. Some authors have indicated that phycocyanin was sometimes encapsulated within the particle, whereas the protein was also located at the interface between the external medium and the particle [56]. There has been no report on the possible degradation of proteins during the formulation processes.

The tested formulation protocols did not modify the antioxidant properties of the protein [56,60,61], and all the reported studies showed enhanced stability of the encapsulated protein with respect to temperature and pH modifications. However, the improved thermostability of phycocyanin in nanofibers was in the same range as that obtained with the use of preservatives [49].

In addition, *in vitro* release studies of polymeric microcapsules encapsulating phycocyanin in conditions mimicking gastric and intestinal fluids were performed. The results demonstrated protection of the protein from acidic gastric fluid and a sustained release profile in simulated intestinal fluid as a function of the used polymer (i.e., alginate, chitosan carrageenan) [58,59]. This emphasized the role of the formulation in protecting the protein from the intestinal fluids.

Although these studies are preliminary, they are interesting in terms of the properties of phycocyanin and for future use in the food industry and other domains such as therapeutics, cosmetics, and sensors.

#### *4.3. Recommendations on the use of preservatives and development of particulate forms*

Research on preservatives is devoted to the study of the intermolecular interactions between the protein and the preservatives [31]. However, apart from this fundamental knowledge, other parameters must be cited. The regulation of additives is complex and differs from one country to another [66]. Many additives are not devoid of side effects. For further information, for example, the reader is referred to the document edited by the European Commission describing the side effects originating from excipients contained in medicines [67]. Similarly, consumers are increasingly questioning additives in food even though they are necessary to ensure the quality of products. Studies that consider this question showed that consumers are generally more sensitive to negative than positive arguments [68–70]. This is balanced by consumers' level of knowledge about additives and the education level of consumers [68–70]. Knowledge regarding food additives has been shown to weaken the risk perception of food safety issues [71]. Other factors must be considered, such as consumers' knowledge of regulation, their trust in regulators, and their preference for natural products [72].

The encapsulation of phycocyanin in different forms (nanofibers, microparticles, or nanoparticles) is another way to prevent its degradation and may open new opportunities for the use of this high added-value protein. The development of formulations for the food, cosmetic, and pharmaceutical purposes is definitely a challenge. Phycocyanin is hydrophilic and highly sensitive to environmental stresses. The best compromise between the intrinsic parameters of the protein, the chosen excipients/additives, and the selected process is not simple [73,74]. The use of benign processes (with no heat, no ultrasound, and no organic solvent) should be prioritized. Future directions using microfluidic [75] or supercritical fluid technology [76] that have already proved their efficiency in formulating sensitive molecules may emerge.

## **5. Conclusion**

Phycocyanin is of particular interest to humans in food, medicine, and cosmetics. However, its sensitivity to environmental conditions may prevent the development of new products. According to previous studies, the optimal conditions required for its stability include a pH between 5.5 and 6.0, low temperature (less than 45 °C), and dark storage conditions. This review suggests the use of separative systems to study the degradation pathways and generated by-products. The addition of stabilizing agents such as mono- or di-saccharides, sodium chloride, or citric acid prevents degradation. Some consideration regarding the use of preservatives is provided towards regulatory aspects and, on more general terms, the acceptance by the population. Finally, as the use of particulate systems appears to be a future direction, some recommendations on the selected processes are included.

## **Declaration of competing interest**

The authors declare that there are no conflicts of interest.

### Acknowledgments

The authors would like to thank Université de Lorraine for the grants provided on the project “manger droit” whose coordinator is François Allard-Huver.

### CRedit author statement

Aida Adjali: Writing- Original draft preparation; Igor Clarot, Zilin, Chen, and Eric Marchioni: Writing- reviewing and editing and Ariane Boudier: supervision and writing - reviewing and editing

### References

- [1]. R.R. Siva Kiran, G.M. Madhu, S.V. Satyanarayana, Spirulina in combating protein energy malnutrition (PEM) and protein energy wasting (PEW) - A review, *J. Nut. Res.* 3 (2016) 62-79.
- [2]. P. Jaouen, B. Lepine, N. Rossignol, et al., Clarification and concentration with membrane technology of a phycocyanin solution extracted from *Spirulina platensis*, *Biotechnol. Tech.* 13 (1999) 877–881.
- [3]. L.V. Venkataraman, *Spirulina platensis (Arthrospira)*: physiology, cell biology and biotechnologym, *J. Appl. Phycol.* 9 (1997) 295–296.
- [4]. P. Bermejo, E. Piñero, À.M. Villar, Iron-chelating ability and antioxidant properties of phycocyanin isolated from a protean extract of *Spirulinaplatensis*, *Food Chem.* 110 (2008) 436–445.
- [5]. L. Fang, C. Zhou, P. Cai, et al., Binding characteristics of copper and cadmium by cyanobacterium *Spirulina platensis*, *J. Hazard. Mater.* 190 (2011) 810–815.
- [6]. R. Chaiklahan, N. Chirasuwan, B. Bunnag, Stability of phycocyanin extracted from *Spirulina* sp.: Influence of temperature, pH and preservatives, *Process Biochem.* 47 (2012) 659–664.
- [7]. S.T. Silveira, J.F.M. Burkert, J.A.V. Costa, et al., Optimization of phycocyanin extraction from *Spirulina platensis* using factorial design, *Bioresour. Technol.* 98 (2007) 1629–1634.
- [8]. Q. Liu, Y. Huang, R. Zhang, et al., Medical application of *Spirulina platensis* derived C-phycocyanin, *Evid. Based Complement. Alternat. Med.* 2016 (2016), 7803846.
- [9]. M.K. Saini, K. Vaiphei, S.N. Sanyal, Chemoprevention of DMH-induced rat colon carcinoma initiation by combination administration of piroxicam and C-phycocyanin, *Mol. Cell. Biochem.* 361 (2012) 217–228.
- [10]. M. Gantar, S. Dhandayuthapani, A. Rathinavelu, Phycocyanin induces apoptosis and enhances the effect of topotecan on prostate cell Line LNCaP, *J. Med. Food.* 15 (2012) 1091–1095.
- [11]. R.P. Nishanth, B.S. Ramakrishna, R.G. Jyotsna, et al., C-Phycocyanin inhibits MDR1 through reactive oxygen species and cyclooxygenase-2 mediated pathways in human hepatocellular carcinoma cell line, *Eur. J. Pharmacol.* 649 (2010) 74–83.



- [12]. J.P. Kehrer, L.-O. Klotz, Free radicals and related reactive species as mediators of tissue injury and disease: implications for Health, *Crit. Rev. Toxicol.* 45 (2015) 765–798.
- [13]. I. Michalak, K. Chojnacka, Algae as production systems of bioactive compounds, *Eng. Life Sci.* 15 (2015) 160–176
- [14]. P. Yu, Y. Wu, G. Wang, et al., Purification and bioactivities of phycocyanin, *Crit. Rev. Food Sci. Nutr.* 57 (2017) 3840–3849.
- [15]. N.T. Eriksen, Production of phycocyanin: a pigment with applications in biology, biotechnology, foods and medicine, *Appl. Microbiol. Biotechnol.* 80 (2008) 1–14.
- [16]. G. Patil, S. Chethana, A.S. Sridevi, et al., Method to obtain C-phycocyanin of high purity, *J. Chromatogr. A.* 1127 (2006) 76–81.
- [17]. P.S. Corrêa, W.G. Morais Jr., A.A. Martins, et al., Microalgae biomolecules: extraction, separation and purification methods, *Processes* 9 (2021), 10.
- [18]. M. Hsieh-Lo, G. Castillo, M. Alberto Ochoa-Becerra, et al., Phycocyanin and phycoerythrin: Strategies to improve production yield and chemical stability, *Algal Res.* 42 (2019), 101600.
- [19]. I. Ílter, S. Akyil, Z. Demirel, et al., Optimization of phycocyanin extraction from *Spirulina platensis* using different techniques, *J. Food Compos. Anal.* 70 (2018) 78–88.
- [20]. D. Berns, R. MacColl, Phycocyanin in physical-chemical studies, *Chem. Rev.* 89 (1989) 807–825.
- [21]. L.M. Colla, C.D. Bertol, D.J. Ferreira, et al., Thermal and photo-stability of the antioxidant potential of *Spirulina platensis* powder, *Braz. J. Biol.* 77 (2017) 332–339.
- [22]. D. Dutta, A. Dutta, U. Raychaudhuri, et al., Rheological characteristics and thermal degradation kinetics of beta-carotene in pumpkin puree, *J. Food Eng.* 76 (2006) 538–546.
- [23]. D. Pez Jaeschke, I. Rocha Teixeira, L. Damasceno Ferreira Marczak, et al., Phycocyanin from *Spirulina*: A review of extraction methods and stability, *Food Res. Int.* 143 (2021), 110314.
- [24]. X.-J. Wu, H. Yang, Y.-T. Chen, et al., Biosynthesis of fluorescent  $\beta$  subunits of C-phycocyanin from *Spirulina subsalsa* in *Escherichia coli*, and their antioxidant properties, *Molecules.* 23 (2018), 1369.
- [25]. M.F. Falkeborg, M.C. Roda-Serrat, K.L. Burnaes, et al., Stabilising phycocyanin by anionic micelles, *Food Chem.* 239 (2018) 771–780.
- [26]. M.C.A. de Amarante, L.C.S. Corrêa Jr., L. Sala, et al., Analytical grade C-phycocyanin obtained by a single-step purification process, *Process. Biochem.* 90 (2020) 215–222.
- [27]. H. Scheer, W. Kufer, Conformational studies on C-phycocyanin from *Spirulina platensis*, *Zeitschrift Für Naturforschung C.* 32 (1977) 513–519
- [28]. F.S. Antelo, J.A.V. Costa, S.J. Kalil, Thermal degradation kinetics of the phycocyanin from *Spirulina platensis*, *Biochem. Eng. J.* 41 (2008) 43–47.
- [29]. L. Böcker, S. Ortmann, J. Surber, et al., Biphasic short time heat degradation of the blue microalgae protein phycocyanin from *Arthrospira platensis*, *Innov. Food Sci. Emerg. Technol.* 52 (2019) 116–121.
- [30]. A. Patel, R. Pawar, S. Mishra, et al., Kinetic studies on thermal denaturation of C-phycocyanin, *Indian J. Biochem. Biophys.* 41 (2004) 254–257.
- [31]. M. Faieta, L. Neri, G. Sacchetti, et al., Role of saccharides on thermal stability of phycocyanin in aqueous solutions, *Food Res. Int.* 132 (2020), 109093.

- [32]. I. Chentir, M. Hamdi, S.M. Li, et al., Stability, bio-functionality and bio-activity of crude phycocyanin from a two-phase cultured *Saharian Arthrospira* sp. strain, *Algal Res.* 35 (2018) 395–406.
- [33]. C. Couteau, S. Baudry, C. Roussakis, et al., Study of thermodegradation of phycocyanin from *Spirulina platensis*, *Sci. Aliments.* 24 (2004) 415–421.
- [34]. V.K. Kannaujiya, R.P. Sinha, Thermokinetic stability of phycocyanin and phycoerythrin in food-grade preservatives, *J. Appl. Phycol.* 28 (2016) 1063–1070.
- [35]. S.K. Mishra, A. Shrivastav, S. Mishra, Effect of preservatives for food grade C-PC from *Spirulina platensis*, *Process Biochem.* 43 (2008) 339–345.
- [36]. H.-L. Wu, G.-H. Wang, W.-Z. Xiang, et al., Stability and Antioxidant Activity of Food-Grade Phycocyanin Isolated from *Spirulina platensis*, *Int. J. Food Prop.* 19 (2016) 2349–2362.
- [37]. J. Aoki, D. Sasaki, M. Asayama, Development of a method for phycocyanin recovery from filamentous cyanobacteria and evaluation of its stability and antioxidant capacity, *BMC Biotechnol.* 21 (2021), 40.
- [38]. W.Y. Choi, H.Y. Lee, Kinetic analysis of stabilizing C-phycocyanin in the *Spirulina platensis* extracts from ultrasonic process associated with effects of light and temperature, *Appl. Sci.* 8 (2018), 1662.
- [39]. F.M.E. Escalante, D.A. Pérez-Rico, J.L. Alarcón-Jiménez, et al., Phycocyanin thermo-photostability: An accelerated life-test analysis, *J. Mex. Chem. Soc.* 64 (2020) 218–229.
- [40]. T. Saito, H. Ishikura, Y. Hada, et al., Photostabilization of phycocyanin and anthocyanin in the presence of biopterin- $\alpha$ -glucoside from *Spirulina platensis* under ultraviolet ray, *Dye. Pigment.* 56 (2003) 203–207.
- [41]. J.B. Moreira, A.L.M. Terra, J.A.V. Costa, et al., Development of pH indicator from PLA/PEO ultrafine fibers containing pigment of microalgae origin, *Int. J. Biol. Macromol.* 118 (2018) 1855–1862.
- [42]. M. Debreczeny, Z. Gombos, V. Csizmadia, et al., Chromophore conformational-analysis in phycocyanin and in related chromopeptides by surface enhanced Raman-spectroscopy, *Biochem. Biophys. Res. Commun.* 159 (1989) 1227–1232.
- [43]. European Medicines Agency, Q1B Photostability Testing of New Active Substances and Medicinal Products, <https://www.ema.europa.eu/en/ich-q1b-photostability-testing-new-active-substances-medicinal-products>. Accessed on December 13<sup>th</sup> 2021.
- [44]. M. Kissoudi, I. Sarakatsianos, V. Samanidou, Isolation and purification of food-grade C-phycocyanin from *Arthrospira platensis* and its determination in confectionery by HPLC with diode array detection, *J. Sep. Sci.* 41 (2018) 975–981.
- [45]. Y. Li, G. Aiello, C. Bollati, et al, Phycobiliproteins from *Arthrospira platensis* (*Spirulina*): A new source of peptides with dipeptidyl peptidase-IV inhibitory activity, *Nutrients.* 12 (2020), 794.
- [46]. C. Simó, M. Herrero, C. Neusüss, et al., Characterization of proteins from *Spirulina platensis* microalga using capillary electrophoresis-ion trap-mass spectrometry and capillary electrophoresis-time of flight-mass spectrometry, *Electrophoresis* 26 (2005) 2674–2683.
- [47]. Z. Zhang, Y. Li, A. Abbaspourrad, Improvement of the colloidal stability of phycocyanin in acidified conditions using whey protein-phycocyanin interactions, *Food Hydrocoll.* 105 (2020), 105747.
- [48]. K. Fukui, T. Saito, Y. Noguchi, et al., Relationship between color development and protein conformation in the phycocyanin molecule, *Dye Pigment.* 63 (2004) 89–94.



- [49]. A.R.C. Braga, F.D.S. Figueira, J.T.D. Silveira, et al., Improvement of thermal stability of C-phycoyanin by nanofiber and preservative agents, *J. Food Process. Preserv.* 40 (2016) 1264–1269.
- [50]. G. Martelli, C. Folli, L. Visai, et al., Thermal stability improvement of blue colorant C-Phycocyanin from *Spirulina platensis* for food industry applications, *Process. Biochem.* 49 (2014) 154–159.
- [51]. Hadiyanto, M. Christwardana, H. Sutanto, et al., Kinetic study on the effects of sugar addition on the thermal degradation of phycocyanin from *Spirulina* sp., *Food Biosci.* 22 (2018) 85–90.
- [52]. Y. Li, H. Yang, F.M. Cao, et al., The stability of C-phycoyanin doped silica biomaterials in UV irradiation, *J. Wuhan Univ. Technol. - Mater. Sci. Ed.* 24 (2009), 852.
- [53]. A. Gustiningtyas, I. Setyaningsih, S.D. Hardiningtyas, et al., Improvement stability of phycocyanin from *Spirulina platensis* encapsulated by water soluble chitosan nanoparticles, *IOP Conf. Ser.: Earth Environ. Sci.* 414 (2020), 012005.
- [54]. A. Chandralekha, H.S. Prashanth, H. Tavanandi, et al., A novel method for double encapsulation of C-phycoyanin using aqueous two phase systems for extension of shelf life, *J. Food Sci. Technol.-Mysore.* 58 (2021) 1750–1763.
- [55]. Z. Zhang, S. Cho, Y. Dadmohammadi, et al., Improvement of the storage stability of C-phycoyanin in beverages by high-pressure processing, *Food Hydrocoll.* 110 (2021), 106055.
- [56]. J.B. Moreira, L.-T. Lim, E.D. Rosa Zavareze, et al., Antioxidant ultrafine fibers developed with microalga compounds using a free surface electrospinning, *Food Hydrocoll.* 93 (2019) 131–136.
- [57]. F. da Silva Figueira, J. Garcia Gettens, J.A. Vieira Costa, et al., Production of nanofibers containing the bioactive compound C-phycoyanin, *J. Nanosci. Nanotechnol.* 16 (2016) 944–949.
- [58]. M. Yan, B. Liu, X. Jiao, et al., Preparation of phycocyanin microcapsules and its properties, *Food Bioprod. Process.* 92 (2014) 89–97.
- [59]. H. Hadiyanto, M. Christwardana, M. Suzery, et al., Effects of carrageenan and chitosan as coating materials on the thermal degradation of microencapsulated phycocyanin from *Spirulina* sp., *Int. J. Food Eng.* 15 (2019), 20180290.
- [60]. W. Pan-utai, S. Iamtham, Enhanced microencapsulation of C-phycoyanin from *Arthrospira* by freeze-drying with different wall materials, *Food Technol. Biotechnol.* 58 (2020) 423–432.
- [61]. D.A. Schmatz, D.J. da Silveira Mastrantonio, J.A. Vieira Costa, et al., Encapsulation of phycocyanin by electrospraying: A promising approach for the protection of sensitive compounds, *Food Bioprod. Process* 119 (2020) 206–215.
- [62]. H.N. Pradeep, C.A. Nayak, Enhanced stability of C-phycoyanin colorant by extrusion encapsulation, *J. Food Sci. Technol.-Mysore.* 56 (2019) 4526–4534.
- [63]. I. İter, M. Koç, Z. Demirel, et al., Improving the stability of phycocyanin by spray dried microencapsulation, *J. Food Process. Preserv.* 45 (2021), e15646.
- [64]. P.V.F. Lemos, L.C.F. Opretzka, L.S. Almeida, et al., Preparation and characterization of C-phycoyanin coated with STMP/ STPP cross-linked starches from different botanical sources, *Int. J. Biol. Macromol.* 159 (2020) 739–750.
- [65]. A.M. Nilamsari, A. Yunanda, H. Hadiyanto, Thermal degradation kinetics of phycocyanin encapsulation as an antioxidant agent, *IOP Conf. Ser.: Earth Environ. Sci.* 102 (2018), 012055

- [66]. L. Wu, C. Zhang, Y. Long, et al., Food additives: From functions to analytical methods, *Crit. Rev. Food Sci. Nutr.* (2021) <http://dx.doi.org/10.1080/10408398.2021.1929823>.
- [67]. Annex to the European Commission guideline on ‘Excipients in the labelling and package leaflet of medicinal products for human use’ (SANTE-2017-11668), [https://www.ema.europa.eu/en/documents/scientific-guideline/annex-european-commission-guideline-excipients-labelling-package-leaflet-medicinal-products-human\\_en.pdf](https://www.ema.europa.eu/en/documents/scientific-guideline/annex-european-commission-guideline-excipients-labelling-package-leaflet-medicinal-products-human_en.pdf). (Accessed 21 September 2021).
- [68]. Y. Zhong, L. Wu, X. Chen, et al., Effects of food-additive-information on consumers’ willingness to accept food with additives, *Int. J. Environ. Res. Public. Heal.* 15 (2018), 2394.
- [69]. P. Varela, S.M. Fiszman, Exploring consumers’ knowledge and perceptions of hydrocolloids used as food additives and ingredients, *Food Hydrocoll.* 30 (2013) 477–484.
- [70]. L. Shan, Q. Zang, L. Xu, et al., A literature review of public perception of food additive safety, *Agro Food Ind. Hi-Tech.* 26 (2015) 49–51.
- [71]. P. Miao, S. Chen, J. Li, et al., Decreasing consumers’ risk perception of food additives by knowledge enhancement in China, *Food Qual. Prefer.* 79 (2020), 103781.
- [72]. A. Bearth, M.-E. Cousin, M. Siegrist, The consumer’s perception of artificial food additives: Influences on acceptance, risk and benefit perceptions, *Food Qual. Prefer.* 38 (2014) 14–23.
- [73]. W. Wang, S. Ohtake, Science and art of protein formulation development, *Int. J. Pharm.* 568 (2019), 118505.
- [74]. G. Manu, N. Firdose, M.K. Jayanthi, et al., Current status and perspectives of oral therapeutic protein and peptide formulations: A review, *J. Pharm. Res. Int.* 33 (2021) 117–137.
- [75]. S. He, N. Joseph, S. Feng, et al., Application of microfluidic technology in food processing, *Food Funct.* 11 (2020) 5726–5737.
- [76]. S.P. Cape, J.A. Villa, E.T.S. Huang, et al., Preparation of active proteins, vaccines and pharmaceuticals as fine powders using supercritical or near-critical fluids, *Pharm. Res.* 25 (2008) 1967–1990.

### **Figure captions**

Fig. 1. Applications of phycocyanin: green represents already marketed products whereas orange represents products under development.

Fig. 2. Structure of (A)  $\alpha\beta$  monomer of phycocyanin and (B) phycocyanobilin.

Table 2. Some extracted results (selected according to compare almost same temperatures) after fitting the data to the first-order kinetic model.

T (°C)	pH 5.0			pH 6.0			pH 7.0			Refs.
	$k$ (min <sup>-1</sup> )	$t_{1/2}$ (min)	Ea (kJ/mol)	$k$ (min <sup>-1</sup> )	$t_{1/2}$ (min)	Ea (kJ/mol)	$k$ (min <sup>-1</sup> )	$t_{1/2}$ (min)	Ea (kJ/mol)	
47	0.006±0.003	116.5±22.3	100.3 <sup>a</sup>	0.0022±0.0001	309.4±12.0	120.32 <sup>a</sup>	0.0054±0.0005	128.6±13.3	116.68 <sup>a</sup>	[6]
55	0.0202±0.0013	34.4±2.2		0.0075±0.0018	92.7±23.2		0.0146±0.0034	47.5±8.9		
69	0.0928±0.0102	7.5±1.7		0.0481±0.0064	14.5±4.2		0.1156±0.0328	6.0±1.7		
74	0.1068±0.0232	6.5±1.4		0.0707±0.0030	9.7±1.6		0.1361±0.0256	5.3±1.0		
50	5.99×10 <sup>-4</sup>	1.155	387.14 <sup>a</sup>	4.79×10 <sup>-4</sup>	1.444	559.96 <sup>a</sup>	1.19×10 <sup>-3</sup>	0.577	202.70 <sup>a</sup>	[28]
55	1.20×10 <sup>-2</sup>	0.057		2.99×10 <sup>-3</sup>	0.231		5.98×10 <sup>-3</sup>	0.115		
57	2.40×10 <sup>-2</sup>	0.028		3.59×10 <sup>-2</sup>	0.019		8.38×10 <sup>-3</sup>	8.250		
60	6.59×10 <sup>-2</sup>	10.500		0.220	3.033		0.01	6.416		
45	9.58×10 <sup>-4</sup>	/	92.048 <sup>b</sup>	3.23×10 <sup>-4</sup>	/	154.48 <sup>b</sup>	1.14×10 <sup>-3</sup>	/	158.99 <sup>b</sup>	[30]
55	2.63×10 <sup>-3</sup>	/		2.51×10 <sup>-3</sup>	/		4.55×10 <sup>-3</sup>	/		
65	8.98×10 <sup>-3</sup>	/		1.14×10 <sup>-2</sup>	/		4.37×10 <sup>-2</sup>	/		
50	0.0022±0.0000	321.3±5.5	78.5 <sup>a</sup>	0.0014±0.0000	495.0±9.9	103.1 <sup>a</sup>	0.0028±0.0001	243.6±10.6	83.3 <sup>a</sup>	[36]
55	0.0036±0.0000	193.0±0.6		0.0027±0.0002	253.9±17.1		0.0058±0.0001	119.1±2.4		
65	0.0116±0.0000	58.8±0.1		0.0133±0.0005	52.0±2.0		0.0210±0.0004	33.0±0.7		
75	0.0165±0.0002	42.1±0.4		0.0194±0.0011	35.7±2.0		0.0242±0.0012	28.6±1.4		

Kinetic constant ( $k$ ), half-life ( $t_{1/2}$ ) and activation energy (Ea) at pH 5.0, 6.0 and 7.0. <sup>a</sup>Ea is calculated using Arrhenius' equation:  $k = A \times e^{-\frac{Ea}{RT}}$  with  $k$  as the kinetic constant,  $A$  as the pre-exponential factor, Ea, as the activation energy,  $R$  as the universal gas constant,  $T$  as the absolute temperature and using the slope of the graph  $\ln K = f\left(\frac{1}{T}\right)$ . For each study a linear relation was obtained with a correlation coefficient  $>0.903$ . <sup>b</sup> Indicated in the article.

Table 3. Chemical compounds used to improve the storage.

Phycocyanin			Conditions		Preservatives	Methods to study the stability	Model to interpret the results	Main result of the study	Refs.
Supplier or extraction method	Purity (ratio $A_{620}/A_{280}$ when indicated)	Initial concentration and medium	pH	T (°C)					
Extraction in phosphate buffer, centrifugation and filtrations	1.43	1 mg/mL Phosphate citrate buffer (pH 5, 6, 7)	7.0	60	Glucose 2.5% to 40% ( <i>m/m</i> ). saccharose 2.5% to 40% ( <i>m/m</i> ). sorbitol 20 % ( <i>m/m</i> ), sodium chloride 2.5% to 20% ( <i>m/m</i> ), ascorbic acid 2.5% ( <i>m/m</i> ), citric acid 2.5% ( <i>m/m</i> ), sodium azide 2.5% ( <i>m/m</i> )	Spectrophotometry UV-vis and scanning electron microscopy	First-order rate law	Glucose. sucrose and NaCl preserve the protein stability	[6]
Filtration	0.46	3.68 mg/mL In water	pH 5, 6, 7	62	Sorbitol 10% to 50 % m/m	Spectrophotometry UV-vis	First-order rate law	Sorbitol improves the stability of phycocyanin even at 10%	[28]
Supplier Sigma-Aldrich and GNT International BV	Reagent grade (1.65) and food grades (1.97 and 0.64)	Phosphate buffer 0.1 M pH 7.0	-	50, 65, 80, 90, 100	Saccharose 20%, 40%, 70% ( <i>m/m</i> ), trehalose 20%, 40% ( <i>m/m</i> )	Spectrophotometry UV-vis and water activity measurement	Weibull model	Sucrose stabilizes more phycocyanin than trehalose <sup>5</sup>	[31]
Precipitation with ammonium sulfate and centrifugations	Reagent grade 1.24	0.5. 2.75 and 5 mg mL <sup>-1</sup> . prepared in 50 mM sodium phosphate buffer (pH 6.0)	6.0	60	Polyethylene glycol-4000 (PEG) or sorbitol and sucrose at a ratio of 1:4 ( <i>m/m</i> ) of phycocyanin for stabilizer	Spectrophotometry UV-vis	None	Stability and antioxidant and antibacterial activity of phycocyanin	[32]

Sonication and centrifugation from <i>Nostoc</i> sp. strain HKAR-2	3.18	0.1 mg/mL in potassium phosphate buffer at pH 7.0	7.0	4, 25, 40	Calcium chloride, ascorbic acid, sucrose, citric acid, and benzoic acid at 0.5, 2.5, and 5 mM	Spectrophotometry UV-vis	First-order rate law	Benzoic acid is the best preservative	[34]
Precipitation with ammonium sulfate and centrifugations	2.25	0.4 mg/mL Phosphate buffer 0.1 M (pH 7.0)	-	0 and 35	Saccharose (4 g/L), calcium chloride (4 g/L), citric acid (4 g/L) and a combination of previous cited compounds at 4 g/L each or 2 g/L	Spectrophotometry UV-vis and differential scanning calorimetry	None	Citric acid is the best preservative	[35]
Precipitation with ammonium sulfate and centrifugations	1.5	0.4 mg/mL Phosphate citrate buffer pH 5.0, 6.0, 7.0, 8.0	5, 6, 7, 8	65	Saccharose. glucose. sodium chloride 20% (m/m)	Spectrophotometry UV-vis	First-order rate law	Sodium chloride stabilizes phycocyanin in a concentration-dependent manner	[36]
Centrifugation	-	-	-	25 to 70 °C in the presence of UV light 254 nm. 40W	Biopterin- $\alpha$ -glucoside	Spectrophotometry UV-vis and spectrofluorimetry	-	Photostabilization of phycocyanin in presence of biopterin- $\alpha$ -glucoside	[40]
Extracted	1.0	-	-	55, 60, 65, 70, 75	Sorbitol (50%), saccharose and glucose (20%), sodium chloride (2.5%), polyethyleneoxide (6%)	Spectrophotometry UV-vis	First-order rate law	Enhancement of phycocyanin stability when using glucose or sorbitol or with nanofibers	[49]
Centrifugation and filtrations	1.4	0.02 to 1.3 mg/mL in phosphate buffer	5, 7, 9	50, 60, 70, 80	Conventional honey. honey from <i>Leptospermum scoparium</i> . fructose (62%, m/V) glucose (37%, m/V) and saccharose (54%, m/V)	Spectrophotometry UV-vis	None	Fructose is the best preservative	[50]
Supplied from CV Neoalgae (Sukoharjo. Indonesia)	-	1 mg/mL in citrate buffer (pH 6.0)	6.0	40, 60, 80	Glucose. sucrose. fructose 10 to 15% (m/m)	Spectrophotometry UV-vis	First-order rate law	Fructose preserves phycocyanin color while	[51]



Table 4. Particulate forms used to improve the storage.

Type of formulation and technology employed	Size of the particles (method used)	Main result concerning the stability of phycocyanin	Refs.
Sodium dodecylsulfate micelles	Not indicated but usually few nanometers	Stabilizing effect of the colour even at acidic pH and at high temperatures	[25]
Complexes with $\alpha$ -lactalbumin, $\beta$ -lactoglobulin, BSA, immunoglobulins, or glycomacropptides	90 to 120 nm	Improvement of phycocyanin stability in acidified solutions	[47]
Nanofibers produced by electrospinning	Average diameter of 295 nm	Better thermostability of phycocyanin in nanofibers (compared to native protein. enhancement by a 2-fold factor) but in the same range compared to the use of preservatives	[49]
Encapsulated in a hydrogel to dope silica materials	30-40 nm	Improvement of the photostability of phycocyanin	[52]
Chitosan nanoparticles	Average diameter of 457 nm	Enhanced thermal stability when encapsulated (90 min at 50 °C)	[53]
Double emulsion by an aqueous two-phase system	8.8 and 380.5 $\mu$ m depending on the experimental conditions	Up to 6 months of stability	[54]
Complexes formed with whey and $\kappa$ -carrageenan	660 to 3925 nm depending on the experimental conditions	Improvement of the photostability of phycocyanin	[55]
Fibers produced by electrospinning	Mean diameters of 196 to 542 nm	Enhancement of the thermal stability when encapsulated (increase of the initial temperature of degradation of the protein measured by thermogravimetry)	[56]
Fibers produced by electrospinning	Average diameter of 295 to 760 nm	Enhanced thermal resistance of phycocyanin	[57]
Microcapsules produced by extrusion	Average diameter of 1.37 to 2.54 mm	Improvement of the stability against temperature increase in microcapsules. No improvement against light degradation	[58]
Microencapsulation using chitosan or $\kappa$ -carrageenan	2 to 4 $\mu$ m	Improvement of phycocyanin stability when microencapsulated	[59]
Microencapsulation	1.5 to 316 $\mu$ m	Better thermal stability when microencapsulated	[60]
Microparticles based on polyvinylalcohol produced by electrospaying	Average diameter of 395 nm	Thermal stability improved in microparticles up to 216 °C	[61]
Microencapsulation using extrusion	1.2 mm	Better stability at high temperatures	[62]
Microencapsulation by emulsion and spray-drying	Not reported	Material used for microencapsulation affects phycocyanin stability	[63]
Cross-linked starches-C-phycocyanin composites	Not reported	Stabilization within the composites	[64]
Microencapsulation using extrusion	Not reported	Better stability of phycocyanin towards heat stress linked to a better antioxidant activity	[65]





Table 1. Experimental conditions used to study phycocyanin degradation.

Phycocyanin from <i>Spirulina</i> except when mentioned			Degradation conditions			Method to study the stability	Model to interpret the results	Focus of the study	Refs.
Supplier or extraction method	Purity (ratio $A_{620}/A_{280}$ when indicated)	Initial concentration and medium	pH	Temperature (°C)	Others				
Extraction in phosphate buffer centrifugation and filtrations	1.43	1 mg/mL Phosphate citrate buffer (pH 5, 6, 7)	5.0, 5.5, 6.0, 6.5, 7.0	26, 31, 35, 39, 43, 47, 51, 55, 59, 64, 69, 74	-	Spectrophotometry UV-vis	First-order rate law	Stability of phycocyanin and the use of preservatives	[6]
Filtration	0.46	3.68 mg/mL Water	5, 6, 7	50, 53, 55, 57, 60, 62, 65	-	Spectrophotometry UV-vis	First-order rate law	Kinetic study of thermal degradation of phycocyanin	[28]
Supplier GNT International BV	Not indicated	1 mg/mL Phosphate buffer 0.1 M, pH 7.3	-	70, 75, 80	-	Spectrophotometry UV-vis	Sixth-order rate law	Thermal color degradation kinetics of a phycocyanin for short time	[29]
Precipitation with ammonium sulfate and centrifugations	>4.0	0.33 mg/mL Phosphate buffer (0.005 M, pH 4.0 to 8.0) Acetate buffer (pH 4.0 to 5.0)	4.0, 8.0	45, 55, 65	-	Spectrophotometry UV-vis	First-order rate law	Kinetic study of phycocyanin degradation as a function of the pH and the temperature	[30]
Supplier Sigma-Aldrich and GNT International BV	Reagent grade (1.65) and food grades (1.97 and 0.64)	Phosphate buffer 0.1 M pH 7.0	7.0	50, 65, 80, 90, 100	-	Spectrophotometry UV-vis and circular dichroism	Weibull model	Saccharides and water play an important role in phycocyanin stability	[31]
Precipitation with ammonium sulfate and centrifugations	Reagent grade 1.24	0.5, 2.75 and 5 mg/mL), prepared in 50 mM sodium phosphate buffer (pH 6.0)	6.0	60	-	Spectrophotometry UV-vis	None	Stability and antioxidant and antibacterial activity of phycocyanin	[32]
Supplier Linablue-A, Dainippon Ink & Chemicals	-	400 mg/mL in water	2.0, 6.5, 8.0	25, 35, 45	-	Spectrophotometry UV-vis	First-order rate law	Thermal degradation of phycocyanin at various pH	[33]

Sonication and centrifugation from <i>Nostoc</i> sp. strain HKAR-2	3.18	0.1 mg/mL in potassium phosphate buffer at pH 7.0	7.0	4, 25, 40	-	Spectrophotometry UV-vis	First-order rate law	Thermal stability in presence of preservatives	[34]
Precipitation with ammonium sulfate and centrifugations	2.25	0.4 mg/mL Phosphate buffer 0.1 M (pH 7.0)	-	0, 35	Presence of urea: 1 to 10 M	Spectrophotometry UV-vis, differential scanning calorimetry	None	Screening of preservatives to improve the stability of phycocyanin	[35]
Precipitation with ammonium sulfate and centrifugations	1.5	0.4 mg/mL Phosphate citrate buffer pH 3.0, 4.0, 5.0, 6.0, 7.0, 8.0	3.0, 4.0, 5.0, 5.5, 6.0, 7.0, 8.0	25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75	Light: intensity: 50 and 100 $\mu\text{mol}/\text{m}^2\cdot\text{s}^1$	Spectrophotometry UV-vis	First-order rate law	Stability and antioxidant activity of phycocyanin	[36]
Extracted from <i>Pseudanabaena</i> sp. ABRG5-3, <i>Limnithrix</i> sp. SK1-2-1, <i>Spirulina platensis</i> NIES-39 via centrifugation and freeze-drying and water addition	3.10, 2.14, or 1.76 depending on the strains	4 mg/mL in deionised water then at 0.8 mg/mL in phosphate-citrate buffer (0.15 M at pH 4, 5, 7)	4.0, 5.0, 7.0	30 or 55	Light: 100 $\mu\text{mol}/\text{m}^2\cdot\text{s}^1$	Spectrophotometry UV-vis	None	Stability and antioxidant capacity of phycocyanin extracted from various cyanobacteria	[37]
Ultra-sound and centrifugation	-	Water	-	4, 25, 40	Light: 20 W white fluorescent lamp	HPLC-Vis	First-order rate law and second-order with light	Kinetic model to reflect the stability of phycocyanin from ultrasonic extraction process	[38]
Supplier Biotecnología Mexicana de Microalgas S.A. de C.V	0.75	333 $\mu\text{g}/\text{mL}$ in distilled water	-	5, 18, 44, 57	Light: 0, 65, 130 $\mu\text{mol}/\text{m}^2\cdot\text{s}^1$	Spectrophotometry UV-vis	Weibull model and others	Thermo-photostability of phycocyanin	[39]
Centrifugation	-	-	-	25 to 70	Light: UV light 254 nm, 40W	Spectrophotometry UV-vis and spectrofluorimetry	-	Photostabilization of phycocyanin in presence of biopterin- $\alpha$ -glucoside	[40]

-: no data.





## Highlights

- Phycocyanin is in great demand for industrial application
- Phycocyanin is sensitive to pH, temperature, and light
- Optimal stability occurs between pH 5.5–6.0 and at temperatures <45 °C in the dark
- The use of preservatives or its encapsulation with polymers enhances its stability