



HAL
open science

Potassium binders for the prevention of hyperkalaemia in heart failure patients: implementation issues and future developments

Faiez Zannad, João Pedro Ferreira, Bertram Pitt

► To cite this version:

Faiez Zannad, João Pedro Ferreira, Bertram Pitt. Potassium binders for the prevention of hyperkalaemia in heart failure patients: implementation issues and future developments. *European Heart Journal Supplements*, 2019, 21 (Supplement_A), pp.A55-A60. 10.1093/eurheartj/suy034. hal-03764606

HAL Id: hal-03764606

<https://hal.univ-lorraine.fr/hal-03764606v1>

Submitted on 30 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Potassium binders for the prevention of hyperkalaemia in heart failure patients: implementation issues and future developments

Faiez Zannad^{1*}, João Pedro Ferreira^{1,2}, and Bertram Pitt³

¹Université de Lorraine INSERM, Centre, d'Investigations Cliniques Plurithématique 1433, INSERM U1116, CHRU de Nancy, F-CRIN INI-CRCT, France;

²Department of Physiology and Cardiothoracic Surgery, Cardiovascular Research and Development Unit, Faculty of Medicine, University of Porto, Portugal; and

³Division of Cardiology, Department of Medicine, University of Michigan School of Medicine, Ann Arbor, MI, USA

KEYWORDS

Potassium binders;
Trials;
Outcomes

New therapeutic options to treat hyperkalaemia, such as potassium binders, have been suggested as potentially beneficial by allowing the maintenance (or increase) of the dose of medications that improve outcomes in several cardiovascular conditions, but which have in common the propensity for raising serum potassium. However, potassium binding drugs have yet to prove their causal association with improvements in patients' prognosis before their widespread use can be recommended. In this review we provided an up-to-date appraisal on potassium binders, their potential clinical applications and directions for future research.

Introduction

Potassium is the most abundant intracellular cation and is essential for normal cellular function. Derangements of potassium regulation often lead to life-threatening neuromuscular, gastrointestinal, and cardiac abnormalities if left untreated.

New therapeutic options to treat hyperkalaemia, such as potassium binders, have been suggested as potentially beneficial by allowing the maintenance (or increase) of the dose of medications that improve outcomes in several cardiovascular conditions, but which have in common the propensity for raising serum potassium. However, potassium-binding drugs will need to be shown to improve clinical outcomes before their widespread use can be recommended.

This article focuses on implementation issues and future directions for the treatment of hyperkalaemia. We use

examples from the heart failure (HF) field which is our area of expertise.

Potassium homeostasis alterations and their association with outcomes

Neuro-hormonal antagonists have driven a remarkable prognostic improvement in patients with heart failure and reduced left ventricular ejection fraction (HF-REF). However, both β -blockers and renin-angiotensin-aldosterone system inhibitors (RAASi) may cause hypotension, and RAASi, in particular, may decrease estimated glomerular filtration rate (eGFR) and increase serum potassium levels. More recently, sacubitril/valsartan has been shown to improve morbidity and mortality compared to enalapril with less frequent hyperkalaemia and eGFR deterioration.¹⁻³

The occurrence of hyperkalaemia has been highlighted in several observational studies and, despite the low-grade evidence, hyperkalaemia has become a major concern for clinicians, particularly in association with the use of mineralocorticoid receptor antagonists (MRAs).⁴ Consequently,

*Corresponding author. Tel: +33 (0) 3 83 15 73 15, Fax: +33 (0) 3 83 15 73 24, Email: zannad@chru-nancy.fr

MRAs have been persistently under-prescribed and under dosed despite their remarkable improvement in clinical outcomes and Class IA guideline indication for patients with HF-REF.⁵⁻¹² Once they are initiated they are often discontinued due to an increase in serum potassium or a decrease in renal function.¹³⁻¹⁵ Furthermore once a MRA is administered patients often do not receive guideline-recommended monitoring of serum potassium or renal function. On the other hand, evidence from prospective randomized trials has consistently demonstrated a low rate of hyperkalaemia without any hyperkalaemia-associated fatal events.^{10,11,16-19} In the large-scale randomized trials of MRAs in patients with HF including RALES, EPHEsus, EMPHASIS-HF, and TOPCAT, the incidence of hyperkalaemia (serum K^+ >5.5 mmol/L) was low (up to a maximum of 12%), and there has not been, to our knowledge, a single death attributable to hyperkalaemia in a patient randomized to a MRA in these trials, including >7000 patients taking MRAs. Therefore, a discrepancy exists between the evidence from randomized clinical trials and daily clinical practice. Awareness and concern about hyperkalaemia may have been enhanced by the recent introduction of new potassium-binding therapies, putting undue emphasis on the hyperkalaemia 'issue'. Actually, while hyperkalaemia is an important issue, it is possible that an increase in serum potassium contributes to the beneficial mechanism of RAASi, particularly reduced rates of sudden cardiac death associated with the administration of a MRA.^{10,16} *Post hoc* analyses of the RALES and EMPHASIS-HF trials show that the outcome benefits of a MRA, compared with placebo, were maintained in patients who developed moderate hyperkalaemia (potassium levels between 5.5 mmol/L and 6.0 mmol/L) during treatment.^{20,21} Presumably, once patients manifest potassium levels in this range, many had study drug and/or concomitant RAASi reduced or discontinued, raising the question whether treatment benefit might have been further enhanced if potassium binders had been used, thereby facilitating preservation of RAASi dosing. Whether the use of potassium binders may improve outcome through enabling optimal use of RAASi is still an open question.^{22,23} In HF, the association of K^+ with mortality has been described as U-shaped i.e. patients with both low- and high-potassium levels have increased death rates.^{24,25} For example, K^+ levels below 4.0 mmol/L are associated with increased death rates with similar risk as K^+ levels above 5.0 mmol/L. Of note, patients with K^+ below 3.0 mmol/L may have higher mortality risk than patients with K^+ above 5.5 mmol/L. U-shaped associations have been described in HF but also in other settings, such as chronic kidney disease (CKD),²⁶ hypertension,²⁷ myocardial infarction,^{28,29} and in the general population.³⁰ Despite the narrow nadir of optimal K^+ levels, clinical attitudes towards serum K^+ levels vary widely, with some clinicians becoming concerned when the serum K^+ is <3.5 or >5.5 mmol/L, whereas others become concerned and consider altering cardiovascular drug therapy at a serum K^+ <4.0 or >5.0 mmol/L.²⁹ Generally, the associations between K^+ levels and clinical outcomes are derived from observational data prone to a number of confounding factors. Reverse causation is one of them, i.e. K^+ alterations arise in consequence of other underlying illnesses or treatments

responsible for the outcomes associations (e.g. hyperkalaemia is frequently caused by haemolysis or renal failure— itself associated with RAASi drug discontinuation, whereas hypokalaemia is often caused by diuretics often used to treat patients with severe signs and symptoms of systemic congestion or high blood pressure).

A rise in potassium level represents a frequent cause for RAASi dose reduction or discontinuation—actions that may deprive patients of therapy proven to improve clinical outcomes.^{7,21} Although optimal dosing of MRAs for outcome benefit has not been explored, for other RAASi antagonists the clinical benefit has been shown to be dose-dependent, despite an increased risk of hyperkalaemia.^{31,32} For these reasons, management of the adverse effects of RAASi, without discontinuing or reducing the dose of the therapeutic agent, may represent an attractive goal. Thus, the recent availability of safe and tolerable oral potassium binders may change the approach to managing hyperkalaemia and RAASi usage.^{22,33} However, without a clear demonstration of outcome improvement with these agents their widespread use cannot be recommended. More research and education about hyperkalaemia may help increase awareness about this issue and promote better clinical practice. This should include: (i) identifying patients at risk of hyperkalaemia, (ii) preventing hyperkalaemia with available 'life-style' changes, including dietary changes, (iii) monitoring serum potassium as per international guidelines, and (iv) treating emerging rise of potassium with dose adjustments of drugs likely to increase serum potassium, and/or using potassium binders. One should use good clinical judgement to mitigate an increase in serum potassium levels before adding another drug to the polypharmacy of a HF patient. Finally, education should be more generally applied to potassium haemostasis. Hypokalaemia is also an often-overlooked cause of iatrogenic mortality among patients with cardiovascular conditions. Focus on hyperkalaemia should not shift the attention to only 'one side of the coin'.

Potassium binders

The majority of potassium is renally excreted, but $\approx 5-10\%$ is secreted in the colon. Two new agents, patiromer and sodium zirconium cyclosilicate (SZC), were developed for the treatment of hyperkalaemia. These agents may offer advantages over existing approaches to hyperkalaemia treatment (e.g. kayexylate). Both patiromer and SZC act to remove potassium by exchanging cations (calcium and sodium for patiromer and SZC, respectively) for potassium in the gastrointestinal tract, binding potassium, and increasing its faecal excretion.³⁴

Patiromer clinical trials

The PEARL-HF study (Evaluation of RLY5016 in Heart Failure Patients) was a multicentre, randomized, double-blind, placebo-controlled parallel-group study to evaluate the effects of patiromer in 120 patients with HF.³⁵ All patients had either an eGFR of <60 mL/min or a documented history of RAASi discontinuation because of hyperkalaemia within 6 months. Spironolactone 25 mg/day was

initiated in all patients. At 28 days, serum potassium was significantly lower in the patiromer group compared with placebo and fewer patients randomized to patiromer experienced serum potassium >5.5 mEq/L (7% vs. 25%; $P=0.015$). However, more patiromer-treated patients had serum potassium <3.5 mEq/L (6% vs. 0%; $P=0.094$) and hypomagnesaemia (<1.8 mg/dL) was reported in 24% of the patiromer patients and 2% of the placebo patients ($P=0.001$). Overall, patiromer was well tolerated.

The AMETHYST-DN (RLY5016 in the Treatment of Hyperkalaemia in Patients With Hypertension and Diabetic Nephropathy) trial³⁶ was a multicentre, randomized, open-label, dose-ranging study, randomizing 304 patients with type 2 diabetes mellitus and CKD (eGFR 15 to <60 mL/min/ 1.73 m²) who developed hyperkalaemia in the setting of RAASi optimization to either patiromer or 'control'. Serum potassium fell with patiromer treatment and rose after patiromer discontinuation. Serum potassium <3.5 mEq/L occurred in 17 patients (5.6%). Hypomagnesaemia (7.2%), and constipation (4.6%) were the most commonly reported treatment-related adverse events.

The OPAL-HK (Study Evaluating the Efficacy and Safety of Patiromer for the Treatment of Hyperkalaemia),²³ examined the effects of patiromer on serum potassium levels of 102 patients with CKD (eGFR 15 to <60 mL/min/ 1.73 m²) and serum potassium of 5.1 to <6.5 mmol/L. In this study, patiromer treatment decreased serum potassium levels and reduced hyperkalaemia recurrence compared with placebo. Hypokalaemia (serum potassium <3.8 mmol/L) was observed in 5% and 2% of the patiromer and placebo groups, respectively. Hypomagnesaemia occurred in 8 (3%) patiromer patients during the initial treatment phase.

Sodium zirconium cyclosilicate clinical trials

The sodium zirconium cyclosilicate in hyperkalaemia³⁷ randomized 753 patients with serum potassium between 5.0 and 6.5 mmol/L to either SZC or placebo for 48 h. No specific requirements for eGFR (excluding dialysis) or RAASi use were specified. The decrease in serum potassium was greater with SZC compared with placebo. Adverse events were rare and similar between SZC and placebo.

The HARMONIZE (Hyperkalaemia Randomized Intervention Multidose SZC Maintenance) trial²² studied the effects of SZC on serum potassium levels in 95 HF patients with baseline $K^+ \geq 5.1$ mmol/L. Among these, 93% achieved the target potassium level of 3.5–5 mmol/L within 48 h of receiving open-label SZC without adjusting RAASi doses. Oedema was reported in 2.4% of the placebo group and up to 14.3% with 15 g SZC dosing. Serum potassium of <3.5 mmol/L was observed in 10% of patients in the SZC 10 g group and 11% of patients in the SZC 15 g group vs. no cases in the 5 g or placebo groups. No clinically significant changes in serum magnesium, phosphate, or bicarbonate were observed.

Taken together, these studies demonstrate that both patiromer and SZC are effective for reducing hyperkalaemia. However, both agents increase the rates of hypokalaemia and patiromer increased rates of hypomagnesaemia, and SZC, in particular, is associated with increased oedema in patients with HF.

Potassium binders for the prevention of hyperkalaemia in heart failure patients: the equipoise

The benefit of potassium-binding agents in controlling hyperkalaemia in patients' at risk, needs to be substantiated with further studies determining whether administration of potassium-binding agents will allow maximizing the use of RAASi in patients at risk of hyperkalaemia, and consequently improving clinical outcomes. The current state of equipoise may be summarized as follows:

- In daily practice, physicians may be less permissive to hyperkalaemia due to concern about its consequences.
- Yet, monitoring of serum potassium in daily practice, including in hyperkalaemia risk situations, is suboptimal and below guideline recommendations.
- Inertia is the likely main driver of suboptimal monitoring of serum potassium.
- Inertia is also one reason for lack of optimal monitoring, because time and resource consuming, HF drug-dose adjustments, including drug re-initiation or re-up titration after drug discontinuation or down titration, in response to rise in potassium.
- An increase in serum creatinine after initiation of a MRA may be due to reversible haemodynamic factors rather than renal injury.
- Down-titration of RAASi in response to hyperkalaemia and or an increase in serum creatinine may be unnecessarily conservative for many patients, provided that potassium levels and kidney function are carefully monitored.³⁸
- Potassium-binding agents may enable using life-saving RAASi at target doses.
- Potassium-binding agents may enable optimal use of life-saving RAASi and therefore may improve clinical outcomes.

RAASi therapy is being refined

In the PARADIGM-HF trial,² sacubitril/valsartan (an angiotensin receptor blocker combined with a neprilysin inhibitor), when used concomitantly with an MRA, resulted in lower hyperkalaemia rates when compared to the concomitant use of enalapril and an MRA. This lower risk of hyperkalaemia was also observed among participants who were not treated with an MRA at baseline but who were initiated on MRA therapy during the course of the study.² These data from the PARADIGM-HF trial suggest that the substitution of an angiotensin-converting enzyme inhibitor (ACEi) or angiotensin receptor blocker (ARB) for sacubitril/valsartan may enhance the potassium safety and tolerability of MRAs in suitable HF-REF patients by moderating the risk of serious hyperkalaemia. Therefore, future studies of potassium binders might be needed specifically within the context of use of HF medications, including sacubitril/valsartan.

Several new non-steroidal MRAs are being developed. The one that has undergone the most extensive clinical evaluation is finerenone.³⁹ Finerenone is tightly bound to

Table 1 Spironolactone dose adjustment proposal

Serum K ⁺ (mmol/L)	Dose adjustment
<4.5	Baseline: eGFR ≥50 mL/min/1.73 m ² → spironolactone dose = 25 mg/day eGFR 30-49 mL/min/1.73 m ² → spironolactone dose = 25 mg every other day Increase dose: If spironolactone dose = 25 mg/day → increase to 50 mg/day If spironolactone dose = 25 mg every other day → increase to 25 mg/day
4.5-5.4	No adjustment recommended
5.5-5.9	Decrease dose: If spironolactone dose = 50 mg/day → decrease to 25 mg/day If spironolactone dose = 25 mg/day → decrease to 25 mg very other day If spironolactone dose = 25 mg every other day → stop treatment
≥6.0	Stop treatment and reintroduce when serum K ⁺ ≤4.5 mmol/L Stop treatment at any case if eGFR ≤30 mL/min/1.73 m ² and reintroduce upon clinical decision

and highly specific for the MRA. In the ARTS trial,⁴⁰ finerenone was compared to spironolactone in patients with HF-REF and CKD. The incidence of hyperkalaemia was less frequent with finerenone compared with spironolactone (5.3% vs. 12.7%, $P=0.048$). In the ARTS-HF study,⁴¹ finerenone was compared to eplerenone in patients with chronic HF-REF plus CKD and/or diabetes mellitus. A potassium level increase to ≥5.6 mmol/L at any time point occurred in 4.3% of patients without differences between finerenone and eplerenone. Finerenone is currently under evaluation in patients with diabetic nephropathy (FIDELIO trial: NCT02540993) and in patients with renal disease at increased risk for cardiovascular events (FIGARO trial: NCT02545049). These patients are also at increased risk of hyperkalaemia. It has to be seen whether finerenone will be 'potassium-safe' and whether potassium binders might be useful in this context of use. Unfortunately, finerenone is not currently being developed for HF.

How to manage the risk of hyperkalaemia?

Most often, hyperkalaemia is predictable, manageable, and reversible. Risk factors for hyperkalaemia are well documented and include: advanced stages of CKD (frequency up to 40-50%),³⁰ chronic HF (frequency up to 30%),²¹ diabetes mellitus (frequency up to 17%),⁴² resistant hypertension with add-on MRA therapy (frequency 8-17%),⁴³ advanced age,⁴⁴ and drugs that interfere with renal potassium excretion (especially when used concomitantly).⁴⁴ In all these conditions, more frequent serum potassium monitoring is advisable, patient education about potassium diet is important, and in case of recurrent episodes of hyperkalaemia the use of potassium binders may be useful in maintaining normokalaemia and optimal RAASi therapy. Monitoring of serum potassium is often overlooked, even in high-risk situations and after an episode of hyperkalaemia.¹⁵ International guidelines suggest that serum potassium should be tested within 120 days before initiation of a RAASi, during the early post-initiation period (Days 1 through 10), and during the extended post-

initiation period (Days 11 through 90), typically every 4 months.⁴⁵ This cycle of monitoring should be repeated at each new RAASi initiation, re-initiation or dose adjustment, and after any change in electrolyte status.

How to manage hyperkalaemia?

Dietary management

Restriction of dietary potassium to <2.4 g/day is recommended in patients with stage 3 (eGFR <60 mL/min/1.73 m²) or higher CKD.⁴⁶ However, specific guideline-directed advice is lacking on dietary potassium intake for other patient groups at risk of hyperkalaemia. In patients with HF in whom sodium restriction is frequently advised, the use of salt substitutes including potassium chloride may expose these patients to the risk of hyperkalaemia, especially if they have concomitant CKD or diabetes mellitus. Although patients are often educated to avoid commonly recognized high-potassium foods, many high-potassium foods may remain unrecognized by patients and healthcare providers. Patients at risk for hyperkalaemia should receive comprehensive dietary potassium education and if necessary consultation with a dietician.

Dose adjustments

The management of hyperkalaemia may require a short-term cessation of potassium-retaining agents and/or RAASi, but this should be minimized and RAASi should be carefully reintroduced as soon as possible while monitoring potassium levels.^{8,9}

One suggested approach is to reduce ACEi/ARB dose by 50% and recheck the serum potassium in 5-7 days until it has returned to baseline. If serum potassium does not return to baseline in ~2-4 weeks, discontinue the ACEi/ARB. If the patient is taking some combination of an ACEi or ARB and a MRA, discontinue one of the agents and recheck the potassium within 5-7 days.⁴⁴

In the case of the MRA spironolactone a detailed algorithm for dose adjustment is provided in *Table 1*.

Future developments and potential clinical applications

The FDA and EMA approved Patiromer and SZC for the treatment of chronic hyperkalaemia in patients receiving RAASi, with the ‘wide’ indication of ‘treatment of hyperkalaemia’.

Physicians will have to figure out how they may take advantage of these agents. Certainly, because of these are safer compared to the ‘older’ potassium binders, physicians may feel more comfortable using Patiromer or SZC to mitigate transient increases in serum potassium in patients receiving RAASi therapy. Further clinical experience will be required, and future studies may help understand when best to use these medications, i.e. at what level of serum potassium they should be initiated and whether or not they should be initiated before or after adjusting the doses of RAASi. The optimal duration of treatment in clinical practice will also need to be better described, i.e. should the K⁺ binder be discontinued after resolution of hyperkalaemia or continued with the aim of preventing a recurrence of hyperkalaemia in high-risk patients? Importantly, Patiromer and SZC are not interchangeable. They have important pharmacokinetic and safety profile differences. Ideally, a head-to-head comparison should help physicians determine the best conditions to use Patiromer and SZC, respectively. Cost and reimbursement are other factors which will influence the clinical use of Patiromer and SZC. Cost-effectiveness studies are required including a comparison to the currently used potassium binder resins (calcium polystyrene sulfonate and sodium polystyrene sulfonate).

The benefit of more aggressive use of RAASi with higher doses might also be investigated if facilitated with concomitant use of potassium binders.

A potential use of the new potassium-lowering agents would be to allow new indications, such as the use of MRAs in patients with an eGFR ≤ 30 mL/min/1.73 m² in whom they are currently contraindicated due to the fear of inducing hyperkalaemia and further renal dysfunction. Another potential use of the new potassium-lowering agents in association with a MRA could be in patients with ESRD on renal dialysis who are at a high risk for HF and sudden cardiac death.

The long-term risks and benefits of strategies using potassium-lowering agents should be thoroughly tested in adequately powered outcome trials.^{29,47}

Conflict of interest: F.Z. has received consultant/scientific advisory board fees from Actelion, Astrazeneca, Amgen, Bayer, Boehringer, Boston Scientific, CEVA, CVRx, EDDH, GE Healthcare, Janssen, Novartis, Pfizer, Quantum Genomics, Relypsa, Resmed, Stealth peptide, Takeda, and ZS Pharma. He is CardioRenal co-founder. J.P.F. has no conflicts of interest to declare. B.P. is a consultant to Bayer, Astra Zeneca, Sanofi, Stealth peptides, Relypsa/Vifor*, Sarfez*, KBP Biosciences*, Tricida*, and Sc Pharmaceuticals*. He has patent—site-specific delivery of eplerenone to the myocardium—US Patent # 9931412 (* denotes stock options).

References

- McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, Rouleau JL, Shi VC, Solomon SD, Swedberg K, Zile MR. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014;**371**:993-1004.
- Desai AS, Vardeny O, Claggett B, McMurray JJ, Packer M, Swedberg K, Rouleau JL, Zile MR, Lefkowitz M, Shi V, Solomon SD. Reduced risk of hyperkalemia during treatment of heart failure with mineralocorticoid receptor antagonists by use of sacubitril/valsartan compared with enalapril: a secondary analysis of the PARADIGM-HF trial. *JAMA Cardiol* 2017;**2**:79-85.
- Damman K, Gori M, Claggett B, Jhund PS, Senni M, Lefkowitz MP, Prescott MF, Shi VC, Rouleau JL, Swedberg K, Zile MR, Packer M, Desai AS, Solomon SD, McMurray JJV. Renal effects and associated outcomes during angiotensin-neprilysin inhibition in heart failure. *JACC Heart Fail* 2018;**6**:489-498.
- Ko DT, Juurlink DN, Mamdani MM, You JJ, Wang JT, Donovan LR, Tu JV. Appropriateness of spironolactone prescribing in heart failure patients: a population-based study. *J Card Fail* 2006;**12**:205-210.
- Juurlink DN, Mamdani MM, Lee DS, Kopp A, Austin PC, Laupacis A, Redelmeier DA. Rates of hyperkalemia after publication of the Randomized Aldactone Evaluation Study. *N Engl J Med* 2004;**351**:543-551.
- Bozkurt B, Agoston I, Knowlton AA. Complications of inappropriate use of spironolactone in heart failure: when an old medicine spirals out of new guidelines. *J Am Coll Cardiol* 2003;**41**:211-214.
- Ferreira JP, Rossignol P, Machu JL, Sharma A, Girerd N, Anker SD, Cleland JG, Dickstein K, Filippatos G, Hillege HL, Lang CC, Ter Maaten JM, Metra M, Ng L, Ponikowski P, Samani NJ, van Veldhuisen DJ, Zwiderman AH, Voors A, Zannad F. Mineralocorticoid receptor antagonist pattern of use in heart failure with reduced ejection fraction: findings from BIOSTAT-CHF. *Eur J Heart Fail* 2017;**19**:1284-1293.
- Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruschitzka F, Rutten FH, van der Meer P. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2016;**37**:2129-2200.
- Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Colvin MM, Drazner MH, Filippatos G, Fonarow GC, Givertz MM, Hollenberg SM, Lindenfeld J, Masoudi FA, McBride PE, Peterson PN, Stevenson LW, Westlake C. 2016 ACC/AHA/HFSA focused update on new pharmacological therapy for heart failure: an update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Am Coll Cardiol* 2016;**68**:1476-1488.
- Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, Palensky J, Wittes J. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med* 1999;**341**:709-717.
- Zannad F, McMurray JJ, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pocock SJ, Pitt B. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med* 2011;**364**:11-21.
- Zannad F, Gattis Stough W, Rossignol P, Bauersachs J, McMurray JJ, Swedberg K, Struthers AD, Voors AA, Ruilope LM, Bakris GL, O'Connor CM, Gheorghiadu M, Mentz RJ, Cohen-Solal A, Maggioni AP, Beygui F, Filippatos GS, Massy ZA, Pathak A, Pina IL, Sabbah HN, Sica DA, Tavazzi L, Pitt B. Mineralocorticoid receptor antagonists for heart failure with reduced ejection fraction: integrating evidence into clinical practice. *Eur Heart J* 2012;**33**:2782-2795.
- Williams EM, Katholi RE, Karambelas MR. Use and side-effect profile of spironolactone in a private cardiologist's practice. *Clin Cardiol* 2006;**29**:149-153.
- Shah KB, Rao K, Sawyer R, Gottlieb SS. The adequacy of laboratory monitoring in patients treated with spironolactone for congestive heart failure. *J Am Coll Cardiol* 2005;**46**:845-849.
- Cooper LB, Hammill BG, Peterson ED, Pitt B, Maciejewski ML, Curtis LH, Hernandez AF. Consistency of laboratory monitoring during initiation of mineralocorticoid receptor antagonist therapy in patients with heart failure. *JAMA* 2015;**314**:1973-1975.
- Pitt B, Williams G, Remme W, Martinez F, Lopez-Sendon J, Zannad F, Neaton J, Roniker B, Hurley S, Burns D, Bittman R, Kleiman J. The

- EPHESUS trial: eplerenone in patients with heart failure due to systolic dysfunction complicating acute myocardial infarction. Eplerenone Post-AMI Heart Failure Efficacy and Survival Study. *Cardiovasc Drugs Ther* 2001;15:79-87.
17. Montalescot G, Pitt B, Lopez de Sa E, Hamm CW, Flather M, Verheugt F, Shi H, Turgonyi E, Orri M, Vincent J, Zannad F. Early eplerenone treatment in patients with acute ST-elevation myocardial infarction without heart failure: the Randomized Double-Blind Reminder Study. *Eur Heart J* 2014;35:2295-2302.
 18. Beygui F, Cayla G, Roule V, Roubille F, Delarche N, Silvain J, Van Belle E, Belle L, Galinier M, Motreff P, Cornillet L, Collet JP, Furber A, Goldstein P, Ecollan P, Legallois D, Lebon A, Rousseau H, Machecourt J, Zannad F, Vicaut E, Montalescot G. Early aldosterone blockade in acute myocardial infarction: the ALBATROSS Randomized Clinical Trial. *J Am Coll Cardiol* 2016;67:1917-1927.
 19. Butler J, Anstrom KJ, Felker GM, Givertz MM, Kalogeropoulos AP, Konstam MA, Mann DL, Margulies KB, McNulty SE, Mentz RJ, Redfield MM, Tang WHW, Whellan DJ, Shah M, Desvigne-Nickens P, Hernandez AF, Braunwald E. Efficacy and safety of spironolactone in acute heart failure: the ATHENA-HF randomized clinical trial. *JAMA Cardiol* 2017;2:950.
 20. Rossignol P, Dobre D, McMurray JJ, Swedberg K, Krum H, van Veldhuisen DJ, Shi H, Messig M, Vincent J, Girerd N, Bakris G, Pitt B, Zannad F. Incidence, determinants, and prognostic significance of hyperkalemia and worsening renal function in patients with heart failure receiving the mineralocorticoid receptor antagonist eplerenone or placebo in addition to optimal medical therapy: results from the Eplerenone in Mild Patients Hospitalization and Survival Study in Heart Failure (EMPHASIS-HF). *Circ Heart Fail* 2014;7:51-58.
 21. Vardeny O, Claggett B, Anand I, Rossignol P, Desai AS, Zannad F, Pitt B, Solomon SD. Incidence, predictors, and outcomes related to hypo- and hyperkalemia in patients with severe heart failure treated with a mineralocorticoid receptor antagonist. *Circ Heart Fail* 2014;7:573-579.
 22. Anker SD, Kosiborod M, Zannad F, Pina IL, McCullough PA, Filippatos G, van der Meer P, Ponikowski P, Rasmussen HS, Lavin PT, Singh B, Yang A, Deedwania P. Maintenance of serum potassium with sodium zirconium cyclosilicate (ZS-9) in heart failure patients: results from a phase 3 randomized, double-blind, placebo-controlled trial. *Eur J Heart Fail* 2015;17:1050-1056.
 23. Pitt B, Bakris GL, Bushinsky DA, Garza D, Mayo MR, Stasiv Y, Christ-Schmidt H, Berman L, Weir MR. Effect of patiromer on reducing serum potassium and preventing recurrent hyperkalemia in patients with heart failure and chronic kidney disease on RAAS inhibitors. *Eur J Heart Fail* 2015;17:1057-1065.
 24. Aldahl M, Jensen A-SC, Davidsen L, Eriksen MA, Møller Hansen S, Nielsen BJ, Krogager ML, Køber L, Torp-Pedersen C, Søgaard P. Associations of serum potassium levels with mortality in chronic heart failure patients. *Eur Heart J* 2017;38:2890-2896.
 25. Nunez J, Bayes-Genis A, Zannad F, Rossignol P, Nunez E, Bodi V, Minana G, Santos E, Chorro FJ, Mollar A, Carratala A, Navarro J, Gorris JL, Lupon J, Husser O, Metra M, Sanchis J. Long-term potassium monitoring and dynamics in heart failure and risk of mortality. *Circulation* 2018;137:1320-1330.
 26. Collins AJ, Pitt B, Reaven N, Funk S, McGaughey K, Wilson D, Bushinsky DA. Association of serum potassium with all-cause mortality in patients with and without heart failure, chronic kidney disease, and/or diabetes. *Am J Nephrol* 2017;46:213-221.
 27. Krogager ML, Torp-Pedersen C, Mortensen RN, Køber L, Gislason G, Søgaard P, Aasbjerg K. Short-term mortality risk of serum potassium levels in hypertension: a retrospective analysis of nationwide registry data. *Eur Heart J* 2016;38:104-112.
 28. Goyal A, Spertus JA, Gosch K, Venkitchalam L, Jones PG, Van den Berghe G, Kosiborod M. Serum potassium levels and mortality in acute myocardial infarction. *JAMA* 2012;307:157-164.
 29. Pitt B, Rossignol P. Serum potassium in patients with chronic heart failure: once we make a U-turn where should we go? *Eur Heart J* 2017;38:2897-2899.
 30. Kovesdy CP, Matsushita K, Sang Y, Brunskill NJ, Carrero JJ, Chodick G, Hasegawa T, Heerspink HL, Hirayama A, Landman GWD, Levin A, Nitsch D, Wheeler DC, Coresh J, Hallan SI, Shalev V, Grams ME, Consortium CKDP. Serum potassium and adverse outcomes across the range of kidney function: a CKD Prognosis Consortium meta-analysis. *Eur Heart J* 2018;39:1535-1542.
 31. Konstam MA, Neaton JD, Dickstein K, Drexler H, Komajda M, Martinez FA, Riegger GA, Malbecq W, Smith RD, Guptha S, Poole-Wilson PA. Effects of high-dose versus low-dose losartan on clinical outcomes in patients with heart failure (HEAAL study): a randomized, double-blind trial. *Lancet* 2009;374:1840-1848.
 32. Packer M, Poole-Wilson PA, Armstrong PW, Cleland JG, Horowitz JD, Massie BM, Ryden L, Thygesen K, Uretsky BF. Comparative effects of low and high doses of the angiotensin-converting enzyme inhibitor, lisinopril, on morbidity and mortality in chronic heart failure. ATLAS Study Group. *Circulation* 1999;100:2312-2318.
 33. Kosiborod M, Rasmussen HS, Lavin P, Qunibi WY, Spinowitz B, Packham D, Roger SD, Yang A, Lerma E, Singh B. Effect of sodium zirconium cyclosilicate on potassium lowering for 28 days among outpatients with hyperkalemia: the HARMONIZE randomized clinical trial. *JAMA* 2014;312:2223-2233.
 34. Tamargo J, Caballero R, Delpon E. New drugs for the treatment of hyperkalemia in patients treated with renin-angiotensin-aldosterone system inhibitors - hype or hope? *Discov Med* 2014;18:249-254.
 35. Pitt B, Anker SD, Bushinsky DA, Kitzman DW, Zannad F, Huang IZ. Evaluation of the efficacy and safety of RLY5016, a polymeric potassium binder, in a double-blind, placebo-controlled study in patients with chronic heart failure (the PEARL-HF) trial. *Eur Heart J* 2011;32:820-828.
 36. Bakris GL, Pitt B, Weir MR, Freeman MW, Mayo MR, Garza D, Stasiv Y, Zawadzki R, Berman L, Bushinsky DA. Effect of patiromer on serum potassium level in patients with hyperkalemia and diabetic kidney disease: the AMETHYST-DN randomized clinical trial. *JAMA* 2015;314:151-161.
 37. Packham DK, Rasmussen HS, Lavin PT, El-Shahawy MA, Roger SD, Block G, Qunibi W, Pergola P, Singh B. Sodium zirconium cyclosilicate in hyperkalemia. *N Engl J Med* 2015;372:222-231.
 38. Konstam MA. Hyperkalemia in heart failure: binding the patient to improved treatment? *Eur J Heart Fail* 2015;17:997-999.
 39. Liu LC, Schutte E, Gansevoort RT, van der Meer P, Voors AA. Finerenone: third-generation mineralocorticoid receptor antagonist for the treatment of heart failure and diabetic kidney disease. *Expert Opin Investig Drugs* 2015;24:1123-1135.
 40. Pitt B, Køber L, Ponikowski P, Gheorghiadu M, Filippatos G, Krum H, Nowack C, Kolkhof P, Kim SY, Zannad F. Safety and tolerability of the novel non-steroidal mineralocorticoid receptor antagonist BAY 94-8862 in patients with chronic heart failure and mild or moderate chronic kidney disease: a randomized, double-blind trial. *Eur Heart J* 2013;34:2453-2463.
 41. Filippatos G, Anker SD, Böhm M, Gheorghiadu M, Køber L, Krum H, Maggioni AP, Ponikowski P, Voors AA, Zannad F, Kim SY, Nowack C, Palombo G, Kolkhof P, Kimmekamp-Kirschbaum N, Pieper A, Pitt B. A randomized controlled study of finerenone vs. eplerenone in patients with worsening chronic heart failure and diabetes mellitus and/or chronic kidney disease. *Eur Heart J* 2016;37:2105-2114.
 42. Uribarri J, Oh MS, Carroll HJ. Hyperkalemia in diabetes mellitus. *J Diabet Complications* 1990;4:3-7.
 43. Khosla N, Kalaitzidis R, Bakris GL. Predictors of hyperkalemia risk following hypertension control with aldosterone blockade. *Am J Nephrol* 2009;30:418-424.
 44. Palmer BF. Managing hyperkalemia caused by inhibitors of the renin-angiotensin-aldosterone system. *N Engl J Med* 2004;351:585-592.
 45. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE, McMurray JJ, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WH, Tsai EJ, Wilkoff BL. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2013;128:1810-1852.
 46. Kidney Disease Outcomes Quality Initiative (K/DOQI). K/DOQI clinical practice guidelines on hypertension and antihypertensive agents in chronic kidney disease. *Am J Kidney Dis* 2004;43:11-290.
 47. Zannad F, Rossignol P, Stough WG, Epstein M, Alonso Garcia M, D L A, Bakris GL, Butler J, Kosiborod M, Berman L, Mebazaa A, Rasmussen HS, Ruilope LM, Stockbridge N, Thompson A, Wittes J, Pitt B. New approaches to hyperkalemia in patients with indications for renin-angiotensin-aldosterone inhibitors: considerations for trial design and regulatory approval. *Int J Cardiol* 2016;216:46-51.