



HAL
open science

Beyond clinical examination and natriuretic peptides: comprehensive quantification of congestion with ultrasound in ambulatory heart failure patients

Nicolas Girerd, Elke Platz

► To cite this version:

Nicolas Girerd, Elke Platz. Beyond clinical examination and natriuretic peptides: comprehensive quantification of congestion with ultrasound in ambulatory heart failure patients. *European Journal of Heart Failure*, 2019, 21 (7), pp.917-920. 10.1002/ejhf.1422 . hal-03765907

HAL Id: hal-03765907

<https://hal.univ-lorraine.fr/hal-03765907v1>

Submitted on 31 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Published in final edited form as:

Eur J Heart Fail. 2019 July ; 21(7): 917–920. doi:10.1002/ejhf.1422.

Beyond clinical examination and natriuretic peptides: comprehensive quantification of congestion with ultrasound in ambulatory heart failure patients

Nicolas Girerd^{1,*}, Elke Platz²

¹Université de Lorraine, INSERM, Centre d'Investigations Cliniques 1433, CHRU de Nancy, Institut Lorrain du Cœur et des Vaisseaux, Nancy, France, and INI-CRCT (Cardiovascular and Renal Clinical Trialists) F-CRIN Network, Nancy, France;

²Department of Emergency Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

Clinical congestion plays a central role in the development of heart failure (HF)-related symptoms and is the cause of the majority of hospitalizations for HF (HHF).^{1,2} Relieving congestion is consequently an important goal of HF management.³ Physical examination in patients with HF has limited accuracy for the detection of congestion,⁴ which supports using additional tools in the evaluation of HF patients to improve risk stratification and possibly guide patient management. In the latest European Society of Cardiology HF guidelines, using ultrasound measurement of inferior vena cava (IVC) diameter for assessment of volume status has a IIbC recommendation, whereas the use of thoracic ultrasound is cited in the text but does not have a formal recommendation.³

The toolbox of sonographic methods that are relevant to congestion evaluation includes pulmonary congestion on lung ultrasound (LUS) (B-lines, pleural effusions), haemodynamic congestion on echocardiography (e.g. E/e' and systolic pulmonary artery pressure) and vascular ultrasound [IVC and jugular vein diameter (JVD) measurement]. LUS is a semi-quantitative, simple, and rapid method to assess pulmonary congestion⁵; this technique produces vertical lines defined as 'B-lines' by scanning along the intercostal spaces using either a phased array or curvilinear transducer. Pulmonary congestion identified by LUS has been shown to be a strong predictor of outcomes following a HHF^{6–8} and in ambulatory patients with HF.^{8,9} E/e' and elevated pulmonary artery pressures measured using echocardiography have been shown to be associated with increased risk for HHF in patients with HF.^{10,11} In addition, both IVC and JVD measurements have been employed in the assessment of haemodynamic congestion in ambulatory patients with HF with heterogeneous results with respect to their prognostic utility.^{12–15} Sonographic JVD evaluation can be either based on the evaluation of JVD at rest or the ratio of JVD during the

*Corresponding author. Centre d'Investigation Clinique Pierre Drouin -INSERM - CHRU de Nancy, Institut Lorrain du Cœur et des Vaisseaux Louis Mathieu, 4, rue du Morvan. 54500 Vandoeuvre-Les-Nancy, France. Tel: +33 3 83157496, Fax: +33 3 83157324, n.girerd@chru-nancy.fr.

The opinions expressed in this article are not necessarily those of the Editors of the *European Journal of Heart Failure* or of the European Society of Cardiology.

Valsalva manoeuvre and at rest. This sonographic measurement has been shown to be associated with adverse clinical outcomes by one research group so far.¹³ Whether each of these sonographic markers individually or combined are associated with important adverse events in ambulatory patients with HF had not been previously evaluated in a large cohort.

In this issue of the Journal, Pellicori and colleagues report the association of most available sonographic markers of congestion with the composite outcome of death or HHF in ambulatory patients with HF, as well as the prevalence of these measures in patients without congestion on physical examination.¹⁶ The key findings of this study are that: (i) nearly 50% of ambulatory patients with HF thought not to be congested based on the traditional evaluation of signs and symptoms actually had sonographic evidence of congestion (either based on B-lines quantification, JVD ratio or IVC diameter), (ii) these three sonographic features of congestion were associated with the aforementioned composite outcome in univariable analysis and improved risk prediction [based on net reclassification improvement (NRI) and integrated discrimination improvement (IDI)] beyond prognostically important clinical variables [namely age, sex, New York Heart Association (NYHA) class, creatinine, haemoglobin and left ventricular ejection fraction], and (iii) adding JVD ratio to similarly prognostically important clinical variables, including N-terminal pro-B-type natriuretic peptide (NT-proBNP), further improved re-classification (by IDI only).

The most striking clinical message of this study is that ultrasound of the lungs, jugular vein and IVC identified 'congestion' in almost half of those ambulatory HF patients who did not have any physical exam findings of congestion. This would suggest that congestion-focused ultrasound could detect and quantify subclinical congestion in patients with HF using simple techniques and widely available technology.

When comparing these results with previous studies, the most surprising finding of Pellicori's study is the weak association of LUS findings with long-term outcomes in multivariable models. This result stands in contrast to three prior observational studies in ambulatory patients with HF. Scali *et al.*¹⁷ demonstrated that > 8 B-lines on 28-zone LUS at rest is prognostically important for the composite of death or HHF at 12 month as demonstrated by an area under the curve of 0.98 [95% confidence interval (CI) 0.93–1.0]. Similarly, in a cohort of 195 NYHA class II–IV HF patients, three or more B-lines (using an 8-zone technique) was independently associated with a higher rate of 6-month death or HHF (adjusted hazard ratio 4.08, 95% CI 1.95–8.54; $P < 0.001$).⁹ Dwyer *et al.*¹⁸ reported similar results in low-risk patients with HF. The discrepancy between these former and the current study may be related to a higher degree of pulmonary congestion in the HF cohort in the current study, differences in LUS technique (e.g. number of zones examined and variable LUS clip duration), longer follow-up period, a lower event rate compared to previous reports (26% over a 6-month period in the report of Platz *et al.*⁹ vs. 18% during a median follow-up of 234 days in the current report of Pellicori *et al.*) or due to adjustment for NT-proBNP.

The work of Pellicori *et al.*¹⁶ should be applauded as it represents an important step forward in our understanding of how ultrasound can aid in the detection of congestion in ambulatory patients with HF. Important strengths of this study are its size (342 ambulatory patients with HF) and the comprehensive sonographic evaluation of congestion, including LUS,

echocardiography, IVC and JVD assessment. However, a few limitations should be considered. First, although an experienced reader analysed ultrasound videos off-line, it is not clear if this reader was blinded to clinical characteristics and subsequent outcomes. Second, the duration of the LUS recordings were based on heart beats (at least five); as shorter clip duration is associated with lower B-line counts, this recording strategy could have resulted in a lower B-line count in sicker patients with higher heart rates. Finally, treating physicians were not blinded to the echocardiogram and the ultrasound of the lungs, jugular vein and IVC. Physicians could have altered therapy based on these findings, which may have impacted the study findings. The reproducibility of these results using appropriate blinding of readers and treating clinicians should be assessed in future investigations.

This study greatly adds to our knowledge regarding the value of sonographic congestion measures in ambulatory patients with HF. It paves the way for a more comprehensive evaluation of congestion in patients with HF by including both sonographic pulmonary and haemodynamic congestion assessment. Congestion-focused ultrasound examinations could complement the clinical examination in ambulatory patients with chronic HF and allow for early detection of subclinical congestion. In Figure 1, we provide a brief overview of the current literature for the three steps of congestion assessment (i.e. clinical, laboratory, and imaging evaluation of congestion); we also detail current gaps in evidence and propose future research directions in the field.^{7,8,10–14,16–25}

Important gaps in knowledge remain. To our knowledge, a comprehensive sonographic congestion assessment including the aforementioned tools has not been evaluated in a randomized trial. In addition, there is a lack of evidence regarding how improved congestion assessment could better inform HF treatment (e.g. based on which sonographic cut-off values specific therapies should be altered). Ultrasound-based congestion parameters could potentially become important diagnostic risk stratification measures. Yet, the degree to which these sonographic congestion parameters can be modified by therapies and whether outcomes can be improved as a result needs to be examined in well-designed clinical trials.

Despite the established use of echocardiography in the assessment of patients with HF and availability of ultrasound equipment, comprehensive sonographic congestion assessment beyond echocardiography is not routinely performed. Demonstrating that acting on subclinical sonographic congestion in the setting of chronic HF does improve patient outcome could prompt its wide spread use and support its formal inclusion in HF guidelines.

Conflict of interest:

N.G. is supported by a public grant overseen by the French National Research Agency (ANR) as part of the second 'Investissements d'Avenir' program FIGHT-HF (reference: ANR-15-RHU-0004) and has received board and consulting fees for Novartis and consulting/lecture fees for Servier and Boehringer Ingelheim. E.P. is supported by a grant from the National Heart Lung and Blood Institute (NHLBI) (grant: K23 HL123533).

References

1. Maggioni AP, Dahlstrom U, Filippatos G, Chioncel O, Leiro MC, Drozdz J, Fruhwald F, Gullestad L, Logeart D, Metra M, Parissis J, Persson H, Ponikowski P, Rauchhaus M, Voors A, Nielsen OW, Zannad F, Tavazzi L, Heart Failure Association of ESC. EURObservational Research Programme:

- the Heart Failure Pilot Survey (ESC-HF Pilot). *Eur J Heart Fail* 2010;12:1076–1084. [PubMed: 20805094]
2. Girerd N, Seronde MF, Coiro S, Chouihed T, Bilbault P, Braun F, Kenizou D, Maillier B, Nazeyrollas P, Roul G, Fillieux L, Abraham WT, Januzzi J Jr, Sebbag L, Zannad F, Mebazaa A, Rossignol P, INI-CRCT Group. Integrative assessment of congestion in heart failure throughout the patient journey. *JACC Heart Fail* 2018;6:273–285. [PubMed: 29226815]
 3. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2016;18:891–975. [PubMed: 27207191]
 4. Wang CS, FitzGerald JM, Schulzer M, Mak E, Ayas NT. Does this dyspneic patient in the emergency department have congestive heart failure? *JAMA* 2005;294:1944–1956. [PubMed: 16234501]
 5. Volpicelli G, Elbarbary M, Blaivas M, Lichtenstein DA, Mathis G, Kirkpatrick AW, Melniker L, Gargani L, Noble VE, Via G, Dean A, Tsung JW, Soldati G, Copetti R, Bouhemad B, Reissig A, Agricola E, Rouby JJ, Arbelot C, Liteplo A, Sargsyan A, Silva F, Hoppmann R, Breitkreutz R, Seibel A, Neri L, Storti E, Petrovic T, International Liaison Committee on Lung Ultrasound for International Consensus Conference on Lung Ultrasound. International evidence-based recommendations for point-of-care lung ultrasound. *Intensive Care Med* 2012;38:577–591. [PubMed: 22392031]
 6. Coiro S, Porot G, Rossignol P, Ambrosio G, Carluccio E, Tritto I, Huttin O, Lemoine S, Sadoul N, Donal E, Zannad F, Girerd N. Prognostic value of pulmonary congestion assessed by lung ultrasound imaging during heart failure hospitalisation: a two-centre cohort study. *Sci Rep* 2016;6:39426. [PubMed: 27995971]
 7. Coiro S, Rossignol P, Ambrosio G, Carluccio E, Alunni G, Murrone A, Tritto I, Zannad F, Girerd N. Prognostic value of residual pulmonary congestion at discharge assessed by lung ultrasound imaging in heart failure. *Eur J Heart Fail* 2015;17:1172–1181. [PubMed: 26417699]
 8. Platz E, Merz AA, Jhund PS, Vazir A, Campbell R, McMurray JJ. Dynamic changes and prognostic value of pulmonary congestion by lung ultrasound in acute and chronic heart failure: a systematic review. *Eur J Heart Fail* 2017;19:1154–1163. [PubMed: 28557302]
 9. Platz E, Lewis EF, Uno H, Peck J, Pivetta E, Merz AA, Hempel D, Wilson C, Frasura SE, Jhund PS, Cheng S, Solomon SD. Detection and prognostic value of pulmonary congestion by lung ultrasound in ambulatory heart failure patients. *Eur Heart J* 2016;37:1244–1251. [PubMed: 26819225]
 10. Carluccio E, Dini FL, Biagioli P, Lauciello R, Simioniu A, Zuchi C, Alunni G, Reboldi G, Marzilli M, Ambrosio G. The ‘Echo Heart Failure Score’: an echocardiographic risk prediction score of mortality in systolic heart failure. *Eur J Heart Fail* 2013;15:868–876. [PubMed: 23512095]
 11. Thavendiranathan P, Yingchoncharoen T, Grant A, Seicean S, Landers SH, Gorodeski EZ, Marwick TH. Prediction of 30-day heart failure-specific readmission risk by echocardiographic parameters. *Am J Cardiol* 2014;113:335–341. [PubMed: 24268036]
 12. Gustafsson M, Alehagen U, Johansson P. Imaging congestion with a pocket ultrasound device: prognostic implications in patients with chronic heart failure. *J Card Fail* 2015;21:548–554. [PubMed: 25725475]
 13. Pellicori P, Kallvikbacka-Bennett A, Dierckx R, Zhang J, Putzu P, Cuthbert J, Boyalla V, Shoaib A, Clark AL, Cleland JG. Prognostic significance of ultrasound-assessed jugular vein distensibility in heart failure. *Heart* 2015;101:1149–1158. [PubMed: 26006717]
 14. Pellicori P, Carubelli V, Zhang J, Castiello T, Sherwi N, Clark AL, Cleland JG. IVC diameter in patients with chronic heart failure: relationships and prognostic significance. *JACC Cardiovasc Imaging* 2013;6:16–28. [PubMed: 23328557]
 15. Pellicori P, Clark AL, Kallvikbacka-Bennett A, Zhang J, Urbinati A, Monzo L, Dierckx R, Anker SD, Cleland JG. Non-invasive measurement of right atrial pressure by near-infrared spectroscopy:

- preliminary experience. A report from the SICA-HF study. *Eur J Heart Fail* 2017;19:883–892. [PubMed: 28387033]
16. Pellicori P, Shah P, Cuthbert J, Urbinati A, Zhang J, Kallvikbacka-Bennett A, Clark AL, Cleland JG. Prevalence, pattern and clinical relevance of ultrasound indices of congestion in outpatients with heart failure. *Eur J Heart Fail* 2019, in press 10.1002/ejhf.1383.
 17. Scali MC, Cortigiani L, Simionuc A, Gregori D, Marzilli M, Picano E. Exercise-induced B-lines identify worse functional and prognostic stage in heart failure patients with depressed left ventricular ejection fraction. *Eur J Heart Fail* 2017;19:1468–1478. [PubMed: 28198075]
 18. Dwyer KH, Merz AA, Lewis EF, Claggett BL, Crousillat DR, Lau ES, Silverman MB, Peck J, Rivero J, Cheng S, Platz E. Pulmonary congestion by lung ultrasound in ambulatory patients with heart failure with reduced or preserved ejection fraction and hypertension. *J Card Fail* 2018;24:219–226. [PubMed: 29499322]
 19. Ambrosy AP, Pang PS, Khan S, Konstam MA, Fonarow GC, Traver B, Maggioni AP, Cook T, Swedberg K, Burnett JC Jr, Grinfeld L, Udelson JE, Zannad F, Gheorghide M, EVEREST Trial Investigators. Clinical course and predictive value of congestion during hospitalization in patients admitted for worsening signs and symptoms of heart failure with reduced ejection fraction: findings from the EVEREST trial. *Eur Heart J* 2013;34:835–843. [PubMed: 23293303]
 20. Rohde LE, Beck-da-Silva L, Goldraich L, Grazziotin TC, Palombini DV, Polanczyk CA, Clausell N. Reliability and prognostic value of traditional signs and symptoms in outpatients with congestive heart failure. *Can J Cardiol* 2004;20:697–702. [PubMed: 15197422]
 21. Welsh P, Kou L, Yu C, Anand I, van Veldhuisen DJ, Maggioni AP, Desai AS, Solomon SD, Pfeffer MA, Cheng S, Gullestad L, Aukrust P, Ueland T, Swedberg K, Young JB, Kattan MW, Sattar N, McMurray JJ. Prognostic importance of emerging cardiac, inflammatory, and renal biomarkers in chronic heart failure patients with reduced ejection fraction and anaemia: RED-HF study. *Eur J Heart Fail* 2018;20:268–277. [PubMed: 28960777]
 22. Masson S, Latini R, Anand IS, Barlera S, Angelici L, Vago T, Tognoni G, Cohn JN, Val-He FTI. Prognostic value of changes in N-terminal pro-brain natriuretic peptide in Val-HeFT (Valsartan Heart Failure Trial). *J Am Coll Cardiol* 2008;52:997–1003. [PubMed: 18786480]
 23. Duarte K, Monnez JM, Albuissou E, Pitt B, Zannad F, Rossignol P. Prognostic value of estimated plasma volume in heart failure. *JACC Heart Fail* 2015;3: 886–893. [PubMed: 26541787]
 24. Brunner-La Rocca HP, Eurlings L, Richards AM, Januzzi JL, Pfisterer ME, Dahlstrom U, Pinto YM, Karlstrom P, Erntell H, Berger R, Persson H, O'Connor CM, Moertl D, Gaggin HK, Frampton CM, Nicholls MG, Troughton RW. Which heart failure patients profit from natriuretic peptide guided therapy? A meta-analysis from individual patient data of randomized trials. *Eur J Heart Fail* 2015;17:1252–1261. [PubMed: 26419999]
 25. Felker GM, Anstrom KJ, Adams KF, Ezekowitz JA, Fiuzat M, Houston-Miller N, Januzzi JL Jr, Mark DB, Pina IL, Passmore G, Whellan DJ, Yang H, Cooper LS, Leifer ES, Desvigne-Nickens P, O'Connor CM. Effect of natriuretic peptide-guided therapy on hospitalization or cardiovascular mortality in high-risk patients with heart failure and reduced ejection fraction: a randomized clinical trial. *JAMA* 2017;318:713–720. [PubMed: 28829876]




| | CONGESTION ASSESSMENT | | |
|--|---|--|--|
| | Clinical markers | Laboratory markers | Imaging markers |
| Parameters |  <p>Dyspnea / Orthopnea JVD Rales Peripheral edema</p> |  <p>Natriuretic peptides Estimated plasma volume (evaluated from hemoglobin and hematocrit) <i>Most conveniently used as point of care measurements</i></p> |  <p>Lung ultrasound JVD and IVC measurements Echocardiographic markers (E/e', pulmonary artery pressures...)</p> |
| Observational data regarding risk stratification | <p>A clinical congestion score to be performed at hospital discharge was validated in the EVEREST trial data: it was strongly associated with all-cause mortality and the risk for HF hospitalization¹⁹</p> <p>In the outpatient setting, the Rohde score, based on the quantification of orthopnea, JVD, rales and edema has been shown to be associated with mortality²⁰</p> | <p>Reported association with HF hospitalization and/or mortality: Natriuretic peptides^{21,22} Estimated plasma volume²³</p> <p>Significantly improved prediction: Natriuretic peptides²¹ Estimated plasma volume²³</p> | <p>Reported association with HF hospitalization and/or mortality: LUS^{8, 12, 17, 18} JVD measurements¹³ IVC measurements¹⁴ Echocardiographic markers^{10, 11}</p> <p>Improved risk prediction for HF hospitalization or mortality: LUS⁷ JVD measurements¹⁶ Echocardiographic markers¹¹</p> |
| Randomized trial-based evidence | <p>Trials unlikely to be performed as clinical markers are part of routine clinical care</p> | <p>A meta-analysis of individual patient data reported that natriuretic peptide-guided compared with symptom-guided therapy resulted in significantly lower mortality and fewer HF admissions in HFrEF patients²⁴.</p> <p>However, the most recent trial, GUIDE-IT²⁵, was neutral; in this trial, NT-ProBNP was mostly use to trigger changes in background treatment rather than treatment targeting congestion.</p> | <p>Trials to be performed</p> |

Figure 1.

The three steps of congestion assessment in ambulatory heart failure (HF) patients. HFrEF, heart failure with reduced ejection fraction; JVD, jugular vein diameter; IVC, inferior vena cava; LUS, lung ultrasound; NT-ProBNP, N-terminal pro-B-type natriuretic peptide.