



HAL
open science

Association of Empagliflozin Treatment With Albuminuria Levels in Patients With Heart Failure

João Pedro Ferreira, Faiez Zannad, Javed Butler, Gerasimos Filippatos, Stuart J Pocock, Martina Brueckmann, Dominik Steubl, Elke Schueler, Stefan D Anker, Milton Packer

► **To cite this version:**

João Pedro Ferreira, Faiez Zannad, Javed Butler, Gerasimos Filippatos, Stuart J Pocock, et al.. Association of Empagliflozin Treatment With Albuminuria Levels in Patients With Heart Failure. JAMA Cardiology, 2022, 7 (11), pp.1148-1159. 10.1001/jamacardio.2022.2924 . hal-03798543

HAL Id: hal-03798543

<https://hal.univ-lorraine.fr/hal-03798543>

Submitted on 5 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Association of Empagliflozin Treatment With Albuminuria Levels in Patients With Heart Failure

A Secondary Analysis of EMPEROR-Pooled

João Pedro Ferreira, MD, PhD; Faiez Zannad, MD, PhD; Javed Butler, MD, MPH; Gerasimos Filippatos, MD, PhD; Stuart J. Pocock, PhD; Martina Brueckmann, MD; Dominik Steubl, MD; Elke Schueller, MSc, Dipl-Math; Stefan D. Anker, MD, PhD; Milton Packer, MD

 Supplemental content

IMPORTANCE Albuminuria, routinely assessed as spot urine albumin-to-creatinine ratio (UACR), indicates structural damage of the glomerular filtration barrier and is associated with poor kidney and cardiovascular outcomes. Sodium-glucose cotransporter-2 (SGLT2) inhibitors have been found to reduce UACR in patients with type 2 diabetes, but its use in patients with heart failure (HF) is less well studied.

OBJECTIVE To analyze the association of empagliflozin with study outcomes across baseline levels of albuminuria and change in albuminuria in patients with HF across a wide range of ejection fraction levels.

DESIGN, SETTING, AND PARTICIPANTS This post hoc analysis included all patients with HF from the EMPEROR-Pooled analysis using combined individual patient data from the international multicenter randomized double-blind parallel-group, placebo-controlled EMPEROR-Reduced and EMPEROR-Preserved trials. Participants in the original trials were excluded from this analysis if they were missing baseline UACR data. EMPEROR-Preserved was conducted from March 27, 2017, to April 26, 2021, and EMPEROR-Reduced was conducted from April 6, 2017, to May 28, 2020. Data were analyzed from January to June 2022.

INTERVENTIONS Randomization to empagliflozin or placebo.

MAIN OUTCOMES AND MEASURES New-onset macroalbuminuria and regression to normoalbuminuria and microalbuminuria.

RESULTS A total of 9673 patients were included (mean [SD] age, 69.9 [10.4] years; 3551 [36.7%] female and 6122 [63.3%] male). Of these, 5552 patients had normoalbuminuria (UACR <30 mg/g) and 1025 had macroalbuminuria (UACR >300 mg/g). Compared with normoalbuminuria, macroalbuminuria was associated with younger age, races other than White, obesity, male sex, site region other than Europe, higher levels of N-terminal pro-hormone brain natriuretic peptide and high-sensitivity troponin T, higher blood pressure, higher New York Heart Association class, greater HF duration, more frequent previous HF hospitalizations, diabetes, hypertension, lower eGFR, and less frequent use of angiotensin-converting enzyme inhibitors or angiotensin receptor blockers and mineralocorticoid receptor antagonists. An increase in events was observed in individuals with higher UACR levels. The association of empagliflozin with cardiovascular mortality or HF hospitalization was consistent across UACR categories (hazard ratio [HR], 0.80; 95% CI, 0.69-0.92 for normoalbuminuria; HR, 0.74; 95% CI, 0.63-0.86 for microalbuminuria; HR, 0.78; 95% CI, 0.63-0.98 for macroalbuminuria; interaction P trend = .71). Treatment with empagliflozin was associated with lower incidence of new macroalbuminuria (HR, 0.81; 95% CI, 0.70-0.94; P = .005) and an increase in rate of remission to sustained normoalbuminuria or microalbuminuria (HR, 1.31; 95% CI, 1.07-1.59; P = .009) but not with a reduction in UACR in the overall population; however, UACR was reduced in patients with diabetes, who had higher UACR levels than patients without diabetes (geometric mean for diabetes at baseline, 0.91; 95% CI, 0.85-0.98 and for no diabetes at baseline, 1.08; 95% CI, 1.01-1.16; interaction P = .008).

CONCLUSIONS AND RELEVANCE In this post hoc analysis of a randomized clinical trial, compared with placebo, empagliflozin was associated with reduced HF hospitalizations or cardiovascular death irrespective of albuminuria levels at baseline, reduced progression to macroalbuminuria, and reversion of macroalbuminuria.

TRIAL REGISTRATION ClinicalTrials.gov Identifiers: [NCT03057977](https://clinicaltrials.gov/ct2/show/study/NCT03057977) and [NCT03057951](https://clinicaltrials.gov/ct2/show/study/NCT03057951)

JAMA Cardiol. doi:10.1001/jamacardio.2022.2924
Published online September 21, 2022.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: João Pedro Ferreira, MD, PhD, Centre d'Investigation Clinique 1433 module Plurithématique, CHRU Nancy-Hopitaux de Brabois, Institut Lorrain du Coeur et des Vaisseaux Louis Mathieu, 4 rue du Morvan, 54500 Vandoeuvre les Nancy, France (j.ferreira@chru-nancy.fr).

Albuminuria, determined as spot urine albumin-to-creatinine ratio (UACR), indicates structural damage of the glomerular filtration barrier and, along with reduced estimated glomerular filtration rate (eGFR), is a key variable in defining chronic kidney disease (CKD).^{1,2} Albuminuria is common in patients with CKD related to diabetes, hypertension, and other cardiovascular diseases that can lead to endothelial dysfunction or increase intraglomerular capillary pressure, and is associated with cardiovascular events, including heart failure (HF) and cardiovascular mortality as well as CKD progression.³⁻⁸

Angiotensin-converting enzyme inhibitors (ACEi) and angiotensin receptor blockers (ARBs) have been shown to reduce albuminuria and the progression to macroalbuminuria.⁹⁻¹¹ In addition, mineralocorticoid receptor antagonists (MRAs) have been found to reduce albuminuria in patients with diabetic kidney disease and HF.¹²⁻¹⁶ More recently, sodium glucose cotransporter-2 inhibitors were found to reduce the progression to macroalbuminuria in patients with diabetes and CKD with and without diabetes.¹⁷⁻²⁰

In patients with HF who participated in the Candesartan in Heart Failure: Assessment of Reduction in Mortality and Morbidity (CHARM) study,²¹ microalbuminuria and macroalbuminuria were associated with an increased risk of HF hospitalization and mortality irrespective of ejection fraction, but treatment with candesartan did not reduce excessive albuminuria. In the Aldosterone Antagonist Therapy for Adults With Heart Failure and Preserved Systolic Function (TOPCAT) trial,¹³ microalbuminuria and macroalbuminuria were associated with an increased risk of HF hospitalizations and mortality and spironolactone with a reduction in albuminuria. In the Study to Evaluate the Efficacy and Safety of LCZ696 Compared to Enalapril on Morbidity and Mortality of Patients With Chronic Heart Failure (PARADIGM-HF)²² and the Study of Renal Effects of the Angiotensin Receptor Neprilysin Inhibitor LCZ696 in Patients With Heart Failure and Preserved Ejection Fraction (PARAMOUNT),²³ sacubitril-valsartan was associated with an increase in albuminuria throughout follow-up compared with enalapril or valsartan. The effect of sodium glucose cotransporter-2 inhibitors in reducing HF hospitalizations or cardiovascular mortality across baseline albuminuria levels and their impact on the progression of albuminuria in HF patients is yet to be reported.

In this post hoc analysis, we studied the association of empagliflozin with the study outcomes across baseline levels of albuminuria and change in albuminuria in patients with HF across a wide range of left ventricular ejection fraction (LVEF) levels using data from the EMPEROR-Pooled analysis²⁴ (ie, Empagliflozin Outcome Trial in Patients With Chronic Heart Failure With Reduced Ejection Fraction [EMPEROR-Reduced]²⁵ and Empagliflozin in Heart Failure with a Preserved Ejection Fraction [EMPEROR-Preserved] trials combined).

Methods

Study Design and Patient Population

The design and primary results of the EMPEROR-Pooled analysis have been previously published.^{24,26} In brief, EMPEROR-

Key Points

Question Is empagliflozin associated with a reduction in albuminuria in patients with heart failure?

Findings In this secondary analysis from EMPEROR-Pooled, treatment with empagliflozin was associated with a reduction in incidence of new macroalbuminuria and an increase in rate of remission to sustained normoalbuminuria or microalbuminuria among patients with macroalbuminuria at baseline. The association of empagliflozin with cardiovascular and kidney outcomes was consistent across albumin-to-creatinine ratio categories.

Meaning Compared with placebo, empagliflozin was associated with a reduction in progression to macroalbuminuria and a reversion from macroalbuminuria as well as a decrease in heart failure hospitalizations or cardiovascular death, irrespective of baseline albuminuria.

Pooled combined individual patient data from EMPEROR-Reduced and EMPEROR-Preserved, 2 phase 3 international multicenter randomized double-blind parallel-group, placebo-controlled trials that enrolled adult patients with chronic HF with New York Heart Association class II to IV symptoms for at least 3 months and elevated natriuretic peptide levels across a wide range of LVEF ($\leq 40\%$ in EMPEROR-Reduced²⁵ and $>40\%$ with no prior measurement $\leq 40\%$ in EMPEROR-Preserved).²⁷

The protocol of each trial complied with the Declaration of Helsinki and was approved by the ethical committees of the participating sites. All patients gave written informed consent to participate in the study.

Randomization, Study Visits, and Event Definition

Patients were randomized in a double-blind manner to receive placebo or empagliflozin, 10 mg daily (1:1 ratio), in addition to their usual therapy. Following entry into the trial, treatments for HF or other medical conditions could be managed at the clinical discretion of the investigator.

Albuminuria was assessed using UACR from a morning void spot urine sample and collected at randomization and each subsequent study visit (weeks 4, 12, 32, and 52 and every 24 weeks thereafter) and analyzed by the central laboratory. Normoalbuminuria was defined as UACR less than 30 mg/g, microalbuminuria as UACR ranging from 30 to 300 mg/g, and macroalbuminuria as UACR greater than 300 mg/g.²⁸

End Points

To analyze the use of empagliflozin in patients with HF across albuminuria categories, the prespecified primary and some secondary end points were studied. The primary outcome in the EMPEROR trials was a composite of cardiovascular death or HF hospitalization. Additionally, the association of empagliflozin with first and recurrent HF hospitalizations, cardiovascular and all-cause death, 2 composite kidney end points (consisting of sustained decline in eGFR $\geq 40\%$ or $\geq 50\%$ from baseline and sustained eGFR <15 or <10 ml/min/1.73 m² for patients with baseline eGFR \geq or <30 ml/min/1.73 m², respectively, long-term dialysis, or kidney transplant), and changes

in annualized eGFR slope (chronic slope) were studied across baseline UACR levels.

In an additional post hoc analysis, the association of empagliflozin with change in albuminuria was studied using progression to macroalbuminuria in patients without macroalbuminuria at baseline and sustained remission to normoalbuminuria or microalbuminuria in patients with macroalbuminuria at baseline. Relative changes in UACR over time were also evaluated in the overall cohort and by UACR and diabetes subgroups at baseline. Lastly, we evaluated treatment safety by baseline UACR categories.

Statistical Analysis

Baseline characteristics were compared across categories of baseline UACR (normoalbuminuria, microalbuminuria, and macroalbuminuria) using ordinal regression likelihood ratio test. Associations between baseline UACR categories and subsequent outcomes were studied by comparing the placebo events rates across categories. The association of treatment (empagliflozin vs placebo) with the study outcomes was assessed using a Cox proportional hazards model including the prespecified baseline covariates of age, sex, geographical region, diabetes, study (EMPEROR-Reduced or EMPEROR-Preserved), LVEF, eGFR, UACR category, and a treatment-by-UACR category interaction term according to the intention-to-treat principle. Race data were reported in accordance with the requirements of the US Food and Drug Administration (FDA) and self-reported according to multiple-choice categories as per FDA guidance, with multiple answers possible. Total number of hospitalizations (first and recurrent) was analyzed using a joint frailty model that accounted for informative censoring because of cardiovascular death. Progression to and remission from macroalbuminuria to microalbuminuria or normoalbuminuria were studied with a Cox proportional hazards model including the prespecified baseline covariates described above (except UACR and the interaction term). The consistency of association of empagliflozin with macroalbuminuria was assessed across a range of clinically relevant participant characteristics, including age, eGFR, LVEF, body mass index, previous HF hospitalization, and diabetes, along with the respective interaction tests. The association of empagliflozin with UACR changes over time was studied using a linear mixed model for repeated measurements with adjustment for the covariates referenced above and treatment-by-visit interaction. *P* values and 95% CIs presented in this report have not been adjusted for multiplicity. All analyses were performed using SAS version 9.4 (SAS Institute). All tests were 2-sided, and *P* < .05 was considered statistically significant.

Results

Patient Characteristics by UACR Categories

A total of 9673 patients were included (mean [SD] age, 69.9 [10.4] years; 3551 [36.7%] female and 6122 [63.3%] male; 1496 [15.5%] Asian, 514 [5.3%] Black, 7130 [73.7%] White, and 476 [4.9%] of another race [including American Indian or Alaska Native, Native Hawaiian or Other Pacific Islander, and multiple races, consolidated owing to low numbers]), 3710 from

EMPEROR-Reduced and 5963 from EMPEROR-Preserved; 45 (0.5%) patients from the total population were excluded due to missing baseline UACR. In the pooled population, 5552 patients had normoalbuminuria (UACR <30 mg/g), and 1025 patients had macroalbuminuria (UACR >300 mg/g). Compared with normoalbuminuria, macroalbuminuria was associated with younger age, races other than White, obesity, male sex, site region other than Europe, higher levels of N-terminal pro-hormone brain natriuretic peptide and high-sensitivity troponin T, higher blood pressure, higher New York Heart Association class, greater HF duration; more frequent previous HF hospitalizations, diabetes (including insulin use), hypertension, lower eGFR, and less frequent use of ACEi or ARBs and MRAs. LVEF was similar across UACR categories (Table).

Risks and Associations of Empagliflozin Across UACR Categories

An increase in events was observed for higher UACR categories. For example, patients receiving placebo with UACR greater than 300 mg/g had a 2.7-fold higher rate of primary outcome events (22.2 vs 8.2 events per 100 person-years) and 2.3-fold higher rate of cardiovascular death events (8.2 vs 3.6 events per 100 person-years) than patients with UACR less than 30 mg/g.

The association of empagliflozin with all analyzed trial outcomes was consistent across UACR categories. For example, the primary outcome was reduced by 20% in patients with UACR less than 30 mg/g (hazard ratio [HR] in patients with UACR <30 mg/g, 0.80; 95% CI, 0.69-0.92; HR in patients with UACR 30-300 mg/g, 0.74; 95% CI, 0.63-0.86; HR in patients with UACR >300 mg/g, 0.78; 95% CI, 0.63-0.98; interaction *P* trend = .71) (Figure 1).

Empagliflozin was associated with a slower decline in annualized eGFR slope. This association was also consistent across UACR categories: UACR <30 mg/g-placebo slope, -2.4 (95% CI, -2.6 to -2.1) mL/min/1.73 m²/y; empagliflozin slope, -0.8 (95% CI, -1.0 to -0.5) mL/min/1.73 m²/y; empagliflozin vs placebo slope difference 1.6 (95% CI, 1.2 to 1.9) mL/min/1.73 m²/y; UACR 30-300 mg/g-placebo slope, -2.5 (95% CI, -2.9 to -2.2) mL/min/1.73 m²/y; empagliflozin slope, -1.4 (95% CI, -1.7 to -1.0) mL/min/1.73 m²/y empagliflozin vs placebo slope difference, 1.2 (95% CI, 0.7 to 1.6) mL/min/1.73 m²/y; UACR >300 mg/g-placebo slope, -4.0 (95% CI, -4.6 to -3.3) mL/min/1.73 m²/y; empagliflozin slope, -2.3 (95% CI, -2.9 to -1.6); empagliflozin vs placebo slope difference, 1.7 (95% CI, 0.8 to 2.6) mL/min/1.73 m²/y (interaction *P* trend = .57).

Regarding safety, we detected higher frequencies of adverse events, adverse events leading to discontinuation, serious adverse events, and acute kidney failure events for patients in higher UACR categories. However, no relevant differences were detected between patients in the placebo vs empagliflozin groups (eTable 1 in the Supplement).

Association of Empagliflozin With Onset of Macroalbuminuria and Remission to Normoalbuminuria or Microalbuminuria

Among the 8648 patients without macroalbuminuria at baseline (4312 empagliflozin and 4336 placebo), treatment with empagliflozin was associated with a reduction in incidence of new

Table. Characteristics of the EMPEROR-Pooled Population by Categories of Urinary Albumin-to-Creatinine Ratio (UACR) at Baseline

Characteristic	UACR categories, No. (%)			P value ^a
	<30 mg/g	30-300 mg/g	>300 mg/g	
No. of patients (total = 9673)	5552	3096	1025	NA
Empagliflozin rand	2765 (49.8)	1547 (50.0)	525 (51.2)	.54
Demographic characteristics				
Age, mean (SD), y	70 (10)	71 (10)	67 (11)	.003
Female	2151 (38.7)	1099 (35.5)	301 (29.4)	<.001
Male	3401 (61.3)	1997 (64.5)	724 (70.6)	
BMI, mean (SD) ^b	29 (6)	29 (6)	30 (6)	.001
BMI categories				
<30	3440 (62.0)	1836 (59.3)	554 (54.0)	<.001
≥30	2112 (38.0)	1260 (40.7)	471 (46.0)	
Race ^c				
Asian	762 (13.7)	535 (17.3)	199 (19.4)	
Black	263 (4.7)	174 (5.6)	77 (7.5)	
White	4240 (76.4)	2214 (71.5)	676 (66.0)	<.001
Other ^d	256 (4.6)	153 (4.9)	67 (6.5)	
Site region				
Asia	586 (10.6)	443 (14.3)	150 (14.6)	
Europe	2607 (47.0)	1120 (36.2)	280 (27.3)	
Latin America	1488 (26.8)	929 (30.0)	381 (37.2)	
North America	550 (9.9)	444 (14.3)	144 (14.0)	
Other ^e	321 (5.8)	160 (5.2)	70 (6.8)	<.001
Laboratory values				
Troponin T, median (IQR), ng/mL ^f	17 (11-26)	22 (14-33)	28 (18-44)	<.001
NT-proBNP, median (IQR), pg/mL ^g	1085 (577-1916)	1568 (824-2890)	1740 (897-3739)	<.001
UACR, median (IQR), mg/g	9 (5-16)	68 (43-125)	793 (450-1637)	NA
eGFR, mean (SD), mL/min/1.73 m ²	63 (20)	60 (21)	54 (22)	<.001
eGFR categories				
≥60	3091 (55.7)	143 (46.3)	389 (38.0)	
45 to <60	1375 (24.8)	832 (26.9)	244 (23.8)	<.001
30 to <45	899 (16.2)	663 (21.4)	235 (22.9)	
<30	187 (3.4)	167 (5.4)	157 (15.3)	
Potassium, mean (SD), mmol/L	4.6 (0.5)	4.6 (0.5)	4.6 (0.6)	.74
Albumin, mean (SD), g/dL	4.4 (0.3)	4.4 (0.3)	4.2 (0.4)	<.001
Hemoglobin, mean (SD), g/dL	13.5 (1.5)	13.4 (1.7)	13.3 (1.9)	<.001
Vital signs, mean (SD)				
Heart rate, bpm	70 (11)	72 (12)	72 (12)	<.001
Blood pressure, mm Hg				
Systolic	126 (16)	128 (17)	137 (17)	<.001
Diastolic	74 (10)	75 (11)	78 (11)	<.001
HF characteristics				
NYHA class III/IV	1012 (18.2)	745 (24.1)	257 (25.1)	<.001
HF diagnosis, mean (SD), y	4.9 (5.6)	5.3 (5.7)	5.2 (5.8)	.02
HHF<12 mo	1360 (24.5)	842 (27.2)	308 (30.0)	<.001
Ischemic HF	2281 (41.1)	1296 (41.9)	443 (43.2)	.21
AFib/flutter	2552 (46.0)	1595 (51.5)	410 (40.0)	.27
LVEF, mean (SD), %	44 (15)	44 (15)	45 (16)	.81
LVEF categories				
≤40% ^f	2078 (37.4)	1236 (39.9)	396 (38.6)	
>40% ^f	3474 (62.6)	1860 (60.1)	629 (61.4)	.053
Comorbidities				
Hypertension	4508 (81.2)	2659 (85.9)	920 (89.8)	<.001
Diabetes	2306 (41.5)	1714 (55.4)	750 (73.2)	<.001

(continued)

Table. Characteristics of the EMPEROR-Pooled Population by Categories of Urinary Albumin-to-Creatinine Ratio (UACR) at Baseline (continued)

Characteristic	UACR categories, No. (%)			P value ^a
	<30 mg/g	30-300 mg/g	>300 mg/g	
Comedication				
Insulin use	465 (8.4)	508 (16.4)	352 (34.3)	<.001
ACEi/ARB	4255 (76.6)	2278 (73.6)	744 (72.6)	<.001
ARNI	457 (8.2)	309 (10.0)	90 (8.8)	.04
β-Blockers	4954 (89.2)	2790 (90.1)	920 (89.8)	.25
Thiazides	912 (16.4)	454 (14.7)	152 (14.8)	.02
Loop diuretics	4010 (72.2)	2349 (75.9)	813 (79.3)	<.001
MRA	2962 (53.4)	1486 (48.0)	435 (42.4)	<.001
CCB	1007 (18.1)	724 (23.4)	369 (36.0)	<.001
Assist devices				
ICD	631 (11.4)	356 (11.5)	80 (7.8)	.052
CRT (CRT-D or -P)	264 (4.8)	161 (5.2)	37 (3.6)	.71

Abbreviations: ACEi, angiotensin-converting enzyme inhibitor; AFib, atrial fibrillation; ARB, angiotensin receptor blocker; ARNI, angiotensin receptor neprilysin inhibitor; BMI, body mass index; CCB, calcium channel blocker; CRT, cardiac resynchronization therapy with or without a defibrillator; eGFR, estimated glomerular filtration rate; FDA, US Food and Drug Administration; HF, heart failure; HHF, hospitalization for heart failure; ICD, implantable cardiac defibrillator with or without cardiac resynchronization therapy; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; NA, not applicable; NT-proBNP, N-terminal pro-hormone brain natriuretic peptide; NYHA, New York Heart Association.

^a P values from ordinal regression likelihood ratio test based on log-transformed data.

^b Calculated as weight in kilograms divided by height in meters squared.

^c Race was recorded for this study in accordance with the requirements of the FDA. Race was self-reported according to multiple-choice categories as per FDA guidance, with multiple answers possible.

^d Included American Indian or Alaska Native, Native Hawaiian or Other Pacific Islander, and multiple races, consolidated owing to small numbers.

^e Included Australia, India, and South Africa, consolidated owing to small numbers.

^f To convert to μg/L, multiply by 1.

^g EMPEROR-Reduced.²⁵

macroalbuminuria (5.7 events per 100 person-years with empagliflozin vs 7.1 events per 100 person-years with placebo, corresponding to a hazard ratio [HR] of 0.81; 95% CI, 0.70-0.94; $P = .005$) (Figure 2). The association was consistent across subgroups (Figure 3).

Among the 1025 patients (525 empagliflozin and 500 placebo) with macroalbuminuria at baseline, treatment with empagliflozin was associated with an increase in the rate of remission to sustained normoalbuminuria or microalbuminuria (HR, 1.31; 95% CI, 1.07-1.59; $P = .009$) (Figure 2). The association was generally consistent across subgroups, except across age subgroups where a trend toward a larger association of empagliflozin with reversion to macroalbuminuria to sustained normoalbuminuria or microalbuminuria was observed for older patients (HR for patients aged <65 years, 1.04; 95% CI, 0.75-1.44; HR for patients aged 56-75 years, 1.30; 95% CI, 0.91-1.86; HR for patients aged >75 years, 1.77; 95% CI, 1.23-2.56; interaction P trend = .03) (Figure 3).

Association of Empagliflozin With Albuminuria Over Time

Empagliflozin was not significantly associated with relative changes in UACR over time in the overall population (baseline to week 52 geometric mean relative change vs placebo, 0.99; 95% CI, 0.95-1.05) (eTable 2 in the Supplement; Figure 4A). However, albuminuria reduction was more pronounced in higher UACR categories (baseline to week 52 geometric mean relative change in UACR with empagliflozin vs placebo in baseline UACR <30 mg/g, 1.04 [95% CI, 0.97-1.11]; in UACR 30-300 mg/g, 0.95 [95% CI, 0.87-1.04]; and in UACR >300 mg/g, 0.88 [95% CI, 0.75-1.03]; interaction P trend, .04) (eTable 2 in the Supplement; Figure 4B-D). Regarding diabe-


tes status at baseline, the geometric mean relative change of UACR from baseline to week 52 was significant in patients with diabetes (0.91; 95% CI, 0.85-0.98) but not in patients without diabetes (1.08; 95% CI, 1.01-1.16); interaction $P = .008$ (eTable 2 in the Supplement).

Discussion

This study confirms that albuminuria above the normal range was frequent (microalbuminuria 32% and macroalbuminuria 11%) and associated with poor prognosis in patients with HF, and showed that empagliflozin was associated with a reduction in new-onset macroalbuminuria and an increase in sustained remission from macroalbuminuria to normoalbuminuria or microalbuminuria in patients with HF, regardless of ejection fraction or diabetes status. Furthermore, empagliflozin was associated with improved cardiovascular and kidney outcomes irrespective of baseline UACR.

The association between macroalbuminuria and age, race, obesity, diabetes, higher N-terminal pro-hormone brain natriuretic peptide and troponin T levels, and poorer kidney function suggests that albuminuria could serve as a marker of disease severity in younger patients with obesity, diabetes, and HF. In our study, patients with macroalbuminuria had a 2- to 3-fold higher risk of HF hospitalizations and cardiovascular mortality compared with patients with normoalbuminuria. The rate of eGFR decline throughout follow-up was also greater in patients with macroalbuminuria. These findings support albuminuria measurement as a cardiovascular risk predictor, associated with HF hospitalizations, mortality from cardiovas-

Figure 1. Association of Empagliflozin With Baseline Urinary Albumin-to-Creatinine Ratio (UACR) Categories



Cox proportional hazard model adjusted for age (continuous), baseline estimated glomerular filtration rate (eGFR; continuous), baseline left ventricular ejection fraction (continuous), study, region, baseline diabetes status, sex, UACR category, treatment, and treatment-by-UACR category. Composite kidney end point defined by sustained decline in eGFR $\geq 40\%$ from baseline and sustained eGFR <15 or <10 mL/min/1.73 m² for patients with baseline eGFR \geq or <30 mL/min/1.73 m², respectively; long-term dialysis; or kidney transplant. Alternative kidney end point defined by sustained decline in eGFR $\geq 50\%$ from baseline and sustained eGFR <15 or <10 mL/min/1.73 m² for patients with baseline eGFR \geq or <30 mL/min/1.73 m², respectively; long-term dialysis;


or kidney transplant. Total hospitalizations for heart failure were analyzed using a joint frailty model accounting for cardiovascular death and adjusting for the same covariates as the Cox model. Hazard ratios (HRs) for the composite kidney end point should be interpreted with caution due to significant heterogeneity across the 2 trials. The subgroup analysis by trial estimated an HR of 0.51 (95% CI, 0.33 to 0.79) in the EMPEROR-Reduced trial²⁵ and 0.95 (95% CI, 0.73 to 1.24) in the EMPEROR-Preserved trial;²⁷ P value for interaction between trials = .02. NA indicates not applicable; NNH, number needed to harm; NNT, number needed to treat; py, person-year.

cular causes, and kidney function decline.^{29,30} Still, treatment with empagliflozin was associated with reduced rate of HF hospitalizations or cardiovascular death irrespective of albuminuria at baseline; the event reduction ranged from 20% to 26% relative reduction, so that patients with macroalbuminuria treated with empagliflozin had event rates similar to patients

with microalbuminuria treated with placebo. The rate of eGFR slope decline was also consistently slower in association with empagliflozin irrespective of baseline albuminuria.

Empagliflozin was associated with reduced incidence of new macroalbuminuria in patients with normoalbuminuria or microalbuminuria at baseline, an association that was consis-

Figure 2. Cumulative Incidence Curves Considering All-Cause Mortality as Competing Risk for Time to Incidence of Macroalbuminuria




tent across all studied subgroups, including patients with and without diabetes and with LVEF above or below 40%. Furthermore, empagliflozin was associated with reversion to normal albuminuria or microalbuminuria among more patients who had macroalbuminuria at baseline.

In some cases, the association of HF therapies with albuminuria has been discordant to the effect of the same drug class reported in patients with diabetes or hypertension and not necessarily in agreement with the association of these therapies with HF hospitalizations and cardiovascular mortality. For

example, in the CHARM study,²¹ the proportion of patients with microalbuminuria and macroalbuminuria was similar to that reported in the present study, and the risk of events increased with excessive albuminuria but candesartan was not associated with a reduction in development or excessive excretion of albumin in urine. However, the CHARM analysis was limited by a high proportion of missing values throughout follow-up. The association of candesartan with albuminuria levels in patients with diabetes was small,³¹ and some studies have suggested that the doses of candesartan required to impact al-


Figure 3. Association of Empagliflozin With Incidence of Macroalbuminuria in Subgroups of Interest



A, Association of empagliflozin with macroalbuminuria in patients with normoalbuminuria or microalbuminuria (urinary albumin-to-creatinine ratio [UACR] ≤300 mg/dl) at baseline. B, Remission to microalbuminuria or normoalbuminuria in patients with macroalbuminuria (UACR >300 mg/dL) at

baseline. Cox proportional hazard model adjusted for age (continuous), baseline estimated glomerular filtration rate (eGFR; continuous), baseline left ventricular ejection fraction (LVEF; continuous), study, region, baseline diabetes status, sex, subgroup of interest, treatment, and treatment-by-subgroup category.

Figure 4. Association of Empagliflozin With Urinary Albumin-to-Creatinine Ratio (UACR) Over Time



A, Overall population. B, Patients with normoalbuminuria (UACR <30 mg/g) at baseline. C, Patients with microalbuminuria (UACR 30-300 mg/g) at baseline. D, Patients with macroalbuminuria (UACR >300 mg/g) at baseline. Models analyzed using a mixed model for repeated measures, including age, baseline estimated glomerular filtration rate, and baseline left ventricular ejection

fraction as linear covariates and study, region, baseline diabetes status, sex, log (baseline UACR) by visit, visit by treatment, and week reachable as fixed effects (overall population). For the subgroup analysis, visit by treatment-by-UACR subgroup was used. Gmean indicates geometric mean.

buminuria are much higher than the doses recommended for clinical use.³² A substudy from the Studies of Left Ventricular Dysfunction (SOLVD)³³ showed an antiproteinuric association with enalapril in patients with HF with reduced ejection fraction and diabetes, but not in those without diabetes.³³ These findings from CHARM and SOLVD suggest that the association of ACEi and ARBs with the proteinuria of patients with HF may be limited and only prevalent in specific subpopulations. Notwithstanding, ACEi and ARBs were associated with a reduction in HF hospitalizations and mortality in patients with HF with reduced ejection fraction and modestly associated with HF hospitalizations in patients with HF with preserved ejection fraction.³⁴⁻³⁶ In patients with HF treated with ACEi or ARBs, MRAs were associated with reduced excessive albuminuria regardless of diabetes status or ejection fraction.^{13,16} MRAs were also associated with improved outcomes in patients with HF with reduced ejection fraction^{37,38} and many patients with HF with preserved ejection fraction.³⁹ Sacubitril-valsartan was associated with an increase in albuminuria in patients with HF with reduced ejection fraction and HF with preserved ejection frac-

tion, but its association with a reduction in HF hospitalizations or cardiovascular death and a slower decline in eGFR were not influenced by albuminuria at baseline or changes during follow-up.^{22,23} It is possible that an increase in natriuretic peptides and other vasoactive substances as result of neprilysin inhibition by sacubitril leads to an increase in glomerular endothelial permeability.^{40,41}

Empagliflozin showed was associated with a reduction in macroalbuminuria, HF hospitalizations or cardiovascular death, and eGFR decline over time in patients with HF. If a reduction in intraglomerular pressure is key for albuminuria control in patients with diabetes,⁴² albuminuria may be influenced by many other factors, such as aldosterone activation, renal venous congestion, low-grade systemic inflammation, and microvascular and endothelial dysfunction, in patients with HF (with and without diabetes), all of which are processes that play a role in the progression of HF and can be mitigated by sodium glucose cotransporter-2 inhibitors.⁴³⁻⁴⁹

In our study, empagliflozin was not associated with a reduction in UACR in the overall population; however,

UACR was reduced in patients with diabetes who had higher UACR levels than patients without diabetes. In addition, a trend toward UACR reduction was observed in patients with higher UACR levels at baseline. These findings support the need for an elevated albuminuria level for a reduction in albuminuria to be observed with empagliflozin and are aligned with the existing data from patients with diabetes and CKD.^{19,20,50,51} In the EMPEROR trials, most patients had a UACR less than 30 mg/g, thus helping to explain why no association was seen in the overall population in this post hoc analysis.

Limitations

This study has limitations. Management of albuminuria was left to the discretion of the treating physician, but between-group variation in treatment approaches is expected to be small due to the randomized and double-blind nature of the study. We could not determine the exact mechanisms by which empagliflozin was associated with a reduction in

macroalbuminuria and dedicated studies should address this question.

Conclusions

In this post hoc analysis of EMPEROR-Pooled, empagliflozin was more frequently associated with a reduction in HF hospitalizations and cardiovascular death irrespective of albuminuria levels at baseline, a reduction in progression to macroalbuminuria, and reversion of macroalbuminuria compared with placebo. Empagliflozin was not associated with a reduction in UACR in the overall population; however, UACR was reduced in patients with diabetes, who had higher UACR levels than patients without diabetes. In addition, a trend toward UACR reduction was observed in patients with higher UACR levels at baseline. These findings support the potential need for an elevated albuminuria level for reduction of albuminuria to be observed with empagliflozin.

ARTICLE INFORMATION

Accepted for Publication: July 19, 2022.

Published Online: September 21, 2022.
doi:10.1001/jamacardio.2022.2924

Open Access: This is an open access article distributed under the terms of the [CC-BY-NC-ND License](https://creativecommons.org/licenses/by-nc-nd/4.0/). © 2022 Ferreira JP et al. *JAMA Cardiology*.

Author Affiliations: Université de Lorraine, Inserm, Centre d'Investigations Cliniques Plurithématique 1433, and Inserm U1116, CHRU, F-CRIN INI-CRCT (Cardiovascular and Renal Clinical Trialists), Nancy, France (Ferreira, Zannad); Department of Surgery and Physiology, Cardiovascular Research and Development Center, University of Porto, Porto, Portugal (Ferreira); Baylor Scott and White Research Institute, Dallas, Texas (Butler, Packer); University of Mississippi Medical Center, Jackson (Butler); National and Kapodistrian University of Athens School of Medicine, Athens, Greece (Filippatos); London School of Hygiene and Tropical Medicine, London, United Kingdom (Pocock); Boehringer Ingelheim International, Ingelheim, Germany (Brueckmann, Steubl); First Department of Medicine, Faculty of Medicine Mannheim, University of Heidelberg, Mannheim, Germany (Brueckmann); Department of Nephrology, Hospital rechts der Isar, Technical University Munich, Munich, Germany (Steubl); mainanalytics, Sulzbach, Germany (Schueler); Department of Cardiology Berlin Institute of Health Center for Regenerative Therapies German Centre for Cardiovascular Research partner site Berlin, Charité Universitätsmedizin Berlin, Berlin, Germany (Anker); Institute of Heart Diseases, Wrocław Medical University, Wrocław, Poland (Anker); Imperial College, London, United Kingdom (Packer).

Author Contributions: Dipl Math Schueler had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Ferreira, Zannad, Butler, Filippatos, Brueckmann, Anker, Packer.

Acquisition, analysis, or interpretation of data: Ferreira, Zannad, Butler, Filippatos, Pocock, Brueckmann, Steubl, Schueler.

Drafting of the manuscript: Ferreira, Butler, Steubl.
Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Ferreira, Pocock, Steubl, Schueler, Packer.

Obtained funding: Brueckmann, Packer.

Administrative, technical, or material support: Steubl, Packer.

Supervision: Ferreira, Zannad, Butler, Brueckmann, Packer.

Conflict of Interest Disclosures: Dr Ferreira reports personal fees from Boehringer Ingelheim during the conduct of the study and personal fees from Boehringer Ingelheim outside the submitted work. Dr Zannad reports personal fees from Acceleron Boehringer Ingelheim during the conduct of the study and personal fees from Janssen, Novartis, Boston Scientific, Amgen, CVRx, AstraZeneca, Vifor Fresenius, Cardior, Cereno Pharmaceutical, Applied Therapeutics, Merck, Bayer, and Cellprothera and other from Cardiovascular Clinical Trialists, Cardiorenal, Novo Nordisk, Servier and G3 Pharmaceuticals outside the submitted work. Dr Butler reports personal fees from Boehringer Ingelheim during the conduct of the study and personal fees from Adrenomed, Amgen, Array, AstraZeneca, Bayer, Bristol Myers Squibb, Boehringer Ingelheim, Cardior, CVRx, Foundry, G3 Pharmaceutical, Imbria, Impulse Dynamics, Innolife, Janssen, LivaNova, Luitpold, Medtronic, Merck, Novartis, Novo Nordisk, Relypsa, Roche, Sanofi, Sequana Medical, V-Wave, and Vifor outside the submitted work. Dr Filippatos reports personal fees from Boehringer Ingelheim during the conduct of the study and personal fees from Medtronic, Vifor, Servier, Novartis, Bayer, Amgen, Windtree, and Boehringer Ingelheim outside the submitted work. Dr Pocock reported personal fees from Boehringer Ingelheim Consultancy outside the submitted work. Dr Brueckmann reports personal fees from Boehringer Ingelheim International and is an employee of Boehringer Ingelheim. Dr Steubl is an employee of Boehringer Ingelheim. Dipl Math Schueler is an employee of mainanalytics, contracted by Boehringer Ingelheim. Dr Pocock reports personal fees from Boehringer Ingelheim during the conduct of the study and personal fees

from Boehringer Ingelheim outside the submitted work. Dr Anker reports personal fees from Boehringer Ingelheim during the conduct of the study and grants and personal fees from Abbott Vascular and Vifor and personal fees from Bayer, Boehringer Ingelheim, Brahms, Cardiac Dimensions, Cordio, Novartis, Occlutech, Servier, and V-Wave outside the submitted work. Dr Packer reports personal fees from Boehringer Ingelheim during the conduct of the study and personal fees from AbbVie, Actavis, Altimune, Amgen, Amarin, AstraZeneca, Boehringer Ingelheim, Bristol Myers Squibb, Caladrius, Casana, CSL Behring, Cytokinetics, Johnson & Johnson, Imara, Eli Lilly & Company, Moderna, Novartis, ParatusRx, Pfizer, Reata, Relypsa, Salamandra, Synthetic Biologics, Theravance, and Casana outside the submitted work. No other disclosures were reported.

Funding/Support: The EMPEROR-Reduced and Preserved trials were funded by Boehringer Ingelheim and Eli Lilly.

Role of the Funder/Sponsor: The sponsors were Boehringer Ingelheim and Eli Lilly. The executive committee, which included representatives of Boehringer Ingelheim, developed the protocol and statistical analysis plan, oversaw the recruitment of patients, and supervised the data analysis.

Additional Information: To ensure independent interpretation of clinical study results and enable authors to fulfill their role and obligations under the International Committee of Medical Journal Editors criteria, Boehringer Ingelheim grants all external authors access to clinical study data pertinent to the development of the publication. In adherence with the Boehringer Ingelheim Policy on Transparency and Publication of Clinical Study Data, scientific and medical researchers can request access to clinical study data when it becomes available on Vivli Center for Global Clinical Research Data, and earliest after publication of the primary manuscript in a peer-reviewed journal, regulatory activities are complete, and other criteria are met. Please visit Medical & Clinical Trials, Clinical Research, MyStudyWindow for further information.

Additional Contributions: Graphical assistance was provided by 7.4 Limited and supported

financially by Boehringer Ingelheim. Editorial assistance was provided by Elevate Scientific Solutions and supported financially by Boehringer Ingelheim.

REFERENCES

- Satchell S. The role of the glomerular endothelium in albumin handling. *Nat Rev Nephrol*. 2013;9(12):717-725. doi:10.1038/nrneph.2013.197
- Korakas E, Ikonomidis I, Markakis K, Raptis A, Dimitriadis G, Lambadiari V. The endothelial glycocalyx as a key mediator of albumin handling and the development of diabetic nephropathy. *Curr Vasc Pharmacol*. 2020;18(6):619-631. doi:10.2174/157016118666191224120242
- Gerstein HC, Mann JF, Yi Q, et al. Albuminuria and risk of cardiovascular events, death, and heart failure in diabetic and nondiabetic individuals. *JAMA*. 2001;286(4):421-426. doi:10.1001/jama.286.4.421
- Blecker S, Matsushita K, Köttgen A, et al. High-normal albuminuria and risk of heart failure in the community. *Am J Kidney Dis*. 2011;58(1):47-55. doi:10.1053/j.ajkd.2011.02.391
- Mogensen CE. Microalbuminuria predicts clinical proteinuria and early mortality in maturity-onset diabetes. *N Engl J Med*. 1984;310(6):356-360. doi:10.1056/NEJM198402093100605
- Ibsen H, Wachtell K, Olsen MH, et al. Albuminuria and cardiovascular risk in hypertensive patients with left ventricular hypertrophy: the LIFE Study. *Kidney Int Suppl*. 2004;(92):S56-S58. doi:10.1111/j.1523-1755.2004.09214.x
- Solomon SD, Lin J, Solomon CG, et al; Prevention of Events With ACE Inhibition (PEACE) Investigators. Influence of albuminuria on cardiovascular risk in patients with stable coronary artery disease. *Circulation*. 2007;116(23):2687-2693. doi:10.1161/CIRCULATIONAHA.107.723270
- Persson F, Bain SC, Mosenson O, et al; LEADER Trial Investigators. Changes in albuminuria predict cardiovascular and renal outcomes in type 2 diabetes: a post hoc analysis of the LEADER trial. *Diabetes Care*. 2021;44(4):1020-1026. doi:10.2337/dc20-1622
- Heeg JE, de Jong PE, van der Hem GK, de Zeeuw D. Reduction of proteinuria by angiotensin converting enzyme inhibition. *Kidney Int*. 1987;32(1):78-83. doi:10.1038/ki.1987174
- Viberti G, Mogensen CE, Groop LC, Pauls JF; European Microalbuminuria Captopril Study Group. Effect of captopril on progression to clinical proteinuria in patients with insulin-dependent diabetes mellitus and microalbuminuria. *JAMA*. 1994;271(4):275-279. doi:10.1001/jama.1994.03510280037029
- Parving HH, Lehnert H, Bröchner-Mortensen J, Gomis R, Andersen S, Arner P; Irbesartan in Patients with Type 2 Diabetes and Microalbuminuria Study Group. The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med*. 2001;345(12):870-878. doi:10.1056/NEJMoa011489
- Davidson MB, Wong A, Hamrahian AH, Stevens M, Siraj ES. Effect of spironolactone therapy on albuminuria in patients with type 2 diabetes treated with angiotensin-converting enzyme inhibitors. *Endocr Pract*. 2008;14(8):985-992. doi:10.4158/EP.14.8.985
- Selvaraj S, Claggett B, Shah SJ, et al. Prognostic value of albuminuria and influence of spironolactone in heart failure with preserved ejection fraction. *Circ Heart Fail*. 2018;11(11):e005288. doi:10.1161/CIRCHEARTFAILURE.118.005288
- Epstein M, Williams GH, Weinberger M, et al. Selective aldosterone blockade with eplerenone reduces albuminuria in patients with type 2 diabetes. *Clin J Am Soc Nephrol*. 2006;1(5):940-951. doi:10.2215/CJN.00240106
- Bakris GL, Agarwal R, Chan JC, et al; Mineralocorticoid Receptor Antagonist Tolerability Study-Diabetic Nephropathy (ARTS-DN) Study Group. Effect of finerenone on albuminuria in patients with diabetic nephropathy: a randomized clinical trial. *JAMA*. 2015;314(9):884-894. doi:10.1001/jama.2015.10081
- Pitt B, Kober L, Ponikowski P, et al. Safety and tolerability of the novel non-steroidal mineralocorticoid receptor antagonist BAY 94-8862 in patients with chronic heart failure and mild or moderate chronic kidney disease: a randomized, double-blind trial. *Eur Heart J*. 2013;34(31):2453-2463. doi:10.1093/eurheartj/ehf187
- Bae JH, Park EG, Kim S, Kim SG, Hahn S, Kim NH. Effects of sodium-glucose cotransporter 2 inhibitors on renal outcomes in patients with type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *Sci Rep*. 2019;9(1):13009. doi:10.1038/s41598-019-49525-y
- Heerspink HJ, Johnsson E, Gause-Nilsson I, Cain VA, Sjöström CD. Dapagliflozin reduces albuminuria in patients with diabetes and hypertension receiving renin-angiotensin blockers. *Diabetes Obes Metab*. 2016;18(6):590-597. doi:10.1111/dom.12654
- Jongs N, Greene T, Chertow GM, et al; DAPA-CKD Trial Committees and Investigators. Effect of dapagliflozin on urinary albumin excretion in patients with chronic kidney disease with and without type 2 diabetes: a prespecified analysis from the DAPA-CKD trial. *Lancet Diabetes Endocrinol*. 2021;9(11):755-766. doi:10.1016/S2213-8587(21)00243-6
- Cherney DZI, Zinman B, Inzucchi SE, et al. Effects of empagliflozin on the urinary albumin-to-creatinine ratio in patients with type 2 diabetes and established cardiovascular disease: an exploratory analysis from the EMPA-REG OUTCOME randomised, placebo-controlled trial. *Lancet Diabetes Endocrinol*. 2017;5(8):610-621. doi:10.1016/S2213-8587(17)30182-1
- Jackson CE, Solomon SD, Gerstein HC, et al. Albuminuria in chronic heart failure: prevalence and prognostic importance. *Lancet*. 2009;374(9689):543-550. doi:10.1016/S0140-6736(09)61378-7
- Damman K, Gori M, Claggett B, et al. Renal effects and associated outcomes during angiotensin-neprilysin inhibition in heart failure. *JACC Heart Fail*. 2018;6(6):489-498. doi:10.1016/j.jchf.2018.02.004
- Voors AA, Gori M, Liu LC, et al; PARAMOUNT Investigators. Renal effects of the angiotensin receptor neprilysin inhibitor LCZ696 in patients with heart failure and preserved ejection fraction. *Eur J Heart Fail*. 2015;17(5):510-517. doi:10.1002/ejhf.232
- Packer M, Butler J, Filippatos G, et al; EMPEROR Trial Committees and Investigators. Design of a prospective patient-level pooled analysis of two parallel trials of empagliflozin in patients with established heart failure. *Eur J Heart Fail*. 2020;22(12):2393-2398. doi:10.1002/ejhf.2065
- Packer M, Anker SD, Butler J, et al; EMPEROR-Reduced Trial Investigators. Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med*. 2020;383(15):1413-1424. doi:10.1056/NEJMoa2022190
- Packer M, Butler J, Zannad F, et al; EMPEROR Study Group. Empagliflozin and major renal outcomes in heart failure. *N Engl J Med*. 2021;385(16):1531-1533. doi:10.1056/NEJMoa2112411
- Anker SD, Butler J, Filippatos G, et al; EMPEROR-Preserved Trial Investigators. Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med*. 2021;385(16):1451-1461. doi:10.1056/NEJMoa2107038
- National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis*. 2002;39(2)(suppl 1):S1-S266.
- Bakris GL, Mollitch M. Microalbuminuria as a risk predictor in diabetes: the continuing saga. *Diabetes Care*. 2014;37(3):867-875. doi:10.2337/dc13-1870
- Laffin LJ, Bakris GL. Intersection between chronic kidney disease and cardiovascular disease. *Curr Cardiol Rep*. 2021;23(9):117. doi:10.1007/s11886-021-01546-8
- Bilous R, Chaturvedi N, Sjølie AK, et al. Effect of candesartan on microalbuminuria and albumin excretion rate in diabetes: three randomized trials. *Ann Intern Med*. 2009;151(1):11-20, W3-4. doi:10.7326/0003-4819-151-1-200907070-00120
- Burgess E, Muirhead N, Rene de Cotret P, Chiu A, Pichette V, Tobe S; SMART (Supra Maximal Atacand Renal Trial) Investigators. Supramaximal dose of candesartan in proteinuric renal disease. *J Am Soc Nephrol*. 2009;20(4):893-900. doi:10.1681/ASN.2008040416
- Capes SE, Gerstein HC, Negassa A, Yusuf S. Enalapril prevents clinical proteinuria in diabetic patients with low ejection fraction. *Diabetes Care*. 2000;23(3):377-380. doi:10.2337/diacare.23.3.377
- McMurray JJ, Ostergren J, Swedberg K, et al; CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin-converting-enzyme inhibitors: the CHARM-Added trial. *Lancet*. 2003;362(9386):767-771. doi:10.1016/S0140-6736(03)14283-3
- Yusuf S, Pitt B, Davis CE, Hood WB, Cohn JN; SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med*. 1991;325(5):293-302. doi:10.1056/NEJM199108013250501
- Yusuf S, Pfeffer MA, Swedberg K, et al; CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved trial. *Lancet*. 2003;362(9386):777-781. doi:10.1016/S0140-6736(03)14283-7
- Pitt B, Zannad F, Remme WJ, et al; Randomized Aldactone Evaluation Study Investigators. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med*. 1999;341(10):709-717. doi:10.1056/NEJM199909023411001

38. Zannad F, McMurray JJ, Krum H, et al; EMPHASIS-HF Study Group. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med*. 2011;364(1):11-21. doi:10.1056/NEJMoa1009492
39. Pfeffer MA, Claggett B, Assmann SF, et al. Regional variation in patients and outcomes in the Treatment of Preserved Cardiac Function Heart Failure With an Aldosterone Antagonist (TOPCAT) trial. *Circulation*. 2015;131(1):34-42. doi:10.1161/CIRCULATIONAHA.114.013255
40. McMurray J, Seidelin PH, Howey JE, Balfour DJ, Struthers AD. The effect of atrial natriuretic factor on urinary albumin and beta 2-microglobulin excretion in man. *J Hypertens*. 1988;6(10):783-786. doi:10.1097/00004872-198810000-00003
41. Lofton CE, Newman WH, Currie MG. Atrial natriuretic peptide regulation of endothelial permeability is mediated by cGMP. *Biochem Biophys Res Commun*. 1990;172(2):793-799. doi:10.1016/0006-291X(90)90744-8
42. Imanishi M, Yoshioka K, Okumura M, et al. Mechanism of decreased albuminuria caused by angiotensin converting enzyme inhibitor in early diabetic nephropathy. *Kidney Int Suppl*. 1997;63: S198-S200.
43. Akiyama E, Sugiyama S, Matsuzawa Y, et al. Incremental prognostic significance of peripheral endothelial dysfunction in patients with heart failure with normal left ventricular ejection fraction. *J Am Coll Cardiol*. 2012;60(18):1778-1786. doi:10.1016/j.jacc.2012.07.036
44. Torre-Amione G, Kapadia S, Lee J, et al. Tumor necrosis factor-alpha and tumor necrosis factor receptors in the failing human heart. *Circulation*. 1996;93(4):704-711. doi:10.1161/01.CIR.93.4.704
45. Blake WD, Wegria R, Keating RP, Ward HP. Effect of increased renal venous pressure on renal function. *Am J Physiol*. 1949;157(1):1-13. doi:10.1152/ajplegacy.1949.157.1.1
46. Butler MJ, Ramnath R, Kadoya H, et al. Aldosterone induces albuminuria via matrix metalloproteinase-dependent damage of the endothelial glycocalyx. *Kidney Int*. 2019;95(1):94-107. doi:10.1016/j.kint.2018.08.024
47. Kuriyama S. A potential mechanism of cardio-renal protection with sodium-glucose cotransporter 2 inhibitors: amelioration of renal congestion. *Kidney Blood Press Res*. 2019;44(4): 449-456. doi:10.1159/000501081
48. Locatelli M, Zoja C, Conti S, et al. Empagliflozin protects glomerular endothelial cell architecture in experimental diabetes through the VEGF-A/caveolin-1/PV-1 signaling pathway. *J Pathol*. 2022;256(4):468-479. doi:10.1002/path.5862
49. Abdollahi E, Keyhanfar F, Delbandi AA, Falak R, Hajimiresmaiel SJ, Shafiei M. Dapagliflozin exerts anti-inflammatory effects via inhibition of LPS-induced TLR-4 overexpression and NF-κB activation in human endothelial cells and differentiated macrophages. *Eur J Pharmacol*. 2022;918:174715. doi:10.1016/j.ejphar.2021.174715
50. Perkovic V, Jardine MJ, Neal B, et al; CREDENCE Trial Investigators. Canagliflozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med*. 2019;380(24): 2295-2306. doi:10.1056/NEJMoa1811744
51. Neuen BL, Ohkuma T, Neal B, et al. Cardiovascular and renal outcomes with canagliflozin according to baseline kidney function. *Circulation*. 2018;138(15):1537-1550. doi:10.1161/CIRCULATIONAHA.118.035901