



**HAL**  
open science

## Assessment of bioavailability and bioaccessibility of persistent organic pollutants in the chain “soil-animal-food”

Farida Amutova, Matthieu Delannoy, G Konuspayeva, Stefan Jurjanz

► **To cite this version:**

Farida Amutova, Matthieu Delannoy, G Konuspayeva, Stefan Jurjanz. Assessment of bioavailability and bioaccessibility of persistent organic pollutants in the chain “soil-animal-food”. Annual seminar of Doctoral School SIRENA, Feb 2020, Vandoeuvre-lès-Nancy, France. hal-03953966

**HAL Id: hal-03953966**

<https://hal.univ-lorraine.fr/hal-03953966v1>

Submitted on 24 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Assessment of bioavailability and bioaccessibility of persistent organic pollutants in the chain «soil-animal-food»

<sup>1,2,3</sup>F. Amutova, <sup>1</sup>M. Delannoy, <sup>2,3</sup>G. Konuspayeva, <sup>1</sup>S. Jurjanz

<sup>1</sup>URAFPA, University of Lorraine-INRA, Vandoeuvre, France

<sup>2</sup>Faculty of Geography and Environmental Sciences, Al-Farabi Kazakh National University, Almaty, Kazakhstan

<sup>3</sup>Antigen LLP, Scientific and Production Enterprise, Almaty region, Kazakhstan

e-mail: amutovafb@gmail.com

## CONTEXT

- Environmental contaminants (metals, POPs) can be stored during centuries in soil, therefore soil is a major exposure matrix
- All free ranged (food producing) animals ingest soil<sup>1;2</sup> at different levels
- Soil is one of the main vector for contaminants in animals and then in food

## OBJECTIVE

Reducing the bioavailability of POPs in the food chain "soil – farm animals – food products of animal origin" using different sequestration materials.

### AIM № 1

Evaluation of POP transfer to food of animal origin

#### Task 1

Literature review (meta-analysis)

### AIM № 2

Characterization of POP mobility in soil

#### Task 2

*In vitro* assessment of POPs availability in soils

### AIM № 3

Reducing the POP transfer from soil to animals by application of sequestration materials

#### Task 3


*In vitro* assessment of the efficiency of POP sequestration

## Preliminary results.



Mean and SDs of carry-over rates (COR) of PCDD/Fs in milk and eggs derived from the literature

Cl <sub>n</sub>	Compound	Transfer to milk		Transfer to eggs	
		COR, %	Transfer level	COR, %	Transfer level
4	2,3,7,8-TCDD	34.5 ± 14.3	high	40.5 ± 0.7	high
5	1,2,3,7,8-PeCDD	36.0 ± 12.4	high	44.5 ± 4.9	high
6	1,2,3,4,7,8-HxCDD	26.8 ± 8.9	high	43.0 ± 1.4	high
6	1,2,3,6,7,8-HxCDD	28.1 ± 13.7	high	44.0 ± 0	high
6	1,2,3,7,8,9-HxCDD	17.8 ± 6.5	medium	33.0 ± 1.4	high
7	1,2,3,4,6,7,8-HpCDD	5.06 ± 3.3	low	21.5 ± 0.7	medium
8	OCDD	0.72 ± 0.6	low	10.5 ± 3.5	low
4	2,3,7,8-TCDF	2.41 ± 2.70	low	40.0 ± 0	high
5	1,2,3,7,8-PeCDF	4.45 ± 4.00	low	43.5 ± 6.4	high
5	2,3,4,7,8-PeCDF	40.6 ± 18.2	high	41.0 ± 4.2	high
6	1,2,3,4,7,8-HxCDF	26.2 ± 11.3	high	43.5 ± 0.7	high
6	1,2,3,6,7,8-HxCDF	25.6 ± 11.6	high	41.0 ± 1.4	high
6	1,2,3,7,8,9-HxCDF	14.3 ± 6.10	medium	33.0 ± 2.8	high
6	2,3,4,6,7,8-HxCDF	19.0 ± 16.9	medium	40.0 ± 1.4	high
7	1,2,3,4,6,7,8-HpCDF	4.63 ± 3.20	low	18.0 ± 0	medium
7	1,2,3,4,7,8,9-HpCDF	5.87 ± 4.10	low	20.5 ± 0.7	medium
8	OCDF	0.54 ± 0.60	low	6.00 ± 1.4	low

Factorial plan (F1, F2) of PCA applied on mean CORs of PCBs for milk (A) and eggs (B), chlorine substitution at none, mono and di-ortho position, number of chlorines, molecular weight (M.W.), n-Octanol/Water Partition Coefficient (K<sub>ow</sub>) and transfer level.



Indication: Numbers correspond to the PCB isomers. Framed congeners are dioxin-like PCBs. Circle with PCBs are the high excreted congeners with the range 37.5-83.5% and 40.5-80.0% for milk and eggs respectively.





## SCIENTIFIC PRODUCTIONS (2018-2021)

N°	Topics of scientific publication	Type of publication	Status	Year	Journal
1	Transfer of persistent organic pollutants in camel milk	Communication/poster	published	Aug., 2019	70th Annual Meeting of the European Federation of Animal Science, Ghent (Belgium), 26 - 30 Aug 2019
2	Evaluation of two contrasted activated carbon based sequestration strategies to reduce soil bound chlordecone bioavailability in piglets	Article	published	Dec., 2019	Environmental Science and Pollution Research (IF = 2.8), In press, doi: 10.1007/s11356-019-06494-z
3	Transfer of persistent organic pollutants in food producing animals (meta-analysis)	Article	In process	2020	Chemosphere (IF=4.4)
4	Assessment of PCDD/Fs and PCBs transfer to milk and eggs	Communication/oral report	In process	2020	DIOXIN 2020, the 40th International Symposium on Halogenated Persistent Organic Pollutants (POPs) from 30 August to 4 September 2020.
Additional articles that were recently published not in the topic of PhD:					
5	Volatile organic compounds profiles in milk fermented by lactic bacteria.	Article	published	Jan., 2018	International Journal of Biology and Chemistry 11, № 2, 57, doi.org/10.26577/ijbch-2018-2-345
6	Comparative study of fatty acid and sterol profiles for the investigation of potential milk fat adulteration.	Article	published	Sept., 2019	Journal of Dairy Science. Volume 102, Issue 9, p. 7723-7733, doi.org/10.3168/jds.2018-15620. (IF = 3.08)

### Task 4

*In vivo* validation of Sequestration strategies



1. Jurjanz, S., Germain, K., Juin, H., Jondreville, C. Plant and soil intake by organic broilers reared in tree- or grass-covered plots as determined by means of n-alkanes and of acid-insoluble ash. 2015, ANIMAL, 9 (5), pp. 888-898.  
2. Jurjanz, S., Collas, C., Lasteil, M.-L., Godard, X., Archimède, H., Rychen, G., Mahieu, M., Feidt, C. Evaluation of soil intake by growing Creole young bulls in common grazing systems in humid tropical conditions. 2017, Animal, 11 (8), pp. 1363-1371.