

Assessment of bioavailability and bioaccessibility of persistent organic pollutants in the chain "soil-animal-food"

Farida Amutova, Matthieu Delannoy, G Konuspayeva, Stefan Jurjanz

▶ To cite this version:

Farida Amutova, Matthieu Delannoy, G Konuspayeva, Stefan Jurjanz. Assessment of bioavailability and bioaccessibility of persistent organic pollutants in the chain "soil-animal-food". Annual seminar of Doctoral School SIReNA, Feb 2020, Vandoeuvre-lès-Nancy, France. hal-03953966

HAL Id: hal-03953966 https://hal.univ-lorraine.fr/hal-03953966

Submitted on 24 Jan 2023 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Assessment of bioavailability and bioaccessibility of persistent organic pollutants in the chain «soil-animal-food»

^{1,2,3}F. Amutova, ¹M. Delannoy, ^{2,3}G.Konuspayeva, ¹S. Jurjanz

¹URAFPA, University of Lorraine-INRA, Vandoeuvre, France

²Faculty of Geography and Environmental Sciences, Al-Farabi Kazakh National University, Almaty, Kazakhstan

³Antigen LLP, Scientific and Production Enterprise, Almaty region, Kazakhstan


e-mail: amutovafb@gmail.com

CONTEXT

- Environmental contaminants (metals, POPs) can be stored during centuries in soil, therefore soil is a major exposure matrix
- All free ranged (food producing) animals ingest soil^{1;2} at different levels
- Soil is one of the main vector for contaminants in animals and then in food

OBJECTIVE


Reducing the bioavailability of POPs in the food chain "soil – farm animals – food products of animal origin" using different sequestration materials.




Preliminary results.

Mean and SDs of carry-over rates (COR) of PCDD/Fs in milk and eggs derived from the literature

Cl_n	Compound	Transfer	Transfer to eggs		
		COR, %	Transfer level	COR,	Transfer
				%	level
4	2,3,7,8-TCDD	34.5 ±14.3	high	$\textbf{40.5} \pm 0.7$	high
5	1,2,3,7,8 -PeCDD	36.0 ± 12.4	high	44.5 ± 4.9	high
6	1,2,3,4,7,8-HxCDD	26.8 ± 8.9	high	43.0 ± 1.4	high
6	1,2,3,6,7,8-HxCDD	28.1 ± 13.7	high	44.0 ±0	high
6	1,2,3,7,8,9-HxCDD	17.8 ± 6.5	medium	33.0 ± 1.4	high
7	1,2,3,4,6,7,8-HpCDD	5.06 ± 3.3	low	21.5 ± 0.7	medium
8	OCDD	0.72 ± 0.6	low	10.5 ± 3.5	low
4	2,3,7,8-TCDF	2.41 ± 2.70	low	40.0± 0	high
5	1,2,3,7,8-PeCDF	4.45 ± 4.00	low	43.5 ± 6.4	high
5	2,3,4,7,8-PeCDF	40.6 ± 18.2	high	41.0 ± 4.2	high
6	1,2,3,4,7,8-HxCDF	26.2 ± 11.3	high	43.5 ± 0.7	high
6	1,2,3,6,7,8-HxCDF	25.6 ± 11.6	high	41.0 ± 1.4	high
6	1,2,3,7,8,9-HxCDF	14.3 ± 6.10	medium	33.0 ± 2.8	high
6	2,3,4,6,7,8-HxCDF	19.0 ± 16.9	medium	40.0 ± 1.4	high
7	1,2,3,4,6,7,8-HpCDF	4.63 ± 3.20	low	18.0 ± 0	medium
7	1,2,3,4,7,8,9-HpCDF	5.87 ± 4.10	low	20.5 ± 0.7	medium
8	OCDF	0.54 ± 0.60	low	6.00 ± 1.4	low



Factorial plan (F1, F2) of PCA applied on mean CORs of PCBs for milk (A) and eggs (B), chlorine substitution at none, mono and di-ortho position, number of chlorines, molecular weight (M.W), n-Octanol/Water Partition Coefficient (K_{ow}) and transfer level.



Indication: Numbers correspond to the PCB isomers. Framed congeners are dioxin-like PCBs. Circle with PCBs are the high excreted congeners with the range 37.5-83.5% and 40.5-80.0% for milk and eggs respectively.

		(1 1		(Dergram), 20 Sorrag 2019
2	Evaluation of two contrasted activated	Article	published	Dec.,	Environmental Science and Pollution
	carbon based sequestration strategies to			2019	Research (IF - 2.8), In press, doi
	reduce soil bound chlordecone				10.1007/s11356-019-06494-z
	bioavailability in piglets				
3	Transfer of persistent organic pollutants	Article	In process	2020	Chemosphere (IF=4.4)
	in food producing animals (meta-				
	analysis)				
4	Assessment of PCDD/Fs and PCBs	Communicati	In process	2020	DIOXIN 2020, the 40th International
	transfer to milk and eggs	on/oral report			Symposium on Halogenated Persistent
					Organic Pollutants (POPs) from 30 August
					to 4 September 2020.
Ac	lditional articles that were recently pub	lished not in t	he topic of l	PhD:	
5	Volatile organic compounds profiles in	Article	published	Jan.,	International Journal of Biology and
•	milk fermented by lactic bacteria.		puolibiliou	2018	Chemistry 11, N_{2} 2, 57
	mink termented by lactic bacteria.			2010	doi.org/10.26577/ijbch-2018-2-345
			1 1 1 1 1	<u> </u>	
-	Comparative study of fatty acid and	Article	published	Sept.,	Journal of Dairy Science. Volume 102
6			_		
6	sterol profiles for the investigation of			2019	Issue 9,, p. 7723-7733
6				2019	Issue 9,, p. 7723-7733 doi.org/10.3168/jds.2018-15620.

1. Jurjanz, S., Germain, K., Juin, H., Jondreville, C. Plant and soil intake by organic broilers reared in tree- or grasscovered plots as determined by means of n-alkanes and of acid-insoluble ash. 2015, ANIMAL, 9 (5), pp. 888-898. 2. Jurjanz, S., Collas, C., Lastel, M.-L., Godard, X., Archimède, H., Rychen, G., Mahieu, M., Feidt, C. Evaluation of soil intake by growing Creole young bulls in common grazing systems in humid tropical conditions. 2017, Animal, 11 (8) pp. 1363-1371



Annual seminar of Doctoral School SIReNA, University of Lorraine, February 13, 2020