



HAL
open science

Patient reported outcome measures and patient engagement in heart failure clinical trials

Faiez Zannad, Jacqueline Alikhaani, Sadegh Alikhaani, Javed Butler, Jason Gordon, Klaus Jensen, Rani Khatib, Lorenzo Mantovani, Robin Martinez, Wanda F Moore, et al.

► To cite this version:

Faiez Zannad, Jacqueline Alikhaani, Sadegh Alikhaani, Javed Butler, Jason Gordon, et al.. Patient reported outcome measures and patient engagement in heart failure clinical trials: multi-stakeholder perspectives. *European Journal of Heart Failure*, 2023, 25 (4), pp.478-487. 10.1002/ejhf.2828 . hal-04033648

HAL Id: hal-04033648

<https://hal.univ-lorraine.fr/hal-04033648>

Submitted on 17 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PATIENT REPORTED OUTCOME MEASURES AND PATIENT ENGAGEMENT IN HEART FAILURE CLINICAL TRIALS: MULTI-STAKEHOLDER PERSPECTIVES

Authors

Faiez Zannad, MD, PhD¹, Jacqueline Alikhaani², Sadegh Alikhaani², Javed Butler, MD³, Jason Gordon, PhD⁴, Klaus Jensen, MD⁵, Rani Khatib, DPharm⁶, Lorenzo Mantovani, DSc⁷, Robin Martinez⁸, Wanda F. Moore⁹, Masahiro Murakami, MD, PhD¹⁰, Lothar Roessig, MD¹¹, John A. Spertus, MD, MPH¹², Norman Stockbridge, MD, PhD¹³, Harriette G.C. Van Spall, MD, MPH¹⁴, Clyde Yancy, MD¹⁵

Affiliations

¹Université de Lorraine, Inserm Clinical Investigation Center at Institut Lorrain du Coeur et des Vaisseaux, University Hospital of Nancy, Nancy, France;

²Los Angeles, USA;

³Department of Medicine, University of Mississippi Medical Center, Jackson, MS, USA;

⁴HEOR- Health Economics and Outcomes Research, Ltd. Cardiff, Wales, UK;

⁵Vifor Pharma; Zürich, Switzerland;

⁶Leeds Institute of Cardiovascular and Metabolic Medicine, University of Leeds, UK; Cardiology Department, Leeds Teaching Hospitals NHS Trust, Leeds, UK;

⁷Value-Based Healthcare Unit, IRCCS Multimedica Hospital, Milan, Italy;

⁸SmartPatients.com;

⁹Sarver Heart Center Women's Heart Health Education Comm., University of Arizona, Tucson, AZ, USA;

¹⁰Diabetes-Cardiovascular, Eli Lilly & Company, Indianapolis, IN, USA;

¹¹Clinical Development Group, Bayer AG, Germany;

¹²Department of Cardiology, Saint Luke's Mid America Heart Institute, University of Missouri-Kansas City, Kansas City, MO, USA;

¹³Division of Cardiology and Nephrology, Food and Drug Administration Center for Drug Evaluation and Research, Silver Spring, MD, USA;

¹⁴Department of Medicine, McMaster University, Hamilton, ON, Canada; Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, ON, Canada; Research Institute of St. Joseph's, Hamilton, ON, Canada; Population Health Research Institute, Hamilton, ON, Canada;

¹⁵Department of Internal Medicine, Division of Cardiology, Northwestern University, Feinberg School of Medicine, Chicago, IL, USA

Corresponding author

Dr. Faiez Zannad, MD, PhD

Université de Lorraine

INSERM, CIC 1439

Institut Lorrain du Coeur et des Vaisseaux, CHU 54500

Vandoeuvre-lès-Nancy, France

Tel: +33 383 15732225, Fax: +33 383 157324

E-mail: f.zannad@chru-nancy.fr

Word count: 3,766 (not including tables, figures, or references)

Abstract

There are many consequences of heart failure (HF), including symptoms, impaired health-related quality of life (HRQoL), and physical and social limitations (functional status). These have a substantial impact on patients' lives, yet are not routinely captured in clinical trials. Patient-reported outcomes (PROs) can quantify patients' experiences of their disease and its treatment.

Steps can be taken to improve the use of PROs in HF trials, in regulatory and payer decisions, and in patient care. Importantly, PRO measures (PROMs) must be developed with involvement of patients, family members, and caregivers from diverse demographic groups and communities. PRO data collection should become more routine not only in clinical trials but also in clinical practice. This may be facilitated by the use of digital tools and interdisciplinary patient advocacy efforts. There is a need for standardisation, not only of the PROM instruments, but also in procedures for analysis, interpretation and reporting PRO data. More work needs to be done to determine the degree of change that is important to patients and that which is associated with increased risks of clinical events. This "minimal clinically important difference (MCID)" requires further research to determine thresholds for different PROMs, to assess consistency across trial populations, and to define standards for improvement that warrant regulatory and reimbursement approvals.

PROs are a vital part of patient care and drug development, and more work should be done to ensure that these measures are both reflective of the patient experience and that they are more widely employed.

Keywords: Patient-reported outcomes, clinical trials, heart failure, minimal clinically important difference, patient partner, health-related quality of life

Introduction

In addition to increased risks of mortality and hospitalisation, heart failure (HF) is associated with symptoms that can result in physical and social limitations, impaired quality of life, and increased rates of depression, anxiety, and stress.^{1,2} While these latter outcomes have an important impact on patients' lives, they are not always captured as primary, or even secondary, endpoints in clinical trials.^{3,4}

One method for quantifying patients' lived experiences is through the formal collection of patient-reported outcomes (PROs) that directly assess the patient's experience, health behaviours, and the impact of the disease and its treatment on patient health status.⁵ There are three general components of patient health status: symptom burden, functional status (physical, mental, emotional, or social functional limitations), and health related quality of life (HRQoL) (Graphical abstract).³ Since each patient views their own health in different ways, HRQoL is measured distinctly from symptoms or functional limitations. Thus, it is essential to assess this aspect with specifically designed tools.³

PROs provide a unique perspective of the patient's experience of their illness and the effects of its treatment. These outcomes are relevant, not just to the patients, their families, and treating clinicians, but also to trialists, regulatory agencies, payers, and industry partners who seek to better understand these patient-centred outcomes. For example, PRO data may be used by trial sponsors to support a new product indication or labelling claim based on patient-reported treatment benefits, while payers may use these data to help determine the comparative value of different therapies.⁶

The Global Cardiovascular Clinical Trialists (CVCT) Workshop brings together CV clinical trialists, patients, clinicians, and regulatory, government and industry representatives to

discuss challenges facing CV clinical research. This paper is based on discussions at the 18th Global CVCT meeting (www.globalcvctforum.com) and aims to look at the benefits and challenges around the use of PROs to improve patient care, to inform drug development, and to aid regulatory and reimbursement decisions. The full session presentations and video recording can be accessed on the CVCT website.

Examples of PROMs used in heart failure

General measures of health status (e.g., 36-Item Short Form Health Survey [SF-36], EuroQol-5 Dimension [EQ-5D]) can provide the clinician and patient with useful information, but are not disease-specific and are designed to quantify total impact of all current health conditions on overall physical and emotional functioning. PROMs are commonly used in HF, and a number of HF-specific PRO measures (PROMs) including the Minnesota Living with Heart Failure Questionnaire (MLHFQ) and Kansas City Cardiomyopathy Questionnaire (KCCQ) (Table 1), are now becoming more commonly used.^{4, 7} These tools not only assess HF-specific health status, but have prognostic utility, with the KCCQ shown to be more sensitive to change^{8,}⁹ and superior to the MLHFQ for the prediction of clinical outcomes (death/transplant/left ventricular assist device implantation and hospitalisation).¹⁰

It is important that PROMs be developed with patients, family members, and caregivers, and many may need ongoing updating as environmental and other issues faced by patients may change over time. It is important to consult patients from diverse demographic groups and communities.

Minimal clinically important difference

While the primary analyses of most clinical trials compare the mean differences across populations of studied patients, these can be difficult to interpret from a clinical perspective. In contrast, categorizing score changes into gradations of clinically important changes can provide deeper insight into how much better (or worse) patients are likely to feel after a particular treatment versus another. It is also necessary to understand the degree of change that is associated with increased risks of clinical events. The minimum difference in score of any outcome that patients can perceive as either beneficial or harmful is referred to as the minimal clinically important difference (MCID).^{11, 12} The MCID provides a method for interpreting changes in scores to help inform treatment decisions, and is also useful when planning clinical trials and calculating sample size.^{11, 12} However, a majority of PROs do not have established MCIDs, and studies typically report whether there is a statistically significant difference in health status between treatment groups, but do not put this in the context of MCID or larger clinically important changes.⁷ Population-average changes often do not differ by thresholds that are greater than the MCID for individual patients, which can make inferences about the magnitude of health status improvement with the interventions difficult to appreciate. Responder analysis, which describes the proportions of patients in each treatment arm who experience changes in their health status of varying clinical magnitudes, is one way of assessing clinical relevance in this setting, although there are arguments against this analytic approach too.^{13, 14} In particular, assessment at a single time point is poor evidence to classify a patient as a responder.

Estimating MCID can be done using either an expert consensus, a distribution-based method, or an anchor-based method. The distribution-based method compares changes in scores and small, moderate, and large changes in patient health status. This method relies on the

statistical properties of the distribution of scores, and is used to estimate whether the magnitude of change in different groups in response to an intervention is more than would be expected from chance alone.^{11, 12, 15} An anchor-based determination compares the patient's numerical change in score for an outcome measure (e.g., changes on KCCQ) with an external subjective, independent evaluation of improvement (e.g., Patient Global Assessment (PGA)).^{11, 16}

The KCCQ is the HF-specific PRO that has undergone the most extensive research to define clinically important changes.^{15, 17-20} These efforts have led to a general consensus that a change in KCCQ total score of >5 points has been considered to be the MCID associated with a small improvement in patient health status.¹⁷ Changes of 10 and 20 points are considered to be moderate to large, and large to very large, clinical changes in patient health status. Such thresholds can transform the mean differences in scores along a 100-point range into the proportions of patients (e.g., number needed to treat) experiencing different magnitudes of improvement with treatment. It is also important to understand the stability and time course of benefit of a given intervention; some interventions may provide benefit early while others may take longer, and similarly, some benefits may be short-term and others more sustained.²¹

Using an anchor-based methodology (based on PGA), and data from the Ferric Carboxymaltose in patients with Heart Failure and Iron Deficiency (FAIR-HF) trial, the MCID threshold for a small positive change was associated with a change in KCCQ summary score of 3.6.¹⁶ This was somewhat lower than the typically accepted 5-point change described above.¹⁷ This demonstrates the utility of anchor-based methodology and supports the concept that even small improvements in HRQoL of a treatment over placebo can be clinically meaningful (Table 2).²²⁻²⁶

In addition, HRQoL may decline over the course of the disease due to deterioration of HF, but also due to other factors that may not be amenable to HF treatment, such as physical and social deterioration related to aging, non-CV comorbid conditions, and psychological factors.^{22,}
²⁷ Therefore, if a treatment can improve or at least stabilise a patient's HRQoL that could be considered beneficial.

Using PROs to improve patient care

PROMs are not yet widely incorporated into routine clinical practice for patients with HF, with clinicians generally placing greater importance on echocardiographic and biomarker responses, or using non-standardised ways to obtain subjective patient feedback and gauge symptomatic treatment response.²⁸ Yet PROs reflect what patients actually perceive and have been shown to improve the accuracy of clinicians' assessments of patients' New York Heart Association (NYHA) classification when available to the assessing physician.²⁹ Moreover, they can be used to assess their experience and behaviours, monitor therapeutic effectiveness, and prompt discussions relating to prognosis or end-of-life care. PROs may also be useful in guiding referrals for invasive cardiac care (e.g., percutaneous coronary intervention or coronary artery bypass grafting), non-cardiac care (e.g., psychiatric care, physiotherapy, or occupational therapy, and for assessing the quality of delivered healthcare. This suggests a potential role for interdisciplinary patient advocacy efforts in support of the widespread collection, standardization and implementation of PROs in clinical practice. Quality of life should be consistently and correctly assessed and managed throughout the patient journey. Incorporating routine measurements of HRQoL in clinical practice using PROMs can help both clinicians and patients to discuss the individual's experiences, and to consider whether the treatment program is meeting

their expectations. Some patients can complete questionnaires while awaiting their appointment, but others may need help completing these assessments. Routine measurements of HRQoL in clinical practice may be facilitated by the use of digital tools and interdisciplinary patient advocacy efforts. Patient organizations can provide coaching and support for patients to self-manage their condition and improve their quality of life. Formal assessments can also highlight unmet needs and areas for potential improvement in care delivery.³⁰

Patient perspective

The priorities and expectations of patients and clinicians often differ. Clinicians frequently are more concerned with cardiac anatomy and physiology, reducing mortality, and specific management issues, while patients just wish to feel better.^{28,31} However, both clinicians and patients generally assign high priority to improving HRQoL. Multiple studies have reported that patients often prefer a healthier life, with the majority willing to accept some decrease in longevity in return for improved health status.^{1,32,33} For example, in a survey of over 500 HF patients by the Pumping Marvellous Foundation, only 3% said that how long they live was the most important aspect of HF treatment, while 97% prioritised quality of life or both longevity and quality of life (Figure 1).²⁸ A survey within a family medicine clinic assessing disability and quality of life among HF patients found that almost 20% had severe disability, and another 28% had moderate disability.¹ The main areas affected were life activities, getting around, and participation in society.

Using PROs to better inform drug development and clinical trial design

Despite there being a significant increase in the inclusion of PROs in HF clinical trials over the past 20 years,⁴ there remains uncertainty around their interpretation and their role in the product development lifecycle. To further optimise study design requires a clearer understanding of the downstream utility of PRO data.

Trialist perspective

The inclusion of PROs in clinical trials was often thought to present multiple scientific and logistical challenges due to their fundamentally subjective nature.^{6, 34} Experience over time, however, has redefined the view of PROs as valid, reliable, and sensitive measures of patients' experiences. Using a PRO as a primary trial outcome complements the conduct of, and potentially reduces the need for large trials to demonstrate the less common outcome of mortality. For example, using the KCCQ overall summary score as the primary outcome measure and collecting data from mobile phones, enabled the recently completed Canagliflozin: Impact on Health Status, Quality of Life and Functional Status in Heart Failure (CHIEF-HF) trial to be conducted during the COVID-19 pandemic without a single in-person visit.^{35, 36} The CHIEF-HF also used PROs in the inclusion/exclusion criteria of the trial to define a more homogeneous study population.^{6, 36} In addition, PROMs, such as KCCQ, can be used to help identify an at risk population who are likely to respond to the treatment being tested. In HF trials, the patient population almost always includes a NYHA classification. However, clinician variability in assigning NYHA class is common,^{37, 38} while PRO scores have demonstrated reliability.^{6, 39}

Compared with HF with reduced ejection fraction (HFrEF), outcomes for HF with preserved ejection fraction (HFpEF) may be best defined by morbidity benefits as there continue to be few therapies that improve survival. Identifying the impact on HRQoL takes on even

greater importance. Because of the unmet need in this area, a drug that improves HRQoL with or without a mortality benefit may represent a beneficial treatment and could be approved by some regulatory bodies.⁴⁰ Using HRQoL and other PROs as primary endpoints in HFpEF clinical trials may be an important option to speed development of new treatments for these patients.

Since a new therapy could conceivably improve HRQoL or symptoms, but increase the risk of death/hospitalisation, or vice versa, it is well understood that PROMs may be primary outcome measures of trials, provided convincing safety data is collected in the same trial and/or in adjacent trials. Hierarchical clinical composites, possibly using the win ratio statistical approach is one way of investigating the HRQoL benefits without compromising data on the risk of death/hospitalisation.

PROs are essential in clinical trials involving patients with chronic diseases, and as the ongoing challenges of their interpretation are being resolved, their use should continue to increase. From the trialist's perspective, there is a need for standards for PRO collection and analyses to help ensure consistency across trials. Notably, the issue of missing PRO data in non-survivors needs appropriate analyses methods, such as hierarchical rank endpoints or imputation methods combining clinical events and PRO assessment.⁴¹

Industry perspective

In HF, there continues to be a need for interventions that provide HRQoL and symptomatic improvements.³¹ HRQoL merits a level of importance similar to traditional “hard” endpoints, but there must be strong evidence to show that an intervention has a beneficial effect on HRQoL and other PROs, especially if it has a neutral effect on survival or disease progression. This is important not just from the regulatory perspective, but also from a value

perspective. PRO can be used as part of the risk:benefit assessment, and help build or refute the value equation.

While the KCCQ is becoming more commonly used in HF, there are still many challenges. It is unknown whether a MCID defined in one trial population will be valid in another, for example, patients with different severities of HF, HF_rEF vs. HF_pEF, or patients with comorbidities such as diabetes, obesity, or a depressive disorder. Choosing a KCCQ subdomain as an endpoint can be a challenge, when the mechanism of drug effects on HRQoL are not known, e.g., via functional improvement and/or symptomatic improvement.

Another challenge is how to implement the KCCQ at the clinical trial sites. The tool has not been widely used by clinicians in clinical practice, and this limits the ability to rapidly screen potentially eligible patients or to supplement clinical trial results with “real-world” comparative effectiveness research. Web- or app-based versions are helping in this regard. Regardless of the measurement method it is important that it remain consistent throughout a trial to minimise within-subject variability.

Overall, from the industry perspective there continues to be a need for agreement among investigators, regulatory authorities, and trial sponsors on standards for instruments, endpoints, analytical approaches, and MCIDs. It is critical for industry to foster partnerships with patients, family, and caregivers from diverse backgrounds as part of their leadership and development teams, to ensure that decisions are relevant.

Using PROs to better inform regulatory and reimbursement decisions

Regulatory perspective

In 2019, the U.S. Food and Drug Administration (FDA) recognised that even in the absence of demonstrated effects on mortality or HF hospitalisations, substantial and persistent symptomatic or functional improvement can be an acceptable endpoint for HF drug approval.⁴⁰

⁴² The FDA has qualified the KCCQ and the MLHFQ as medical device development tools and the KCCQ for drug development, making approval and labelling claims more attainable with demonstration of clinically important improvements. The EMA guidance for chronic HF trials has not yet taken this step, but recognises PROs as a surrogate for exercise capacity for patient who are unable to undergo exercise testing.⁴³

While trialists and industry sponsors want standards for determining a MCID for a given PROM, regulatory agencies have not yet established these thresholds. For mortality or hospitalisation outcomes there is no concept of MCID, and while payers, clinicians, and patients might question the value of a small improvement, regulators only assess whether the benefits exceed the risks for approval of a treatment. However, for functional and symptomatic claims, the magnitude of change does matter. The rationale for this discrepant view may be the idea that society may benefit from a 1% decline in hospitalisation, but not from a 1% improvement in global symptoms.

It is often difficult to interpret the various analyses of PROMs from clinical trials. There can be substantial variability between patient populations. Despite a high intra-class correlation coefficient in stable patients, even within a given individual, results can vary with patients showing improvement at one time point but not at another. Disease trajectories can lead to changes even in PROs. Randomization and placebo control, as well as double blinding, are

critical in distinguishing benefit of an interventional treatment from the usual course of disease under standard management. This makes determining response rates a challenge. While there is no absolute threshold for clinically relevant finding, any effect larger than zero is taken into consideration. Statistically significant small changes may not be necessarily clinically meaningful, which argues for responder analyses that can clarify the proportions of patients that experience different magnitudes of clinical change.

Overall, there is a need to increase the understanding of the importance of specific HRQoL benefits. Answering questions such as: When is a new therapy approvable if the benefit versus existing care is solely an improvement in HRQoL? What if this therapy has less safety documentation and certainty than established therapies with years of experience? Will patients and clinicians find the documented HRQoL measures convincing enough to prefer the new therapy? Providing clear answers to industry decision-makers will increase the likelihood that better and lower-risk programs can be designed from early drug development and facilitate prioritisation and investment into HRQoL-oriented therapies.

Health economics/payer perspective

Assessing the incremental health benefits of an intervention relative to its incremental net long-term costs is often used by payers to determine treatment value, with PROs being an important component of the numerator in the value equation.⁴⁴ Payers make decisions that maximise both patient and population health outcomes, while operating within the context of finite resources. It is important that appropriate endpoints for capturing clinical, PRO, and economic outcomes are incorporated at the trial stage to subsequently inform reimbursement decisions.

Given the established association between clinical outcomes (e.g., survival, hospitalisation) and HF-specific PROMs, such as the KCCQ, such measures can be used to quantify clinical risk. The HF-specific PROMs allow quantification for decisions within HF, but the results of generic assessment tools are also crucial for payers who have to consider reimbursement decisions not just for HF but other CV conditions as well. For example, the FAIR-HF trial included not only the HF-specific self-reported patient global assessment and the KCCQ, but also the generic EQ-5D, which helps in the determination of treatment value across disease states.⁴⁵ However, in studies that do not collect such utility assessments, the KCCQ can be mapped to predict EQ-5D utilities, which can be used to support cost effectiveness analyses.⁴⁶ However, at least in HF, we are yet to see payers willing to pay a premium for a new therapy based only on improvement in PROs.⁴⁷

PROs are used both before, during, and after the introduction of treatments to support regulatory and reimbursement approvals. For example, the DAPA-HF trial (Dapagliflozin and Prevention of Adverse-Outcomes in Heart Failure) showed a reduction in the risks of mortality and hospitalisation for HF, and improvements in symptoms.⁴⁸ An economic evaluation was conducted using KCCQ and other outcomes to capture the progression of disease and the impact of different treatment modalities on both the clinical outcomes and the patient-relevant outcomes.⁴⁹ Results suggested that dapagliflozin did represent value for money for the treatment of HFrEF in the UK, German and Spanish healthcare systems.

In health economic evaluation of treatment value, PROs can play an important role in assessing the real-world relationship between patient health status, medication adherence, treatment effectiveness and clinical outcomes. In HF, lower HRQoL and symptoms of depression may be associated with non-adherence with HF medications.⁵⁰ Non-adherence may be associated

with poorer patient outcomes (e.g., HF mortality) and increased health system burden (e.g., increased frequency of hospitalisation for HF).⁵¹

Incorporating PROs into clinical practice guidelines

Despite recognition of the importance of functional capacity and HRQoL for patients with HF, recommendations in HF guidelines continue to be based predominantly on morbidity and mortality outcomes.^{52, 53} Recommendations based on PRO data are generally not class I, even when the data fulfil the criterion of “data derived from multiple randomised clinical trials or meta-analyses”. For example, in the 2021 ESC HF guidelines the recommendations relating to iron supplementation continue to be class IIa (“should be considered”) rather than class I (“is recommended” or “is indicated”).⁵²

This is likely related to ongoing issues around standards for reporting and interpreting PRO data. In HF clinical trials the NYHA class is not reported as a mean for each treatment arm, but rather as the proportion of patients within each NYHA class. This categorization facilitates understanding of the applicability of the trial to a certain patient population. Yet this approach is not used for HRQoL variables, with each trial reporting their own mean scores for the PROM used. An established and agreed-upon range for PROMs such as KCCQ or MLWHF scores would enable more standardised reporting and interpretation of PRO data.¹⁵

Relevant questions to be considered include how to assess PRO fluctuations to identify responders and how to convert patient- and population-improvement concepts into interpretable results for action by the clinician, including how to define the MCID for within-patient changes in different patient populations. Until these questions are answered, guideline recommendations will continue to receive a lower-class rating despite a high level of evidence. One approach would be to issue a class I recommendation where warranted especially in rare diseases, but one

that is accompanied with a specific statement acknowledging the lack of long-term mortality/morbidity data. Such a shift in guidance would further acknowledge the importance of improvements in PROs.

Beyond PROs: Involving patients in clinical trial design and execution

The focus of PROMs has been on developing and validating instruments that have optimal test characteristics. However, this often fails to consider the user-friendliness of these tools. Although using PROMs is a step in the right direction, it is important to ensure that they are being designed, and this type of data is being collected, from a diverse population that is representative of individuals with the disease. The burden of disease extends to family members and caregivers, and their perspective should also be solicited when designing PROMs. Having comprehensive representation of patients in clinical trials will ensure that HRQoL, efficacy and safety data are more relevant, and generalisable to broader populations.

“Patient-centric”, the concept of having the patient at the centre of clinical research and care, underscores that the goal of generating new knowledge is to improve patient care and outcomes. However, for some people this implies patients in the middle with things going on around them. Patient-centred research is meant to anchor research on the values and priorities of patients and to include them as partners, but true patient partnership continues to be underused. Even in Patient-Centered Outcomes Research Institute (PCORI) funded studies, engagement often included only passive consultants, while true collaboration and shared leadership occurred in only 37% of trials. A report from the PCORI funded ADAPTABLE (Aspirin Dosing: A Patient-centric Trial Assessing Benefits and Long-term Effectiveness) study concluded that patients must be an active part of the steering committee, and that true patient engagement

requires not just one or two “token” patients but a group of at least five or six diverse patients to produce effective, shared ideas and confidence.⁵⁴ It is critical that patients are seen as partners in their healthcare management and in clinical trials from development through trial design, execution, and dissemination.

The CardioVascular Clinical Trialists (CVCT) Forum has a long history of integrating patients as speakers and panellists during its annual meetings (www.globalcvctforum.com/patient-trialists). Each patient brings their own perspectives and priorities, their own experiences and knowledge about their disease to the table, and thus they are one of the most valuable resources in the development of improved treatments for HF.

Summary

Steps are needed to address the challenges around the standardisation and use of PROs in CV trials, regulatory and payers’ decisions and patient care. One of the most important aspects is to ensure that patients, family members, and caregivers from diverse demographic groups and communities are involved in the development of PROMs, as well as the design, analysis and interpretation of PROs in clinical trials. There is a need for standardisation not only of the PROM instruments, but also in procedures for analysis, interpretation and reporting of PRO data. A summary of recommendation is shown in Table 3.

ACKNOWLEDGEMENTS

This article was generated from discussions at the 18th Global Cardiovascular Clinical Trialists (CVCT) Forum held online in December 2021 (<https://www.globalcvctforum.com/>). The CVCT Forum is a strategic workshop for high level dialogue between clinical trialists, industry representatives, regulatory authorities, and patients. The authors would like to thank Pauline Lavigne and Steven Portelance (unaffiliated, supported by the CVCT Forum) for contributions to writing, and editing the manuscript.

Disclosures

Zannad: Personal fees for trial oversight committees, advisory boards, or speakers' bureau from Acceleron, Amgen, Applied Therapeutics, AstraZeneca, Bayer, Biopeutics, Boehringer, Cardior, Cereno Scientific, Cellprothera, CVRx, G3 Pharmaceuticals, Merck, Merck AG, Novartis, NovoNordisk, Pfizer, Roche, Servier, and Vifor Fresenius; is co-founder of CardioRenal and CVCT.

J. Alikhaani: None.

S. Alikhaani: None.

Butler: Consulting fees from Abbott, Adrenomed, Amgen, Applied Therapeutics, Array, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol Myers Squibb, CVRx, G3 Pharma, Impulse Dynamics, Innolife, Janssen, LivaNova, Luitpold, Medtronic, Merck, Novartis, NovoNordisk, Relypsa, Roche, Sequana Medical, and Vifor; and payment or honoraria for lectures, presentations, speakers bureaus from Novartis, Boehringer Ingelheim-Lilly, AstraZeneca, and Janssen.

Gordon: None.

Mantovani: Support from the Italian Ministry of Health Ricerca Corrente IRCCS Multimedica; participation on data safety monitoring boards or advisory boards for Pfizer, Bayer, Roche Diagnostics, and MSD.

Martinez: Salary support from SmartPatients.com.

Moore: None

Jensen: Employment with Vifor Pharma.

Khatib: Grants or contracts (jointly with institution) from Boehringer Ingelheim and AstraZeneca, and an educational grant paid directly to the institution from Novartis; speaker fees from Boehringer Ingelheim, Sanofi, Amarin, and AstraZeneca; leadership roles on various committees for the ESC (Science Committee, Allied Professions Task Force), UK Clinical Pharmacy Association (Cardiology Committee Chair), Pumping Marvelous Charity (Clinical Board Member), and National Institute for Health and Care Excellence (Heart Failure Quality Standards).

Roessig: Employment with Bayer AG; stock or stock options from Bayer AG.

Spertus: Grants or contracts from Janssen, Myokardia/Bristol Meyers Squibb, Abbott Vascular, and the National Institutes of Health; holds the copyright to the Kansas City Cardiomyopathy

Questionnaire, Seattle Angina Questionnaire, and the Peripheral Artery Questionnaire; consulting fees from Novartis, Bristol Meyers Squibb, Imbria Pharmaceuticals, Janssen, Merck, Terumo, Bayer, and AstraZeneca; and a leadership or fiduciary role with Blue Cross/Blue Shield of Kansas City.

Stockbridge: None.


Van Spall: Grants or contracts from Canadian Institutes of Health Research, Heart and Stroke Foundation of Canada.

Yancy: Participation on various NIH and NHLBI sponsored clinical trial and clinical research network oversight committees; served as President (past) of the American Heart Association (AHA); prior spousal employment with Abbott Labs, Inc.

Murakami: Employment with Eli Lilly and Company; stock or stock options from Eli Lilly and Company.

Graphical abstract

Components of patient-reported health status



Symptoms and functional status can be impacted by both the disease and its treatments

Tables

Table 1: Examples of types of heart failure PRO measures (PROMs)

Heart-failure specific questionnaires	Generic questionnaires
Minnesota Living With Heart Failure Questionnaire (MLHFQ)*	Self-Reported Dyspnea Scale
Kansas City Cardiomyopathy Questionnaire (KCCQ)*	EuroQol-5 Dimensions (EQ-5D)
Chronic Heart Failure Questionnaire*	Short Form Survey 36 or 12
Care-Related Quality of Life survey for Chronic Heart Failure (CaReQoL CHF)	Patient Global Assessment
Chronic Heart Failure Patient-Reported Outcome Measure (CHF-PROM)	Generic Quality of Life
Heart Failure Somatic Awareness Scale (HFSPS)	
Knowledge, Attitude, Self-care Practice & Health-related Quality of Life in Heart Failure (KAPQ-HF)	
Left Ventricular Dysfunction Questionnaire (LVD-36)	
Memorial Symptom Assessment Scale-Heart Failure (MSAS-HF)	
Patient-Reported Outcomes Measurement Information System-Plus-Heart Failure (PROMIS-Plus-HF)	
Quality of Life Questionnaire for Severe Heart Failure (QLQ-SHF)	

From references.^{4,7*} Adhere to validation and reporting of all FDA-recommended psychometric properties.⁷

Table 2: Treatment effect on HRQoL and morbidity-mortality outcomes after 6-12 months in HF trials

Trial²²	Difference in KCCQ OSS with active treatment vs. control	Clinical outcomes HF hospitalisation or CV death
FAIR-HF	+7 (SE ± 2)	Event rate decreased with treatment*
DAPA-HF	+2.8 (N/A)	HR 0.74 (0.65-0.85)
SHIFT	+2.4 (95% CI 0.91-3.85)	HR 0.82 (0.75-0.90)
EMPEROR-HFrEF ^{24, 25}	+1.5 (95% CI 0.29-2.74)	HR 0.75 (0.65-0.86)
HFpEF ^{23, 26}	+1.6 (95% CI 0.76-2.44)	HR 0.77 (0.67-0.87)
PARADIGM-HF	+1.3 (95% CI 0.6-2.0)	HR 0.80 (0.73-0.87)
TOPCAT	+1.3 (SE ± 0.4)	HR 0.89 (0.77-1.04)
PARAGON-HF [†]	+1.0 (95% CI 0.0-2.1)	RR 0.87 (0.75-1.01)

Adapted from references.²²⁻²⁶

*HRs not reported because study was underpowered for morbidity-mortality outcomes; [†]Results for the KCCQ clinical summary score. CI, confidence interval; HF, hospitalisation; HR, time to first event hazard ratio; OSS, overall summary score, RR, total event risk ratio; SE, standard error


Table 3: Recommendation for improving the use of PROs in cardiovascular trials, regulatory and payers' decisions and care

PROMs	<ul style="list-style-type: none"> • Involve patients, family members, and caregivers from diverse demographic groups and communities in the development of PROMs • Improve PROM and data collection using Web- or app-based versions
PRO use in clinical care	<ul style="list-style-type: none"> • Incorporate routine measurements of HRQoL in clinical practice • Develop interdisciplinary patient advocacy efforts in support of the widespread collection, standardisation and implementation of PROMs in clinical practice
PROs in clinical trials	<ul style="list-style-type: none"> • Early assessment and inclusion of appropriate PROs in trial design and endpoint selection to support value-based evidence assessment and translation in regulatory, reimbursement, and clinical practice decision-making
PRO data analysis	<ul style="list-style-type: none"> • Standardize PRO data analyses to help ensure consistency across trials • Develop consensual methods for analysis of missing PRO data in non-survivors • Develop methods to analyse PRO fluctuations to identify responders
PRO results reporting and interpretation	<ul style="list-style-type: none"> • Adhere to standards for validation and reporting, and develop guidelines for interpreting PRO data • Determine the degree of change that is important to patients, or that is associated with increased risks of clinical events to support responder analyses and interpretability of study results • Investigate the issue of consistency of MCID across various trial populations
International regulatory PRO guidance	<ul style="list-style-type: none"> • Develop standards for clinically meaningful thresholds of PRO improvement • Define approvable PRO benefit ratios • Develop criteria to assess whether a new therapy is approvable if the benefit vs. existing care is solely an improvement in HRQoL
International payer valuation methods of PRO benefits	<ul style="list-style-type: none"> • Reach agreement on when coverage for a new therapy is warranted based solely on improvement in PROs

HRQoL, health-related quality of life; MCID, minimal clinically important difference; PRO, Patient-reported outcome; PROM, PRO measures

Figures

Figure 1: Survey of patients with HF: What is more important to you in terms of HF drugs and interventions? Quality of life, how long you live, or both?



Adapted from reference.²⁸ Data from a survey of over 500 patients conducted by the Pumping Marvellous Foundation.

References

1. Garcia-Olmos L, Batlle M, Aguilar R, Porro C, Carmona M, Alberquilla A, Sanchez-Gomez LM, Monge E, Lopez-Rodriguez AB, Benito L, Banos N, Simon A, Martinez-Alvarez MA, Luque EM, Garcia-Benito C. Disability and quality of life in heart failure patients: a cross-sectional study. *Fam Pract* 2019;**36**(6):693-698.
2. Tsabedze N, Kinsey JH, Mpanya D, Mogashoa V, Klug E, Manga P. The prevalence of depression, stress and anxiety symptoms in patients with chronic heart failure. *Int J Ment Health Syst* 2021;**15**(1):44.
3. Rumsfeld JS, Alexander KP, Goff DC, Jr., Graham MM, Ho PM, Masoudi FA, Moser DK, Roger VL, Slaughter MS, Smolderen KG, Spertus JA, Sullivan MD, Treat-Jacobson D, Zerwic JJ, American Heart Association Council on Quality of C, Outcomes Research CoC, Stroke Nursing CoE, Prevention CoPVD, Stroke C. Cardiovascular health: The importance of measuring patient-reported health status: a scientific statement from the American Heart Association. *Circulation* 2013;**127**(22):2233-49.
4. Eliya Y, Averbuch T, Le N, Xie F, Thabane L, Mamas MA, Van Spall HGC. Temporal trends and factors associated with the inclusion of patient-reported outcomes in heart failure randomized controlled trials: A systematic review. *J Am Heart Assoc* 2021;**10**(21):e022353.
5. Kluzek S, Dean B, Wartolowska KA. Patient-reported outcome measures (PROMs) as proof of treatment efficacy. *BMJ Evid Based Med* 2022;**27**(3):153-155.
6. Coles TM, Hernandez AF, Reeve BB, Cook K, Edwards MC, Boutin M, Bush E, Degboe A, Roessig L, Rudolph A, McNulty P, Patel N, Kay-Mugford T, Vernon M, Woloschak M, Buchele G, Spertus JA, Roe MT, Bury D, Weinfurt K. Enabling patient-reported outcome measures in clinical trials, exemplified by cardiovascular trials. *Health Qual Life Outcomes* 2021;**19**(1):164.
7. Chew DS, Whitelaw S, Vaduganathan M, Mark DB, Van Spall HGC. Patient-reported outcome measures in cardiovascular disease: An evidence map of the psychometric properties of health status instruments. *Ann Intern Med* 2022;**175**(10):1431-1439.
8. Kitzman DW, Brubaker P, Morgan T, Haykowsky M, Hundley G, Kraus WE, Eggebeen J, Nicklas BJ. Effect of caloric restriction or aerobic exercise training on peak oxygen consumption and quality of life in obese older patients with heart failure with preserved ejection fraction: A randomized clinical trial. *JAMA* 2016;**315**(1):36-46.
9. Green CP, Porter CB, Bresnahan DR, Spertus JA. Development and evaluation of the Kansas City Cardiomyopathy Questionnaire: a new health status measure for heart failure. *J Am Coll Cardiol* 2000;**35**(5):1245-55.
10. Yee D, Novak E, Platts A, Nassif ME, LaRue SJ, Vader JM. Comparison of the Kansas City Cardiomyopathy Questionnaire and Minnesota Living With Heart Failure Questionnaire in predicting heart failure outcomes. *Am J Cardiol* 2019;**123**(5):807-812.
11. McGlathlin AE, Lewis RJ. Minimal clinically important difference: defining what really matters to patients. *JAMA* 2014;**312**(13):1342-3.
12. Apaza J, Franco J, Meza N, Madrid E, Loezar C, Garegnani L. Minimal clinically important difference: The basics. *Medwave* 2021;**21**:e8149.
13. Snapinn SM, Jiang Q. Responder analyses and the assessment of a clinically relevant treatment effect. *Trials* 2007;**8**:31.
14. Psootka MA, Abraham WT, Fiuzat M, Filippatos G, Lindenfeld J, Ahmad T, Felker GM, Jacob R, Kitzman DW, Leifer ES, Lewis EF, Mentz RJ, Nkulikiyinka R, Ni W, Schaber DE,

- Sharma A, Solomon SD, Stockbridge N, Teerlink JR, Unger EF, Whellan DJ, Wittes J, Anker SD, O'Connor CM. Functional and symptomatic clinical trial endpoints: The HFC-ARC Scientific Expert Panel. *JACC Heart Fail* 2022;**10**(12):889-901.
15. Spertus JA, Jones PG, Sandhu AT, Arnold SV. Interpreting the Kansas City Cardiomyopathy Questionnaire in clinical trials and clinical care: JACC state-of-the-art review. *J Am Coll Cardiol* 2020;**76**(20):2379-2390.
 16. Butler J, Khan MS, Mori C, Filippatos GS, Ponikowski P, Comin-Colet J, Roubert B, Spertus JA, Anker SD. Minimal clinically important difference in quality of life scores for patients with heart failure and reduced ejection fraction. *Eur J Heart Fail* 2020;**22**(6):999-1005.
 17. Spertus J, Peterson E, Conard MW, Heidenreich PA, Krumholz HM, Jones P, McCullough PA, Pina I, Tooley J, Weintraub WS, Rumsfeld JS, Cardiovascular Outcomes Research Consortium. Monitoring clinical changes in patients with heart failure: a comparison of methods. *Am Heart J* 2005;**150**(4):707-15.
 18. Pokharel Y, Khariton Y, Tang Y, Nassif ME, Chan PS, Arnold SV, Jones PG, Spertus JA. Association of serial Kansas City Cardiomyopathy Questionnaire assessments with death and hospitalization in patients with heart failure with preserved and reduced ejection fraction: A secondary analysis of 2 randomized clinical trials. *JAMA Cardiol* 2017;**2**(12):1315-1321.
 19. Flynn KE, Lin L, Moe GW, Howlett JG, Fine LJ, Spertus JA, McConnell TR, Pina IL, Weinfurt KP. Relationships between changes in patient-reported health status and functional capacity in outpatients with heart failure. *Am Heart J* 2012;**163**(1):88-94 e3.
 20. Dreyer RP, Jones PG, Kutty S, Spertus JA. Quantifying clinical change: Discrepancies between patients' and providers' perspectives. *Qual Life Res* 2016;**25**(9):2213-20.
 21. Khan MS, Butler J. Stability of changes in health status: Next step in comprehensively assessing patient-reported outcomes. *JAMA* 2022;**328**(10):923-924.
 22. Ferreira JP. Health-related quality of life scores: ending the minimum 5-point difference as the clinically meaningful threshold. *Eur J Heart Fail* 2020;**22**(6):1006-1008.
 23. Packer M, Butler J, Zannad F, Filippatos G, Ferreira JP, Pocock SJ, Carson P, Anand I, Doehner W, Haass M, Komajda M, Miller A, Pehrson S, Teerlink JR, Schnaidt S, Zeller C, Schnee JM, Anker SD. Effect of empagliflozin on worsening heart failure events in patients with heart failure and preserved ejection fraction: EMPEROR-Preserved trial. *Circulation* 2021;**144**(16):1284-1294.
 24. Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, Januzzi J, Verma S, Tsutsui H, Brueckmann M, Jamal W, Kimura K, Schnee J, Zeller C, Cotton D, Bocchi E, Bohm M, Choi DJ, Chopra V, Chuquiure E, Giannetti N, Janssens S, Zhang J, Gonzalez Juanatey JR, Kaul S, Brunner-La Rocca HP, Merkely B, Nicholls SJ, Perrone S, Pina I, Ponikowski P, Sattar N, Senni M, Seronde MF, Spinar J, Squire I, Taddei S, Wanner C, Zannad F, EMPEROR-Reduced Trial Investigators. Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med* 2020;**383**(15):1413-1424.
 25. Butler J, Anker SD, Filippatos G, Khan MS, Ferreira JP, Pocock SJ, Giannetti N, Januzzi JL, Pina IL, Lam CSP, Ponikowski P, Sattar N, Verma S, Brueckmann M, Jamal W, Vedin O, Peil B, Zeller C, Zannad F, Packer M, EMPEROR-Reduced Trial Committees Investigators. Empagliflozin and health-related quality of life outcomes in patients with heart failure with reduced ejection fraction: the EMPEROR-Reduced trial. *Eur Heart J* 2021;**42**(13):1203-1212.
 26. Butler J, Filippatos G, Jamal Siddiqi T, Brueckmann M, Bohm M, Chopra VK, Pedro Ferreira J, Januzzi JL, Kaul S, Pina IL, Ponikowski P, Shah SJ, Senni M, Vedin O, Verma S, Peil B, Pocock SJ, Zannad F, Packer M, Anker SD. Empagliflozin, health status, and quality of life in

- patients with heart failure and preserved ejection fraction: The EMPEROR-Preserved trial. *Circulation* 2022;**145**(3):184-193.
27. Chandra A, Lewis EF, Claggett BL, Desai AS, Packer M, Zile MR, Swedberg K, Rouleau JL, Shi VC, Lefkowitz MP, Katova T, McMurray JJV, Solomon SD. Effects of sacubitril/valsartan on physical and social activity limitations in patients with heart failure: A secondary analysis of the PARADIGM-HF trial. *JAMA Cardiol* 2018;**3**(6):498-505.
 28. Hooper J, Hartshorne-Evans N, Cunnington C, Ahmed FZ. Iron deficiency—the invisible comorbidity in HF: prioritising QoL as a target for treatment. *Br J Cardiol* 2021;**28**(suppl 1):S15-S18.
 29. Sandhu AT, Zheng J, Kalwani N, Gupta A, Calma J, Skye M, Lan R, Yu B, Spertus J, Heidenreich P. Impact of patient-reported outcome measurement in heart failure clinic on clinician health status assessment and patient experience: A sub-study of the PRO-HF trial. *Circ Heart Fail* 2022.
 30. Mercieca-Bebber R, King MT, Calvert MJ, Stockler MR, Friedlander M. The importance of patient-reported outcomes in clinical trials and strategies for future optimization. *Patient Relat Outcome Meas* 2018;**9**:353-367.
 31. Taylor CJ, Huntley AL, Burden J, Gadoud A, Gronlund T, Jones NR, Wicks E, McKelvie S, Byatt K, Lehman R, King A, Mumford B, Feder G, Mant J, Hobbs R, Johnson R. Research priorities in advanced heart failure: James Lind alliance priority setting partnership. *Open Heart* 2020;**7**(1).
 32. Kraai IH, Vermeulen KM, Luttik ML, Hoekstra T, Jaarsma T, Hillege HL. Preferences of heart failure patients in daily clinical practice: quality of life or longevity? *Eur J Heart Fail* 2013;**15**(10):1113-21.
 33. Lewis EF, Johnson PA, Johnson W, Collins C, Griffin L, Stevenson LW. Preferences for quality of life or survival expressed by patients with heart failure. *J Heart Lung Transplant* 2001;**20**(9):1016-24.
 34. Calvert M, King M, Mercieca-Bebber R, Aiyegbusi O, Kyte D, Slade A, Chan AW, Basch E, Bell J, Bennett A, Bhatnagar V, Blazeby J, Bottomley A, Brown J, Brundage M, Campbell L, Cappelleri JC, Draper H, Dueck AC, Ells C, Frank L, Golub RM, Griebisch I, Haywood K, Hunn A, King-Kallimanis B, Martin L, Mitchell S, Morel T, Nelson L, Norquist J, O'Connor D, Palmer M, Patrick D, Price G, Regnault A, Retzer A, Revicki D, Scott J, Stephens R, Turner G, Valakas A, Velikova G, von Hildebrand M, Walker A, Wenzel L. SPIRIT-PRO Extension explanation and elaboration: Guidelines for inclusion of patient-reported outcomes in protocols of clinical trials. *BMJ Open* 2021;**11**(6):e045105.
 35. Spertus JA, Birmingham MC, Butler J, Lingvay I, Lanfear DE, Abbate A, Kosiborod ML, Fawcett C, Burton P, Damaraju CV, Januzzi JL, Whang J. Novel trial design: CHIEF-HF. *Circ Heart Fail* 2021;**14**(3):e007767.
 36. Spertus JA, Birmingham MC, Nassif M, Damaraju CV, Abbate A, Butler J, Lanfear DE, Lingvay I, Kosiborod MN, Januzzi JL. The SGLT2 inhibitor canagliflozin in heart failure: The CHIEF-HF remote, patient-centered randomized trial. *Nat Med* 2022;**28**(4):809-813.
 37. Raphael C, Briscoe C, Davies J, Ian Whinnett Z, Manisty C, Sutton R, Mayet J, Francis DP. Limitations of the New York Heart Association functional classification system and self-reported walking distances in chronic heart failure. *Heart* 2007;**93**(4):476-82.
 38. Taichman DB, McGoon MD, Harhay MO, Archer-Chicko C, Sager JS, Murugappan M, Chakinali MM, Palevsky HI, Gallop R. Wide variation in clinicians' assessment of New York

Heart Association/World Health Organization functional class in patients with pulmonary arterial hypertension. *Mayo Clin Proc* 2009;**84**(7):586-92.

39. Spertus JA, Jones PG. Development and validation of a short version of the Kansas City Cardiomyopathy Questionnaire. *Circ Cardiovasc Qual Outcomes* 2015;**8**(5):469-76.
40. US Department of Health Human Services FDA Center for Drug Evaluation Research, US Department of Health Human Services FDA Center for Biologics Evaluation Research. *Treatment for heart failure: Endpoints for drug development - draft guidance*. 2019, <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/treatment-heart-failure-endpoints-drug-development-guidance-industry>.
41. Spertus JV, Hatfield LA, Cohen DJ, Arnold SV, Ho M, Jones PG, Leon M, Zuckerman B, Spertus JA. Integrating quality of life and survival outcomes in cardiovascular clinical trials. *Circ Cardiovasc Qual Outcomes* 2019;**12**(6):e005420.
42. Khan MS, Butler J, Greene SJ. Patient-reported outcomes for heart failure with preserved ejection fraction: conducting quality studies on quality of life. *Eur J Heart Fail* 2020;**22**(6):1019-1021.
43. European Medicines Agency. *Guideline on clinical investigation of medicinal products for the treatment of chronic heart failure*. CPMP/EWP/235/95, Rev.2. 2017, https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-clinical-investigation-medicinal-products-treatment-chronic-heart-failure-revision-2_en.pdf (March 17 2022).
44. Anderson JL, Heidenreich PA, Barnett PG, Creager MA, Fonarow GC, Gibbons RJ, Halperin JL, Hlatky MA, Jacobs AK, Mark DB, Masoudi FA, Peterson ED, Shaw LJ, ACC AHA Task Force on Performance Measures, ACC AHA Task Force on Practice Guidelines. ACC/AHA statement on cost/value methodology in clinical practice guidelines and performance measures: A report of the American College of Cardiology/American Heart Association Task Force on Performance Measures and Task Force on Practice Guidelines. *Circulation* 2014;**129**(22):2329-45.
45. Anker SD, Comin Colet J, Filippatos G, Willenheimer R, Dickstein K, Drexler H, Luscher TF, Bart B, Banasiak W, Niegowska J, Kirwan BA, Mori C, von Eisenhart Rothe B, Pocock SJ, Poole-Wilson PA, Ponikowski P, Fair-HF Trial Investigators. Ferric carboxymaltose in patients with heart failure and iron deficiency. *N Engl J Med* 2009;**361**(25):2436-48.
46. Parizo JT, Goldhaber-Fiebert JD, Salomon JA, Khush KK, Spertus JA, Heidenreich PA, Sandhu AT. Cost-effectiveness of dapagliflozin for treatment of patients with heart failure with reduced ejection fraction. *JAMA Cardiol* 2021;**6**(8):926-935.
47. Thomas M, Jones PG, Cohen DJ, Suzanne AV, Magnuson EA, Wang K, Thourani VH, Fonarow GC, Sandhu AT, Spertus JA. Predicting the EQ-5D utilities from the Kansas City Cardiomyopathy Questionnaire in patients with heart failure. *Eur Heart J Qual Care Clin Outcomes* 2021;**7**(4):388-396.
48. Kosiborod MN, Jhund PS, Docherty KF, Diez M, Petrie MC, Verma S, Nicolau JC, Merkely B, Kitakaze M, DeMets DL, Inzucchi SE, Kober L, Martinez FA, Ponikowski P, Sabatine MS, Solomon SD, Bengtsson O, Lindholm D, Niklasson A, Sjostrand M, Langkilde AM, McMurray JJV. Effects of dapagliflozin on symptoms, function, and quality of life in patients with heart failure and reduced ejection fraction: Results from the DAPA-HF trial. *Circulation* 2020;**141**(2):90-99.
49. McEwan P, Darlington O, McMurray JJV, Jhund PS, Docherty KF, Bohm M, Petrie MC, Bergenheim K, Qin L. Cost-effectiveness of dapagliflozin as a treatment for heart failure with

reduced ejection fraction: a multinational health-economic analysis of DAPA-HF. *Eur J Heart Fail* 2020;**22**(11):2147-2156.

50. Rasmussen AA, Wiggers H, Jensen M, Berg SK, Rasmussen TB, Borregaard B, Thrysoe L, Thorup CB, Mols RE, Larsen SH, Johnsen SP. Patient-reported outcomes and medication adherence in patients with heart failure. *Eur Heart J Cardiovasc Pharmacother* 2021;**7**(4):287-295.

51. Fitzgerald AA, Powers JD, Ho PM, Maddox TM, Peterson PN, Allen LA, Masoudi FA, Magid DJ, Havranek EP. Impact of medication nonadherence on hospitalizations and mortality in heart failure. *J Card Fail* 2011;**17**(8):664-9.

52. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumach A, Bohm M, Burri H, Butler J, Celutkiene J, Chioncel O, Cleland JGF, Coats AJS, Crespo-Leiro MG, Farmakis D, Gilard M, Heymans S, Hoes AW, Jaarsma T, Jankowska EA, Lainscak M, Lam CSP, Lyon AR, McMurray JJV, Mebazaa A, Mindham R, Muneretto C, Francesco Piepoli M, Price S, Rosano GMC, Ruschitzka F, Kathrine Skibelund A, Group. ESD. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;**42**(36):3599-3726.

53. Maddox TM, Januzzi JL, Jr., Allen LA, Breathett K, Butler J, Davis LL, Fonarow GC, Ibrahim NE, Lindenfeld J, Masoudi FA, Motiwala SR, Oliveros E, Patterson JH, Walsh MN, Wasserman A, Yancy CW, Youmans QR. 2021 Update to the 2017 ACC Expert Consensus Decision Pathway for Optimization of Heart Failure Treatment: Answers to 10 Pivotal Issues About Heart Failure With Reduced Ejection Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol* 2021;**77**(6):772-810.

54. Faulkner M, Alikhaani J, Brown L, Cruz H, Davidson D, Gregoire K, Berdan L, Rorick T, Jones WS, Pletcher MJ. Exploring meaningful patient engagement in ADAPTABLE (Aspirin Dosing: A Patient-centric Trial Assessing Benefits and Long-term Effectiveness). *Med Care* 2018;**56 Suppl 10 Suppl 1**:S11-S15.