



**HAL**  
open science

## Sodium-glucose co-transporter 2 inhibitors in heart failure: an updated evidence-based practical guidance for clinicians

Luca Monzo, Ilaria Ferrari, Francesco Cicogna, Claudia Tota, Gennaro Cice, Nicolas Girerd, Leonardo Calò

### ► To cite this version:

Luca Monzo, Ilaria Ferrari, Francesco Cicogna, Claudia Tota, Gennaro Cice, et al.. Sodium-glucose co-transporter 2 inhibitors in heart failure: an updated evidence-based practical guidance for clinicians. European Heart Journal Supplements, 2023, 25 (Supplement C), pp.C309 - C315. 10.1093/eurheartj-sup/suad055 . hal-04087732

**HAL Id: hal-04087732**

**<https://hal.univ-lorraine.fr/hal-04087732>**

Submitted on 3 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# Sodium-glucose co-transporter 2 inhibitors in heart failure: an updated evidence-based practical guidance for clinicians

Luca Monzo<sup>1,2\*</sup>, Ilaria Ferrari<sup>2</sup>, Francesco Cicogna<sup>2</sup>, Claudia Tota<sup>2</sup>,  
Gennaro Cice<sup>2</sup>, Nicolas Girerd<sup>1</sup>, and Leonardo Calò<sup>2</sup>

<sup>1</sup>Centre d'Investigation Clinique 1433 module Plurithématique, CHRU Nancy - Hôpitaux de Brabois, Institut Lorrain du Cœur et des Vaisseaux Louis Mathieu, Université de Lorraine INSERM, 4 rue du Morvan, 54500 Vandoeuvre les Nancy, France; and <sup>2</sup>Department of Cardiology, Policlinico Casilino, Rome 00169, Italy

## KEYWORDS

Heart failure;  
SGLT2 inhibitors;  
Cardiovascular outcome trials;  
Clinical practical guide

The sodium-glucose co-transporter 2 (SGLT2) inhibitors have been shown to reduce risks of clinical events in patients with heart failure (HF), with early and sustained benefits regardless of ejection fraction, diabetic status, and care setting. As part and parcel of the modern foundational HF therapy, clinicians should be familiar with these drugs, in order to implement their use and limit the potential adverse effects. We present an up-to-date review of current evidence and a practical guide for the prescription of SGLT2 inhibitors in patients with HF, highlighting important elements for patient selection, treatment initiation, dosing, and problem solving.

## Introduction

Over the last 5 years, sodium-glucose co-transporter 2 (SGLT2) inhibitors have been tested in dedicated heart failure (HF) trials with outstanding positive results (Figure 1). A recent meta-analysis including 21,947 participants from five HF trials showed that SGLT2 inhibitors significantly reduced the risk of composite cardiovascular death or hospitalization for HF (hazard ratio [HR] 0.77; 95% confidence interval [CI], 0.72-0.82), cardiovascular death for HF (HR 0.87; 95% CI, 0.79-0.95), first hospitalization for HF (HR 0.72; 95% CI, 0.67-0.78), and all-cause mortality (HR 0.92; 95% CI, 0.86-0.99), agnostic of left ventricular ejection fraction (LVEF) and diabetic status.<sup>1</sup> In light of the impressive results from recent trials, clinical practice guidelines recommend the use of SGLT2 inhibitors in HF patients across all the spectrum of ejection fraction and in a wide range of care settings.<sup>2,3</sup>


We provide an up-to-date review of current evidence and a practical guide for the prescription of SGLT2

inhibitors in patients with HF, highlighting important elements for patient selection, treatment initiation, dosing, and anticipated adverse effects.

## SGLT2 inhibitors in patients with chronic heart failure

The SGLT2 inhibitors are, together with the renin-angiotensin-system inhibitors (RASi), angiotensin receptor-neprilysin inhibitors (ARNi),  $\beta$ -blockers, and mineralocorticoid receptor antagonists (MRA), the evidence-based backbone therapy in HF patients with reduced ejection fraction (HFrEF).<sup>3</sup> Indeed, a recent meta-analysis including 95,444 participants from 75 trials showed that the association of ARNi,  $\beta$ -blockers, MRA, and SGLT2 inhibitors was the best combination for reducing all-cause death (HR 0.39; 95% CI, 0.31-0.49) and the composite outcome of cardiovascular death or first hospitalization for HF (HR 0.36; 95% CI, 0.29-0.46).<sup>10</sup> Currently, there are two major outcome trials assessing the effect of SGLT2 inhibitors in patients with chronic stable HFrEF with or without diabetes: DAPA-HF<sup>8</sup> (assessing dapagliflozin) and EMPEROR-Reduced<sup>11</sup> (assessing empagliflozin) trials (Figure 1). Overall, in these two studies, SGLT2 inhibition with

\*Corresponding author. Tel: +33 383157306, Fax: +33 383157324, Email: [l.monzo@chru-nancy.fr](mailto:l.monzo@chru-nancy.fr)



**Figure 1** Timeline of SGLT2 inhibitor trials targeting patients with heart failure (HF). Large and narrow boxes show enrolment and follow-up period, respectively. The result for the primary outcome is reported below the trial name. Modified from Tromp *et al.*<sup>4</sup> Abbreviations: CHF, chronic heart failure; HR, hazard ratio; IV, intravenous; WR, win ratio. The primary endpoint was defined as (A) EMPULSE<sup>4</sup>—clinical benefit, defined as a hierarchical composite of death from any cause, number of HF events and time to first HF event, or a  $\geq 5$  point difference in change from baseline KCCQ score at 90 days, as assessed using a WR; (B) SOLOIST-WHF<sup>5</sup>—composite of deaths from cardiovascular causes, HF hospitalizations and urgent visits for HF (first and subsequent events); (C) EMPEROR-Reduced<sup>6</sup> and EMPEROR-Preserved<sup>7</sup>—composite of cardiovascular death or hospitalizations for HF; (D) DAPA-HF<sup>8</sup> and DELIVER<sup>9</sup>—composite of worsening HF (defined as either an unplanned hospitalization for HF or an urgent visit for HF) or cardiovascular death. \*minimum time to randomization from the withdrawal of intravenous HF therapy in acute decompensated HF patients [EMPULSE—off inotropes for 24 h and no vasodilators or escalating diuretics for 6 h; SOLOIST-WHF—off IV inotropes or IV vasodilators (except for nitrates) for 48 h and having transitioned from IV to oral diuretic therapy; DELIVER—off intravenous HF therapy (including diuretics) for at least 24 h]. <sup>#</sup>enrolled only type 2 diabetic patients.

empagliflozin or dapagliflozin—when added to guideline-recommended therapy for HF—reduced all-cause and cardiovascular death, hospitalizations for HF, and serious adverse renal outcomes in patients with a broad spectrum of severity of HF<sub>rEF</sub>.<sup>12</sup> The benefit of empagliflozin and dapagliflozin on the primary endpoint of both trials—the combined risk of cardiovascular death or hospitalization for HF—was primarily driven by an approximately 30% relative reduction in the risk of hospitalization for HF. A benefit on hospitalizations for HF was observed whether the analysis was confined to first events or to all events (first and recurrent).<sup>12</sup> The modest size of the cardiovascular death benefit might explain why it is observed inconsistently in individual trials. Specifically, the relative reduction in cardiovascular death was 18% in DAPA-HF and 8% in EMPEROR-Reduced.<sup>8,11</sup>

In chronic stable HF patients with mildly reduced (HF<sub>m</sub>rEF) or preserved (HF<sub>p</sub>rEF) ejection fraction, two large outcome trials evaluated the effect of SGLT2 inhibitors in patients with or without diabetes: DELIVER<sup>9</sup> (assessing dapagliflozin) and EMPEROR-Preserved<sup>7</sup> (assessing empagliflozin) trials (Figure 1). In both these studies, SGLT2 inhibition similarly and strongly reduced the combined risk of cardiovascular death or hospitalization for HF (primary endpoint) in a broad range of patients. Similar to HF<sub>rEF</sub> trials on SGLT2 inhibitors, risk reductions in the primary composite endpoint were driven by substantial treatment effects on hospitalizations for HF

(average 26% relative risk reduction), with modest effects on cardiovascular death. The main finding of the DELIVER trial that distinguished it from the EMPEROR-Preserved trial (and also other neurohormonal blockade trials) was the benefit among patients with an LVEF of 60% or more and among those with an LVEF that had improved to more than 40% (namely HF with improved LVEF—HF<sub>imp</sub>EF). Indeed, DELIVER demonstrated no hint of therapeutic heterogeneity in the higher range of EF and in patients with HF<sub>imp</sub>EF, where dapagliflozin was as effective as it was at the lower end of the EF spectrum.<sup>9</sup>

Providing SGLT2 inhibition in HF patients not only reduces hard outcomes but also improves the quality of life. Indeed, a large meta-analysis of five trials (DAPA-HF, DELIVER, EMPEROR-Reduced, EMPEROR-Preserved, and SOLOIST-WHF) showed that more participants in the SGLT2 inhibitor groups than in placebo groups had clinically meaningful improvements and fewer participants had clinically meaningful deterioration in each of the three KCCQ summary scores by 8 months, without evidence of heterogeneity by trial.<sup>1</sup>

### SGLT2 inhibitors in patients with acute heart failure

Patients with acutely decompensated and acute heart failure (AHF) constitute a particularly vulnerable cohort since

they show a high post-discharge readmission and mortality rates,<sup>13</sup> together with limited evidence regarding outcome-modifying treatments.<sup>3</sup>

To date, SGLT2 inhibitors have been tested in AHF in two dedicated outcome trials: the SOLOIST-WHF and the EMPULSE trials (Figure 1). In the SOLOIST-WHF trial,<sup>5</sup> 1222 diabetic patients with an acute or recent episode of HF (49% of patients initiated therapy in-hospital and 51% within a median of 2 days after discharge) were randomized to the SGLT1/SGLT2 inhibitor sotagliflozin or placebo. After a median follow-up of 9 months, patients randomized to sotagliflozin showed a lower risk of total CV deaths, HF hospitalizations, or urgent HF visits than the placebo group (HR 0.67; 95% CI, 0.52-0.85).

Differently from SOLOIST-WHF trial, the EMPULSE trial<sup>14</sup> included AHF patients ( $n = 530$ ) with and without diabetes, randomized in-hospital (median 3 days after admission) to either empagliflozin or placebo, and continuing treatment during the 90 days post-discharge (i.e. 'vulnerable phase'). EMPULSE used a hierarchical clinical composite endpoint of all-cause death, number of HF events (HF hospitalizations, urgent visits for HF, and unplanned outpatient visits), time to first HF event, and health-related quality of life ( $\geq 5$ -point difference in change from baseline in Kansas City Cardiomyopathy Questionnaire Total Symptom Score at 90 days) using the win ratio (WR), a method that considers the order and clinical importance of the events, with death as the most important event.<sup>4</sup> The results showed that more patients treated with empagliflozin had a clinical benefit compared with placebo (stratified WR 1.36; 95% CI, 1.09-1.68), both in the acute *de novo* and decompensated chronic HF group and regardless of EF and diabetic status. The SOLOIST-WHF and the EMPULSE trials showed that SGLT2 inhibitors are also safe in the AHF setting without excess risk compared with a placebo.<sup>5,14</sup> Recently, a small clinical trial (EMPAG-HF)<sup>15</sup> demonstrated that the early addition of empagliflozin (25 mg daily) to standard diuretic therapy increases urine output (25% increase over 5 days vs. placebo) without affecting renal function in patients hospitalized for AHF. A phase 3 trial evaluating dapagliflozin in HF patients hospitalized for decompensated HF (DAPA ACT HF-TIMI 68; NCT04363697) is now ongoing and slated to report in 2023.<sup>16</sup>

### SGLT2 inhibitors in patients with heart failure and chronic kidneys disease

Chronic kidney disease (CKD) is one of the most common comorbidities in HF and is a powerful predictor of morbidity and mortality.<sup>17</sup> SGLT2 inhibitors reduce the risk of serious adverse renal outcomes in type 2 diabetes,<sup>18</sup> but the renal effects of these drugs in patients with HF remain uncertain. Empagliflozin and dapagliflozin have been reported—after an initial decrement (early dip) in the estimated glomerular filtration rate (eGFR) due to secondary changes in intraglomerular pressures (i.e. restoration of tubular-glomerular feedback)<sup>19</sup>—to slow the rate of decline in renal function (Table 1).<sup>7-9,11</sup> However, changes in the eGFR slope may not predict the effects of these drugs on major renal outcomes. A significant reduction of serious adverse renal events (i.e. profound and sustained decreases in eGFR or renal-replacement therapy) was demonstrated only in the EMPEROR-Reduced trial (empagliflozin vs. placebo: HR 0.50, 95% CI, 0.32-0.77),<sup>11</sup> while

in all other HF outcomes trials testing this hypothesis the effect on renal outcomes was modest (Table 1).

Pre-specified analysis of SGLT2 inhibitor trials demonstrated similar reductions in the primary outcome of cardiovascular death or hospitalization for HF across the spectrum of ejection fraction regardless of baseline eGFR and albuminuria.<sup>20-23</sup> DAPA-HF enrolled patients with  $eGFR \geq 30$  mL/min/1.73 m<sup>2</sup>,<sup>8</sup> while the eGFR threshold was  $\geq 25$  mL/min/1.73 m<sup>2</sup> in DELIVER<sup>9</sup> and  $\geq 20$  mL/min/1.73 m<sup>2</sup> in both EMPEROR-Reduced<sup>11</sup> and EMPEROR-Preserved<sup>7</sup> (in all trials eGFR was calculated by CKD-EPI formula). Based on the safety data from these studies, as well as on trials specifically targeting the CKD population (i.e. DAPA-CKD and EMPA-Kidney),<sup>24,25</sup> current labelling encourages treatment when eGFR is  $\geq 25$  mL/min/1.73 m<sup>2</sup> for dapagliflozin and  $\geq 20$  mL/min/1.73 m<sup>2</sup> for empagliflozin (Figure 2).

### Timing for initiation of SGLT2 inhibitors

Clinical practice guidelines recommend (class IA) an early initiation of dapagliflozin and empagliflozin together with the classical foundational therapy (betablockers, RASi/ARNI, and MRA) in HF patients in order to reduce mortality and HF hospitalizations.<sup>2,3</sup> For patients with HFmrEF and HFpEF, recent AHA guidelines recommend treatment with SGLT2 inhibitors in class IIA (the higher among the other outcome-modifying treatments),<sup>2</sup> basing their recommendation only on EMPEROR-Preserved data (but published data from DELIVER trial are anticipated to strengthen current treatment recommendations in this population). Both empagliflozin and dapagliflozin can be taken at the dose of 10 mg once daily, irrespective of food.

Recent evidence suggests that SGLT2 inhibition should be implemented as soon as possible. Indeed, in patients with HFpEF, a sustained statistically significant reduction in clinical events (primary endpoint) was observed by 28 days after randomization (HR 0.51; 95% CI, 0.28-0.94) in DAPA-HF,<sup>26</sup> and after only 12 days (HR 0.76; 95% CI, 0.67-0.87) in EMPEROR-Reduced.<sup>6</sup> Similarly, in HFmrEF/HFpEF patients, dapagliflozin reduced the risk for the primary endpoint within 2 weeks of treatment initiation (HR 0.45; 95% CI, 0.20-0.99),<sup>27</sup> while empagliflozin takes 18 days from randomization to induce a sustained reduction of clinical events (HR 0.41; 95% CI, 0.17-0.99).<sup>28</sup> Early SGLT2 inhibition is even more crucial in patients at higher risk, such as those with a recent episode of AHF. Indeed, in a post-hoc analysis of the DAPA-HF trial patients in closer proximity to a prior HF hospitalization experienced greater relative and absolute risk reductions of the composite of death and HF hospitalization when treated with dapagliflozin.<sup>26</sup>

Patients hospitalized for AHF without symptomatic hypotension or need for diuretic escalation, inotropic drugs or IV vasodilators within the previous 6 h may be eligible for SGLT2 inhibitors (Figure 2). In this setting, an early introduction (within the first 5 days) of these drugs has been demonstrated safe and effective in improving clinical outcomes.<sup>29</sup>

Taken together these results reinforce early sustained clinical benefits with SGLT2 inhibitors in patients with HF, underscoring the need for timely initiation of therapy. Moreover, SGLT2 inhibitors can be easily implemented in the complex HF therapy (single dose, low impact on blood

**Table 1** Effect of SGLT2 inhibitors on the change of renal function and kidney outcomes

	Mean baseline eGFR (mL/min/1.73 m <sup>2</sup> )	Mean absolute eGFR dip in the SGLT2 inhibitor group (mL/min/1.73 m <sup>2</sup> )	Mean slope of eGFR change (mL/min/1.73 m <sup>2</sup> per year)	Between-groups difference in eGFR slope (95% CI)	Renal adverse outcomes
DAPA-HF	66	-4.2	Dapagliflozin, -1.09 vs. placebo, -2.85	1.73 (1.10; 2.37) mL/min/1.73 m <sup>2</sup> per year; <i>P</i> < 0.001	HR 0.71 (95% CI, 0.44-1.16) <sup>a</sup>
EMPEROR-Reduced	62	-3.8	Empagliflozin, -0.55 vs. placebo, -2.28	1.73 (1.10; 2.37) mL/min/1.73 m <sup>2</sup> per year; <i>P</i> < 0.001	HR 0.50 (95% CI, 0.32-0.77) <sup>b</sup>
EMPEROR-Preserved	61	-	Empagliflozin, -1.25 vs. placebo, -2.62	1.36 (1.06; 1.66) mL/min/1.73 m <sup>2</sup> per year; <i>P</i> < 0.001	HR 0.95 (95% CI, 0.73-1.24) <sup>b</sup>
DELIVER	61	-3.7	Dapagliflozin, 1.0 vs. placebo, -1.5	1.4 (1.0; 1.8) mL/min/1.73 m <sup>2</sup> per year; <i>P</i> < 0.001	HR 1.08, (95% CI, 0.79-1.49) <sup>a</sup>
SOLOIST-WHF	50 (median)	-	Sotagliflozin, -0.34 vs. placebo, -0.18	-0.16 (-1.30; 0.98) mL/min/1.73 m <sup>2</sup> per year; <i>P</i> = NS	-
EMPULSE	52 (median)	-	-	-	Not possible to fit a model due to low event rate <sup>c</sup>






The dash stands for not reported or not investigated data.

Abbreviations: CI, confidence interval; eGFR, estimated glomerular filtration rate; NS, non-significant; HR, hazard ratio.

<sup>a</sup>Composite of sustained reduction of  $\geq 50\%$  in the eGFR for  $\geq 28$  days or sustained eGFR < 15 mL/min/1.73m<sup>2</sup> for  $\geq 28$  days or long-term dialysis treatment or renal transplantation or death from renal causes.

<sup>b</sup>Composite of  $\geq 40\%$  sustained decline eGFR or end-stage renal disease (defined as: 1) chronic dialysis; 2) renal transplantation; or 3) a sustained eGFR < 15 mL/min/1.73 m<sup>2</sup> [for patients with baseline eGFR  $\geq 30$ ] or sustained eGFR < 10 mL/min/1.73m<sup>2</sup> [for patients with baseline eGFR < 30 mL/min/1.73 m<sup>2</sup>]).

<sup>c</sup>Occurrence of chronic dialysis or renal transplant or sustained reduction of  $\geq 40\%$  eGFR or sustained eGFR < 15 mL/min/1.73 m<sup>2</sup> [for patients with baseline eGFR < 10 mL/min/1.73 m<sup>2</sup>] or sustained eGFR < 10 mL/min/1.73 m<sup>2</sup> [for patients with baseline eGFR < 30 mL/min/1.73 m<sup>2</sup>]).

<b>Eligible patients</b> <ul style="list-style-type: none"> <li>All symptomatic HF patients, regardless of LVEF, diabetic status and care setting</li> </ul>	
<b>Contraindications</b> <ul style="list-style-type: none"> <li>Type 1 diabetes mellitus or history of ketoacidosis</li> <li>Hypotension (caution if SBP &lt;100 mmHg)</li> <li>Severe CKD (dapagliflozin: eGFR &lt;25 ml/min/1.73m<sup>2</sup>; empagliflozin: eGFR &lt;20 ml/min/1.73m<sup>2</sup>)<sup>a</sup></li> <li>Pregnancy/risk of pregnancy and breastfeeding period</li> <li>Caution in patients with history of recurrent genital or urinary tract infections</li> <li>In AHF, use of inotropes within the last 24h or use of IV vasodilators or LD escalation within the last 6h</li> </ul>	
<b>Dose</b> <ul style="list-style-type: none"> <li>10 mg once daily for both dapagliflozin and empagliflozin (irrespective of food)</li> </ul>	
<b>Monitoring</b> <ul style="list-style-type: none"> <li>Check renal function when starting the therapy and then after 1-2 weeks<sup>a,b</sup></li> <li>Blood glucose (if SGLT2 inhibitors are used in association with anti-diabetic drugs – mainly insulin and insulin secretagogues)</li> <li>Acute illness or major surgery<sup>c</sup></li> </ul>	
<b>Patient/caregiver counselling</b> <ul style="list-style-type: none"> <li>Ensure adequate daily genital hygiene</li> <li>Watch for symptoms of volume depletion<sup>d</sup>, uro-genital infections<sup>e</sup> and diabetic ketoacidosis<sup>f</sup></li> <li>Avoid dehydration, low carbohydrate (ketogenic) diet and excessive alcohol consumption</li> </ul>	

**Figure 2** Practical guide to initiation of SGLT2 inhibitors in patients with heart failure. Abbreviations: AHF, acute heart failure; CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate; HF, heart failure; IV, intravenous; LD, loop diuretics; LVEF, left ventricular ejection fraction; SBP, systolic blood pressure. <sup>a</sup>eGFR calculated by CKD-EPI formula. <sup>b</sup>Initial decline in eGFR of 3-4 mL/min/1.73 m<sup>2</sup> or 10-15% from baseline (mainly in patients with initial low eGFR) is common/expected, does not reflex acute kidney injury and therapy should be continued unless major fall in eGFR. <sup>c</sup>FDA recommend to withhold SGLT2 inhibitors in case of prolonged fasting or 3 days before major surgery. The treatment can be restarted only once the patient's oral intake is restored and any other risk factors for ketoacidosis are resolved. <sup>d</sup>Symptoms of volume depletion are weakness, orthostatic hypotension, weight decrease >1 kg over 24 h or >2 kg in 1 week. <sup>e</sup>Symptoms of uro-genital infections are pain or burning on urination, redness, swelling, itching in the genital area, nasty-smelling vaginal or penile secretion, and fever. <sup>f</sup>Symptoms of diabetic ketoacidosis are excessive thirst, sweet-smelling breath, a change in urine or sweat odour, nausea, vomiting, abdominal pain, confusion, weakness, and fever. Note that it could also manifest with relatively normal glucose levels (euglycemic diabetic ketoacidosis).

pressure, no impact on heart rate) fitting almost all treatment strategies.<sup>30</sup> Finally, clinical inertia should be avoided considering the broad eligibility to these drugs,<sup>31,32</sup> and the consequent expected strong clinical benefit in the treated population.

### Management of concomitant treatments

Heart failure treatment and its comorbidities are met with significant polypharmacy, which is associated in turn with an increased risk of drug-drug interactions and lower patient compliance.

Subgroup analyses of the DAPA-HF and EMPEROR-Reduce trials demonstrated the efficacy of dapagliflozin and empagliflozin regardless of concomitant MRA or ARNi use,<sup>33-36</sup> supporting the combined use of SGLT2 inhibitors with these drugs. Interestingly, SGLT2 inhibitors showed to reduce the incidence of clinically meaningful hyperkalaemia in HF patients, possibly enabling the concomitant use or up-titration of renin-angiotensin-aldosterone system inhibitors to target doses.<sup>37</sup>

For stable patients with HF, loop diuretics dose should be adapted on longitudinal evaluation of clinical signs and symptoms of congestion. None of the SGLT2 inhibitor

trials had a routine protocolized adjustment of loop diuretics when initiating study therapy. In general, the prevalence of subjects who experienced volume depletion across the trials was low (i.e. ~7% in DAPA-HF and ~11% in EMPEROR-Reduced) and on average not significantly different compared with placebo.<sup>8,11</sup> Notably, in the DAPA-HF trial volume depletion was slightly more common with dapagliflozin than with placebo (8.0% vs. 4.9%) in the higher-dose diuretic group (≥40 mg of furosemide equivalent per day).<sup>38</sup> Accordingly, it is important to regularly assess the patient's volume status and correct hypovolemia-modulating loop diuretic doses as deemed appropriate. Patients and caregivers should be educated to recognize the warning symptoms of volume depletion (excessive reduction in body weight, low blood pressure, dizziness, etc.) (Figure 2).

SGLT2 inhibitors show a modest effect on blood pressure lowering across the published HF trials, with an average reduction of -1.32 (-2.19; -0.45) mmHg and -1.06 (-3.20; 1.08) mmHg in systolic and diastolic blood pressure, respectively.<sup>39</sup> Accordingly, SGLT2 inhibitors seem rather neutral from a blood pressure standpoint and their initiation should not raise any issue about other HF treatments impacting on blood pressure. Discontinuing these

drugs in patients with hypotension should probably be perceived as questionable, given that patients with low blood pressure are intrinsically more at risk of events and experience a strong benefit from SGLT2 inhibitors.<sup>40</sup>

Given the expected mild reduction in glycated haemoglobin (0.5–1.2%) and the low incidence of hypoglycaemia in patients with type 2 diabetes, the dose adjustment of glucose-lowering medications is usually not required. A notable exception is the concomitant therapy with insulin and insulin secretagogues—such as sulfonylureas—where the incidence of hypoglycaemic events slightly increases (although remaining rare). In these cases, a down-titration of insulin dose could be required after consulting a diabetologist.<sup>41</sup>

### Adverse effects and problem solving

Although SGLT2i are generally well tolerated, several adverse effects may limit patient compliance and the overall effectiveness of the medication (Figure 2).

The most common major safety concern of SGLT2 inhibition is the increased risk of genital mycotic infections. The incidence of genital infections is between 5% and 15% and is four to five times more frequent in women—as mycotic vulvovaginitis—than in men—who usually develop a mycotic balanitis.<sup>42</sup> A history of genital infection, obesity, and premenopausal state are all risk factors for the development of mycotic infections during SGLT2 inhibitors treatment.<sup>43</sup> The incidence of this complication is the highest in the first few months of treatment followed by an attenuation in frequency. The infections are mostly symptomatic and typically respond to topical or single-dose oral antifungal therapy, usually without the need for interruption of SGLT2 inhibitors.<sup>44</sup> An adequate genital hygiene may reduce the incidence of this complication.

SGLT2 inhibitors should be withheld in diabetic patients during periods of anticipated or unexpected poor oral intake, including the preparation for elective surgeries, due to the increased risk of diabetic ketoacidosis. Although uncommon, it is a potentially life-threatening condition that could manifest at relatively lower glucose levels (euglycemic diabetic ketoacidosis) and with vague symptoms (i.e. polyuria, abdominal pain, nausea/vomiting, and confusion).<sup>42</sup> The U.S. Food and Drug Administration released multiple warnings on this issue, recommending to discontinue SGLT2 inhibitors 3 days (4 days for ertugliflozin) before scheduled surgery and restarting the treatment only once the patient's oral intake is restored and any other risk factors for ketoacidosis are resolved.<sup>45</sup> Patients and their caregivers should be also counselled to avoid very low carbohydrates, ketogenic diets, and excessive alcohol consumption.

Osmotic diuresis with SGLT2 inhibitors may lead to orthostatic hypotension (frequency of 1.2–1.5%), especially among the elderly and in the first weeks of treatment. Prevention of hypovolemia ensuring an adequate hydration and modulating diuretic treatment might reduce the occurrence of this event.<sup>46</sup>

### Conclusions

Sodium-glucose co-transporter 2 inhibitors reduce the risk of clinical events in a broad range of patients with HF, irrespective of ejection fraction, diabetic status, or care

setting. Their clinical benefit has been demonstrated early and sustained, without any major safety concern. Therefore, unless contraindicated, SGLT2 inhibitors should be rapidly initiated as part of the foundational therapy in all patients with HF. Clinicians should be confident in handling these drugs, in order to implement their use and limit the potential adverse effects.

### Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

**Conflict of interest:** L.M. received speaker fees from AstraZeneca and Vifor Pharma. N.G. received honoraria from AstraZeneca, Bayer, Boehringer, Lilly, Novartis, Roche Diagnostics, and Vifor Pharma. All other authors declare no conflict of interest.

### Data availability

DAS: Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

### References

1. Vaduganathan M, Docherty KF, Claggett BL, Jhund PS, de Boer RA, Hernandez AF *et al.* SGLT-2 inhibitors in patients with heart failure: a comprehensive meta-analysis of five randomised controlled trials. *Lancet* 2022;**400**:757–767.
2. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM *et al.* 2022 AHA/ACC/HFSA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2022;**145**:e876–e894.
3. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumback A, Bohm M *et al.* 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;**42**:3599–3726.
4. Tromp J, Ponikowski P, Salsali A, Angermann CE, Biegus J, Blatchford J *et al.* Sodium-glucose co-transporter 2 inhibition in patients hospitalized for acute decompensated heart failure: rationale for and design of the EMPULSE trial. *Eur J Heart Fail* 2021;**23**:826–834.
5. Bhatt DL, Szarek M, Steg PG, Cannon CP, Leiter LA, McGuire DK *et al.* Sotagliflozin in patients with diabetes and recent worsening heart failure. *N Engl J Med* 2021;**384**:117–128.
6. Packer M, Anker SD, Butler J, Filippatos G, Ferreira JP, Pocock SJ *et al.* Effect of empagliflozin on the clinical stability of patients with heart failure and a reduced ejection fraction: the EMPEROR-Reduced trial. *Circulation* 2021;**143**:326–336.
7. Anker SD, Butler J, Filippatos G, Ferreira JP, Bocchi E, Bohm M *et al.* Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med* 2021;**385**:1451–1461.
8. McMurray JJV, Solomon SD, Inzucchi SE, Kober L, Kosiborod MN, Martinez FA *et al.* Dapagliflozin in patients with heart failure and reduced ejection fraction. *N Engl J Med* 2019;**381**:1995–2008.
9. Solomon SD, McMurray JJV, Claggett B, de Boer RA, DeMets D, Hernandez AF *et al.* Dapagliflozin in heart failure with mildly reduced or preserved ejection fraction. *N Engl J Med* 2022;**387**:1089–1098.
10. Tromp J, Ouwerkerk W, van Veldhuisen DJ, Hillege HL, Richards AM, van der Meer P *et al.* A systematic review and network meta-analysis of pharmacological treatment of heart failure with reduced ejection fraction. *JACC Heart Fail* 2022;**10**:73–84.
11. Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P *et al.* Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med* 2020;**383**:1413–1424.
12. Zannad F, Ferreira JP, Pocock SJ, Anker SD, Butler J, Filippatos G *et al.* SGLT2 inhibitors in patients with heart failure with reduced ejection fraction: a meta-analysis of the EMPEROR-Reduced and DAPA-HF trials. *Lancet* 2020;**396**:819–829.

13. Ambrosy AP, Fonarow GC, Butler J, Chioncel O, Greene SJ, Vaduganathan M *et al*. The global health and economic burden of hospitalizations for heart failure: lessons learned from hospitalized heart failure registries. *J Am Coll Cardiol* 2014;**63**:1123-1133.
14. Voors AA, Angermann CE, Teerlink JR, Collins SP, Kosiborod M, Biegus J *et al*. The SGLT2 inhibitor empagliflozin in patients hospitalized for acute heart failure: a multinational randomized trial. *Nat Med* 2022;**28**:568-574.
15. Schulze PC, Bogoviku J, Westphal J, Aftanski P, Haertel F, Grund S *et al*. Effects of early empagliflozin initiation on diuresis and kidney function in patients with acute decompensated heart failure (EMPAG-HF). *Circulation* 2022;**146**:289-298.
16. Dapagliflozin and Effect on Cardiovascular Events in Acute Heart Failure - Thrombolysis in Myocardial Infarction 68 (DAPA ACT HF-TIMI 68). <https://clinicaltrials.gov/ct2/show/NCT04363697> (18 November 2022).
17. Hillege HL, Nitsch D, Pfeffer MA, Swedberg K, McMurray JJ, Yusuf S *et al*. Candesartan in heart failure: assessment of reduction in M. Morbidity I. Renal function as a predictor of outcome in a broad spectrum of patients with heart failure. *Circulation* 2006;**113**:671-678.
18. Giorgino F, Vora J, Fenici P, Solini A. Renoprotection with SGLT2 inhibitors in type 2 diabetes over a spectrum of cardiovascular and renal risk. *Cardiovasc Diabetol* 2020;**19**:196.
19. Cice G, Calò L, Monzo L. Sodium-glucose co-transporter 2 inhibitors for the treatment of cardio-renal syndrome. *Eur Heart J Suppl* 2022;**24**:168-171.
20. Ferreira JP, Zannad F, Butler J, Filippatos G, Pocock SJ, Brueckmann M *et al*. Association of empagliflozin treatment with albuminuria levels in patients with heart failure: a secondary analysis of EMPEROR-Pooled. *JAMA Cardiol* 2022;**7**:1148-1159.
21. Jhund PS, Solomon SD, Docherty KF, Heerspink HJL, Anand IS, Bohm M *et al*. Efficacy of dapagliflozin on renal function and outcomes in patients with heart failure with reduced ejection fraction: results of DAPA-HF. *Circulation* 2021;**143**:298-309.
22. Zannad F, Ferreira JP, Pocock SJ, Zeller C, Anker SD, Butler J *et al*. Cardiac and kidney benefits of empagliflozin in heart failure across the spectrum of kidney function: insights from EMPEROR-Reduced. *Circulation* 2021;**143**:310-321.
23. Causland FR M, Claggett BL, Vaduganathan M, Desai AS, Jhund P, de Boer RA *et al*. Dapagliflozin and kidney outcomes in patients with heart failure with mildly reduced or preserved ejection fraction: a prespecified analysis of the DELIVER randomized clinical trial. *JAMA Cardiol* 2023;**8**:56-65.
24. Heerspink HJL, Stefansson BV, Correa-Rotter R, Chertow GM, Greene T, Hou FF *et al*. Dapagliflozin in patients with chronic kidney disease. *N Engl J Med* 2020;**383**:1436-1446.
25. The EMPA-KIDNEY Collaborative Group, Herrington WG, Staplin N, Wanner C, Green JB, Hauske SJ *et al*. Empagliflozin in patients with chronic kidney disease. *N Engl J Med* 2023;**388**:117-127.
26. Berg DD, Jhund PS, Docherty KF, Murphy SA, Verma S, Inzucchi SE *et al*. Time to clinical benefit of dapagliflozin and significance of prior heart failure hospitalization in patients with heart failure with reduced ejection fraction. *JAMA Cardiol* 2021;**6**:499-507.
27. Vaduganathan M, Claggett BL, Jhund P, de Boer RA, Hernandez AF, Inzucchi SE *et al*. Time to clinical benefit of dapagliflozin in patients with heart failure with mildly reduced or preserved ejection fraction: a prespecified secondary analysis of the DELIVER randomized clinical trial. *JAMA Cardiol* 2022;**7**:1259-1263.
28. Butler J, Siddiqi TJ, Filippatos G, Ferreira JP, Pocock SJ, Zannad F *et al*. Early benefit with empagliflozin in heart failure with preserved ejection fraction: insights from the EMPEROR-Preserved trial. *Eur J Heart Fail* 2022;**24**:245-248.
29. Ferreira JP, Pimenta J, Moura B, Aguiar C, Franco F. Use of sodium glucose co-transporter 2 inhibitors in acute heart failure: a practical guidance. *ESC Heart Fail* 2022;**9**:4344-4347.
30. Rosano GMC, Moura B, Metra M, Bohm M, Bauersachs J, Ben Gal T *et al*. Patient profiling in heart failure for tailoring medical therapy. A consensus document of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2021;**23**:872-881.
31. Monzo L, Ferrari I, Cicogna F, Tota C, Calò L. What proportion of patients with heart failure and preserved ejection fraction are eligible for empagliflozin? *J Cardiovasc Med (Hagerstown)* 2022;**23**:567-569.
32. Monzo L, Ferrari I, Cicogna F, Tota C, Calò L. Sodium-glucose co-transporter-2 inhibitors eligibility in patients with heart failure with reduced ejection fraction. *Int J Cardiol* 2021;**341**:56-59.
33. Solomon SD, Jhund PS, Claggett BL, Dewan P, Kober L, Kosiborod MN *et al*. Effect of dapagliflozin in patients with HFrEF treated with sacubitril/valsartan: the DAPA-HF trial. *JACC Heart Fail* 2020;**8**:811-818.
34. Packer M, Anker SD, Butler J, Filippatos G, Ferreira JP, Pocock SJ *et al*. Influence of neprilysin inhibition on the efficacy and safety of empagliflozin in patients with chronic heart failure and a reduced ejection fraction: the EMPEROR-Reduced trial. *Eur Heart J* 2021;**42**:671-680.
35. Shen L, Kristensen SL, Bengtsson O, Bohm M, de Boer RA, Docherty KF *et al*. Dapagliflozin in HFrEF patients treated with mineralocorticoid receptor antagonists: an analysis of DAPA-HF. *JACC Heart Fail* 2021;**9**:254-264.
36. Ferreira JP, Butler J, Zannad F, Filippatos G, Schueler E, Steubl D *et al*. Mineralocorticoid receptor antagonists and empagliflozin in patients with heart failure and preserved ejection fraction. *J Am Coll Cardiol* 2022;**79**:1129-1137.
37. Ferreira JP, Zannad F, Butler J, Filippatos G, Ritter I, Schuler E *et al*. Empagliflozin and serum potassium in heart failure: an analysis from EMPEROR-Pooled. *Eur Heart J* 2022;**43**:2984-2993.
38. Jackson AM, Dewan P, Anand IS, Belohlavek J, Bengtsson O, de Boer RA *et al*. Dapagliflozin and diuretic use in patients with heart failure and reduced ejection fraction in DAPA-HF. *Circulation* 2020;**142**:1040-1054.
39. Li M, Yi T, Fan F, Qiu L, Wang Z, Weng H *et al*. Effect of sodium-glucose cotransporter-2 inhibitors on blood pressure in patients with heart failure: a systematic review and meta-analysis. *Cardiovasc Diabetol* 2022;**21**:139.
40. Bohm M, Anker SD, Butler J, Filippatos G, Ferreira JP, Pocock SJ *et al*. Empagliflozin improves cardiovascular and renal outcomes in heart failure irrespective of systolic blood pressure. *J Am Coll Cardiol* 2021;**78**:1337-1348.
41. Tsushima Y, Lansang MC, Makin V. The role of SGLT-2 inhibitors in managing type 2 diabetes. *Cleve Clin J Med* 2021;**88**:47-58.
42. Halimi S, Verges B. Adverse effects and safety of SGLT-2 inhibitors. *Diabetes Metab* 2014;**40**:S28-S34.
43. Engelhardt K, Ferguson M, Rosselli JL. Prevention and management of genital mycotic infections in the setting of sodium-glucose cotransporter 2 inhibitors. *Ann Pharmacother* 2021;**55**:543-548.
44. Johnsson KM, Ptaszynska A, Schmitz B, Sugg J, Parikh SJ, List JF. Vulvovaginitis and balanitis in patients with diabetes treated with dapagliflozin. *J Diabetes Complications* 2013;**27**:479-484.
45. FDA revises labels of SGLT2 inhibitors for diabetes to include warnings about too much acid in the blood and serious urinary tract infections (fda.gov). 2022. <https://www.fda.gov/drugs/drug-safety-and-availability/fda-revises-labels-sglt2-inhibitors-diabetes-include-warnings-about-too-much-acid-blood-and-serious>.
46. Vardeny O, Vaduganathan M. Practical guide to prescribing sodium-glucose cotransporter 2 inhibitors for cardiologists. *JACC Heart Fail* 2019;**7**:169-172.