



**HAL**  
open science

## Defining myocardial infarction in trials of people receiving hemodialysis: consensus report from the SONG-HD MI Expert Working group

Emma O'loné, Fred Apple, James Burton, Fergus Caskey, Jonathan Craig, Christopher Defilippi, Derek Forfang, Karen Hicks, Vivekanand Jha, Kenneth Mahaffey, et al.

### ► To cite this version:

Emma O'loné, Fred Apple, James Burton, Fergus Caskey, Jonathan Craig, et al.. Defining myocardial infarction in trials of people receiving hemodialysis: consensus report from the SONG-HD MI Expert Working group. *Kidney International*, 2023, 103 (6), pp.1028-1037. 10.1016/j.kint.2023.02.033 . hal-04094804

**HAL Id: hal-04094804**

**<https://hal.univ-lorraine.fr/hal-04094804v1>**

Submitted on 11 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Defining Myocardial Infarction in trials of people receiving hemodialysis: consensus report from the SONG-HD MI Expert Working group**

Brief title: Defining MI in patients on hemodialysis.

O'Lone E<sup>1</sup>, Apple FS<sup>2</sup>, Burton JO<sup>3</sup>, Caskey FJ<sup>4</sup>, Craig JC<sup>5</sup>, de Filippi CR<sup>6</sup>, Forfang D<sup>7</sup>, Hicks KA<sup>8§</sup>, Jha V<sup>9,10,11</sup>, Mahaffey KW<sup>12</sup>, Mark PB<sup>13</sup>, Rossignol P<sup>14,15</sup>, Scholes-Robertson N<sup>5</sup>, Jaure A<sup>1,16</sup>, Viecelli AK<sup>17</sup>, Wang AY<sup>18</sup>, Wheeler DC<sup>19</sup>, White D<sup>20</sup>, Winkelmayr WC<sup>21</sup>, Herzog CA<sup>22, 23</sup>

1. The University of Sydney, Camperdown, Sydney, Australia
2. Departments of Laboratory Medicine and Pathology, Hennepin Healthcare/Hennepin County Medical Center and University of Minnesota, Minneapolis, Minnesota
3. Department of Cardiovascular Sciences, University of Leicester and NIHR Leicester Cardiovascular Biomedical Research Unit, Glenfield Hospital Leicester, Leicester, UK
4. Population Health Sciences, University of Bristol, Southmead Hospital, Bristol, UK
5. College of Medicine and Public Health, Flinders University, Adelaide, South Australia, Australia
6. Inova Heart and Vascular Institute, Falls Church, VA, USA
7. The National Forum of ESRD Networks, Kidney Patient Advisory Council (KPAC) WI USA
8. Division of Cardiology and Nephrology, Office of Cardiology, Hematology, Endocrinology, and Nephrology, Center for Drug Evaluation and Research (CDER), United States Food and Drug Administration, Silver Spring, Maryland, USA
9. George Institute of Global Health, UNSW, New Delhi, India
10. School of Public Health, Imperial College, London, UK
11. Prasanna School of Public Health, Manipal Academy of Higher Education, Manipal, India
12. The Stanford Center for Clinical Research, Department of Medicine, Stanford University School of Medicine, Stanford, CA
13. University of Glasgow, Institute of Cardiovascular and Medical Sciences, Glasgow, UK
14. Université de Lorraine, Centre d'Investigation Clinique Plurithématique 1433 -INSERM-CHRU de Nancy, Inserm U1116 & FCRIN INI-CRCT (Cardiovascular and Renal Clinical Trialists), Vandoeuvre-les-Nancy, France
15. Medical specialties and nephrology -hemodialysis departments, Princess Grace Hospital, and Monaco Private Hemodialysis Centre, Monaco, Monaco
16. Centre for Kidney Research, Children's Hospital at Westmead, Westmead, NSW, Australia
17. Department of Nephrology, Princess Alexandra Hospital, Brisbane, Australia
18. Queen Mary Hospital, The University of Hong Kong, Hong Kong, China
19. University College London, London, United Kingdom
20. American Association of Kidney Patients, Tampa, Florida
21. Selzman Institute for Kidney Health, Section of Nephrology, Department of Medicine, Baylor College of Medicine, Houston, Texas

22. Chronic Disease Research Group, Hennepin Healthcare Research Institute, Minneapolis, Minnesota
23. Division of Cardiology, Department of Medicine, Hennepin Healthcare and University of Minnesota, Minneapolis, Minnesota

§This publication reflects the views of the author and should not be construed to represent FDA's views or policies.

Manuscript word count: 4330

Funding: This work was supported by the National Health and Medical Research Council Australia (NHMRC; 1098815).

Corresponding author: Dr Emma O'Lone, Centre for Kidney Research, Centre for Kidney Research Locked Bag 4001

The Children's Hospital at Westmead, Sydney NSW 2006 Australia. Email:

[Eolo0909@uni.sydney.edu.au](mailto:Eolo0909@uni.sydney.edu.au) Twitter handle: @emma\_olone. [Tel:+61298451469](tel:+61298451469), Fax: +61 2 9845 1491

Abstract:

Cardiovascular disease is the leading cause of death in patients receiving hemodialysis. Currently there is no standardized definition of myocardial infarction (MI) for patients receiving hemodialysis. Through an international consensus process MI was established as the core CVD measure for this population in clinical trials. The Standardised Outcomes in Nephrology Group – Hemodialysis (SONG-HD) initiative convened a multidisciplinary, international working group to address the definition of MI in this population. Based on current evidence, the working group recommends using the 4th Universal Definition of MI with specific caveats with regard to the interpretation of “ischemic symptoms” and performing a baseline 12-lead electrocardiogram to facilitate interpretation of acute changes on subsequent tracings. The working group do not recommend obtaining baseline cardiac troponin values, nor do we recommend obtaining serial cardiac biomarkers in settings where ischemia is suspected. Application of an evidence-based uniform definition should increase the reliability and accuracy of trial results.

Key words: hemodialysis, myocardial infarction, outcome, definition, trials, recommendations

One sentence summary. An international expert working group support the use of the 4<sup>th</sup> Universal definition of myocardial infarction for use in trials in people requiring hemodialysis.

## INTRODUCTION

Cardiovascular disease (CVD) is the leading cause of death in patients with kidney failure requiring replacement therapy<sup>1</sup>. The incidence of myocardial infarction (MI) in patients receiving hemodialysis is at least 4-times higher than in the general population and is associated with substantially poorer outcomes<sup>1-3</sup>. One-year mortality after a MI in patients receiving hemodialysis is also 60% compared to less than 10% in the general population<sup>4-7</sup>.

The higher prevalence of MI in the hemodialysis population is multifactorial. Traditional cardiovascular (CV) risk factors, including hypertension and diabetes, are more common. In addition, there are risk factors that are unique to patients receiving hemodialysis, including dysregulation of bone and mineral metabolism leading to increased vascular calcification and uremic toxins. Dialysis also results in rapid hemodynamic changes, heightened inflammation, endothelial and immune dysfunction<sup>8-11</sup>.

Patients receiving hemodialysis are usually excluded from large-scale CV interventional trials<sup>12</sup> but when included the most frequently measured and reported CV outcomes are surrogate endpoints such as serum biomarkers which may be of uncertain clinical significance and of little relevance to patients<sup>13, 14</sup>. Composite CV outcomes in this population are frequently used to achieve adequate statistical power however, across different trials use different components (e.g. stroke, MI, heart failure) to form each composite outcome<sup>13</sup>. MI, which in a recent international survey including patients and clinicians has been shown to be of the highest importance to patients receiving hemodialysis<sup>15</sup>, is frequently a component of a CV composite endpoint but is defined inconsistently<sup>13, 16</sup>. A review of five large CV trials in patients receiving hemodialysis demonstrated four different definitions of MI (Table 1) and different adjudication processes. Applying an alternative definition or adjudication process to each trial potentially lead a to

clinically significant difference in number of reported events. Although not specific to trials in this population, the lack of a standardized and validated definition for MI in patients receiving hemodialysis and the heterogeneous measurement and reporting of MI limit the comparison of interventions across trials.

The Standardised Outcomes in Nephrology (SONG) Initiative, a National Health and Medical Research Council (NHMRC) funded project, has established core outcome sets across the spectrum of kidney disease since 2014. A core outcome set is an agreed standardized set of outcomes that should be measured and reported, as a minimum, in all clinical trials in specific areas of health<sup>17</sup>. The core outcomes are based on priorities defined by patients, caregivers, and health professionals. Through the SONG-HD (hemodialysis) consensus process involving over 1500 patients, caregivers, and health professionals from more than 70 countries, CVD was identified as a core outcome, with MI established as the core outcome measure<sup>15, 18</sup>. To use MI as a core outcome measure in trials involving patients receiving hemodialysis, consensus on a uniform definition for MI in this population is needed.

The SONG-HD initiative convened an international working group of experts at a roundtable meeting in Washington, D.C. on November 8, 2019. The expert working group was formed using purposive sampling to represent a broad range of countries and experience. It included cardiologists, nephrologists, clinical trialists, a clinical chemist, representatives from regulatory bodies and registries, and patients with experience of hemodialysis, to recommend a uniform definition of MI for use in clinical trials evaluating patients receiving hemodialysis as well as potential for use in clinical practice. We began by considering whether the 4<sup>th</sup> Universal Definition of Myocardial Infarction (UDMI)<sup>19</sup>, a standard definition formulated for the general population, was applicable to patients receiving hemodialysis. This definition has not been specifically validated in the dialysis population although it is felt to be applicable to all patients with

the caveat that there may be a greater percentage with chronic myocardial injury. There remain several limitations to the definition in people receiving hemodialysis. Specific considerations raised in the SONG-HD CVD consensus workshop regarding criteria required for an appropriate definition of MI informed the deliberations<sup>18</sup>. These considerations included consistency, applicability and specificity of the definition to patients receiving hemodialysis, the importance of the type of MI, variability in MI symptoms in patients receiving hemodialysis, and the uncertainty in the clinical utility of biomarkers specific to hemodialysis. This report summarizes the meeting discussion and the resulting recommendations.

## THE 4<sup>TH</sup> UNIVERSAL DEFINITION OF MYOCARDIAL INFARCTION

In 2018 the European Society of Cardiology (ESC)/ American College of Cardiology (ACC)/the American Heart Association (AHA)and the World Heart Federation (WHF) jointly published an expert consensus document designated the Fourth UDMI<sup>19</sup> (Box 1). This definition was based on studies of MI in the general population. Although the document briefly discusses myocardial injury/infarction in patients with chronic kidney disease (CKD), including those with kidney failure, the publication does not address patients receiving hemodialysis specifically. Box 2 summarizes the limitations of the 4<sup>th</sup> UDMI and its applicability to Type 1 MI in people with kidney failure receiving hemodialysis.

## TYPES OF MI

The Fourth UDMI classifies MI into five types. Criteria for the types of MI as laid out by the 4<sup>th</sup> UDMI are summarised in Box 1. The criteria required for diagnosing type 1 and 2 MIs in people requiring hemodialysis are outlined below. In brief, MI is defined as follows: (1) a rise and/or fall in

cTn with at least 1 value >99<sup>th</sup> percentile URL in patients with an initial cTn ≤99<sup>th</sup> percentile URL; and (2) a >20% rise and/or fall in cTn in patients with an initial cTn >99<sup>th</sup> percentile URL.

Patients receiving hemodialysis have both a high prevalence of acute myocardial injury and are predisposed to acute MI, due to underlying pathophysiological features. Diagnosis of a type 2 MI requires consideration of both the context and mechanism leading to the imbalance of oxygen supply and demand. This is of particular importance to patients receiving hemodialysis, a process which has been shown to have significant hemodynamic effects which increase myocardial oxygen demand<sup>20</sup>. Hemodialysis also induces significant global and segmental reductions in myocardial blood flow<sup>21</sup>. Underlying pathophysiological changes related to kidney failure, including left ventricular hypertrophy and associated capillary/myocyte mismatch, reduced peripheral arterial compliance, endothelial dysfunction, anemia, and reduced coronary flow reserve, predispose patients requiring hemodialysis to demand ischemia. The prevalence of type 2 MIs is high in the hemodialysis population<sup>69</sup>. Estimating prevalence is difficult because differentiation between MI types 1 and 2 often requires expert adjudication in large clinical cohorts and ideally includes coronary angiography to definitively exclude coronary thrombosis<sup>22</sup>.

To establish appropriate treatment according to current guidelines<sup>27,28</sup>, it is important to classify MI into ST-elevation MI (STEMI) or non-ST-elevation MI (NSTEMI). In the setting of a STEMI, primary percutaneous coronary intervention or thrombolytic therapy may be required and in the setting of a NSTEMI, coronary angiography may be indicated. Further classification into a Type 1 or 2 MI is not as well defined and there are limited quantitative data on the efficacy of coronary angiography in differentiating a type 1 from type 2 MI. Short and long-term mortality rates for type 2 MI are higher than for type 1 MI, although it is unclear whether there are differences in attributable cause-specific mortality<sup>23-26</sup>. Treatment of type 2 MI largely consists of addressing the underlying supply and demand imbalance noninvasively. In the case of hemodialysis, reducing



“demand” may be possible through reducing ultrafiltration rates but there are often patient or centre based limitations to moving away from standard short intermittent dialysis sessions. Even in the general population long-term treatment strategies for type 2 MI in the absence of coronary artery disease lack trial data or guidelines <sup>29</sup>. Differentiating type 1 from type 2 MIs is challenging for adjudication experts as well as clinicians. Recent published guidelines advocate that in the absence of a clear alternative cause, the initial working diagnosis for most patients with evidence of acute myocardial injury and signs and symptoms consistent with ischemia should prompt classification and management according to established guidelines for type 1 MI <sup>29</sup>. Although treatment based on the type of MI may differ, the working group considered MI, regardless of type, to be the most important outcome captured in clinical trials. Evidence-based protocols for the management of type 2 MI remain limited.

**Differentiating types 1 and 2 MI can be challenging. Improving our ability to recognize and treat MIs is a priority for future research.**

**MI should be a core outcome reported in clinical trials involving people with kidney failure receiving hemodialysis and a uniform definition should be used.**

#### CRITERIA USED TO DEFINE TYPES 1, 2 AND 3 MI

##### 1. Ischemic symptoms

Studies indicate that patients with CKD, and particularly those receiving hemodialysis, often do not describe typical symptoms of MI. The classic triad of chest discomfort, arm/jaw pain and sweating is experienced by less than 50% of patients with CKD <sup>30</sup>. In patients receiving

hemodialysis, the most common “ischemic symptom” is shortness of breath, experienced by nearly 50% of patients receiving dialysis<sup>30,31</sup>. Chest pain or discomfort associated with MI is experienced by less than 20% of patients receiving dialysis compared to over 35% of patients with normal kidney function<sup>31</sup>. Patient receiving dialysis described a background level of pain and discomfort and felt that non-specific symptoms or a change in sensation or degree of unwellness should also raise suspicion as an “ischemic symptom.” It is important to note that highlighting more non-specific ischemic symptoms may result in increased diagnosis of myocardial injury as well as MI<sup>32</sup>.

**In a patient receiving hemodialysis, atypical symptoms or any changes in symptoms should raise a high index of suspicion for a potential MI and prompt further investigation and treatment.**

## 2. ELECTROCARDIOGRAM

Fluid and electrolyte changes during hemodialysis affect ECG waveforms. The removal of fluid over the course of a dialysis session has been shown to augment the P wave as well as the QRS amplitude and duration<sup>33</sup>. Similarly, electrolyte shifts during dialysis have been shown to affect the P wave, QRS, and QTc<sup>33,34</sup>. Whether an ECG is acquired during, before or after dialysis should be considered in its interpretation; persistent changes such as left bundle-branch-block (BBB) are unlikely to be influenced by variations in dialysis.

Patients receiving hemodialysis often have abnormal baseline ECGs making it difficult to determine acute change. In one series approximately 30% of such patients were found to have electrical conduction abnormalities including left and right BBB on a baseline ECG. ST elevation occurs in less than 20% of patients with an MI on dialysis compared to over 35% of patients who are not on dialysis<sup>35</sup>. A non-specific ECG change is the most common finding in patients presenting with MI on hemodialysis<sup>35</sup>. Patients with CKD are significantly less likely to develop a pathological Q wave than patients without CKD (19% compared to 34%)<sup>30</sup>. It is important that non-specific ECG findings are taken in the context of the clinical presentation and in combination with troponin findings to help rule-in an MI though may not help ruling out MI.

Although patients with CKD, and particularly patients receiving hemodialysis, have a higher prevalence of silent MI compared to the general population<sup>36,37</sup>, there is no consensus regarding whether routine (annual or more frequent) collection of ECGs in this population confer incremental value in the absence of other clinical or biochemical abnormalities. Hence, for the purposes of clinical trials, the working group recommends obtaining ECGs in association with acute events but recommends where relevant discussing with regulators prior to study conduct, to consider whether

baseline ECG and any subsequent changes consistent with silent MI events should be considered as an MI in a trial.

Patients indicated that an additional ECG at baseline “would not be a burden because it is non-invasive” and clinicians thought a baseline ECG would better inform trial endpoint definitions and patient care.

**A baseline ECG in all patients receiving hemodialysis when stable and asymptomatic may aid in the interpretation of acute ECG changes on subsequent tracings in the setting of acute clinical symptoms or biomarker changes suggestive of MI. On trial entry, we suggest performing a single baseline ECG on each patient. We also recommend obtaining serial ECGs during an acute event followed by another ECG when the patient is clinically stable.**

### 3. Cardiac Troponin

Troponin is a complex of three regulatory proteins (troponin I, C and T). During myocardial injury cTnI and cTnT are released as individual subunits as well as non-covalent ternary and binary complexes<sup>38,39</sup>. cTnT and cTnI are now the preferred biomarkers of myocardial injury. Troponin is a biomarker for myocardial injury; the 4<sup>th</sup> UDMI includes clinical evidence of acute myocardial ischemia with detection of a rise and/or fall of cTn values with at least one value above the 99th percentile upper reference limit (URL) plus additional criteria as summarized in Box 1.

#### *Assay variability*

High sensitivity cardiac troponin (hs-cTn) assays need to meet two criteria: a coefficient of variation (total imprecision) of  $\leq 10\%$  at the 99th percentile URL for both men and women and

measurable concentrations below the 99th percentile URL need to be detectable above the assay's limit of detection for  $\geq 50\%$  of healthy individuals in the population of interest<sup>40-42</sup>. Sex-specific 99<sup>th</sup> percentile URLs for hs-cTn assays have been derived and validated in the general population based on healthy individuals<sup>42</sup>. It is not possible to achieve disease specific thresholds due to the heterogeneity within each specific disease population.

hs-Tn assays are now in widespread use and accurately and precisely measure five to 100-fold lower concentrations of cTn in blood than older, contemporary assays<sup>43,44</sup>. The various hs-Tn assays use monoclonal antibodies to a number of different epitopes along the cTnT or cTnI protein<sup>45</sup>. On account of the large variations in epitopes targeted by the different antibodies used in each assay, it is not possible to standardize URLs across assays.

#### *Biological variability*

Biological variability describes the fluctuation of biomarker levels around a homeostatic set-point in healthy individuals or those with stable disease and which is of no clinical significance. Biological variability is low, with intra-patient coefficients of variation quoted as 7.9% for weekly measurements and 12.6% for monthly measurements. Although, there may be increased biological variability when using hs-cTn<sup>63</sup>, over the course of a year this biological variability is minimal and if an acute event occurs, cTn subsequently returns to the individual patient's baseline<sup>47</sup>. Inter-patient variability is high in patients on hemodialysis<sup>46</sup>.

#### *Effect of dialysis on cardiac troponin:*

To date only relatively small studies have been conducted to evaluate the effect of dialysis on cTn. The troponin complex (52 kDa), as well as the subunits (cTnI is 24 kDa and cTnT is

37 kDa) are classified as middle molecules by size. Older dialyzer membranes predominantly filtered out small, water-soluble molecules and not the “middle molecules”, now thought to be the cause of much of the historical morbidity and mortality. New synthetic membranes and the increase in convective therapies have improved the clearance of a number of these molecules. There remains a lack of consensus regarding the effect of dialysis on cTnI and cTnT levels however there is some evidence that cTnI is adsorbed onto the surface of the dialyzer membrane<sup>49-51</sup>. High flux dialyzers may increase cTn clearance more than low flux dialyzers<sup>52,53</sup> and clearance is potentially increased further still with hemodiafiltration<sup>53</sup>. The increasing use of nocturnal dialysis and prolonged hours dialysis may have significant effect on both the production and handling of troponin. To date, changes in levels of both cTnT and cTnI are relatively small and there is insufficient evidence at this time to suggest that the effect of standard dialysis on troponin is significant enough to alter the diagnosis of MI in patients receiving hemodialysis.

#### *Elevated baseline troponin*

Levels of cTnT and cTnI over the sex specific 99<sup>th</sup> percentile URL have been demonstrated in up to 80% for cTnT but < 20% for cTnI of patients requiring hemodialysis<sup>46, 54, 55</sup>. Reduced kidney clearance is not the main driver of an elevated cTn in this population<sup>56</sup>. The exact etiology/mechanism of elevated baseline cTn, or why cTnT remains increased longer than cTnI, is not entirely clear but is likely to be multifactorial including increased instability of the cardiac myocyte membrane, microinfarctions, and myocardial necrosis as well as increased left ventricular hypertrophy and heart failure causing myocyte strain and apoptotic cell death<sup>57-61</sup>.

In the absence of an acute event, elevated baseline cTn in patients receiving hemodialysis is a strong predictor of adverse outcomes. It has been shown that increased cTnT and cTnI are both predictive of CV and all-cause mortality in ESKD<sup>47, 62-65</sup>. Identifying risk has an effect on the

individual patient and the cost to healthcare. There is currently insufficient evidence to suggest a pathway in response to the identified increased risk.

Therefore, although a determinant of risk, an historical baseline troponin should not contribute to the diagnosis of acute MI and we do not support the concept of acquiring a cTnI or cTnT on entry into a trial in HD dialysis patients. We acknowledge that a baseline level of cTnI or cTnT in an individual patient is a useful tool in identifying chronic myocardial injury and that the collection of a baseline troponin may offer future opportunities for further biomarker investigation and clarification of the role of troponin. However, in the assessment of a symptomatic patient presenting to the hospital, serial troponin monitoring should be used for ruling in or ruling out an MI.

**Currently we do not suggest collecting a baseline (trial entry) cTn in a trial setting.**

**Performing baseline cTn in stable, asymptomatic patients receiving hemodialysis may identify patients at increased risk for adverse outcome but with no currently proven treatment to reduce risk, resulting in unwanted concerns for patients without contributing to the diagnosis of acute MI.**

## *Delta Troponin*

The US National Academy of Clinical Biochemistry (NACB) (redesignated as, Academy of the American Association of Clinical Chemistry [AACC]) recommended a  $\delta$  in standard assays for cTn of  $> 50\%$  if cTn is less than the 99<sup>th</sup> percentile URL and  $\geq 20\%$  once values are elevated above the 99<sup>th</sup> percentile URL. These values are calculated to distinguish a true change from one that could be attributed to biological variability alone and yet maintain sensitivity<sup>44</sup>.

The clinical sensitivity of hs-cTn assays to detect myocardial injury remain high in patients on hemodialysis, reported as 100%<sup>31</sup> however high prevalence of elevated baseline cTn can reduce specificity to as low as 40%. Hence, in a hemodialysis patient with an initial cTn value greater than the 99<sup>th</sup> percentile URL, a rise and/or fall of more than 20% is suggestive of an acute MI and should be included in the endpoint definitions in a trial setting. Short-term intra-patient biological and analytical variability is minimal in the absence of an acute event, however, a one to two hour follow up sample after initial testing may not be sufficient to rule out MI. Patients presenting early after an MI are unlikely to be missed using a two-hour sample but for patients who present with atypical symptoms, it may be harder to know where on the cTn kinetic curve (Figure 1) they are at a given time point. Any dynamic change in cTn or strong clinical suspicion should prompt further cTn samples so as not to miss a significant delta. This may require samples to be taken after up to 6 hours to ensure an MI is not missed.

**Current evidence suggests that, in the context of an initial cTn above the sex specific 99<sup>th</sup> percentile URL, a  $\delta$  cTn  $> 20\%$  in cTn in addition to the clinical criteria should be an**



**accepted rise and/or fall to diagnose acute MI in the patients requiring hemodialysis. An early rule-out sample may not be sufficient to exclude MI in patients requiring hemodialysis.**

## SUMMARY

The (SONG-HD) Initiative convened an expert working group to discuss MI definitions most appropriate for use in clinical trials enrolling patients requiring hemodialysis. Although all definitions have limitations, the working group recommends using the 4<sup>th</sup> UDMI and suggests that trialists and clinicians maintain a broad interpretation of “ischemic symptoms” in this population. The working group also recommends obtaining a baseline ECG to aid in the interpretation of acute changes noted on subsequent tracings. Serial cardiac biomarkers, optimally measured by hs-cTn assays, are needed to evaluate potential events of myocardial ischemia and myocardial injury. In addition, a greater than 20% delta in cTn is required to define an acute MI event in patients on hemodialysis with a baseline (at time of presentation) cTn that exceeds 99<sup>th</sup> percentile URL. Our future work will include validation of the use of the 4<sup>th</sup> UDMI in patients receiving hemodialysis. Box 3 summarizes the recommendations for clinical end point committee adjudication criteria for diagnosis of myocardial infarction in trials including people receiving hemodialysis. Supplementary material 1 outlines the recommended elements to incorporate into a Case Report Form for suspected or confirmed acute myocardial infarction events in trials including people receiving hemodialysis.

The working group has identified a number of directions for future research. At this time there are limited data in the hemodialysis population to create evidence-based guidelines. We recommend evaluating diagnostic methods for MI and MI type in patients receiving hemodialysis as well as improving the prevention and treatment of type 2 MI. We recommend investigating the reported lower percentage of STEMIs in this population and whether this is due to fewer acute coronary occlusions noted on angiography or confounders related to baseline ECG abnormalities. We recommend determining whether these findings impact reperfusion therapy in this population.

Consistent definitions and standardized reporting should improve trial quality, reproducibility, and comparability which will assist in endeavours to improve outcomes for this high-risk population.



1 Disclosures:

2 EO, JC, DF, KH, KS, NS, AT, AV, DW, WW have no disclosures.

3 FA is on the Board of Directors for HyTest Ltd, is the Associate Editor of Clinical Chemistry. He  
4 is also on Advisory Boards at: Instrumentation Laboratory, Siemens Healthcare, and Qurvo. He has  
5 received Honorariums for Speaking at Industry Sponsored Conferences for Siemens Healthcare and  
6 Abbott Diagnostics. He is the PI on Industry Funded Grants (non-salaried) on cardiac biomarkers  
7 through hospital research institute (HHRI) with Abbott Diagnostics, Abbott POC, BD, Beckman  
8 Coulter, Ortho-Clinical Diagnostics, Roche Diagnostics, Siemens Healthcare, ET Healthcare and  
9 Quidel.

10 JB has a research grant from Yakult (Honsha) to investigate the effect of probiotics on markers of  
11 gut permeability and cardiovascular risk in HD patients and have received speaker honoraria from  
12 NAPP, BMS and Pharmacosmos on the topics of diabetic kidney disease, cardiovascular risk and  
13 anaemia.

14 FC receives honoraria from Baxter. Research funding from NIHR and Kidney Research UK

15 C de F receives grants to my institution and advisory board from Roche diagnostics: Ortho  
16 diagnostics: consulting and endpoint education, Siemens Healthineers: consulting, FujiRebio:  
17 consulting and Up-to-date: royalties.

18 VJ reports personal fees from GSK, personal fees from Bayer, personal fees from Boehringer  
19 Ingelheim, personal fees from Baxter Healthcare, grants from NephroPlus, grants and personal fees  
20 from Zydus Cadilla, outside the submitted work; under the policy of all payments being made to the  
21 organization

22 KM has received research support and/or consulting fees from the following companies: Abbott,  
23 Afferent, Amgen, Anthos, Apple, Inc, AstraZeneca, Baim Institute, Boehringer Ingelheim, Cardiva  
24 Medical, Inc, CSL Behring, Elsevier, Ferring, Google (Verily), Intermountain Health, Johnson &

1 Johnson, Luitpold, Medscape, Medtronic, Merck, Mount Sinai, Mundi Pharma, Myokardia, NIH,  
2 Novartis, Novo Nordisk, Portola, Regeneron, Sanofi, SmartMedics, St. Jude, Theravance.

3 PM reports personal fees and non-financial support from Vifor, personal fees from Astrazeneca,  
4 grants from Boehringer Ingelheim, personal fees and non-financial support from Pharmacosmos,  
5 personal fees from Janssen, personal fees from Novartis, personal fees from Pfizer, personal fees  
6 from Bristol Myers Squibb, personal fees and non-financial support from Napp, outside the  
7 submitted work.

8 PR receives consulting fees and travel support from Novartis, consulting fees from Novo Nordisk,  
9 AstraZeneca, Grünenthal, and Corvidia, consulting fees, lecture fees, fees for serving on a steering  
10 committee, and travel support from Relypsa/Vifor/Vifor Fresenius Medical Care, fees for serving  
11 on a steering committee and fees for serving on a critical event committee from Idorsia, lecture fees  
12 and travel support from Bayer and Servier, owning stock options in G3 Pharmaceuticals, and fees  
13 for serving as co-founder and owning stock in CardioRenal.

14 AW has received speaker honorarium from Sanofi Renal and Fresenius Kabi, and research grants  
15 from Sanofi Renal and Baxter Corporation.

16 DCW has received honoraria and/or consultancy fees from Amgen, AstraZeneca, Astellas, Bayer,  
17 Boehringer Ingelheim, GlaxoSmithKline, Janssen, Merck Sharpe and Dohme, Mundipharma, Napp,  
18 Reata, Tricida, and Vifor Fresenius

19 CH reports grants from NIDDK/NIH (USRDS Contract), during the conduct of the study; personal  
20 fees from Abbvie, grants and personal fees from Amgen, personal fees from AstraZeneca, personal  
21 fees from Corvidia, personal fees from Diamedica, personal fees from FibroGen, personal fees from  
22 Janssen, personal fees from NxStage, personal fees from Pfizer, grants and personal fees from  
23 Relypsa, personal fees from Sanifit, personal fees from University of Oxford, grants from Bristol-  
24 Myers Squibb, grants from University of British Columbia, personal fees from UpToDate, other

1 from Boston Scientific, other from General Electric, other from Johnson & Johnson, personal fees  
2 and other from Merck, other from Hennepin Healthcare, grants and personal fees from National  
3 Heart Lung and Blood Institute (NHLBI/NIH), outside the submitted work; .

4

5 Acknowledgements:

6 Kimberly Smith and Aliza Thompson from the Division of Cardiology and Nephrology, Office of  
7 Cardiology, Hematology, Endocrinology, and Nephrology, Center for Drug Evaluation and  
8 Research (CDER), United States Food and Drug Administration who attended and contributed to  
9 the consensus workshop. David Charytan who reviewed the manuscript.

10

1   References

- 2   1.     Herzog CA. Sudden cardiac death and acute myocardial infarction in dialysis patients:  
3   perspectives of a cardiologist. *Semin Nephrol* 2005;**25**(6):363-6.
- 4   2.     Iseki K, Fukiyama K. Long-term prognosis and incidence of acute myocardial infarction in  
5   patients on chronic hemodialysis. The Okinawa Dialysis Study Group. *Am J Kidney Dis*  
6   2000;**36**(4):820-5.
- 7   3.     O'Lone E, Kelly PJ, Masson P, Kotwal S, Gallagher M, Cass A, Craig JC, Webster AC.  
8   Incidence of Ischaemic Heart Disease in Men and Women With End-Stage Kidney Disease: A  
9   Cohort Study. *Heart Lung Circ* 2020;**29**(10):1517-1526.
- 10  4.     Charytan D, Mauri L, Agarwal A, Servoss S, Scirica B, Kuntz RE. The use of invasive  
11  cardiac procedures after acute myocardial infarction in long-term dialysis patients. *Am Heart J*  
12  2006;**152**(3):558-64.
- 13  5.     Herzog CA, Ma JZ, Collins AJ. Poor long-term survival after acute myocardial infarction  
14  among patients on long-term dialysis. *N Engl J Med* 1998;**339**(12):799-805.
- 15  6.     Montalescot G, Dallongeville J, Van Belle E, Rouanet S, Baulac C, Degrandart A, Vicaut  
16  E, Investigators O. STEMI and NSTEMI: are they so different? 1 year outcomes in acute  
17  myocardial infarction as defined by the ESC/ACC definition (the OPERA registry). *Eur Heart J*  
18  2007;**28**(12):1409-17.
- 19  7.     Armstrong PW, Fu Y, Chang WC, Topol EJ, Granger CB, Betriu A, Van de Werf F, Lee  
20  KL, Califf RM. Acute coronary syndromes in the GUSTO-IIb trial: prognostic insights and impact  
21  of recurrent ischemia. The GUSTO-IIb Investigators. *Circulation* 1998;**98**(18):1860-8.
- 22  8.     Horl WH, Cohen JJ, Harrington JT, Madias NE, Zusman CJ. Atherosclerosis and uremic  
23  retention solutes. *Kidney Int* 2004;**66**(4):1719-31.



- 1 9. Rostand SG. Coronary heart disease in chronic renal insufficiency: some management  
2 considerations. *J Am Soc Nephrol* 2000;**11**(10):1948-56.
- 3 10. London GM, Guerin AP, Marchais SJ, Metivier F, Pannier B, Adda H. Arterial media  
4 calcification in end-stage renal disease: impact on all-cause and cardiovascular mortality. *Nephrol  
5 Dial Transplant* 2003;**18**(9):1731-40.
- 6 11. Goodman WG, Goldin J, Kuizon BD, Yoon C, Gales B, Sider D, Wang Y, Chung J,  
7 Emerick A, Greaser L, Elashoff RM, Salusky IB. Coronary-artery calcification in young adults with  
8 end-stage renal disease who are undergoing dialysis. *N Engl J Med* 2000;**342**(20):1478-83.
- 9 12. Charytan D, Kuntz RE. The exclusion of patients with chronic kidney disease from clinical  
10 trials in coronary artery disease. *Kidney Int* 2006;**70**(11):2021-30.
- 11 13. O'Lone E, Viecelli AK, Craig JC, Tong A, Sautenet B, Roy D, Herrington WG, Herzog CA,  
12 Jafar T, Jardine M, Krane V, Levin A, Malyszko J, Rocco MV, Strippoli G, Tonelli M, Wang  
13 AYM, Wanner C, Zannad F, Winkelmayer WC, Webster AC, Wheeler DC. Cardiovascular  
14 Outcomes Reported in Hemodialysis Trials. *J Am Coll Cardiol* 2018;**71**(24):2802-2810.
- 15 14. Yudkin JS, Lipska KJ, Montori VM. The idolatry of the surrogate. *BMJ* 2011;**343**:d7995.
- 16 15. O'Lone E, Howell M, Viecelli AK, Craig JC, Tong A, Sautenet B, Herrington WG, Herzog  
17 CA, Jafar TH, Jardine M, Krane V, Levin A, Malyszko J, Rocco MV, Strippoli G, Tonelli M, Wang  
18 AY, Wanner C, Zannad F, Winkelmayer WC, Wheeler DC. Identifying critically important  
19 cardiovascular outcomes for trials in hemodialysis: an international survey with patients, caregivers  
20 and health professionals. *Nephrol Dial Transplant* 2020;**35**(10):1761-1769.
- 21 16. Cholesterol Treatment Trialists C, Herrington WG, Emberson J, Mihaylova B, Blackwell L,  
22 Reith C, Solbu MD, Mark PB, Fellstrom B, Jardine AG, Wanner C, Holdaas H, Fulcher J, Haynes  
23 R, Landray MJ, Keech A, Simes J, Collins R, Baigent C. Impact of renal function on the effects of

- 1 LDL cholesterol lowering with statin-based regimens: a meta-analysis of individual participant data  
2 from 28 randomised trials. *Lancet Diabetes Endocrinol* 2016;**4**(10):829-39.
- 3 17. Williamson PR, Altman DG, Bagley H, Barnes KL, Blazeby JM, Brookes ST, Clarke M,  
4 Gargon E, Gorst S, Harman N, Kirkham JJ, McNair A, Prinsen CAC, Schmitt J, Terwee CB, Young  
5 B. The COMET Handbook: version 1.0. *Trials* 2017;**18**(Suppl 3):280.
- 6 18. O'Lone E, Viecelli AK, Craig JC, Tong A, Sautenet B, Herrington WG, Herzog CA, Jafar  
7 TH, Jardine M, Krane V, Levin A, Malyszko J, Rocco MV, Strippoli G, Tonelli M, Wang AYM,  
8 Wanner C, Zannad F, Winkelmayr WC, Wheeler DC, Investigators S-HCCW. Establishing Core  
9 Cardiovascular Outcome Measures for Trials in Hemodialysis: Report of an International  
10 Consensus Workshop. *Am J Kidney Dis* 2020;**76**(1):109-120.
- 11 19. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, White HD, Executive  
12 Group on behalf of the Joint European Society of Cardiology /American College of Cardiology  
13 /American Heart Association /World Heart Federation Task Force for the Universal Definition of  
14 Myocardial I. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol*  
15 2018;**72**(18):2231-2264.
- 16 20. Bos WJ, Bruin S, van Olden RW, Keur I, Wesseling KH, Westerhof N, Krediet RT, Arisz  
17 LA. Cardiac and hemodynamic effects of hemodialysis and ultrafiltration. *Am J Kidney Dis*  
18 2000;**35**(5):819-26.
- 19 21. McIntyre CW, Burton JO, Selby NM, Leccisotti L, Korsheed S, Baker CS, Camici PG.  
20 Hemodialysis-induced cardiac dysfunction is associated with an acute reduction in global and  
21 segmental myocardial blood flow. *Clin J Am Soc Nephrol* 2008;**3**(1):19-26.
- 22 22. Shroff GR, Li S, Herzog CA. Trends in Discharge Claims for Acute Myocardial Infarction  
23 among Patients on Dialysis. *J Am Soc Nephrol* 2017;**28**(5):1379-1383.

- 1 23. Sandoval Y, Thordsen SE, Smith SW, Schulz KM, Murakami MM, Pearce LA, Apple FS.  
2 Cardiac troponin changes to distinguish type 1 and type 2 myocardial infarction and 180-day  
3 mortality risk. *Eur Heart J Acute Cardiovasc Care* 2014;**3**(4):317-25.
- 4 24. Saaby L, Poulsen TS, Diederichsen AC, Hosbond S, Larsen TB, Schmidt H, Gerke O,  
5 Hallas J, Thygesen K, Mickley H. Mortality rate in type 2 myocardial infarction: observations from  
6 an unselected hospital cohort. *Am J Med* 2014;**127**(4):295-302.
- 7 25. Bonaca MP, Wiviott SD, Braunwald E, Murphy SA, Ruff CT, Antman EM, Morrow DA.  
8 American College of Cardiology/American Heart Association/European Society of  
9 Cardiology/World Heart Federation universal definition of myocardial infarction classification  
10 system and the risk of cardiovascular death: observations from the TRITON-TIMI 38 trial (Trial to  
11 Assess Improvement in Therapeutic Outcomes by Optimizing Platelet Inhibition With Prasugrel-  
12 Thrombolysis in Myocardial Infarction 38). *Circulation* 2012;**125**(4):577-83.
- 13 26. Sandoval Y, Smith SW, Sexter A, Thordsen SE, Bruen CA, Carlson MD, Dodd KW, Driver  
14 BE, Hu Y, Jacoby K, Johnson BK, Love SA, Moore JC, Schulz K, Scott NL, Apple FS. Type 1 and  
15 2 Myocardial Infarction and Myocardial Injury: Clinical Transition to High-Sensitivity Cardiac  
16 Troponin I. *Am J Med* 2017;**130**(12):1431-1439 e4.
- 17 27. Amsterdam EA, Wenger NK, Brindis RG, Casey DE, Jr., Ganiats TG, Holmes DR, Jr., Jaffe  
18 AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine  
19 MS, Smalling RW, Zieman SJ. 2014 AHA/ACC Guideline for the Management of Patients with  
20 Non-ST-Elevation Acute Coronary Syndromes: a report of the American College of  
21 Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*  
22 2014;**64**(24):e139-e228.
- 23 28. O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Jr., Chung MK, de Lemos JA, Ettinger  
24 SM, Fang JC, Fesmire FM, Franklin BA, Granger CB, Krumholz HM, Linderbaum JA, Morrow

- 1 DA, Newby LK, Ornato JP, Ou N, Radford MJ, Tamis-Holland JE, Tommaso JE, Tracy CM, Woo  
2 YJ, Zhao DX, Force CAT. 2013 ACCF/AHA guideline for the management of ST-elevation  
3 myocardial infarction: executive summary: a report of the American College of Cardiology  
4 Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*  
5 2013;**127**(4):529-55.
- 6 29. DeFilippis AP, Chapman AR, Mills NL, de Lemos JA, Arbab-Zadeh A, Newby LK,  
7 Morrow DA. Assessment and Treatment of Patients With Type 2 Myocardial Infarction and Acute  
8 Nonischemic Myocardial Injury. *Circulation* 2019;**140**(20):1661-1678.
- 9 30. Sosnov J, Lessard D, Goldberg RJ, Yarzebski J, Gore JM. Differential symptoms of acute  
10 myocardial infarction in patients with kidney disease: a community-wide perspective. *Am J Kidney*  
11 *Dis* 2006;**47**(3):378-84.
- 12 31. Gunsolus I, Sandoval Y, Smith SW, Sexter A, Schulz K, Herzog CA, Apple FS. Renal  
13 Dysfunction Influences the Diagnostic and Prognostic Performance of High-Sensitivity Cardiac  
14 Troponin I. *J Am Soc Nephrol* 2018;**29**(2):636-643.
- 15 32. Chapman AR, Sandoval Y. Type 2 Myocardial Infarction: Evolving Approaches to  
16 Diagnosis and Risk-Stratification. *Clin Chem* 2021;**67**(1):61-69.
- 17 33. Poulikakos D, Malik M. Challenges of ECG monitoring and ECG interpretation in dialysis  
18 units. *J Electrocardiol* 2016;**49**(6):855-859.
- 19 34. Morris ST, Galiatsou E, Stewart GA, Rodger RS, Jardine AG. QT dispersion before and  
20 after hemodialysis. *J Am Soc Nephrol* 1999;**10**(1):160-3.
- 21 35. Herzog CA, Littrell K, Arko C, Frederick PD, Blaney M. Clinical characteristics of dialysis  
22 patients with acute myocardial infarction in the United States: a collaborative project of the United  
23 States Renal Data System and the National Registry of Myocardial Infarction. *Circulation*  
24 2007;**116**(13):1465-72.

- 1 36. Rizk DV, Gutierrez O, Levitan EB, McClellan WM, Safford M, Soliman EZ, Warnock DG,  
2 Muntner P. Prevalence and prognosis of unrecognized myocardial infarctions in chronic kidney  
3 disease. *Nephrol Dial Transplant* 2012;**27**(9):3482-8.
- 4 37. Ohtake T, Kobayashi S, Moriya H, Negishi K, Okamoto K, Maesato K, Saito S. High  
5 prevalence of occult coronary artery stenosis in patients with chronic kidney disease at the initiation  
6 of renal replacement therapy: an angiographic examination. *J Am Soc Nephrol* 2005;**16**(4):1141-8.
- 7 38. Park KC, Gaze DC, Collinson PO, Marber MS. Cardiac troponins: from myocardial  
8 infarction to chronic disease. *Cardiovasc Res* 2017;**113**(14):1708-1718.
- 9 39. Szczykowska J, Hryszko T, Naumnik B. Cardiac troponins in chronic kidney disease  
10 patients with special emphasis on their importance in acute coronary syndrome. *Adv Med Sci*  
11 2019;**64**(1):131-136.
- 12 40. Jaffe AS, Ravkilde J, Roberts R, Naslund U, Apple FS, Galvani M, Katus H. It's time for a  
13 change to a troponin standard. *Circulation* 2000;**102**(11):1216-20.
- 14 41. Apple FS, Collinson PO, Biomarkers ITFoCAoC. Analytical characteristics of high-  
15 sensitivity cardiac troponin assays. *Clin Chem* 2012;**58**(1):54-61.
- 16 42. Apple FS, Wu AHB, Sandoval Y, Sexter A, Love SA, Myers G, Schulz K, Duh SH,  
17 Christenson RH. Sex-Specific 99th Percentile Upper Reference Limits for High Sensitivity Cardiac  
18 Troponin Assays Derived Using a Universal Sample Bank. *Clin Chem* 2020;**66**(3):434-444.
- 19 43. Apple FS. A new season for cardiac troponin assays: it's time to keep a scorecard. *Clin*  
20 *Chem* 2009;**55**(7):1303-6.
- 21 44. Vasile VC, Jaffe AS. High-Sensitivity Cardiac Troponin for the Diagnosis of Patients with  
22 Acute Coronary Syndromes. *Curr Cardiol Rep* 2017;**19**(10):92.

- 1 45. Apple FS, Sandoval Y, Jaffe AS, Ordonez-Llanos J, Bio-Markers ITFoCAoC. Cardiac  
2 Troponin Assays: Guide to Understanding Analytical Characteristics and Their Impact on Clinical  
3 Care. Clin Chem 2017;**63**(1):73-81.
- 4 46. Fahim MA, Hayen AD, Horvath AR, Dimeski G, Coburn A, Tan KS, Johnson DW, Craig  
5 JC, Campbell SB, Hawley CM. Biological variation of high sensitivity cardiac troponin-T in stable  
6 dialysis patients: implications for clinical practice. Clin Chem Lab Med 2015;**53**(5):715-22.
- 7 47. Hassan HC, Howlin K, Jefferys A, Spicer ST, Aravindan AN, Suryanarayanan G, Hall BM,  
8 Cleland BD, Wong JK, Suranyi MG, Makris A. High-sensitivity troponin as a predictor of cardiac  
9 events and mortality in the stable dialysis population. Clin Chem 2014;**60**(2):389-98.
- 10 49. Wayand D, Baum H, Schatzle G, Scharf J, Neumeier D. Cardiac troponin T and I in end-  
11 stage renal failure. Clin Chem 2000;**46**(9):1345-50.
- 12 50. Donnino MW, Karriem-Norwood V, Rivers EP, Gupta A, Nguyen HB, Jacobsen G,  
13 McCord J, Tomlanovich MC. Prevalence of elevated troponin I in end-stage renal disease patients  
14 receiving hemodialysis. Acad Emerg Med 2004;**11**(9):979-81.
- 15 51. Gaze DC, Collinson PO. Cardiac troponin I but not cardiac troponin T adheres to  
16 polysulfone dialyser membranes in an in vitro hemodialysis model: explanation for lower serum  
17 cTnI concentrations following dialysis. Open Heart 2014;**1**(1):e000108.
- 18 52. Lippi G, Tessitore N, Montagnana M, Salvagno GL, Lupo A, Guidi GC. Influence of  
19 sampling time and ultrafiltration coefficient of the dialysis membrane on cardiac troponin I and T.  
20 Arch Pathol Lab Med 2008;**132**(1):72-6.
- 21 53. Laveborn E, Lindmark K, Skagerlind M, Stegmayr B. NT-proBNP and troponin T levels  
22 differ after hemodialysis with a low versus high flux membrane. Int J Artif Organs 2015;**38**(2):69-  
23 75.

- 1 54. van Berkel M, Dekker MJE, Bogers H, Geerse DA, Konings C, Scharnhorst V. Diagnosis of  
2 acute myocardial infarction in hemodialysis patients may be feasible by comparing variation of  
3 cardiac troponins during acute presentation to baseline variation. *Clin Chim Acta* 2016;**456**:36-41.
- 4 55. Huang HL, Zhu S, Wang WQ, Nie X, Shi YY, He Y, Song HL, Miao Q, Fu P, Wang LL, Li  
5 GX. Diagnosis of Acute Myocardial Infarction in Hemodialysis Patients With High-Sensitivity  
6 Cardiac Troponin T Assay. *Arch Pathol Lab Med* 2016;**140**(1):75-80.
- 7 56. van der Linden N, Cornelis T, Kimenai DM, Klinkenberg LJJ, Hilderink JM, Luck S,  
8 Litjens EJR, Peeters F, Streng AS, Breidhardt T, van Loon LJC, Bekers O, Kooman JP,  
9 Westermark PO, Mueller C, Meex SJR. Origin of Cardiac Troponin T Elevations in Chronic  
10 Kidney Disease. *Circulation* 2017;**136**(11):1073-1075.
- 11 57. Abbas NA, John RI, Webb MC, Kempson ME, Potter AN, Price CP, Vickery S, Lamb EJ.  
12 Cardiac troponins and renal function in nondialysis patients with chronic kidney disease. *Clin Chem*  
13 2005;**51**(11):2059-66.
- 14 58. Masson S, Anand I, Favero C, Barlera S, Vago T, Bertocchi F, Maggioni AP, Tavazzi L,  
15 Tognoni G, Cohn JN, Latini R, Valsartan Heart Failure T, Gruppo Italiano per lo Studio della  
16 Sopravvivenza nell'Insufficienza Cardiaca-Heart Failure I. Serial measurement of cardiac troponin  
17 T using a highly sensitive assay in patients with chronic heart failure: data from 2 large randomized  
18 clinical trials. *Circulation* 2012;**125**(2):280-8.
- 19 59. Hamwi SM, Sharma AK, Weissman NJ, Goldstein SA, Apple S, Canos DA, Pinnow EE,  
20 Lindsay J. Troponin-I elevation in patients with increased left ventricular mass. *Am J Cardiol*  
21 2003;**92**(1):88-90.
- 22 60. Di Lullo L, Gorini A, Russo D, Santoboni A, Ronco C. Left Ventricular Hypertrophy in  
23 Chronic Kidney Disease Patients: From Pathophysiology to Treatment. *Cardiorenal Med*  
24 2015;**5**(4):254-66.

- 1 61. Ooi DS, Isotalo PA, Veinot JP. Correlation of antemortem serum creatine kinase, creatine  
2 kinase-MB, troponin I, and troponin T with cardiac pathology. *Clin Chem* 2000;**46**(3):338-44.
- 3 62. Apple FS, Murakami MM, Pearce LA, Herzog CA. Predictive value of cardiac troponin I  
4 and T for subsequent death in end-stage renal disease. *Circulation* 2002;**106**(23):2941-5.
- 5 63. Sandoval Y, Herzog CA, Love SA, Cao J, Hu Y, Wu AH, Gilbertson D, Brunelli SM,  
6 Young A, Ler R, Apple FS. Prognostic Value of Serial Changes in High-Sensitivity Cardiac  
7 Troponin I and T over 3 Months Using Reference Change Values in Hemodialysis Patients. *Clin*  
8 *Chem* 2016;**62**(4):631-8.
- 9 64. Deegan PB, Lafferty ME, Blumsohn A, Henderson IS, McGregor E. Prognostic value of  
10 troponin T in hemodialysis patients is independent of comorbidity. *Kidney Int* 2001;**60**(6):2399-  
11 405.
- 12 65. Khan NA, Hemmelgarn BR, Tonelli M, Thompson CR, Levin A. Prognostic value of  
13 troponin T and I among asymptomatic patients with end-stage renal disease: a meta-analysis.  
14 *Circulation* 2005;**112**(20):3088-96.
- 15 66. Willeit P, Welsh P, Evans JDW, Tschiderer L, Boachie C, Jukema JW, Ford I, Trompet S,  
16 Stott DJ, Kearney PM, Mooijaart SP, Kiechl S, Di Angelantonio E, Sattar N. High-Sensitivity  
17 Cardiac Troponin Concentration and Risk of First-Ever Cardiovascular Outcomes in 154,052  
18 Participants. *J Am Coll Cardiol* 2017;**70**(5):558-568.
- 19 67. Sandoval Y, Smith SW, Love SA, Sexter A, Schulz K, Apple FS. Single High-Sensitivity  
20 Cardiac Troponin I to Rule Out Acute Myocardial Infarction. *Am J Med* 2017;**130**(9):1076-1083  
21 e1.
- 22 68. Shah AS, Anand A, Sandoval Y, Lee KK, Smith SW, Adamson PD, Chapman AR, Langdon  
23 T, Sandeman D, Vaswani A, Strachan FE, Ferry A, Stirzaker AG, Reid A, Gray AJ, Collinson PO,  
24 McAllister DA, Apple FS, Newby DE, Mills NL, High Si. High-sensitivity cardiac troponin I at



1 presentation in patients with suspected acute coronary syndrome: a cohort study. *Lancet*  
2 2015;**386**(10012):2481-8.

3 69. Petrie MC, Jhund PS, Connolly E, Mark PB, MacDonald MR, Robertson M, Anker SD,  
4 Bhandari S, Farrington K, Kalra PA, Wheeler DC, Tomson CRV, Ford I, McMurray JJV,  
5 Macdougall IC; PIVOTAL Investigators and Committees. High-dose intravenous iron reduces  
6 myocardial infarction in patients on hemodialysis. *Cardiovasc Res.* 2021 Dec 7:cvab317. Epub  
7 ahead of print. PMID: 34875022.

8

9

1 **TABLES AND FIGURES**

2

3 **BOX 1 4<sup>TH</sup> UNIVERSAL DEFINITION OF MYOCARDIAL INFARCTION, TYPE 1 MI**

4

5 **BOX 2 SUMMARY OF THE LIMITATIONS OF THE 4<sup>TH</sup> UNIVERSAL DEFINITION IN DIAGNOSING TYPE**  
6 **1 MI IN PATIENTS RECEIVING HEMODIALYSIS THERAPY**

7

8 **BOX 3 RECOMMENDATIONS FOR CLINICAL END POINT COMMITTEE ADJUDICATION CRITERIA FOR**  
9 **DIAGNOSIS OF MYOCARDIAL INFARCTION IN TRIALS INCLUDING PEOPLE RECEIVING**  
10 **HEMODIALYSIS.**

11

12

13 **TABLE 1 COMPARISON OF DEFINITIONS FOR MYOCARDIAL INFARCTION USED IN LARGE**  
14 **CARDIOVASCULAR TRIALS IN PATIENTS RECEIVING HEMODIALYSIS**

15 **FIGURE 1 EARLY CARDIAC TROPONIN KINETICS IN PATIENTS AFTER ACUTE MYOCARDIAL**  
16 **INJURY INCLUDING ACUTE MYOCARDIAL INFARCTION**

17

1 Box

Type 1	<p>Detection of a rise and/or fall of cTn values with at least one value above the sex-specific 99th percentile URL, and at least one of the following:</p> <ul style="list-style-type: none"> <li>○ Symptoms of acute myocardial ischemia;</li> <li>○ New ischemic ECG changes;</li> <li>○ Development of pathological Q waves;</li> <li>○ Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality in a pattern consistent with an ischemic etiology;</li> <li>○ Identification of a coronary thrombus by angiography including intracoronary imaging or by autopsy</li> </ul>
Type 2	<p>Detection of a rise and/or fall of cardiac cTn values with at least one value above the sex-specific 99th percentile URL and evidence of an imbalance between myocardial oxygen supply and demand unrelated to acute coronary atherothrombosis in addition to the symptoms, ECG findings, and imaging criteria discussed above for a type 1 MI</p>
Type 3	<p>Patients who suffer a cardiac death, with symptoms suggestive of myocardial ischemia accompanied by presumed new ischemic ECG changes or ventricular fibrillation, but die before blood samples for biomarkers (preferably cTn) can be obtained, or before increases in cardiac biomarkers can be identified, or in whom MI is detected by autopsy examination</p>
Type 4	<p>Cardiac procedural myocardial injury is arbitrarily defined by increases of cTn values (&gt;99th percentile URL) in patients with normal baseline values (=99th percentile URL) or a rise of cTn values &gt;20% of the baseline value when it is above the 99th percentile URL but it is stable or falling. Type 4a is =48 hours PCI, type 4b PCI-related MI is stent/scaffold thrombosis, type 4c is restenosis post balloon angioplasty.</p>
Type 5	<p>CABG-related MI is arbitrarily defined as elevation of cTn values &gt;10 times the 99th percentile URL in patients with normal baseline cTn values. In patients with elevated preprocedure cTn in whom cTn levels are stable (=20% variation) or falling, the postprocedure cTn must rise by &gt;20%. However, the absolute postprocedural value still must be &gt;10 times the 99th percentile URL. In addition, 1 of the following elements is required:</p> <ul style="list-style-type: none"> <li>• Development of new pathological Q waves*;</li> <li>• Angiographic documented new graft occlusion or new native coronary artery occlusion;</li> <li>• Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality in a pattern consistent with an ischemic etiology.</li> </ul>

cTn – cardiac troponin, URL – upper reference limit, ECG – electrocardiogram, MI – myocardial infarction, PCI – percutaneous coronary intervention

2

3

1 Box 2

<b>4<sup>th</sup> Universal Definition: Criteria for Type 1 Myocardial Infarction</b>		<b>Limitation in patients receiving hemodialysis</b>
<b>cTn &gt;99<sup>th</sup> percentile URL</b>		In a stable population of patients receiving hemodialysis, 50% to 90% of hs-cTnT concentrations are above the 99th percentile URL compared to 5-25% for hs-cTnI assays (46,54,55)
<b>Plus one of these criteria</b>	Symptoms of acute myocardial ischemia	Typical ischemic symptoms are >50% less likely in patients receiving dialysis; <20% of these patients present with chest discomfort (30, 31)
	New ischemic ECG changes	About 30% of patients receiving hemodialysis have conduction abnormalities at baseline (35). >40% of patients receiving kidney replacement therapy present with non-specific changes and <20% present with typical ST elevation. (35)
	Development of pathological Q waves	>5% of patients receiving hemodialysis already have Q waves on baseline ECG and are less likely to develop a Q wave MI (30, 35)
	Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality in a pattern consistent with an ischemic etiology	
	Identification of a coronary thrombus by angiography including intracoronary imaging or by post mortem examination	<10% of patients receiving hemodialysis with MI undergo reperfusion/angiography

Hs-cTnT – high sensitivity cardiac troponin T, URL – Upper reference limit, ECG - electrocardiogram, cTn – cardiac troponin



### Recommendations for Clinical end point committee adjudication criteria for diagnosis of myocardial infarction in trials including people receiving haemodialysis.

Acute Myocardial infarction in people receiving haemodialysis is a syndrome defined as acute myocardial injury plus clinical evidence of acute myocardial ischemia (MI). An acute myocardial infarction may or may not result in death (fatal or non-fatal MI). Acute MI may be further sub classified by type as defined by the Fourth Universal Definition of Myocardial Infarction (2018)(4<sup>th</sup> UDMI).

Acute non-fatal myocardial infarction criteria Types 1 and 2 (adapted from 4<sup>th</sup> UDMI):

	Criteria	Notes
Required	Cardiac enzyme	<ul style="list-style-type: none"> <li>If index cTn is &lt;99<sup>th</sup> percentile URL: Detection of a rise and/or fall of cTn values with at least 1 value above the 99th percentile URL</li> <li>If index cTn is &gt;99<sup>th</sup> percentile URL: Detection of a rise and/or fall of cTn values with at least 1 value above the 99th percentile URL AND a rise or fall of &gt;20% cTn</li> </ul>
Plus one or more of these additional criteria	1. Clinical presentation	Symptoms suspicious of acute myocardial ischemia including: <ul style="list-style-type: none"> <li>Chest pain/discomfort</li> <li>Sweating</li> <li>Arm/throat/neck/jaw pain or discomfort</li> <li>Shortness of breath</li> <li>Non-specific pain, discomfort or nausea, over and above or different to, usual background pain, discomfort or nausea.</li> </ul>
	2. ECG  Comparison should be made to trial entry ECG or ECG performed prior to event when patient was stable and asymptomatic.	<p><b>New</b> ischemic ECG changes or <b>new</b> development of pathological Q waves:</p> <ul style="list-style-type: none"> <li>New ST-elevation at the J-point in 2 contiguous leads with the cut-point: =1 mm in all leads other than leads V2–V3 where the following cutpoints apply: =2 mm in men =40 years; =2.5 mm in men &lt;40 years, or =1.5 mm in women regardless of age.</li> <li>New horizontal or downsloping ST-depression =0.5 mm in 2 contiguous leads and/or T inversion &gt;1 mm in 2 contiguous leads with prominent R wave or R/S ratio &gt;1.</li> <li>In patients with pre-existing LBBB, ST-segment elevation =1 mm concordant with the QRS complex in any lead may be an indicator of acute myocardial ischemia.</li> <li>Presumed new LBBB or RBBB that is not rate-related may be an indicator of acute myocardial ischaemia</li> <li>New Q wave development: ECG Changes Associated With Prior Myocardial Infarction (In the Absence of Left Ventricular Hypertrophy and LBBB) Any Q wave in leads V2–V3 &gt;0.02 s or QS complex in leads V2–V3. Q wave ?0.03 s and ?1 mm deep or QS complex in leads I, II, aVL, aVF or V4–V6 in any 2 leads of a contiguous lead grouping (I, aVL; V1–V6; II, III, aVF).* R wave &gt;0.04 s in V1–V2 and R/S &gt;1 with a concordant positive T wave in absence of conduction defect. *The same criteria are used for supplemental leads V7–V9. s indicates seconds.</li> </ul>
	3. Imaging	Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality in a pattern consistent with an ischemic etiology
	4. Acute thrombus	Identification of a coronary thrombus by angiography including intracoronary imaging or by post mortem examination

URL-upper reference limit, ECG–electrocardiogram, LBBB- left bundle branch block, RBBB- right bundle branch block

1 Figure 1




Figure taken from Thygesen et al. J Am Coll Cardiol 2018;72(18):2231-64 (with permission).

2

3