



A class of fractional parabolic reaction-diffusion systems with control of total mass

Maha Daoud, El-Haj Laamri, Azeddine Baalal

► To cite this version:

Maha Daoud, El-Haj Laamri, Azeddine Baalal. A class of fractional parabolic reaction-diffusion systems with control of total mass: theory and numerics. 2023. hal-04123268

HAL Id: hal-04123268

<https://hal.univ-lorraine.fr/hal-04123268>

Preprint submitted on 9 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A class of fractional parabolic reaction-diffusion systems with control of total mass: theory and numerics

Maha Daoud ¹, El-Haj Laamri ^{*2}, and Azeddine Baalal¹

¹Département de Mathématiques et Informatique, Faculté des Sciences Ain-chock,
Université Hassan II, Casablanca 20100, Maroc

²Institut Elie Cartan, Université de Lorraine, 54 506 Vandoeuvre-les-Nancy, France

mahaadaoud@gmail.com, el-haj.laamri@univ-lorraine.fr, abaalal@gmail.com

June 9, 2023

Abstract

In this paper, we prove global-in-time existence of strong solutions to a class of fractional parabolic reaction-diffusion systems posed in a bounded domain of \mathbb{R}^N . The nonlinear reactive terms are assumed to satisfy natural structure conditions which provide non-negativity of the solutions and uniform control of the total mass. The diffusion operators are of type $u_i \mapsto d_i(-\Delta)^s u_i$ where $0 < s < 1$. Global existence of strong solutions is proved under the assumption that the nonlinearities are at most of polynomial growth. Our results extend previous results obtained when the diffusion operators are of type $u_i \mapsto -d_i \Delta u_i$. On the other hand, we use numerical simulations to examine the global existence of solutions to systems with exponentially growing right-hand sides, which remains so far an open theoretical question even in the case $s = 1$.

Mathematics Subject Classification (2020): 35R11, 35J62, 30G50, 47H10, 35B45.

Keywords. Reaction-diffusion system, fractional diffusion, strong solution, global existence, numerical simulation.

1 Introduction

The purpose of this work is the study of global existence in time of nonnegative strong solutions to fractional parabolic reaction-diffusion systems of the form:

$$(S) \quad \begin{cases} \forall i = 1, \dots, m, \\ \partial_t u_i(t, \mathbf{x}) + d_i(-\Delta)^s u_i(t, \mathbf{x}) = f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})), & (t, \mathbf{x}) \in (0, T) \times \Omega, \\ u_i(t, \mathbf{x}) = 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ u_i(0, \mathbf{x}) = u_{0i}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases}$$

where Ω is a bounded regular open subset of \mathbb{R}^N where $N \geq 1$, $0 < s < 1$, $m \geq 2$ and for each $i = 1, \dots, m$, the diffusion coefficient $d_i > 0$ and f_i is locally Lipschitz continuous.

Here, we mean by $(-\Delta)^s$ the classical fractional Laplacian operator of order s defined by

$$(1.1) \quad (-\Delta)^s \phi(\mathbf{x}) := a_{N,s} \text{P.V.} \int_{\mathbb{R}^N} \frac{\phi(\mathbf{x}) - \phi(\mathbf{y})}{\|\mathbf{x} - \mathbf{y}\|^{N+2s}} d\mathbf{y},$$

*Corresponding author: el-haj.laamri@univ-lorraine.fr

where $\|\cdot\|$ is the euclidean norm of \mathbb{R}^N , P.V. stands for the Cauchy principal value and $a_{N,s}$ is a normalization constant such that the following pair of identities :

$$\lim_{s \rightarrow 0^+} (-\Delta)^s \phi = \phi \quad \text{and} \quad \lim_{s \rightarrow 1^-} (-\Delta)^s \phi = -\Delta \phi$$

holds (see, *e.g.*, [26, Proposition 4.4] and [25, Proposition 2.1]). We refer readers not familiar with fractional laplacians to [25, 58, 26] and the references therein.

Parabolic reaction-diffusion systems appear in various disciplines such as chemistry, biology and environment science (see [66, 74, 70, 61] and their bibliographies). When governed by the classical Laplacian, such systems are related to normal diffusion processes. They have been extensively studied in the literature, see, for instance, [66, 47, 30, 49, 50, 40, 69, 76, 20, 75, 31, 31, 2] and the references therein.

However, in order to describe nonlocal diffusive processes, the classical Laplacian is no longer a well-suited model. Hence the interest of fractional Laplacian operators, which allow one to derive more advanced models, especially anomalous diffusion processes. Additional insights can be found in [1, 78, 18, 79, 80, 42, 56, 25]. More generally, nonlocal partial differential equations (NPDEs) have come out for some years, and play a significant role in numerous scientific fields, as illustrated in [71, 45] and their references. The fractional Laplacians are the prototypical example of nonlocal operators appearing in NPDEs.

Recently, many papers have investigated the existence of solutions to fractional reaction-diffusion problems consisting of a single parabolic equation ($m = 1$) on open subsets of \mathbb{R}^N . We refer the reader, for instance, to [7, 37, 39, 54, 13, 14, 35].

On the other hand, there are few works dealing with *systems* ($m \geq 2$) governed by fractional Laplacians, see [8, 9, 10, 11] and [4, 5, 6] in the case where $\Omega = \mathbb{R}^N$. In this work, we are going to discuss the extensions of results known for $s = 1$ or $m = 1$ to the more general situation $0 < s < 1$ and $m \geq 2$.

Going back to System (S), we are only interested in nonnegative solutions, as in applications the unknown $\mathbf{u} = (u_1, \dots, u_m)$ represents for example concentrations of chemical species or population densities. Therefore, the initial data have to be chosen nonnegative ; $u_{0i} \geq 0$ for each $i \in \{1, \dots, m\}$. It is also well-known that the solutions (as long as they exist) remain nonnegative, provided that the reaction terms f_i satisfy the so-called “quasi-positivity” property, namely:

$$(P) \quad \forall 1 \leq i \leq m, \quad f_i(r_1, \dots, r_{i-1}, 0, r_{i+1}, \dots, r_m) \geq 0, \quad \forall \mathbf{r} = (r_1, \dots, r_m) \in [0, +\infty)^m.$$

Furthermore, in many cases a control (sometimes even the preservation) of the total mass naturally follows the model, namely:

$$(1.2) \quad \sum_{i=1}^m \int_{\Omega} u_i(t, \mathbf{x}) d\mathbf{x} \leq C \sum_{i=1}^m \int_{\Omega} u_{0i}(\mathbf{x}) d\mathbf{x} \quad \text{for some } C > 0.$$

The control (1.2) is fulfilled if

$$(M) \quad \exists (a_1, \dots, a_m) \in (0, +\infty)^m \text{ such that } \sum_{i=1}^m a_i f_i \leq 0,$$

or, more generally, if this sum grows at most linearly in its variables, *i.e.*

$$(M') \quad \forall \mathbf{r} \in [0, +\infty)^m, \quad \sum_{i=1}^m a_i f_i(\mathbf{r}) \leq C \left[1 + \sum_{i=1}^m r_i \right] \quad \text{for some } C \geq 0.$$

Let us emphasize that properties (P) and (M) are satisfied in many applications, *e.g.* in models describing evolution phenomena involving both spatial diffusion and chemical reactions. Nevertheless, there are some instances where the total mass of the solution is not controlled; for example : $m = 2$, $f_1(u_1, u_2) = u_2^3$ and $f_2(u_1, u_2) = u_1^2$.

The question of global existence for reaction-diffusion systems has been a classical topic since the seventies, yet there are still many open and challenging problems. There are two major obstructions to the construction of global solutions. *Firstly*, when the right-hand sides f_i have quadratic, even faster, growth for large values of the u_i 's. A typical example for three species is :

$$f_1 = \alpha_1 g, \quad f_2 = \alpha_2 g, \quad f_3 = -\alpha_3 g \quad \text{where } g = u_3^{\alpha_3} - u_1^{\alpha_1} u_2^{\alpha_2} \quad \text{and } \alpha_1, \alpha_2, \alpha_3 \geq 1.$$

See System (1.3) below for more precisions. *Secondly*, when the diffusion coefficients d_i are very different, there is no comparison principle. Thus, no a priori estimates are available besides the L^1 control (1.2).

To put our work in context and highlight the novelty of this paper, let us briefly review the existing literature.

• Case where the diffusions are driven by classical Laplacian (i.e. $s = 1$)

It turns out that the structure **(P)**+ **(M)** does not keep the solution from blowing up in L^∞ -norm, even in finite time. More precisely, the paper [68] gives an explicit example of a two-species system fulfilling both **(P)** and **(M)**, with $d_1 \neq d_2$ and strictly superquadratic polynomial nonlinearities, and such that

$$\exists T^* < +\infty, \quad \lim_{t \nearrow T^*} \|u_1(t, \cdot)\|_{L^\infty(\Omega)} = \lim_{t \nearrow T^*} \|u_2(t, \cdot)\|_{L^\infty(\Omega)} = +\infty.$$

Thus, in addition to the structure **(P)**+ **(M)**, some growth restrictions and extra structure on the nonlinearities are needed if one expects global existence of strong (or even weak) solutions.

— **Strong solutions:** This issue has been intensively studied, especially when the initial data are bounded. Let us review some sufficient conditions on the f_i 's guaranteeing the global existence of a *strong* solution (see Definition 2.1):

- (i) *triangular structure* (see (1.6)) and polynomial growth. For more details, we refer to [66] and references therein. A prototypical example of triangular structure when $m = 2$ is $f_1 \leq 0$ and $f_1 + f_2 \leq 0$;
- (ii) quadratic growth. See, for instance, [40, 69, 20, 75, 31, 32]);
- (iii) diffusion coefficients close to one another, see, e.g. [19, 30] (sometimes called *quasi-uniform* in some references, for example in [31, 32]);
- (iv) when the growth of the f_i 's is slightly stronger than polynomial, few results are known, and only for $m = 2$. The most-studied model by far is $f_2(u_1, u_2) = -f_1(u_1, u_2) = u_1 e^{u_2^\beta}$. In this case, the existence of strong solutions is established in [41] for $\beta < 1$, and [12] for $\beta = 1$. The article [73] gives some extensions of those results. Surprisingly, in the three previous works, the global existence is established under a restrictive assumption on the size of the initial data u_{01} . On the other hand, the problem is still open for $\beta > 1$.

— **Weak solutions:** By a *weak* solution, we mean a solution in the sense of distributions or, equivalently here, in the sense of the variation-of-constant formula with the suitable semigroup (see Definition 2.2). Such weak L^1 -solution had already been considered in [63, 46, 17] to handle initial data in L^1 . However, an extra condition of *triangular structure* of the nonlinearities was required. Furthermore [64] (see also [66, theorem 5.9] for $m \geq 2$), if for some reason the f_i 's are bounded in $L^1((0, T) \times \Omega)$ for any T , then there exists a global weak solution for *any* initial data in $L^1(\Omega)$, without any further assumption beyond non-negativity. This L^1 bound can be established under any one of the following assumptions:

1. there are m independent inequalities between the f_i 's, and the u_{0i} 's are just assumed to be in $L^1(\Omega)$. For example, System (S) where $m = 2$ and $f_2(u_1, u_2) = -f_1(u_1, u_2) = u_1 e^{u_2^\beta}$ has a weak solution for any $\beta > 0$ and any initial data in $L^1(\Omega)$. We refer to [64, 66] (see also [50]);

2. the f_i 's are of quadratic growth, and the u_{0i} 's belong to $L^2(\Omega)$, see [66, theorem 5.11];
3. the f_i 's are super-quadratic, as it occurs in complex chemical reactions, and the $u_{0,i}$'s are in $L^1(\Omega) \cap H^{-1}(\Omega)$. For more details, see [49] (see also [22]).

Concerning the elliptic case, the interested reader is referred to [50, 52].

To conclude this short summary, we would like to point out that the above references do not exhaust the rich literature on the subject. The interested reader can find an exhaustive review of the results known before 2010 in the extensive survey [66], which also gives a general presentation of the problem, further references, and many deep comments on the mathematical difficulties raised by such systems. For some more recent results, see [47, 30, 19, 69, 72, 20, 75, 31, 32] and their references.

• Case where the diffusions are driven by the fractional Laplacian (*i.e.* $0 < s < 1$):

Unlike the case $s = 1$, relatively little is known. To our knowledge, the existing works fall into two broad categories: either the domain Ω is bounded and the right-hand sides are of potential-gradient or gradient-gradient type; or $\Omega = \mathbb{R}^N$ and the r.h.s. are of polynomial or exponential growth.

- ◊ Ω bounded: in [8], the first two authors and their coworkers investigated the case of a 2×2 System where the right-hand sides are $f_1(u_1, u_2) = \|\nabla u_2\|^q + h_1$ and $f_2(u_1, u_2) = \|\nabla u_1\|^p + h_2$ with $h_1, h_2 \geq 0$. Obviously, f_1 and f_2 do not belong to the framework of (S), nor do they satisfy (M). They also studied the elliptic version of the system in the following two cases :

$$(S_1) \quad (-\Delta)^s u_1 = \|\nabla u_2\|^q + \lambda h_1, \quad (-\Delta)^s u_2 = \|\nabla u_1\|^p + \mu h_2,$$

$$(S_2) \quad (-\Delta)^s u_1 = u_2^q + \lambda h_1, \quad (-\Delta)^s u_2 = \|\nabla u_1\|^p + \mu h_2,$$

where $\lambda, \mu > 0$. For more details, we refer the reader to [9] for (S_1) and [10] for (S_2) ; see also [11] for a generalized System (S_1) with two different diffusions *i.e.* $s_1 \neq s_2$.

- ◊ $\Omega = \mathbb{R}^N$: the only works that we know are those of [4, 5, 6].

As far as we know, the question of global existence of solutions to System (S), set in a bounded domain Ω , with the reaction terms satisfying (P)+(M) and of polynomial or exponential growth, has not been addressed so far. This is the main goal of this work. More specifically:

1. we will extend to the fractional case ($0 < s < 1$) two main known results in the classical case (*i.e.* $s = 1$) and the reaction terms are polynomial growing. The first one deals with the typical case of reversible chemical reactions for three species (see Theorem 4.1). The second one deals with the case where the number of equations $m \geq 2$ and the f_i 's have a triangular structure (see Theorem 4.2).
2. we present some numerical simulations to examine the global existence of solutions to System (S) in the case where $m = 2$, $f_2(u_1, u_2) = -f_1(u_1, u_2) = u_1 e^{u_2^\beta}$ with $\beta > 1$. See Section 5.

In order to prove our main theorems, we will extend several tools known in the classical case to the fractional case such as the maximal regularity theorem (see Theorem 3.1), a Lambertion-type estimation in L^p (see Theorem 3.3), the so-called Pierre's duality Lemma (see Lemma 4.2). Such results are interesting in themselves. Nevertheless, it should be noted that it is not always possible to extend any result from the classical case to the fractional case; and when it is possible, it is usually far from being trivial. See, for instance, [8, 9, 10, 11] in a different context.

For the ease of reader and the sake of completeness, let us now recall those two results that we will extend.

• **First known result.** Let us consider the following System

$$(1.3) \quad \begin{cases} \partial_t u_1(t, \mathbf{x}) - d_1 \Delta u_1(t, \mathbf{x}) = f_1(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ \partial_t u_2(t, \mathbf{x}) - d_2 \Delta u_2(t, \mathbf{x}) = f_2(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ \partial_t u_3(t, \mathbf{x}) - d_3 \Delta u_3(t, \mathbf{x}) = f_3(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ u_1(t, \mathbf{x}) = u_2(t, \mathbf{x}) = u_3(t, \mathbf{x}) = 0, & (t, \mathbf{x}) \in (0, T) \times \partial\Omega, \\ u_1(0, \mathbf{x}) = u_{01}(\mathbf{x}) \geq 0, & \mathbf{x} \in \Omega, \\ u_2(0, \mathbf{x}) = u_{02}(\mathbf{x}) \geq 0, & \mathbf{x} \in \Omega, \\ u_3(0, \mathbf{x}) = u_{03}(\mathbf{x}) \geq 0, & \mathbf{x} \in \Omega, \end{cases}$$

where

$$f_1 = \alpha_1 g, \quad f_2 = \alpha_2 g, \quad f_3 = -\alpha_3 g \quad \text{with } g = u_3^{\alpha_3} - u_1^{\alpha_1} u_2^{\alpha_2} \text{ and } \alpha_1, \alpha_2, \alpha_3 \geq 1.$$

That system naturally arises in chemical kinetics when modeling the following reversible reaction

$$(1.4) \quad \alpha_1 U_1 + \alpha_2 U_2 \rightleftharpoons \alpha_3 U_3$$

where u_1, u_2, u_3 stand for the density of U_1, U_2 and U_3 respectively, and $\alpha_1, \alpha_2, \alpha_3$ are the stoichiometric coefficients.

Let us notice that in addition to fulfilling **(P)**, f_1, f_2 and f_3 also satisfy **(M)**, namely

$$\alpha_2 \alpha_3 f_1 + \alpha_1 \alpha_3 f_2 + 2\alpha_1 \alpha_2 f_3 = 0.$$

Now, we are ready to state the first known result, (see [47, Theorems 1 et 2]).

Theorem 1.1. *Assume that $(u_{01}, u_{02}, u_{03}) \in (L^\infty(\Omega))^3$. System (1.3) admits a unique nonnegative global strong solution in the following cases*

- (i) $\alpha_1 + \alpha_2 < \alpha_3$;
- (ii) $\alpha_3 = 1$ whatever are α_1 and α_2 ;
- (iii) $d_1 = d_3$ or $d_2 = d_3$ whatever are α_1, α_2 and α_3 ;
- (iv) $d_1 = d_2 \neq d_3$ for any $(\alpha_1, \alpha_2, \alpha_3) \in [1, +\infty)^2 \times (1, +\infty)$ such that $\alpha_1 + \alpha_2 \neq \alpha_3$.

Comments:

Assume that the diffusion coefficients d_1, d_2 and d_3 are very different. The question of global existence of a strong (even weak) solution to System (1.3) in the case $2 < \alpha_3 < \alpha_1 + \alpha_2$ is widely open since at least 2011. This is very surprising from a chemical point of view because the reaction (1.4) is supposed to be reversible and therefore a strong solution should exist and be global as in the case $\alpha_1 + \alpha_2 < \alpha_3$. More generally, that question is far from being understood. For example, the global existence of a strong solution of the system modeling a reversible reaction of the form $\alpha_1 U_1 + \alpha_2 U_2 \rightleftharpoons \alpha_3 U_3 + \alpha_4 U_4$ has only recently been proved and in the sole case $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = 1$, see [20, 75, 31, 32]).

• **Second known result.** Introducing the classical counterpart of System (S), namely:

$$(1.5) \quad \begin{cases} \forall i = 1, \dots, m, \\ \partial_t u_i(t, \mathbf{x}) - d_i \Delta u_i(t, \mathbf{x}) = f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})), & (t, \mathbf{x}) \in (0, T) \times \Omega, \\ u_i(t, \mathbf{x}) = 0, & (t, \mathbf{x}) \in (0, T) \times \partial\Omega, \\ u_i(0, \mathbf{x}) = u_{0i}(\mathbf{x}), & \mathbf{x} \in \Omega. \end{cases}$$

It holds that (see [66, Theorem 3.5] and its references, especially [59])

Theorem 1.2. *Assume that for each $i \in \{1, \dots, m\}$, $u_{0i} \in (L^\infty(\Omega))^+$ and f_i is at most of polynomial growth and satisfy **(P)**. Moreover, assume that*

$$(1.6) \quad \begin{cases} \text{there exist a vector } \mathbf{b} \in \mathbb{R}^m \text{ and a lower triangular invertible matrix } Q \in M_m(\mathbb{R}^+) \text{ s.t.} \\ \forall \mathbf{r} = (r_1, \dots, r_m) \in [0, +\infty)^m, \quad Q\mathbf{f}(\mathbf{r}) \leq \left[1 + \sum_{1 \leq i \leq m} r_i\right] \mathbf{b}. \end{cases}$$

Then, System (1.5) admits a unique nonnegative global strong solution.

The rest of this article is set out in the following manner. In Section 2, we present the necessary background for the remaining sections. First, we set up some basic definitions about the notion of semigroups, and some required results concerning the associated evolution problem. Then, we recall some fundamental properties of the fractional laplacian within the framework of semigroup theory. In Section 3, we focus on the fractional evolution equation and examine the conditions that ensure the boundedness of the solution. Furthermore, we concentrate on significant concepts such as the dual problem, the maximal regularity and the comparison principle. Section 4 is divided into three subsections. In subsection 4.1, we first investigate the local existence of solutions to System (S), then we prove the global existence in a particular case and we end it by extending Pierre's duality lemma to the fractional case. Subsections 4.2 and 4.3 are devoted respectively to the proofs of our two main theorems. In Section 5, we present some numerical simulations to examine the global existence of solutions to a 2×2 System with exponential growth (see (S_{exp}) in page 19), which remains a theoretical open question even in the case $s = 1$. At last, we introduce the proofs of Theorems 3.1 and 3.2 in the Appendix.

Prior to ending this section, let us fix some notations.

Notations. Throughout this paper, we will use the following standard notations :

- Ω is a bounded regular open subset of \mathbb{R}^N with $N \geq 1$.
- For any $T > 0$, $Q_T := (0, T) \times \Omega$.
- $\|\cdot\|$ is the Euclidean norm of \mathbb{R}^N .
- For any $p \in [1, +\infty)$, $\|\phi\|_{L^p(\Omega)} = \left(\int_{\Omega} |\phi(\mathbf{x})|^p d\mathbf{x} \right)^{\frac{1}{p}}$ and $\|\psi\|_{L^p(Q_T)} = \left(\int_0^T \int_{\Omega} |\psi(t, \mathbf{x})|^p dt d\mathbf{x} \right)^{\frac{1}{p}}$.
- $\|\phi\|_{L^\infty(\Omega)} = \text{ess sup}_{\mathbf{x} \in \Omega} |\phi(\mathbf{x})|$ and $\|\psi\|_{L^\infty(Q_T)} = \text{ess sup}_{(t, \mathbf{x}) \in Q_T} |\psi(t, \mathbf{x})|$.
- $\mathbb{H}_0^s(\Omega) := \{\phi \in L^2(\mathbb{R}^N) ; \|\phi\|_{\mathbb{H}_0^s(\Omega)} < +\infty \text{ and } \phi = 0 \text{ in } \mathbb{R}^N \setminus \Omega\}$, where

$$\|\phi\|_{\mathbb{H}_0^s(\Omega)} := \left(\iint_{\mathbb{R}^N \times \mathbb{R}^N} \frac{|\phi(\mathbf{x}) - \phi(\mathbf{y})|^2}{\|\mathbf{x} - \mathbf{y}\|^{N+2s}} d\mathbf{x} d\mathbf{y} \right)^{\frac{1}{2}}.$$

2 Preliminaries

In this section, we briefly review some relevant results for later use. First, we recall existence results of solutions to evolution problems associated with semigroups, along with a Lamberton-type estimation based on semigroups on L^2 . Then, we discuss the main properties of the fractional Laplacian and the semigroup generated by it on L^2 .

2.1 Evolution problems associated with semigroups

Let (E, Σ, μ) be a measure space. Let A be the infinitesimal generator of a strongly continuous semigroup $\{S_A(t)\}_{t \geq 0}$ on $L^2(E, \Sigma, \mu)$ (or simply $L^2(E)$). We refer readers not familiar with the notion of semigroups to standard works such as [62, Chapter 1] and [81, Chapter 7].

Now, let us associate with the operator A the following evolution problem :

$$(2.1) \quad \begin{cases} \frac{dw}{dt} - Aw &= h, & \text{in } (0, T), \\ w(0) &= w_0, \end{cases}$$

where $h(t)$ and w_0 belong to $L^p(E)$, $p \geq 1$.

Before stating the existence theorems, let us make precise what we mean by *strong* solution and *weak* solution to Problem (2.1).

Definition 2.1 (Strong solution). *A function $w : [0, T] \rightarrow L^p(E)$ is called a strong solution to (2.1) if :*

- (i) $w \in \mathcal{C}([0, T]; L^p(E)) \cap \mathcal{C}^1(0, T; L^p(E))$;
- (ii) for any $t \in (0, T)$, $w(t) \in D(A)$;
- (iii) Problem (2.1) is satisfied a.e. on $[0, T]$.

Definition 2.2 (Weak solution). *Let $h \in L^1(0, T; L^p(E))$. A function $w \in \mathcal{C}([0, T]; L^p(E))$ is called a weak solution to (2.1) if it satisfies the following integral equation :*

$$(2.2) \quad w(t) = S_A(t)w_0 + \int_0^t S_A(t - \tau)h(\tau)d\tau, \quad \forall t \in [0, T].$$

The first existence Theorem reads as (see, for instance, [62, Chapter 4]):

Theorem 2.1. *Let $h \in L^1(0, T; L^p(E))$ and $w_0 \in L^p(E)$. Then, Problem (2.1) admits a unique weak solution.*

Now, let us assume that $\{S_A(t)\}_{t \geq 0}$ satisfies the two additional assumptions :

(H1) $\{S_A(t)\}_{t \geq 0}$ is a bounded analytic semigroup on $L^2(E)$;

(H2) for any $p \in [1, +\infty]$ and any $\phi \in L^p(E) \cap L^2(E)$, we have

$$\forall t \geq 0, \quad \|S_A(t)\phi\|_{L^p(E)} \leq \|\phi\|_{L^p(E)}.$$

The following theorem provides a well-known result for estimating the weak solution to Problem (2.1) in L^p when $w_0 = 0$, see [53, Theorem 1].

Theorem 2.2 (Lamberton-type estimation). *Assume that $w_0 = 0$ and $h \in L^p((0, T) \times E)$ with $p \in (1, +\infty)$. Let w be the weak solution to Problem (2.1). Then, there exists a constant $C_p := C(p) > 0$ such that*

$$(2.3) \quad \left\| \frac{dw}{dt} \right\|_{L^p((0, T) \times E)} + \|Aw\|_{L^p((0, T) \times E)} \leq C_p \|h\|_{L^p((0, T) \times E)}.$$

We end this subsection by noting that when $h \equiv 0$, we have $w(t) = S_A(t)w_0$ as the weak solution to (2.1). Hence, it is clear that not every weak solution to (2.1) becomes a strong solution. However, the two notions of solution coincide if h satisfies a further condition, namely if $h \in \mathcal{C}^1(0, T; L^p(E))$. More precisely, one has the next result (see, e.g., [62, Corollary 3.3]).

Theorem 2.3. *Assume that $h \in L^1(0, T; L^p(E)) \cap \mathcal{C}^1(0, T; L^p(E))$. Then, the weak solution to Problem (2.1) is a strong solution.* \square

From now on, we set $E = \Omega$. As stated in the introduction, Ω is assumed to be a bounded regular open subset of \mathbb{R}^N .

2.2 Semigroup generated by the Fractional Laplacian

It is well-known that the operator $A = -(-\Delta)^s$ with domain

$$(2.4) \quad D(A) = \{\phi \in \mathbb{H}_0^s(\Omega), (-\Delta)^s \phi \in L^2(\Omega)\}$$

generates a strongly continuous *submarkovian*¹ semigroup $\{S_A(t)\}_{t \geq 0}$ on $L^2(\Omega)$, see [39, Proposition 2.14] and [21, Theorem 3.3].

The following theorem will be used throughout the rest of this paper. We refer, for instance, to [21, Theorem 2.10] or [39, Theorem 2.16] for further details.

¹A strongly continuous semigroup $\{S_A(t)\}_{t \geq 0}$ is said submarkovian if it satisfies the two following properties:
– if $\phi \in L^2(\Omega)$ and $\phi \geq 0$, then $S_A(t)\phi \geq 0 \forall t \geq 0$;
– if $\phi \in L^\infty(\Omega)$, then $\|S_A(t)\phi\|_{L^\infty(\Omega)} \leq \|\phi\|_{L^\infty(\Omega)}$.

Theorem 2.4. Let $\{S_A(t)\}_{t \geq 0}$ be the submarkovian semigroup generated by A on $L^2(\Omega)$ with domain given by (2.4). Then:

(i) the semigroup $\{S_A(t)\}_{t \geq 0}$ can be extended to a contraction semigroup on $L^p(\Omega)$ for any $p \in [1, +\infty]$. Moreover, each contraction semigroup is strongly continuous if $p \in [1, +\infty)$ and bounded analytic if $p \in (1, +\infty)$;

(ii) the semigroup $\{S_A(t)\}_{t \geq 0}$ is ultracontractive, i.e., for any $\phi \in L^p(\Omega)$ and $1 \leq p \leq q \leq +\infty$, there exists a constant $C > 0$ such that

$$(2.5) \quad \|S_A(t)\phi\|_{L^q(\Omega)} \leq Ct^{-\frac{N}{2s}(\frac{1}{p}-\frac{1}{q})} \|\phi\|_{L^p(\Omega)}, \quad \forall t > 0.$$

Moreover, if $p = q = +\infty$ and $C = 1$, the semigroup $\{S_A(t)\}_{t \geq 0}$ is said L^∞ -contractive. \square

3 Fractional evolution problem

In order to study the existence and regularity of solutions to our System, we will rely on properties of solutions to the equation. To do so, let us consider the following problem :

$$(3.1) \quad \begin{cases} \partial_t w(t, \mathbf{x}) + d(-\Delta)^s w(t, \mathbf{x}) &= h(t, \mathbf{x}), & (t, \mathbf{x}) \in Q_T, \\ w(t, \mathbf{x}) &= 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ w(0, \mathbf{x}) &= w_0(\mathbf{x}) \geq 0, & \mathbf{x} \in \Omega, \end{cases}$$

where $d > 0$, $w_0 \in L^p(\Omega)$ and $h \in L^p(Q_T)$, $p > 1$.

First, we recall the conditions under which the solution to Problem (3.1) exists and is bounded. Then, we consider the dual problem of (3.1) in order to prove a Lamberton-type estimation in L^p .

3.1 Existence and Boundedness

Now, let us denote $A := -d(-\Delta)^s$. According to Theorem 2.4, the operator A verifies the assumptions **(H1)** and **(H2)**. Hence, by Theorem 2.1, Problem (3.1) admits a unique weak solution $w \in \mathcal{C}([0, T], L^p(\Omega))$ which fulfills

$$(3.2) \quad w(t, \cdot) = S_A(t)w_0 + \int_0^t S_A(t - \tau)h(\tau, \cdot)d\tau.$$

Furthermore by Theorem 2.2, when $w_0 = 0$ there exists a constant $C_p > 0$ such that

$$(3.3) \quad \|\partial_t w\|_{L^p(Q_T)} + d \|(-\Delta)^s w\|_{L^p(Q_T)} \leq C_p \|h\|_{L^p(Q_T)}.$$

The regularity of the weak solution to Problem (3.1) relies on the two Theorems presented below.

— The first Theorem outlines the necessary conditions for the solution to Problem (3.1) to have a maximal regularity property in L^∞ .

Theorem 3.1. Assume that $w_0 \in L^\infty(\Omega)$ and $h \in L^p(Q_T)$ with $p > 1$. Let w be the weak solution to Problem (3.1). Then, for any $p > \frac{N + 2s}{2s}$, there exists a constant $C > 0$ such that

$$\|w\|_{L^\infty(Q_T)} \leq \|w_0\|_{L^\infty(\Omega)} + C \|h\|_{L^p(Q_T)}.$$

— The second Theorem is related to the comparison principle for Problem (3.1).

Theorem 3.2. Assume that $w_0 \in L^\infty(\Omega)$ and $h \in L^p(Q_T)$ with $p > 1$. Let w be the weak solution to Problem (3.1). If $h \leq 0$, then $w \in L^\infty(Q_T)$ and

$$\|w\|_{L^\infty(Q_T)} \leq \|w_0\|_{L^\infty(\Omega)}.$$

The two previous results are classical when $s = 1$. Surprisingly, we did not find, in the literature, their equivalents in the fractional case. Therefore, we have decided to introduce their proofs in this paper although they are close to those of the $s = 1$ case. Nevertheless, for reading fluency we preferred to postpone these proofs in the appendix.

3.2 Fractional dual problem of Problem (3.1)

In this subsection, we state and prove the existence and regularity result of solutions to the dual problem of (3.1). Such result will be useful to prove the global existence of solutions to our System.

The dual problem of Problem (3.1) is given by

$$(P_\varphi) \quad \begin{cases} -\partial_t \mathcal{Z}(t, \mathbf{x}) + d(-\Delta)^s \mathcal{Z}(t, \mathbf{x}) &= \varphi(t, \mathbf{x}), & (t, \mathbf{x}) \in Q_T, \\ \mathcal{Z}(t, \mathbf{x}) &= 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ \mathcal{Z}(T, \mathbf{x}) &= 0, & \mathbf{x} \in \Omega, \end{cases}$$

where φ is a regular function.

The following theorem provides the conditions for the solution of the dual problem (P_φ) to exist and verify a Lambertion-type estimation in L^p .

Theorem 3.3. *Assume that $\varphi \in L^p(Q_T)$ with $p > 1$. Then, Problem (P_φ) admits a unique weak solution. Moreover, there exists a constant $C := C(p, T) > 0$ such that*

$$(3.4) \quad \|\partial_t \mathcal{Z}\|_{L^p(Q_T)} + \|\mathcal{Z}\|_{L^p(Q_T)} + d \|(-\Delta)^s \mathcal{Z}\|_{L^p(Q_T)} + \|\mathcal{Z}_0\|_{L^p(\Omega)} \leq C \|\varphi\|_{L^p(Q_T)},$$

where $\mathcal{Z}_0 := \mathcal{Z}(0, \cdot)$.

Proof. In Problem (P_φ) , let us make the following change of variable : $t = T - \tau \Rightarrow \tau = T - t \in (0, T)$. Furthermore, let us denote $w(\tau, \cdot) := \mathcal{Z}(T - \tau, \cdot)$ and $h(\tau, \cdot) := \varphi(T - \tau, \cdot)$.

Then, Problem (P_φ) becomes

$$(3.5) \quad \begin{cases} \partial_\tau \mathcal{Z}(T - \tau, \mathbf{x}) + d(-\Delta)^s \mathcal{Z}(T - \tau, \mathbf{x}) &= \varphi(T - \tau, \mathbf{x}), & (\tau, \mathbf{x}) \in Q_T, \\ \mathcal{Z}(T - \tau, \mathbf{x}) &= 0, & (\tau, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ \mathcal{Z}(T, \mathbf{x}) &= 0, & \mathbf{x} \in \Omega, \end{cases}$$

which is equivalent to

$$(3.6) \quad \begin{cases} \partial_\tau w(\tau, \mathbf{x}) + d(-\Delta)^s w(\tau, \mathbf{x}) &= h(\tau, \mathbf{x}), & (\tau, \mathbf{x}) \in Q_T, \\ w(\tau, \mathbf{x}) &= 0, & (\tau, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ w(0, \mathbf{x}) &= 0, & \mathbf{x} \in \Omega. \end{cases}$$

Hence, the existence of a unique weak solution to (P_φ) follows from Theorem 3.1. In addition, there exists a constant $C_1 := C_1(p) > 0$ such that

$$(3.7) \quad \|\partial_t \mathcal{Z}\|_{L^p(Q_T)} + d \|(-\Delta)^s \mathcal{Z}\|_{L^p(Q_T)} \leq C_1 \|\varphi\|_{L^p(Q_T)}.$$

Now, let us integrate the first equation of (P_φ) over $(0, T)$. We obtain

$$(3.8) \quad -\int_0^T \partial_t \mathcal{Z}(t, \mathbf{x}) dt = \int_0^T (\varphi(t, \mathbf{x}) - d(-\Delta)^s \mathcal{Z}(t, \mathbf{x})) dt.$$

For any $\mathbf{x} \in \Omega$, $\mathcal{Z}(T, \mathbf{x}) = 0$. Then,

$$(3.9) \quad \mathcal{Z}_0(\mathbf{x}) = \int_0^T (\varphi(t, \mathbf{x}) - d(-\Delta)^s \mathcal{Z}(t, \mathbf{x})) dt.$$

Therefore, we get

$$(3.10) \quad \begin{cases} \|\mathcal{Z}_0\|_{L^p(\Omega)} &= \left\| \int_0^T (\varphi(t, \cdot) - d(-\Delta)^s \mathcal{Z}(t, \cdot)) dt \right\|_{L^p(\Omega)} \\ &\leq \int_0^T \|\varphi(t, \cdot) - d(-\Delta)^s \mathcal{Z}(t, \cdot)\|_{L^p(\Omega)} dt \\ &\leq \int_0^T \|\varphi(t, \cdot)\|_{L^p(\Omega)} dt + \int_0^T \|d(-\Delta)^s \mathcal{Z}(t, \cdot)\|_{L^p(\Omega)} dt \\ &\leq T^{1-\frac{1}{p}} (\|\varphi\|_{L^p(Q_T)} + d \|(-\Delta)^s \mathcal{Z}\|_{L^p(Q_T)}) \\ &\leq T^{1-\frac{1}{p}} \max(1, C_1) \|\varphi\|_{L^p(Q_T)}. \end{cases}$$

The last inequality follows by the estimate (3.7). Hence,

$$(3.11) \quad \|\mathcal{Z}_0\|_{L^p(\Omega)} \leq C_2 \|\varphi\|_{L^p(Q_T)},$$

where $C_2 := C_2(p, T) > 0$.

As \mathcal{Z} is the weak solution to Problem (P_φ) , thus it verifies the following integral equation :

$$(3.12) \quad \mathcal{Z}(t, \mathbf{x}) = \int_t^T S_A(\tau - t) \varphi(\tau, \mathbf{x}) d\tau, \quad (t, \mathbf{x}) \in [0, T] \times \Omega.$$

Thus, we have

$$(3.13) \quad \left\{ \begin{array}{l} \|\mathcal{Z}(t, \cdot)\|_{L^p(\Omega)} \leq \left\| \int_t^T S_A(\tau - t) \varphi(\tau, \cdot) d\tau \right\|_{L^p(\Omega)} \\ \leq \int_t^T \|S_A(\tau - t) \varphi(\tau, \cdot)\|_{L^p(\Omega)} d\tau \\ \leq \int_0^T \|S_A(\tau - t) \varphi(\tau, \cdot)\|_{L^p(\Omega)} d\tau \\ \leq C \int_0^T \|\varphi(\tau, \cdot)\|_{L^p(\Omega)} d\tau \\ \leq CT^{1-\frac{1}{p}} \|\varphi\|_{L^p(Q_T)}. \end{array} \right.$$

Consequently,

$$(3.14) \quad \|\mathcal{Z}\|_{L^\infty(0, T; L^p(\Omega))} \leq CT^{1-\frac{1}{p}} \|\varphi\|_{L^p(Q_T)}.$$

As $L^\infty(0, T; L^p(\Omega)) \hookrightarrow L^p(Q_T)$, we obtain the estimate

$$(3.15) \quad \|\mathcal{Z}\|_{L^p(Q_T)} \leq C_3 \|\varphi\|_{L^p(Q_T)},$$

where $C_3 := C_3(p, T) > 0$.

Finally, let us denote $C := \max(C_1, C_2, C_3) > 0$. From the estimates (3.7), (3.11) and (3.15), we obtain (3.4). \square

Remark 3.1. Let us mention another notion of solution often found in the PDE literature.

Let $\beta \in (0, 1)$. It is well known (see [55, 33]) that if $\varphi \in L^\infty(Q_T) \cap \mathcal{C}^{0, \beta}(Q_T)$, Problem (P_φ) admits a regular solution $\mathcal{Z} \in L^\infty(Q_T)$ and the equation $-\partial_t \mathcal{Z} + (-\Delta)^s \mathcal{Z} = \varphi$ is satisfied in a pointwise sense.

Let us multiply (3.2) by \mathcal{Z} and integrate over Q_T . Then, we obtain

$$(3.16) \quad \iint_{Q_T} w(t, \mathbf{x}) (-\partial_t \mathcal{Z}(t, \mathbf{x}) + (-\Delta)^s \mathcal{Z}(t, \mathbf{x})) d\mathbf{x} dt = \iint_{Q_T} h(t, \mathbf{x}) \mathcal{Z}(t, \mathbf{x}) d\mathbf{x} dt + \int_\Omega w_0(\mathbf{x}) \mathcal{Z}_0(\mathbf{x}) d\mathbf{x}.$$

A solution $\mathcal{Z} \in \mathcal{C}([0, T]; L^1(\Omega))$ that satisfies (3.16) is called a solution in the sense of distributions to Problem (3.1). The existence and uniqueness criteria for such solutions have been established in [55, Theorem 28] by using an approximation argument. Due to the properties of the semigroup generated by the fractional Laplacian, this notion coincides with that of a weak solution. \square

4 Main results of global existence for Fractional Reaction-Diffusion Systems

Let us recall System

$$(S) \quad \left\{ \begin{array}{ll} \forall i = 1, \dots, m, \\ \partial_t u_i(t, \mathbf{x}) + d_i (-\Delta)^s u_i(t, \mathbf{x}) = f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ u_i(t, \mathbf{x}) = 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ u_i(0, \mathbf{x}) = u_{0i}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{array} \right.$$

where $m \geq 2$, $0 < s < 1$ and for each $i = 1, \dots, m$, $d_i > 0$, $u_{0i} \in L^\infty(\Omega)$ and f_i is locally Lipchitz continuous.

As mentioned in the introduction, our main contribution is to extend Theorem 1.1 and Theorem 1.2 to the fractional case. That is the purpose of this section. Moreover, for the sake of clarity, this section will be divided into three subsections. In the first one, we will treat the local existence and extend Pierre's duality lemma to the fractional case. Subsections 4.2 and 4.3 will be devoted respectively to the proofs of our main Theorems 4.1 and 4.2.

One should not conclude this short introduction without mentioning that once we have established the necessary tools adapted to the fractional Laplacian, the demonstrations are similar to those for the case where the diffusions are governed by the classical Laplacian. However, for the convenience of the reader and for the completeness of the article, we will give the proofs in detail.

4.1 Local existence and extension of Pierre's duality lemma to the fractional case

• **Local existence.** The local existence lemma is classical. Let us recall here its statement. As the proof is straightforward, we omit it.

Lemma 4.1. *Assume that $\mathbf{u}_0 := (u_{01}, \dots, u_{0m}) \in (L^\infty(\Omega))^m$, the f_i 's are locally Lipschitz continuous. Then, there exists $T_{\max} > 0$ and $\Phi = (\varphi_1, \dots, \varphi_m) \in \mathcal{C}([0, T_{\max}], [0, +\infty)^m)$ such that :*

- (i) *System (S) has a unique nonnegative strong solution $\mathbf{u} = (u_1, \dots, u_m)$ in $(0, T_{\max}) \times \Omega$;*
- (ii) *for each $i \in \{1, \dots, m\}$*

$$(4.1) \quad \|u_i(t, \cdot)\|_{L^\infty(\Omega)} \leq \varphi_i(t) \text{ for any } t \in (0, T_{\max});$$

- (iii) *if $T_{\max} < +\infty$, then $\lim_{t \nearrow T_{\max}} \sum_{i=1}^m \|u_i(t, \cdot)\|_{L^\infty(\Omega)} = +\infty$;*

- (iv) *if, in addition, the f_i 's satisfy **(P)**, then*

$$(\forall i \in \{1, \dots, m\}, u_{0i}(\cdot) \geq 0) \implies (\forall i \in \{1, \dots, m\}, u_i(t, \cdot) \geq 0 \forall t \in [0, T_{\max})).$$

Remark 4.1. *According to (iii), in order to prove global existence of strong solutions to System (S), it is sufficient to prove an a priori estimate of the form*

$$(4.2) \quad \forall t \in [0, T_{\max}), \sum_{i=1}^m \|u_i(t, \cdot)\|_{L^\infty(\Omega)} < \psi(t),$$

where $\psi : [0, +\infty) \rightarrow [0, +\infty)$ is a continuous function.

• Proof of global existence in a specific case

It turns out that an estimate like (4.2) is far from being obvious for our System except in the trivial case where $d_1 = \dots = d_m = d$ and the f_i 's satisfy **(P)**.

Then, suppose that for some $(a_1, \dots, a_m) \in (0, +\infty)^m$ such that

$$(4.3) \quad \forall \mathbf{r} \in [0, +\infty)^m, \sum_{i=1}^m a_i f_i(\mathbf{r}) \leq C \left[1 + \sum_{i=1}^m r_i \right], \quad \text{where } C \geq 0.$$

Let $T \in (0, T_{\max})$, $t \in (0, T]$ and $\mathbf{x} \in \Omega$. We denote $W(t, \mathbf{x}) := \sum_{i=1}^m a_i u_i(t, \mathbf{x})$ and $W_0(\mathbf{x}) := W(0, \mathbf{x})$. Let us multiply each i th equation of (S) by a_i and sum over i . Thus, we get

$$(4.4) \quad \partial_t W(t, \mathbf{x}) + d(-\Delta)^s W(t, \mathbf{x}) = \sum_{i=1}^m a_i f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})).$$

— If $C = 0$ in (4.3) (i.e. we have the assumption **(M)**), we deduce from Theorem 3.2 that:

$$\|W\|_{L^\infty(Q_T)} \leq \|W_0\|_{L^\infty(\Omega)}$$

which yields, thanks the nonnegativity of the u_i 's, to

$$(4.5) \quad \sum_{i=1}^m \|u_i\|_{L^\infty(Q_T)} \leq \frac{1}{a_{i_0}} \|W_0\|_{L^\infty(\Omega)} \text{ where } a_{i_0} := \min\{a_1, \dots, a_m\}.$$

Given that (4.5) implies (4.2), this concludes the proof in this case.

— If $C > 0$ in (4.3) (i.e. we have the assumption **(M')**), introduce w the weak solution to

$$(4.6) \quad \begin{cases} \partial_t w(t, \mathbf{x}) + d(-\Delta)^s w(t, \mathbf{x}) &= C \left[1 + \sum_{i=1}^m u_i(t, \mathbf{x}) \right], & (t, \mathbf{x}) \in Q_T, \\ w(t, \mathbf{x}) &= 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ w(0, \mathbf{x}) &= 0, & \mathbf{x} \in \Omega. \end{cases}$$

Then, w satisfies

$$(4.7) \quad w(t, \mathbf{x}) = C \int_0^t S_i(t - \tau) \left[1 + \sum_{i=1}^m u_i(\tau, \mathbf{x}) \right] d\tau, \quad (t, \mathbf{x}) \in [0, T) \times \Omega.$$

Let us multiply each i th equation of (S) by a_i and sum over i . Thus, we get

$$(4.8) \quad \partial_t W(t, \mathbf{x}) + d(-\Delta)^s W(t, \mathbf{x}) = \sum_{i=1}^m a_i f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})).$$

By substracting the first equation of (4.6) from (4.8), we obtain

$$(4.9) \quad \partial_t (W - w)(t, \mathbf{x}) + d(-\Delta)^s (W - w)(t, \mathbf{x}) \leq 0.$$

By Theorem 3.2, we get

$$(4.10) \quad \|W - w\|_{L^\infty(Q_T)} \leq \|W_0\|_{L^\infty(\Omega)}.$$

Therefore,

$$(4.11) \quad \|W\|_{L^\infty(Q_T)} \leq \|W_0\|_{L^\infty(\Omega)} + \|w\|_{L^\infty(Q_T)}.$$

By taking the $L^\infty(\Omega)$ -norm of (4.7) and using the L^∞ -contractivity of $\{S_A(t)\}_{t \geq 0}$, we obtain

$$a_{i_0} \left\| \sum_{i=1}^m u_i(t, \cdot) \right\|_{L^\infty(\Omega)} \leq \left\| \sum_{i=1}^m a_i u_{0i} \right\|_{L^\infty(\Omega)} + C \int_0^t \left[1 + \left\| \sum_{i=1}^m u_i(\tau, \cdot) \right\|_{L^\infty(\Omega)} \right] d\tau.$$

Now, by applying Gronwall's Lemma, we deduce that

$$\left\| \sum_{i=1}^m u_i(t, \cdot) \right\|_{L^\infty(\Omega)} \leq g(t),$$

where $g : [0, +\infty) \rightarrow [0, +\infty)$ is a continuous function. As **(P)** assures us the nonnegativity of u_i for any $i \in \{1, \dots, m\}$, then $u_i(t, \cdot)$ is bounded in $L^\infty(\Omega) \forall t \in [0, T_{\max})$. This implies that $T_{\max} = +\infty$. \square

At this point, the following remark is in order.

Remark 4.2. *As we have just proved in the case $d_1 = \dots = d_m$, System (S) admits a global strong solution **whatever the growth** of the f_i 's. However, the situation is quite more complicated if the diffusion coefficients are different from each other. Some additional assumptions on the growth of the source terms are needed if one expects global existence of strong solutions even if $s = 1$ as already specified in the introduction.*

• **Extension of Pierre's duality lemma to the fractional case**

One of the main tools that we will use to establish our global existence theorems is the following lemma. This is nothing other than an extension of Pierre's duality lemma to the fractional case.

Lemma 4.2. *Given $T > 0$. Assume that $m = 2$, f_1 and f_2 satisfy (P) and (M). Let (u_1, u_2) be the strong solution to (S) in Q_T . Then, there exists a constant $C = C(p, T, \Omega) > 0$ such that*

$$\|u_2\|_{L^p(Q_T)} \leq C (1 + \|u_1\|_{L^p(Q_T)}) \quad \forall 1 < p < +\infty.$$

Pierre's duality Lemma was first stated in [44] and then widely exploited, see for example [46, 47, 30, 19, 66] and the references therein.

Proof. Without loss of generality, we will assume that $a_1 = a_2 = 1$.

Let $T \in (0, T_{\max})$ and $t \in (0, T]$. By summing the two equations of u_1 and u_2 , we obtain

$$(4.12) \quad \partial_t(u_1(t, \mathbf{x}) + u_2(t, \mathbf{x})) + (-\Delta)^s(d_1 u_1(t, \mathbf{x}) + d_2 u_2(t, \mathbf{x})) \leq 0, \quad (t, \mathbf{x}) \in Q_T.$$

Let φ be a nonnegative regular function and let \mathcal{Z} be the solution to Problem (P_φ) with $d = d_2$. Now, we multiply (4.12) by \mathcal{Z} and integrate over Q_T . We get

$$(4.13) \quad \begin{cases} - \iint_{Q_T} (u_1(t, \mathbf{x}) + u_2(t, \mathbf{x})) \partial_t \mathcal{Z}(t, \mathbf{x}) d\mathbf{x} dt + d_2 \iint_{Q_T} u_2(t, \mathbf{x}) (-\Delta)^s \mathcal{Z}(t, \mathbf{x}) d\mathbf{x} dt \\ \leq \int_{\Omega} (u_{01} - u_{02})(\mathbf{x}) \mathcal{Z}_0(\mathbf{x}) d\mathbf{x} - d_1 \iint_{Q_T} u_1(t, \mathbf{x}) (-\Delta)^s \mathcal{Z}(t, \mathbf{x}) d\mathbf{x} dt. \end{cases}$$

Then,

$$(4.14) \quad \begin{cases} \iint_{Q_T} u_2(t, \mathbf{x}) [-\partial_t \mathcal{Z}(t, \mathbf{x}) + d_2 (-\Delta)^s \mathcal{Z}(t, \mathbf{x})] d\mathbf{x} dt - \iint_{Q_T} u_1(t, \mathbf{x}) \partial_t \mathcal{Z}(t, \mathbf{x}) d\mathbf{x} dt \\ \leq \int_{\Omega} (u_{01} - u_{02})(\mathbf{x}) \mathcal{Z}_0(\mathbf{x}) d\mathbf{x} - d_1 \iint_{Q_T} u_1(t, \mathbf{x}) (-\Delta)^s \mathcal{Z}(t, \mathbf{x}) d\mathbf{x} dt. \end{cases}$$

Therefore,

$$(4.15) \quad \iint_{Q_T} u_2(t, \mathbf{x}) \varphi(t, \mathbf{x}) d\mathbf{x} dt \leq \int_{\Omega} (u_{01} - u_{02})(\mathbf{x}) \mathcal{Z}_0(\mathbf{x}) d\mathbf{x} + \iint_{Q_T} u_1(t, \mathbf{x}) [\partial_t \mathcal{Z}(t, \mathbf{x}) - d_1 (-\Delta)^s \mathcal{Z}(t, \mathbf{x})] d\mathbf{x} dt.$$

Let $p > 1$. By using Hölder's inequality and the estimate (3.4), we obtain

$$(4.16) \quad \iint_{Q_T} u_2(t, \mathbf{x}) \varphi(t, \mathbf{x}) d\mathbf{x} dt \leq C [\|u_{01} - u_{02}\|_{L^p(\Omega)} \|\mathcal{Z}_0\|_{L^{p'}(\Omega)} + (1 + d_1) \|u_1\|_{L^p(Q_T)} \|\varphi\|_{L^{p'}(Q_T)}],$$

where $p' = \frac{p}{p-1}$ and $C := C(p, T, \Omega) > 0$. As φ is regular, we deduce by duality that

$$\|u_2\|_{L^p(Q_T)} \leq C \left(\|u_{01} + u_{02}\|_{L^p(\Omega)} + (1 + d_1) \|u_1\|_{L^p(Q_T)} \right).$$

Hence,

$$\|u_2\|_{L^p(Q_T)} \leq C \max \left\{ \|u_{01} + u_{02}\|_{L^p(\Omega)}, 1 + d_1 \right\} (1 + \|u_1\|_{L^p(Q_T)}),$$

which completes the proof. \square

4.2 First result of global existence : case of reversible chemical reaction

Introducing the fractional counterpart of System (1.3), namely:

$$(4.17) \quad \begin{cases} \partial_t u_1(t, \mathbf{x}) + d_1(-\Delta)^s u_1(t, \mathbf{x}) = f_1(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ \partial_t u_2(t, \mathbf{x}) + d_2(-\Delta)^s u_2(t, \mathbf{x}) = f_2(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ \partial_t u_3(t, \mathbf{x}) + d_3(-\Delta)^s u_3(t, \mathbf{x}) = f_3(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ u_1(t, \mathbf{x}) = u_2(t, \mathbf{x}) = u_3(t, \mathbf{x}) = 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ u_1(0, \mathbf{x}) = u_{01}(\mathbf{x}), & \mathbf{x} \in \Omega, \\ u_2(0, \mathbf{x}) = u_{02}(\mathbf{x}), & \mathbf{x} \in \Omega, \\ u_3(0, \mathbf{x}) = u_{03}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases}$$

where $0 < s < 1$, $f_1 = \alpha_1 g$, $f_2 = \alpha_2 g$, $f_3 = -\alpha_3 g$ with $g = u_3^{\alpha_3} - u_1^{\alpha_1} u_2^{\alpha_2}$ and $\alpha_1, \alpha_2, \alpha_3 \geq 1$.

Our first result of global existence reads as

Theorem 4.1. *Assume that $(u_{01}, u_{02}, u_{03}) \in (L^\infty(\Omega)^+)^3$. System (4.17) admits a unique non-negative global strong solution in the following cases:*

- (i) $\alpha_1 + \alpha_2 < \alpha_3$;
- (ii) $\alpha_3 = 1$ whatever are α_1 and α_2 ;
- (iii) $d_1 = d_3$ or $d_2 = d_3$ whatever are α_1, α_2 and α_3 ;
- (iv) $d_1 = d_2 \neq d_3$ for any $(\alpha_1, \alpha_2, \alpha_3) \in [1, +\infty)^2 \times (1, +\infty)$ such that $\alpha_1 + \alpha_2 \neq \alpha_3$.

Before proceeding to the proof of the previous theorem, it should be mentioned that it is similar to those of [47, Theorems 1 and 3].

Proof. Let $T \in (0, T_{\max})$ and $t \in (0, T]$. Moreover, let us denote by C a positive constant which may be different from line to another.

(i) The case $\alpha_1 + \alpha_2 < \alpha_3$.

Let us begin by noting that f_3 and f_1 (resp. f_2) satisfy **(M)**. In fact, $\alpha_3 f_1 + \alpha_1 f_3 = 0$ and $\alpha_3 f_2 + \alpha_2 f_3 = 0$. Therefore, we deduce from Lemma 4.2 that for any $1 < p < +\infty$

$$(4.18) \quad \|u_1\|_{L^p(Q_T)} \leq C(1 + \|u_3\|_{L^p(Q_T)})$$

and

$$(4.19) \quad \|u_2\|_{L^p(Q_T)} \leq C(1 + \|u_3\|_{L^p(Q_T)}).$$

Therefore, in order to prove the global existence, it is sufficient to prove that $u_3 \in L^p(Q_T)$ for p large enough.

Let $q > 1$. By multiplying the third equation of System (4.17) by $(u_3(t, \mathbf{x}))^q$ and integrating over Q_T , we get

$$(4.20) \quad \begin{cases} \frac{1}{q+1} \int_{\Omega} (u_3(T, \mathbf{x}))^{q+1} d\mathbf{x} + d_3 \iint_{Q_T} [(-\Delta)^s u_3(t, \mathbf{x})] (u_3(t, \mathbf{x}))^q d\mathbf{x} dt + \alpha_3 \iint_{Q_T} (u_3(t, \mathbf{x}))^{q+\gamma} d\mathbf{x} dt \\ = \alpha_3 \iint_{Q_T} (u_1(t, \mathbf{x}))^{\alpha_1} (u_2(t, \mathbf{x}))^{\alpha_2} (u_3(t, \mathbf{x}))^q d\mathbf{x} dt + \frac{1}{q+1} \int_{\Omega} (u_{03}(\mathbf{x}))^{q+1} d\mathbf{x}. \end{cases}$$

By using the the so-called Stroock-Varopoulos inequality (see, for instance, [15, Formula (B7)]), we have

$$(4.21) \quad \iint_{Q_T} [(-\Delta)^s u_3(t, \mathbf{x})] (u_3(t, \mathbf{x}))^q d\mathbf{x} dt \geq \frac{4q}{(q+1)^2} \iint_{Q_T} \left| (-\Delta)^{\frac{s}{2}} (u_3(t, \mathbf{x}))^{\frac{q+1}{2}} \right|^2 d\mathbf{x} dt \geq 0.$$

Furthermore, by using Hölder's inequality, we have

$$(4.22) \quad \iint_{Q_T} (u_1(t, \mathbf{x}))^{\alpha_1} (u_2(t, \mathbf{x}))^{\alpha_2} (u_3(t, \mathbf{x}))^q d\mathbf{x}dt \leq \|u_1\|_{L^{\alpha_1\beta}(Q_T)}^{\alpha_1} \|u_2\|_{L^{\alpha_2\gamma}(Q_T)}^{\alpha_2} \|u_3\|_{L^{\alpha_3+q}(Q_T)}^q,$$

where

$$\frac{1}{\beta} + \frac{1}{\gamma} + \frac{q}{q + \alpha_3} = 1.$$

Since $\alpha_1 + \alpha_2 < \alpha_3$, we can choose β such that $\beta\alpha_1 \leq q + \alpha_3$ and γ such that $\gamma\alpha_2 \leq q + \alpha_3$. Then, $L^{q+\alpha_3}(Q_T) \subset L^{\alpha_1\beta}(Q_T)$ and $L^{q+\alpha_3}(Q_T) \subset L^{\alpha_2\gamma}(Q_T)$. Consequently, we have

$$(4.23) \quad \iint_{Q_T} (u_1(t, \mathbf{x}))^{\alpha_1} (u_2(t, \mathbf{x}))^{\alpha_2} (u_3(t, \mathbf{x}))^q d\mathbf{x}dt \leq C \|u_1\|_{L^{\alpha_3+q}(Q_T)}^{\alpha_1} \|u_2\|_{L^{\alpha_3+q}(Q_T)}^{\alpha_2} \|u_3\|_{L^{\alpha_3+q}(Q_T)}^q.$$

By using the same concept of duality as of Lemma 4.2, we obtain

$$(4.24) \quad \|u_1\|_{L^{\alpha_3+q}(Q_T)} \leq C \left(1 + \|u_3\|_{L^{\alpha_3+q}(Q_T)}\right)$$

and

$$(4.25) \quad \|u_2\|_{L^{\alpha_3+q}(Q_T)} \leq C \left(1 + \|u_3\|_{L^{\alpha_3+q}(Q_T)}\right).$$

Thanks to (4.24) and (4.25), Formula (4.23) implies that

$$(4.26) \quad \begin{cases} \iint_{Q_T} (u_1(t, \mathbf{x}))^{\alpha_1} (u_2(t, \mathbf{x}))^{\alpha_2} (u_3(t, \mathbf{x}))^q d\mathbf{x}dt \\ \leq C \left(1 + \|u_3\|_{L^{\alpha_3+q}(Q_T)}\right)^{\alpha_1} \left(1 + \|u_3\|_{L^{\alpha_3+q}(Q_T)}\right)^{\alpha_2} \left(1 + \|u_3\|_{L^{\alpha_3+q}(Q_T)}\right)^q. \end{cases}$$

If $\|u_3\|_{L^{\alpha_3+q}(Q_T)} \leq 1$ then the proof ends up. Otherwise,

$$(4.27) \quad \iint_{Q_T} (u_1(t, \mathbf{x}))^{\alpha_1} (u_2(t, \mathbf{x}))^{\alpha_2} (u_3(t, \mathbf{x}))^q d\mathbf{x}dt \leq C \|u_3\|_{L^{\alpha_3+q}(Q_T)}^{q+\alpha_1+\alpha_2}.$$

Hence, we have by (4.20)

$$(4.28) \quad \iint_{Q_T} (u_3(t, \mathbf{x}))^{q+\alpha_3} \leq C \|u_3\|_{L^{\alpha_3+q}(Q_T)}^{q+\alpha_1+\alpha_2} + \frac{1}{q+1} \int_{\Omega} (u_{03}(\mathbf{x}))^{q+1} d\mathbf{x}.$$

Therefore, by using Young's inequality, we get

$$(4.29) \quad \|u_3\|_{L^{\alpha_3+q}(Q_T)} \leq C.$$

By choosing $\frac{\alpha_3 + q}{\alpha_3} > \frac{N + 2s}{2s}$, we have by Theorem 3.1

$$(4.30) \quad \|u_1\|_{L^\infty(Q_T)} \leq C$$

and

$$(4.31) \quad \|u_2\|_{L^\infty(Q_T)} \leq C.$$

Now going back to the third equation of (4.17), we deduce from (4.30) and (4.31) that

$$(4.32) \quad \|u_3\|_{L^\infty(Q_T)} \leq C,$$

which implies that $T_{\max} = +\infty$.

(ii) **The case $\alpha_3 = 1$.** From the semigroup property and thanks to comparison principle, it holds for any $p > 1$

$$(4.33) \quad \|u_1(t, \cdot)\|_{L^p(\Omega)} \leq \|u_0\|_{L^p(\Omega)} + \int_0^t \|u_3(\tau, \cdot)\|_{L^p(\Omega)} d\tau.$$

By applying Hölder's inequality and the same duality arguments as (4.18), we obtain

$$(4.34) \quad \int_0^t \|u_3(\tau, \cdot)\|_{L^p(\Omega)} d\tau \leq Ct^{1/p'} \left[1 + \left(\int_0^t \int_{\Omega} (u_1(\tau, \mathbf{x}))^p d\tau d\mathbf{x} \right)^{1/p} \right],$$

where $p' = \frac{p}{p-1}$. For any $t \in (0, T]$, let us set $h(t) := \int_{\Omega} |u_1(t, \mathbf{x})|^p d\mathbf{x}$. Then, (4.33) implies that

$$(4.35) \quad h(t)^{\frac{1}{p}} \leq C \left[1 + \left(\int_0^t h(\tau) d\tau \right)^{\frac{1}{p}} \right].$$

By taking the p th power of (4.35), we obtain

$$(4.36) \quad h(t) \leq C \left[1 + \int_0^t h(\tau) d\tau \right].$$

Therefore, by applying Gronwall's Lemma, we get

$$(4.37) \quad \|u_1\|_{L^p(Q_T)} \leq C.$$

Repeating the method above with u_2 instead of u_1 , we obtain

$$(4.38) \quad \|u_2\|_{L^p(Q_T)} \leq C.$$

Estimates (4.37) and (4.38) imply that for some $q > \frac{N+2s}{2s}$

$$(4.39) \quad \|u_1^{\alpha_1} u_2^{\alpha_2}\|_{L^q(Q_T)} \leq C.$$

Going back to the third equation of System (4.17), we have by Theorem 3.1

$$(4.40) \quad \|u_3\|_{L^\infty(Q_T)} \leq C.$$

Hence, $T_{\max} = +\infty$ which completes the proof in this case.

(iii) and (iv). The proofs in these two cases are straightforward and left to the reader. □

4.3 Second result of global existence : case of triangular structure

Our second result of global existence reads as

Theorem 4.2. *Let $(u_{01}, \dots, u_{0m}) \in (L^\infty(\Omega)^+)^m$ where $m > 2$. Besides **(P)**, assume that f_i is at most polynomial for each $i = 1, \dots, m$. Moreover, assume that there exist a vector $\mathbf{b} \in [0, +\infty)^m$, and a lower triangular invertible matrix $Q \in \mathcal{M}_m([0, +\infty))$, such that*

$$(4.41) \quad \forall \mathbf{r} = (r_1, \dots, r_m) \in [0, +\infty)^m, \quad Q\mathbf{f}(\mathbf{r}) \leq \left[1 + \sum_{i=1}^m r_i \right] \mathbf{b}.$$

Then, System (S) admits a unique nonnegative global strong solution.

Proof. Let us denote $Q = (q_{ij})_{i,j=1,\dots,m}$, $\mathbf{b} = (b_1, \dots, b_m)^T$ and $\mathbf{f}(\mathbf{r}) = (f_1(\mathbf{r}), \dots, f_m(\mathbf{r}))^T$. By Lemma 4.1, System (S) admits a unique strong solution \mathbf{u} on $[0, T_{\max}) \times \Omega$. In what follows, we prove that this solution is global, *i.e.* $T_{\max} = +\infty$. Let us denote by C a positive constant which may be different from line to another. Moreover, let $T \in (0, T_{\max})$ and $t \in (0, T]$. For each $i = 1, \dots, m$, we denote by w_i the weak solution to

$$(4.42) \quad \begin{cases} \partial_t w_i(t, \mathbf{x}) + d_i(-\Delta)^s w_i(t, \mathbf{x}) &= \left[1 + \sum_{i=1}^m u_i(t, \mathbf{x})\right] \frac{b_i}{q_{ii}}, & (t, \mathbf{x}) \in Q_T, \\ w_i(t, \mathbf{x}) &= 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ w_i(0, \mathbf{x}) &= 0, & \mathbf{x} \in \Omega. \end{cases}$$

Then, w_i fulfills

$$(4.43) \quad w_i(t, \mathbf{x}) = \int_0^t S_i(t - \tau) \left[1 + \sum_{i=1}^m u_i(\tau, \mathbf{x})\right] \frac{b_i}{q_{ii}} d\tau, \quad (t, \mathbf{x}) \in [0, T) \times \Omega.$$

According to (4.41), we have

$$(4.44) \quad \sum_{j=1}^i q_{ij} f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})) \leq \left[1 + \sum_{j=1}^m u_j(t, \mathbf{x})\right] b_i.$$

• The case $i = 1$. We subtract the first equation of (4.42) from the first equation of (S) multiplied by q_{11} . Then,

$$(4.45) \quad q_{11} \left[\partial_t (u_1 - w_1)(t, \mathbf{x}) + d_1(-\Delta)^s (u_1 - w_1)(t, \mathbf{x}) \right] \leq 0.$$

Then, by Theorem 3.2, we get

$$\|u_1 - w_1\|_{L^\infty(Q_T)} \leq \|u_{01}\|_{L^\infty(\Omega)}.$$

Therefore,

$$\|u_1\|_{L^\infty(Q_T)} \leq C[1 + \|w_1\|_{L^\infty(Q_T)}].$$

By taking the $L^\infty(\Omega)$ -norm of (4.43) and using the L^∞ -contractivity of $\{S_A(t)\}_{t \geq 0}$, we get

$$(4.46) \quad \|u_1(t, \cdot)\|_{L^\infty(\Omega)} \leq C \left[1 + \int_0^t \left\| \sum_{i=1}^m u_i(\tau, \cdot) \right\|_{L^\infty(\Omega)} d\tau \right].$$

Hence,

$$(4.47) \quad \|u_1(t, \cdot)\|_{L^\infty(\Omega)} \leq C \left[1 + \int_0^t \|u_1(\tau, \cdot)\|_{L^\infty(\Omega)} d\tau + \int_0^t \left\| \sum_{i=2}^m u_i(\tau, \cdot) \right\|_{L^\infty(\Omega)} d\tau \right].$$

• The case $i \geq 2$. By subtracting the i th equation of (4.42) from the i th equation of (S) multiplied by q_{ii} , we obtain

$$(4.48) \quad q_{ii} \left[\partial_t (u_i - w_i)(t, \mathbf{x}) + d_i(-\Delta)^s (u_i - w_i)(t, \mathbf{x}) \right] \leq - \sum_{j=1}^{i-1} q_{ij} \left[\partial_t u_j(t, \mathbf{x}) + d_j(-\Delta)^s u_j(t, \mathbf{x}) \right].$$

Now, we use the same concept of duality as of Lemma 4.2. More precisely, let φ be a nonnegative regular function and let \mathcal{Z} be the solution to Problem (P_φ) with $d = \frac{d_i}{q_{ii}}$. Then, we multiply the two sides of the inequality (4.48) by \mathcal{Z} and integrate over Q_t . We obtain

$$(4.49) \quad \begin{cases} \iint_{Q_t} q_{ii} \partial_t (u_i - w_i)(\tau, \mathbf{x}) \mathcal{Z}(\tau, \mathbf{x}) d\tau d\mathbf{x} + \iint_{Q_t} d_i(-\Delta)^s (u_i - w_i)(\tau, \mathbf{x}) \mathcal{Z}(\tau, \mathbf{x}) d\tau d\mathbf{x} := I_1 \\ \leq - \sum_{j=1}^{i-1} q_{ij} \iint_{Q_t} \partial_t u_j(\tau, \mathbf{x}) \mathcal{Z}(\tau, \mathbf{x}) d\tau d\mathbf{x} - \sum_{j=1}^{i-1} q_{ij} d_j \iint_{Q_t} (-\Delta)^s u_j(\tau, \mathbf{x}) \mathcal{Z}(\tau, \mathbf{x}) d\tau d\mathbf{x} := I_2. \end{cases}$$

Now, let us simplify the terms I_1 and I_2 .
We have

$$(4.50) \quad I_1 = q_{ii} \iint_{Q_t} (u_i - w_i)(\tau, \mathbf{x}) \varphi(\tau, \mathbf{x}) d\tau d\mathbf{x} - q_{ii} \int_{\Omega} u_{0i}(\mathbf{x}) \mathcal{Z}_0(\mathbf{x}) d\mathbf{x}.$$

In addition, we have

$$(4.51) \quad I_2 = \sum_{j=1}^{i-1} q_{ij} \iint_{Q_t} u_j(\tau, \mathbf{x}) (\partial_t \mathcal{Z}(\tau, \mathbf{x}) - d_j(-\Delta)^s \mathcal{Z}(\tau, \mathbf{x})) d\tau d\mathbf{x} + \sum_{j=1}^{i-1} q_{ij} \int_{\Omega} u_{0j}(\mathbf{x}) \mathcal{Z}_0(\mathbf{x}) d\mathbf{x}.$$

Then, (4.49) implies

$$(4.52) \quad \begin{cases} q_{ii} \iint_{Q_t} (u_i - w_i)(\tau, \mathbf{x}) \varphi(\tau, \mathbf{x}) d\tau d\mathbf{x} \\ \leq \sum_{j=1}^i q_{ij} \int_{\Omega} u_{0j}(\mathbf{x}) \mathcal{Z}_0(\mathbf{x}) d\mathbf{x} + \sum_{j=1}^{i-1} q_{ij} \iint_{Q_t} u_j(\tau, \mathbf{x}) (\partial_t \mathcal{Z}(\tau, \mathbf{x}) - d_j(-\Delta)^s \mathcal{Z}(\tau, \mathbf{x})) d\tau d\mathbf{x}. \end{cases}$$

Let $p > 1$ large enough and $p' = \frac{p}{p-1}$. By using Hölder's inequality and the estimate (3.4), we get

$$(4.53) \quad \begin{cases} \iint_{Q_t} (u_i - w_i)(\tau, \mathbf{x}) \varphi(\tau, \mathbf{x}) d\tau d\mathbf{x} \\ \leq C \sum_{j=1}^i \|u_{0j}\|_{L^p(\Omega)} \|\mathcal{Z}_0\|_{L^{p'}(\Omega)} + C \sum_{j=1}^{i-1} \|u_j\|_{L^p(Q_t)} \left(\|\partial_t \mathcal{Z}\|_{L^{p'}(Q_t)} + d_j \|(-\Delta)^s \mathcal{Z}\|_{L^{p'}(Q_t)} \right) \\ \leq C \sum_{j=1}^i \|u_{0j}\|_{L^p(\Omega)} \|\varphi\|_{L^{p'}(Q_t)} + C \sum_{j=1}^{i-1} \|u_j\|_{L^p(Q_t)} \|\varphi\|_{L^{p'}(Q_t)} \\ \leq C \left(1 + \sum_{j=1}^{i-1} \|u_j\|_{L^p(Q_t)} \right) \|\varphi\|_{L^{p'}(Q_t)}. \end{cases}$$

As φ is regular, we obtain by duality that

$$(4.54) \quad \|u_i - w_i\|_{L^p(Q_t)} \leq C \left(1 + \sum_{j=1}^{i-1} \|u_j\|_{L^p(Q_t)} \right).$$

Hence, by an elementary induction on i , we obtain

$$(4.55) \quad \|u_i\|_{L^p(Q_t)} \leq C \left(1 + \sum_{j=1}^i \|u_j\|_{L^p(Q_t)} \right).$$

Consequently, by summing over i and taking the p th power, we obtain

$$(4.56) \quad \left\| \sum_{i=2}^m u_i \right\|_{L^p(Q_t)}^p \leq C \left(1 + \sum_{i=2}^m \|u_i\|_{L^p(Q_t)}^p \right).$$

Now, let us take the $L^p(\Omega)$ -norm of (4.43). Then, by using the estimate (2.5) with $q = p$ and Hölder's inequality, we get

$$(4.57) \quad \|w_i(t, \cdot)\|_{L^p(\Omega)}^p \leq C \left[1 + \left\| \sum_{i=2}^m u_i \right\|_{L^p(Q_t)}^p \right].$$

Hence, we obtain by (4.56)

$$(4.58) \quad \sum_{i=2}^m \|w_i(t, \cdot)\|_{L^p(\Omega)}^p \leq C \left[1 + \int_0^t \sum_{i=2}^m \|w_i(\tau, \cdot)\|_{L^p(\Omega)}^p d\tau \right].$$

By applying Gronwall's Lemma, we get

$$\sum_{i=2}^m \|w_i(t, \cdot)\|_{L^p(\Omega)}^p < g(t),$$

where $g(t) : [0, +\infty) \rightarrow [0, +\infty)$ is a non-decreasing and continuous function. Thus, for each $i = 2, \dots, m$, $w_i(t, \cdot)$ is bounded in $L^\infty(\Omega)$, and so is $\sum_{i=2}^m w_i(t, \cdot)$ thanks to (4.56). By hypothesis, f_i is at most polynomial. Then, we deduce that $f_i \in L^q(Q_T)$ where q is large enough. Then, we can choose $q > \frac{N+2s}{2s}$ so that we have by Theorem 3.1:

$$\|u_i\|_{L^\infty(Q_T)} \leq C \quad \forall i = 2, \dots, m.$$

Going back to (4.47), we also apply Gronwall's lemma to get $\|u_1\|_{L^\infty(Q_T)} \leq C$. Consequently, $T_{\max} = +\infty$. \square

5 Reaction-Diffusion System with exponential growth : Numerical simulations

The main objective of this section is to examine numerically the existence of strong solutions to the following system :

$$(S_{\text{exp}}) \quad \begin{cases} \partial_t u(t, \mathbf{x}) + d_1(-\Delta)^s u(t, \mathbf{x}) &= -u(t, \mathbf{x})e^{(v(t, \mathbf{x}))^\beta}, & (t, \mathbf{x}) \in Q_T, \\ \partial_t v(t, \mathbf{x}) + d_2(-\Delta)^s v(t, \mathbf{x}) &= u(t, \mathbf{x})e^{(v(t, \mathbf{x}))^\beta}, & (t, \mathbf{x}) \in Q_T, \\ u(t, \mathbf{x}) = v(t, \mathbf{x}) &= 0, & (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ u(0, \mathbf{x}) &= u_0(\mathbf{x}), & \mathbf{x} \in \Omega, \\ v(0, \mathbf{x}) &= v_0(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases}$$

where $d_1, d_2 > 0$, $u_0, v_0 \in (L^\infty(\Omega))^+$ in the case where $\beta > 1$.

As stated in the introduction, the global existence of strong solutions to (S_{exp}) remains an open problem even for the classical case ($s = 1$). In [48], the second author and his coworkers have investigated numerically this open problem in the classical case (see also [57]). In the present paper, we use a different approach adapted to the fractional Laplacian in order to treat System (S_{exp}) . More precisely, we perform our numerical simulations for 2D-case by considering the unit ball $\Omega = B(0, 1) \subset \mathbb{R}^2$. Furthermore, we carry out these simulations using :

- the Crank–Nicolson scheme to approximate the time derivative;
- the finite element method (FEM) to discretise in space and approximate the solution. To realize this, we rely on [3] where the authors proposed a FEM adapted to the fractional case as well as a Matlab code in 2D-case. Notably, we use this code and modify it to adapt it to our problem;
- the fixed-point iteration method to solve the resulting nonlinear equations at each discrete time.

As in [3], we consider an admissible triangulation of $\bar{\Omega}$ and we denote by h its mesh size. Regarding the temporal discretization, let $0 = t_0 < t_1 < t_2 < \dots < t_{\bar{N}} = T$ be a partition of the interval $(0, T)$ with time step $h_t = t_{n+1} - t_n$. As for the fixed-point iterative processes, they are performed under the control of a tolerance level equal to 10^{-6} .

Before going further, we would like to point out that we use numerical simulations to build a conjecture about the existence of solutions to System (S_{exp}) , without conducting any theoretical study of errors. That being said, it should be recalled that the Crank-Nicolson scheme is a second-order numerical method (see, for instance, [60, Chapter 2]). Moreover, according to [3], the order of convergence of the FEM in the linear case (Poisson equation for example) depends on the value of s . Namely, for $0 < s \leq \frac{1}{2}$, the expected order of convergence with respect to h is $s + \frac{1}{2}$, while for $\frac{1}{2} < s < 1$, the expected order of convergence is 1. On the other hand, let us comment on the convergence of the fixed-point iterations. Although its theoretical analysis is limited in our case, we are able to numerically estimate the constants of contraction at each iteration and verify that they remain < 1 . Furthermore, it should be stressed that we present here a partial numerical investigation of System (S_{exp}) . For more details, we refer the interested reader to [24].

Now, let us present some numerical simulations that illustrate the performance of our numerical scheme. To do so, let us consider the function $\rho(\mathbf{x}) = (1 - \|\mathbf{x}\|^2)^s$. It is known that $(-\Delta)^s \rho(\mathbf{x}) = \lambda(s)$, where $\lambda(s) := 4^s \Gamma(s+1)^2$ (see [28]). We use this fact to construct a solution to System (S_{exp}) with additional terms on the right-hand sides (r.h.s.). Specifically, the couple $(u(t, \mathbf{x}), v(t, \mathbf{x})) = (1, 1/2)\rho(\mathbf{x})e^{-t}$ solves the system with the following r.h.s. terms:

$$\begin{cases} \text{r.h.s.1} &= -u(t, \mathbf{x})e^{v(t, \mathbf{x})^\beta} + (d_1 \lambda(s) - \rho(\mathbf{x}) + \rho(\mathbf{x}) \exp[2^{-\beta} e^{-\beta t} \rho(\mathbf{x})^\beta]) e^{-t} \\ \text{r.h.s.2} &= u(t, \mathbf{x})e^{v(t, \mathbf{x})^\beta} + (1/2 d_2 \lambda(s) - 1/2 \rho(\mathbf{x}) - \rho(\mathbf{x}) \exp[2^{-\beta} e^{-\beta t} \rho(\mathbf{x})^\beta]) e^{-t}. \end{cases}$$

To assess the accuracy of our scheme, we compare the approximated solution (denoted with the subscript “ h ”) to the exact one (denoted with the subscript “ ex ”). Furthermore, we denote by U_h and V_h the last vectors obtained at a certain time after performing the necessary fixed-point iterations. We also denote by U_{ex} and V_{ex} the vectors which contain the exact solution at the same nodes as U_h and V_h respectively. In what follows, we use these notations.

Moreover, we consider different values of s , d_1 , d_2 and β . In fact, for $s = 0.25$ we fix $(d_1, d_2, \beta) = (1, 3, 2)$, for $s = 0.5$ we fix $(d_1, d_2, \beta) = (2, 1, 3)$, for $s = 0.75$ we fix $(d_1, d_2, \beta) = (1, 2, 3)$ and for $s = 0.9$ we fix $(d_1, d_2, \beta) = (3, 4, 2)$.

In **Figure 1**, we have plotted the evolution in 1D of U_h and V_h for $s = 0.75$ over time for $t = 1, 2, 3, 4, 5$ by fixing the y -axis at 0.

Figures 2 and **3** show the comparison graphs between the numerical and exact solutions for $s = 0.75$ at $t = 5$.




Figure 1. Evolution in time of U_h and V_h for $s = 0.75$ and $h = \frac{1}{48}$ with the y -axis fixed at 0




Figure 2. Comparison between U_h and U_{ex} for $s = 0.75$ and $h = \frac{1}{48}$




Figure 3. Comparison between V_h and V_{ex} for $s = 0.75$ and $h = \frac{1}{48}$ at $t = 5$

In order to verify the order of convergence of the FEM in the presented scheme, we have plotted in **Figure 4** the evolution of the L^2 -error in Ω between the numerical and exact solutions for different values of s .




Figure 4. Order of convergence for $h = \frac{1}{8}, \frac{1}{12}, \frac{1}{16}, \frac{1}{24}, \frac{1}{32}, \frac{1}{48}$

As shown in **Figure 4**, our computed convergence orders for different values of s match the theoretical expectations reported in [3].

Now, after testing the performance of our scheme, let us go back to our System (S_{exp}) with $u_0(\mathbf{x}) = \rho(\mathbf{x})$ and $v_0(\mathbf{x}) = 1/2\rho(\mathbf{x})$. Our main focus is to investigate the existence of a nonnegative numerical solution to System (S_{exp}) that may be globally time. To this end, we aim to numerically solve the system until a sufficiently large final time ($t = 5 \times 10^{10}$ in our case).

Figures 5 and **6** show the graphs of the numerical solutions obtained at $t = 5 \times 10^{10}$ for $s = 0.75$ and different values of d_1, d_2 and β by fixing $h_t = 10^{-2}$ and $h = \frac{1}{48}$.




Figure 5. Plots of U_h and V_h for $s = 0.75$ and $(d_1, d_2, \beta) = (1, 2, 3)$ at $t = 5 \times 10^{10}$




Figure 6. Plots of U_h and V_h for $s = 0.75$ and $(d_1, d_2, \beta) = (4, 3, 5)$ at $t = 5 \times 10^{10}$

In addition to the numerical solution plots, we also compute the l^∞ -norm of U_h and V_h at $t = 5 \times 10^{10}$ for different values of h . More precisely, we have fixed the time step h_t to 10^{-2} and varied the mesh size h . The resulting norms are summarized in **Table 1**. The purpose of this table is to ensure that the solution does not change significantly with respect to the mesh size. As shown in the table, the norms remain relatively constant as the mesh size is varied. This indicates that the numerical solution is consistent.

	$d_1 = 1, d_2 = 2, \beta = 3$		$d_1 = 4, d_2 = 3, \beta = 5$	
h	$\ U_h\ _\infty$	$\ V_h\ _\infty$	$\ U_h\ _\infty$	$\ V_h\ _\infty$
1/8	0.43189	0.42518	0.09623	0.15000
1/12	0.43221	0.42623	0.09630	0.15007
1/16	0.43246	0.42701	0.09635	0.15012
1/24	0.43271	0.42772	0.09637	0.15016
1/32	0.43285	0.42887	0.09643	0.15022
1/48	0.43292	0.42853	0.09645	0.15028

Table 1. Table of ℓ^∞ -errors of U_h and V_h for $s = 0.75$ at $t = 5 \times 10^{10}$

In conclusion, we have studied the global existence of nonnegative numerical solutions to the fractional reaction-diffusion system (S_{exp}) . Our numerical experiments have showed that the system has nonnegative numerical solutions that can be computed for a large final time. This indicates that a global solution in time to System (S_{exp}) seems to exist. However, it is not clear

whether this solution will exist for all times. In future work, we aim at investigating this issue theoretically.

6 Appendix

In this section, we give the proofs of Theorems 3.1 and 3.2. For the convenience of the reader, we recall their statements.

Theorem 3.1. *Assume that $w_0 \in L^\infty(\Omega)$ and $h \in L^p(Q_T)$ with $p > 1$. Let w be the weak solution to Problem (3.1). Then, for any $p > \frac{N+2s}{2s}$, there exists a constant $C > 0$ such that*

$$\|w\|_{L^\infty(Q_T)} \leq \|w_0\|_{L^\infty(\Omega)} + C\|h\|_{L^p(Q_T)}.$$

Proof. For any $t \geq 0$, the unique weak solution to Problem (3.1) reads as :

$$(3.4) \quad w(t, \cdot) = S_A(t)w_0 + \int_0^t S_A(t-\tau)f(\tau, \cdot)d\tau.$$

As $\{S_A(t)\}_{t \geq 0}$ is L^∞ -contractive, we obtain

$$\|w(t, \cdot)\|_{L^\infty(\Omega)} \leq \|w_0\|_{L^\infty(\Omega)} + \int_0^t \|S_A(t-\tau)f(\tau, \cdot)\|_{L^\infty(\Omega)}d\tau.$$

According to Theorem 2.4, the semigroup $\{S_A(t)\}_{t \geq 0}$ is ultracontractive. Thus, for $q = +\infty$ and for any $p > 1$, there exists a constant $C > 0$ such that

$$(3.5) \quad \|S_A(t-\tau)f(\tau, \cdot)\|_{L^\infty(\Omega)} \leq C(t-\tau)^{-\frac{N}{2sp}}\|f(\tau, \cdot)\|_{L^p(\Omega)}, \quad \text{for any } \tau \in (0, t).$$

Therefore, we get

$$\|w(t, \cdot)\|_{L^\infty(\Omega)} \leq \|w_0\|_{L^\infty(\Omega)} + C \int_0^t (t-\tau)^{-\frac{N}{2sp}}\|f(\tau, \cdot)\|_{L^p(\Omega)}d\tau.$$

Now, by using Hölder's inequality, we obtain

$$\|w(t, \cdot)\|_{L^\infty(\Omega)} \leq \|w_0\|_{L^\infty(\Omega)} + C \left(\int_0^t (t-\tau)^{-\frac{N}{2s(p-1)}}d\tau \right)^{\frac{p-1}{p}} \|f\|_{L^p(Q_T)}.$$

The quantity $\int_0^t (t-\tau)^{-\frac{N}{2s(p-1)}}d\tau$ is finite if and only if $p > \frac{N+2s}{2s}$. Hence , we get the desired result. \square

Theorem 3.2. *Assume that $w_0 \in L^\infty(\Omega)$ and $h \in L^p(Q_T)$ with $p > 1$. Let w be the weak solution to Problem (3.1). If $h \leq 0$, then $w \in L^\infty(Q_T)$ and*

$$\|w\|_{L^\infty(Q_T)} \leq \|w_0\|_{L^\infty(\Omega)}.$$

Proof. The idea of the proof is similar to the proof of [34, Proposition 3.18].

Let us define $v := \|w_0\|_{L^\infty(\Omega)} - w$ and write $v = v^+ + v^-$, where $v^+ = \max\{v, 0\}$ and $v^- = \min\{v, 0\}$.

We will proceed by contradiction. Suppose that the set $\{\mathbf{x} \in \mathbb{R}^N : v^-(t, \mathbf{x}) \neq 0 \ \forall t \in (0, T)\}$ is nonempty.

Let us multiply the first equation of Problem (3.1) by v^- and integrate over \mathbb{R}^N then over $(0, T)$. Then, we obtain

$$\begin{aligned} \int_0^T \int_{\Omega} \partial_t w(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt + \frac{a_{N,s}}{2} \int_0^T \int \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{(w(t, \mathbf{x}) - w(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y}))}{\|\mathbf{x} - \mathbf{y}\|^{N+2s}} d\mathbf{x} d\mathbf{y} dt \\ = \int_0^T \int_{\Omega} f(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt. \end{aligned}$$

For any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^N$, we have

$$\begin{aligned} (w(t, \mathbf{x}) - w(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y})) \\ = (w(t, \mathbf{x}) - \|w\|_{L^\infty(\Omega)} + \|w\|_{L^\infty(\Omega)} - w(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y})) \\ = (-v(t, \mathbf{x}) + v(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y})) \\ = (-v^+(t, \mathbf{x}) + v^-(t, \mathbf{x}) + v^+(t, \mathbf{y}) - v^-(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y})) \\ = (v^-(t, \mathbf{x}) - v^-(t, \mathbf{y}))^2 - (v^+(t, \mathbf{x}) - v^+(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y})). \end{aligned}$$

Moreover, we know that if $v^- > 0$ then $v^+ = 0$, and if $v^- = 0$ then $v^+ > 0$. Therefore,

$$v^+(t, \mathbf{y})(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y})) \geq 0,$$

This implies

$$(w(t, \mathbf{x}) - w(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y})) \geq 0.$$

Thence, we get

$$\frac{a_{N,s}}{2} \int_0^T \int \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{(w(t, \mathbf{x}) - w(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y}))}{\|\mathbf{x} - \mathbf{y}\|^{N+2s}} d\mathbf{x} d\mathbf{y} dt \geq 0.$$

As $w = \|w_0\|_{L^\infty(\Omega)} - v$, then $\partial_t w = -\partial_t v = -\partial_t v^+ + \partial_t v^-$ and

$$\int_0^T \int_{\Omega} \partial_t w(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt = \int_0^T \int_{\Omega} \partial_t v^-(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt - \int_0^T \int_{\Omega} \partial_t v^+(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt.$$

Furthermore, we have $\int_0^T \int_{\Omega} \partial_t v^+(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt = 0$, then

$$\begin{aligned} \int_0^T \int_{\Omega} \partial_t w(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt &= \frac{1}{2} \int_{\Omega} \int_0^T \partial_t (v^-(t, \mathbf{x}))^2 dt d\mathbf{x} \\ &= \frac{1}{2} \int_{\Omega} ((v^-(T, \mathbf{x}))^2 - (v^-(0, \mathbf{x}))^2) d\mathbf{x} \\ &= \frac{1}{2} \int_{\Omega} (v^-(T, \mathbf{x}))^2 d\mathbf{x} \geq 0, \end{aligned}$$

as $v^-(0, \mathbf{x}) = 0$.

Now, we have

$$\int_0^T \int_{\Omega} \partial_t w(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt + \frac{a_{N,s}}{2} \int_0^T \int \int_{\mathbb{R}^N \times \mathbb{R}^N} \frac{(w(t, \mathbf{x}) - w(t, \mathbf{y}))(v^-(t, \mathbf{x}) - v^-(t, \mathbf{y}))}{\|\mathbf{x} - \mathbf{y}\|^{N+2s}} d\mathbf{x} d\mathbf{y} dt \geq 0,$$

which is a contradiction as $f \leq 0$ and $\int_0^T \int_{\Omega} f(t, \mathbf{x}) v^-(t, \mathbf{x}) d\mathbf{x} dt \leq 0$. This completes the proof. \square

References

- [1] N. Abatangelo, E. Valdinoci: *Getting Acquainted with the Fractional Laplacian*. In: Dipierro, S. (eds) *Contemporary Research in Elliptic PDEs and Related Topics*. Springer INdAM Series 33. Springer, 2019.
- [2] B. Abdellaoui, A. Attar, R. Bentifour, E.-H. Laamri: *Existence results to a class of non-linear parabolic systems involving potential and gradient terms*. *Mediterranean Journal of Mathematics* 17(119) (2020), 1–30.
- [3] G. Acosta, F.M. Bersetche, J.P. Borthagaray: *A short FE implementation for a 2d homogeneous Dirichlet problem of a fractional Laplacian*. *Comput. Math. Appl.* 74(4) (2017), 784–816.
- [4] B. Ahmad, M.S. Alhothuali, H.H. Alsulami, M. Kirane, S. Timoshin: *On nonlinear nonlocal systems of reaction diffusion equations*. *Abstr. Appl. Anal.* 2 (2014), 1–6.
- [5] B. Ahmad, A. Alsaedi, D. Hnaien, M. Kirane: *On a semi-linear system of nonlocal time and space reaction diffusion equations with exponential nonlinearities*. *J. Integral Equations and Applications* 30 (2018), 17–40.
- [6] A. Alsaedi, M. Al-Yami, M. Kirane, F. Momenkhan: *A triangular nonlinear reaction-fractional diffusion system with a balance law*. *Math. Methods Appl. Sci.* 41 (2018), 1825–1830.
- [7] F. Andreu-Vaillo, J.M. Mazón, J.D. Rossi, J.J. Toledo-Melero: *Nonlocal Diffusion Problems*. *Mathematical Surveys and Monographs*, 165, American Mathematical Society, Providence, RI; Real Sociedad Matemática Española, Madrid, 2010.
- [8] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri: *On some nonlocal parabolic reaction-diffusion systems with gradient source terms*. Submitted.
- [9] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri: *On some nonlocal elliptic systems with gradient source terms*. *Acta Appl. Math.* 181(9) (2022). Doi.org/10.1007/s10440-022-00528-4.
- [10] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri: *On a class of some fractional parabolic systems with potential-gradient source terms*. Submitted.
- [11] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri: *Fractional elliptic reaction-diffusion systems with coupled gradient terms and different diffusion*. Submitted.
- [12] A. Barabanova: *On the global existence of solutions of a reaction-diffusion system with exponential nonlinearity*. *Proc. Amer. Math. Soc.* 122 (1994), 827–831.
- [13] U. Biccari, M. Warma, E. Zuazua: *Local Regularity for Fractional Heat Equations*. *Recent Advances in PDEs: Analysis, Numerics and Control*, SEMA SIMAI Springer Ser. 17, Springer, Cham, 2018, 233–249.
- [14] U. Biccari, M. Warma, E. Zuazua: *Control and numerical approximation of fractional diffusion equations*. *Handbook of Numerical Analysis XXIII. Numerical Control: Part A*, 23 (2022), 1–58.
- [15] M. Bonforte, A. Figalli, X. Ros-Oton: *Infinite Speed of Propagation and Regularity of Solutions to the Fractional Porous Medium Equation in General Domains*. *Commun. Pure Appl. Math.* 70(8) (2017), 1472–1508.
- [16] A. Bonito, W. Lei, J.E. Pasciak: *Numerical approximation of the integral fractional Laplacian*. *Numerische Mathematik* 142 (2019), 235–278.

- [17] N. Boudiba: *Existence globale pour des systèmes de réaction-diffusion avec contrôle de masse*. Ph.D. thesis, Université de Rennes 1, France, 1999.
- [18] C. Bucur, E. Valdinoci: *Nonlocal diffusion and applications*. ISBN 978-3-319-28738-6. Lecture Notes of the Unione Matematica Italiana, 2016.
- [19] J.A. Cañizo, L. Desvillettes, K. Fellner: *Improved Duality Estimates and Applications to Reaction-Diffusion Equations*. Commun. Partial Differ. Equ. 39(6) (2014), 1185–1204.
- [20] C. Caputo, T. Goudon, A. Vasseur: *Solutions of the 4-species quadratic reaction-diffusion system are bounded and C^∞ -smooth, in any space dimension*. Anal. PDE 12(7) (2019), 1773–1804.
- [21] B. Claus, M. Warma: *Realization of the fractional Laplacian with nonlocal exterior conditions via forms method*. J. Evol. Equ. 20 (2020), 1597–1631.
- [22] G. Ciavolella, B. Perthame: *Existence of a global weak solution for a reaction–diffusion problem with membrane conditions*. J. Evol. Equ. 21 (2021), 1513–1540.
- [23] A. Cotsiolis, N.K. Tavoularis: *Best constants for Sobolev inequalities for higher order fractional derivatives*. J. Math. Anal. Appl. 295(1) (2004), 225–236.
- [24] M. Daoud: *Nonlinear elliptic and parabolic reaction-diffusion systems governed by fractional laplacians : Analysis and Numerics*. Ph.D. thesis, Hassan II University of Casablanca, 2023.
- [25] M. Daoud, E.-H. Laamri: *Fractional Laplacians : A short survey*. Discrete Contin. Dyn. Syst. - S 15(1) (2022), 95–116.
- [26] E. Di Nezza, G. Palatucci, E. Valdinoci: *Hitchhiker’s guide to the fractional Sobolev spaces*. Bull. Sci. Math., 136(5) (2012), 521–573.
- [27] S. Duo, H. Wang, Y. Zhang : *A comparative study on nonlocal diffusion operators related to the fractional Laplacian*. Discrete Contin. Dyn. Syst. -B 24 (2019), 231–256.
- [28] B. Dyda: *Fractional calculus for power functions and eigenvalues of the fractional Laplacian*. Fract. Calc. Appl. Anal. 15(4) (2012), 536–555.
- [29] D.E. Edmunds, W.D. Evans: *Fractional Sobolev Spaces and Inequalities*. Cambridge University Press. 2023. ISBN 978-1-009-25463-2.
- [30] K. Fellner, E.-H. Laamri: *Exponential decay towards equilibrium and global classical solutions for nonlinear reaction-diffusion systems*. J. Evol. Equ. 16(3) (2016), 681–704.
- [31] K. Fellner, J. Morgan, B.Q. Tang: *Global classical solutions to quadratic systems with mass control in arbitrary dimensions*. Ann. Inst. H. Poincaré Anal. Non Linéaire 37(2) (2020), 281–307.
- [32] K. Fellner, J. Morgan, B.Q. Tang: *Uniform-in-time for quadratic reaction-diffusion systems with mass dissipation in higher dimensions*. Discrete Contin. Dyn. Syst. -S 14(2) (2021), 635–651.
- [33] M. Felsinger, M. Kassmann: *Local regularity of parabolic nonlocal operators*. Commun. Partial Differ. Equ. 38(9) (2013), 1539–1573.
- [34] X. Fernandez-Real: *Boundary regularity for the fractional heat equation*. Bachelor’s degree thesis, Polytechnic University of Catalonia, September 2014.
- [35] X. Fernandez-Real, X. Ros-Oton: *Boundary regularity for the fractional heat equation*. Rev. Acad. Cienc. Ser. A Math. 110 (2016), 49–64.

- [36] A. Fiscella, R. Servadei, E. Valdinoci: *Density properties for fractional Sobolev spaces*. Annales Academiae Scientiarum Fennicae. Mathematica 40(1) (2015), 235–253.
- [37] C.G. Gal, M. Warma: *Reaction-diffusion equations with fractional diffusion on non-smooth domains with various boundary conditions*. Discrete Contin. Dyn. Syst. 36(3) (2016), 1279–1319.
- [38] C.G. Gal, M. Warma: *Long-term behavior of reaction-diffusion equations with nonlocal boundary conditions on rough domains*. Z. Angew. Math. Phys. 67(4) (2016), 1–42.
- [39] C.G. Gal, M. Warma: *Nonlocal transmission problems with fractional diffusion and boundary conditions on nonsmooth interfaces*. Commun. Partial Differ. Equ. 42(4) (2017), 579–625.
- [40] T. Goudon, A. Vasseur: *Regularity Analysis for Systems of Reaction-Diffusion Equations*. Annales Sci. ENS (4) 43(1) (2010), 117–142.
- [41] A. Haraux, A. Youkana: *On a result of K. Masuda concerning reaction-diffusion equations*. Tôhoku Math. J. 40 (1988), 159–163 .
- [42] B.I. Henry, T.A.M. Langlands, P. Straka: *An Introduction to Fractional Diffusion*. Complex Physical, Biophysical and Econophysical Systems (2010), 37–89.
- [43] M.A. Herrero, A.A. Lacey, J.L. Velázquez: *Global existence for reaction-diffusion systems modelling ignition*. Arch. Rat. Mech. Anal. 142 (1998), 219–251.
- [44] S.L. Hollis, R.H. Martin, M. Pierre: *Global existence and boundedness in reaction-diffusion systems*. SIAM J. Math. Ana. 18 (1987), 744–761.
- [45] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo : *Theory and Applications of Fractional Differential Equations*. Elsevier, Amsterdam, 2006.
- [46] E.-H. Laamri: *Existence globale pour des systèmes de réaction-diffusion dans L^1* , Ph.D thesis, Université de Nancy 1, France, 1988.
- [47] E.-H. Laamri: *Global existence of classical solutions for a class of reaction-diffusion systems*. Acta Appl. Math. 115(2) (2011), 153–165.
- [48] E.-H. Laamri, R. Malek, C. Ziti: *Global existence for parabolic reaction-diffusion systems with exponential growth: Numerical study*. Submitted.
- [49] E.-H. Laamri, B. Perthame: *Reaction-diffusion systems with initial data of low regularity*. J. Differ. Equ. 269(11) (2020), 9310–9335.
- [50] E.-H. Laamri, M. Pierre: *Global existence for reaction-diffusion systems with nonlinear diffusion and control of mass*. Ann. Inst. H. Poincaré Anal. Non Linéaire 34(3) (2017), 571–591.
- [51] E.-H. Laamri, M. Pierre: *Stationary reaction-diffusion systems in L^1* . M3AS 28(11) (2018), 2161–2190.
- [52] E.-H. Laamri, M. Pierre: *Stationary reaction-diffusion systems in L^1 revisited*. Discrete Contin. Dyn. Syst. -S 14(2) (2021), 455–464.
- [53] D. Lamberton: *Equations d'évolution linéaires associées à des semi-groupes de contractions dans les espace L^p* . J. Funct. Anal. 72 (1987), 252–262.
- [54] A. Léculier, S. Mirrahimi, J.M. Roquejoffre: *Propagation in a fractional reaction-diffusion equation in a periodically hostile environment*. J. Dyn. Diff. Equat. 33 (2021), 863–890.

- [55] T. Leonori, I. Peral, A. Primo, F. Soria: *Basic estimates for solutions of a class of nonlocal elliptic and parabolic equations*. Discrete Contin. Dyn. Syst. 35(12) (2015), 6031–6068.
- [56] A. Lischke, G. Pang, M. Gulian, F. Song, C. Glusa, X. Zheng, Z. Mao, W. Cai, M.M. Meerschaert, M. Ainsworth, G.E. Karniadakis: *What is the Fractional Laplacian?* Journal of Computational Physics 404 (2020), 109009.
- [57] R. Malek: *Étude et simulations numériques de quelques problèmes concrets présentant ou non des singularités*. Ph.D thesis, Université Moulay Ismail de Meknès, Maroc, May 2022.
- [58] G. Molica Bisci, V.D. Radulescu, R. Servadei: *Variational Methods For Nonlocal Fractional Problems*. Cambridge University Press, 2016.
- [59] J. Morgan: *Global existence for semilinear parabolic systems*. SIAM J. Math. Ana. 20 (1989), 1128–1144.
- [60] K.W. Morton, D. Mayers : *Numerical Solution of Partial Differential Equations: An Introduction*. Second Edition, Cambridge University Press, 2005.
- [61] J.D. Murray: *Mathematical biology : I. An introduction*. Springer-Verlag, Berlin, Heidelberg, 2002.
- [62] A. Pazy: *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer, New York, 1983.
- [63] M. Pierre: *An L^1 method to prove global existence in some reaction-diffusion systems*. In “Contributions to Nonlinear Partial Differential Equations”, Vol.II, Pitman Research notes, J.I. Diaz and P.L. Lions ed., 155 (1987), 220–231.
- [64] M. Pierre: *Weak solutions and supersolutions in L^1 for reaction-diffusion systems*. J. Evol. Equ. 3 (2003), 153–168.
- [65] M. Pierre: *Systèmes de réaction-diffusion*. École de printemps, équations aux dérivées partielles non-linéaires, Marrakech, 2008.
- [66] M. Pierre: *Global Existence in Reaction-Diffusion Systems with Control of Mass : a Survey*. Milan J. Math. 78 (2010), 417–455.
- [67] M. Pierre, G. Rolland: *Global existence for a class of quadratic reaction-diffusion system with nonlinear diffusion and L^1 initial data*. Nonlinear Anal. TMA 138 (2017) 369–387.
- [68] M. Pierre, D. Schmitt: *Blowup in Reaction-Diffusion Systems with Dissipation of Mass*. SIAM Rev. 42(1) (2000), 93–106.
- [69] M. Pierre, T. Suzuki, Y. Yamada : *Dissipative reaction diffusion systems with quadratic growth*. Indiana Univ. Math. J. 68(1) (2019), 291–322.
- [70] L.M. Pismen: *Patterns and interfaces in dissipative dynamics*. Springer Science & Business Media, 2006.
- [71] I. Podlubny: *Fractional Differential Equations*. Academic Press, San Diego, 1999.
- [72] P. Quittner, Ph. Souplet: *Superlinear Parabolic Problems: Blow-up, Global Existence and Steady States*. Advanced Texts, Birkhäuser, (2019).
- [73] B. Rebiai, S. Benachour: *Global classical solutions for reaction-diffusion systems with nonlinearities of exponential growth*. J. Evol. Equ. 10(3) (2010), 511–527.

- [74] P. Scherer, S.F. Fischer: *Reaction-Diffusion Systems*. In : Theoretical Molecular Biophysics. Biological and Medical Physics, Biomedical Engineering. Springer, Berlin, Heidelberg, 2010.
- [75] Ph. Souplet: *Global existence for reaction-diffusion systems with dissipation of mass and quadratic growth*. J. Evol. Equ. 18(4) (2018), 1713–1720.
- [76] T. Suzuki, Y. Yamada : *Global-in-time behavior of Lotka-Volterra system with diffusion: skew-symmetric case*. Indiana Univ. Math. J. 64(1) (2015), 181–216.
- [77] E. Trélat, E. Zuazua: *Numerical Control: Part A*. First Edition, Elsevier Science & Technology. Handbook of Numerical Analysis XXIII 23 (2022), 1–558,
- [78] E. Valdinoci: *From the long jump random walk to the fractional Laplacian*. Bol. Soc. Esp. Mat. Apl. 49 (2009), 33–44.
- [79] J.L. Vázquez: *Nonlinear Diffusion with Fractional Laplacian Operators*. In: Holden, H., Karlsen, K. (eds) Nonlinear Partial Differential Equations. Abel Symposia 7. Springer, Berlin, Heidelberg, 2012.
- [80] J.L. Vázquez: *The mathematical theories of diffusion. Nonlinear and fractional diffusion*. Springer Lecture Notes in Mathematics, CIME Subseries, 2017.
- [81] I.I. Vrabie: *C_0 -Semigroups and Application*. Elsevier Science B.V., 2003.