

Supplemental Table 1. Baseline characteristics of patients with HFrEF and controls.

	HFrEF (N=148)	Control (N=36)	p-value
Clinical characteristics			
Age, years	54 ± 12	54 ± 10	0.923
Male, n (%)	127 (84)	22 (61)	0.004
Body mass index, kg/m ²	27.4 ± 4.5	28.3 ± 4.8	0.282
NYHA class, n (%): I-II; III-IV	34 (23); 114 (77)	NA	NA
Ischemic HF aetiology, n (%)	65 (44)	NA	NA
Diabetes mellitus, n (%)	50 (33)	3 (8)	0.002
Ischemic heart disease, n (%)	69 (46)	4 (11)	0.002
Haemoglobin, mg/L	137.0 ± 18.3	139.1 ± 17.3	0.525
BNP, ng/L	1019 [509; 1824]	29 [9; 46]	<0.001
Echocardiography*			
LV end-diastolic diameter, mm	69.7 ± 8.5	46.7 ± 4.1	<0.001
LV mass index, gr/m ²	128.5 ± 31.7	75.9 ± 14.2	<0.001
LV ejection fraction, %	22.0 ± 6.7	60.0 ± 5.0	<0.001
Mitral regurgitation, n (%): 0-2 grade; 3-4 grade	72 (48); 78 (52)	36 (100); 0 (0)	<0.001
Tricuspid regurgitation, n (%): 0-2 grade; 3-4 grade	98 (65); 52 (35)	34 (97); 1 (3)	<0.001
Right ventricular basal dimensions, mm	45.0 ± 3.4	36.0 ± 4.2	<0.001
Tricuspid annular plane systolic excursion, mm	15.2 ± 4.2	23.8 ± 4.8	<0.001
Electrocardiography			
Heart rate, bpm	80 ± 16	68 ± 12	<0.001
Atrial rhythm, n (%): Sinus; Paced; AFib or AFlu	128 (85); 3 (2); 19 (13)	36 (0); 0 (0); 0 (0)	<0.001
QRS morphology, n (%): LBBB; RBBB; normal; other alterations*	31 (21); 13 (9); 36 (24); 68 (45)	1 (3); 2 (5); 30 (83); 3 (8)	<0.001
QRS duration, ms	120.4 ± 24.8	93.2 ± 18.3	<0.001
Therapy			
Furosemide, n (%)	143 (97)	4 (12)	<0.001
ACE, ARB or ARNI, n (%)	103 (70)	13 (36)	<0.001
MRA, n (%)	128 (86)	4 (11)	<0.001
Betablocker, n (%)	124 (84)	9 (25)	<0.001
Devices, n (%): ICD; CRT; none	111 (75); 12 (8); 25 (17)	NA	NA

*Mitral and tricuspid regurgitation were assessed semiquantitatively and reported using a 4-stage grading (i.e., 0) none-trace; 1) mild; 2) moderate; 3) severe).

Abbreviations: ACE – Angiotensin Converting Enzyme Inhibitor; AFlu – Atrial Flutter; AFib – Atrial Fibrillation; ARB – Angiotensin Receptor Blocker; ARNI – Angiotensin Receptor Neprilysin Inhibitor; BNP – Brain Natriuretic Peptide; CRT – Cardiac Resynchronization Therapy; HFrEF – Heart Failure with Reduced Ejection Fraction; ICD – Implantable Cardioverter-Defibrillator; LBBB – Left Bundle Branch Block; MRA – Mineralocorticoid Receptor Antagonist; NA – Not Applicable; NYHA – New York Heart Association; RBBB – Right Bundle Branch Block. *Other alterations include: left anterior fascicular block, left posterior fascicular block, incomplete LBBB and nonspecific intraventricular conduction delay.

Supplemental Table 2. Imaging and hemodynamic data in patients with HFrEF and controls

	HF patients (N=148)	Controls (N=36)	p-value
GSPECT			
RV phase SD, %	11.2 ± 4.9	7.6 ± 2.5	<0.001
RV end-diastolic volume – mL	256.4 ± 97.9	204.4 ± 61.1	<0.001
RV end-diastolic volume index – mL/m ²	138.0 ± 47.9	103.1 ± 24.3	<0.001
RV end-systolic volume – mL	165.5 ± 82.1	96.5 ± 32.1	<0.001
RV end-systolic volume index – mL/m ²	85.8 ± 41.9	48.1 ± 14.0	<0.001
RV ejection fraction - %	37.7 ± 14.9	53.1 ± 5.4	<0.001
Right heart catheterization			
RA mean pressure - mmHg	9.6 ± 5.8	5.0 ± 2.9	<0.001
RV systolic elastance (Ees) - mmHg/mL	0.32 [0.23; 0.49]	0.32 [0.23; 0.47]	0.177
RV diastolic compliance - mL/mmHg	0.03 [0.02; 0.05]	0.04 [0.03; 0.06]	0.003
Cardiac output, L/m	3.8 ± 0.9	5.6 ± 1.4	<0.001
Stroke volume, mL	48.8 ± 14.7	76.9 ± 23.8	<0.001
PA mean pressure - mmHg	34.3 ± 10.8	16.3 ± 6.5	<0.001
PA compliance - mL/mmHg	1.8 [2.3; 2.6]	4.8 [3.5; 6.5]	<0.001
PA elastance (Ea _{PA}) - mmHg/mL	1.00 [0.73; 1.53]	0.36 [0.26; 0.44]	<0.001
PA wedge pressure - mmHg	22.8 ± 8.5	8.2 ± 4.2	<0.001
PVR - WU	2.9 [1.9; 4.2]	1.4 [1.0; 2.1]	<0.001
RV ventricular-arterial coupling ratio	0.29 [0.21; 0.51]	0.92 [0.61; 1.37]	<0.001
Systolic blood pressure - mmHg	109.4 ± 16.2	128.5 ± 20.9	<0.001
Diastolic blood pressure - mmHg	75.0 ± 10.0	77.1 ± 9.9	0.258


Abbreviations: GSPECT - Gated Single-Photon Emission Computed Tomography; HFrEF – Heart Failure with Reduced Ejection Fraction; PA – Pulmonary Artery; PVR – Pulmonary Vascular Resistance; RA – Right Atrial; RV – Right Ventricular; SD – Standard deviation; SVR – Systemic Vascular Resistance; WU - Wood Units

Supplemental Table 3 Effect of afterload reduction (sildenafil) and preload increase (leg rise) on right ventricular dyssynchrony, hemodynamics and cardiac volumes derived from SPECT

	Baseline	Afterload reduction (N=15)	p-value	Baseline	Preload modulation (N=135)	p-value
RV dyssynchrony, degrees	44.9 ± 19.2	29.6 ± 13.9	0.004	40.5 ± 17.6	37.7 ± 16.4	0.080
<i>Right heart catheterization</i>						
RA mean pressure, mmHg	7.9 ± 3.5	4.3 ± 2.5	<0.001	9.3 ± 5.7	11.7 ± 6.4	<0.001
RV end-diastolic pressure, mmHg	12.0 ± 4.3	6.3 ± 3.7	<0.001	11.5 ± 6.1	14.4 ± 6.0	<0.001
PA mean pressure, mmHg	44.0 ± 8.6	31.2 ± 9.9	<0.001	33.6 ± 10.4	38.5 ± 9.7	<0.001
PA wedge pressure mean, mmHg	25.3 ± 5.9	19.6 ± 8.7	0.005	22.3 ± 8.1	26.3 ± 7.8	<0.001
PA elastance, mmHg/mL	1.7 [1.1; 1.9]	0.9 [0.6; 1.1]	<0.001	1.0 [0.7; 1.5]	1.2 [0.8; 1.6]	<0.001
RV systolic elastance (Ees) - mmHg/mL	0.48 [0.29; 0.79]	0.50 [0.25; 0.90]	0.889	0.31 [0.22; 0.49]	0.31 [0.22; 0.49]	0.489
RV ventricular-arterial coupling ratio	0.43 [0.15; 0.54]	0.57 [0.34; 0.90]	0.002	0.29 [0.20; 0.51]	0.28 [0.17; 0.47]	0.001
Pulmonary vascular resistance - WU	4.9 [4.3; 6.6]	2.7 [1.8; 3.8]	<0.001	2.8 [1.8; 3.9]	3.1 [2.1; 4.4]	0.031
Cardiac output, L/min	3.7 ± 0.8	4.4 ± 0.8	0.003	3.8 ± 0.9	3.7 ± 0.9	0.411
<i>SPECT-derived data</i>						
RV end-diastolic volume, mL	256.1 ± 77.7	214.3 ± 75.4	0.002	278.2 ± 98.4	289.8 ± 94.1	0.001
RV end-systolic volume, mL	152.7 ± 75.5	107.1 ± 66.1	<0.001	173.1 ± 86.9	178.9 ± 84.0	0.023
RV ejection fraction, %	43.1 ± 12.6	54.1 ± 14.1	<0.001	40.8 ± 14.9	41.0 ± 14.0	0.867


Abbreviations: PA – Pulmonary Artery; PVR – Pulmonary Vascular Resistance; RA – Right Atrial; RV – Right Ventricular; WU - Wood Units.

Supplemental Figure 1




Supplemental Figure 2


A




B




Supplemental Figure 3




Supplemental Figure 4





Supplemental Figure 5

A**B**