



**HAL**  
open science

## **7-Ketocholesterol: Effects on viral infections and hypothetical contribution in COVID-19**

Imen Ghzaiel, Khoulood Sassi, Amira Zarrouk, Thomas Nury, Mohamed Ksila, Valerio Leoni, Balkiss Bouhaouala-Zahar, Sonia Hammami, Mohamed Hammami, John Mackrill, et al.

► **To cite this version:**

Imen Ghzaiel, Khoulood Sassi, Amira Zarrouk, Thomas Nury, Mohamed Ksila, et al.. 7-Ketocholesterol: Effects on viral infections and hypothetical contribution in COVID-19. *Journal of Steroid Biochemistry and Molecular Biology*, 2021, 212, pp.105939. 10.1016/j.jsbmb.2021.105939 . hal-04842319

**HAL Id: hal-04842319**

**<https://hal.univ-lorraine.fr/hal-04842319v1>**

Submitted on 19 Dec 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## 7-ketocholesterol: effects on viral infections and hypothetical contribution in COVID-19

Imen Ghzaïel<sup>1,2,3</sup>, Khoulood Sassi<sup>1,4</sup>, Amira Zarrouk<sup>2,5</sup>, Thomas Nury<sup>1</sup>, Mohamed Ksila<sup>6</sup>, Valerio Leoni<sup>7</sup>, Balkiss Bouhaouala-Zahar<sup>8</sup>, Sonia Hammami<sup>2</sup>, Mohamed Hammami<sup>2</sup>, John J. Mackrill<sup>9</sup>, Mohammad Samadi<sup>10</sup>, Taoufik Ghraïri<sup>6</sup>, Anne Vejux<sup>1</sup>, Gérard Lizard<sup>1\*</sup>

1: University Bourgogne Franche-Comté, Team 'Biochemistry of the Peroxisome, Inflammation and Lipid Metabolism' EA 7270 / Inserm, 21000 Dijon, France; [gerard.lizard@u-bourgogne.fr](mailto:gerard.lizard@u-bourgogne.fr), [thomas.nury@u-bourgogne.fr](mailto:thomas.nury@u-bourgogne.fr), [anne.vejux@u-bourgogne.fr](mailto:anne.vejux@u-bourgogne.fr)

2: University of Monastir, Faculty of Medicine, LR12ES05, Lab-NAFS 'Nutrition - Functional Food & Vascular Health', 5000, Monastir, Tunisia; [zarroukamira@gmail.com](mailto:zarroukamira@gmail.com), [imenghzaïel93@gmail.com](mailto:imenghzaïel93@gmail.com), [sonia.hammami@fmm.rnu.tn](mailto:sonia.hammami@fmm.rnu.tn), [mohamed.hammami@fmm.rnu.tn](mailto:mohamed.hammami@fmm.rnu.tn)

3: University Tunis-El Manar, Faculty of Sciences of Tunis, 2092 Tunis, Tunisia; [imenghzaïel93@gmail.com](mailto:imenghzaïel93@gmail.com)

4: University Tunis El Manar, Laboratory of Onco-Hematology (LR05ES05), Faculty of Medicine, 1007 Tunis, Tunisia; [sassikhouloud@hotmail.com](mailto:sassikhouloud@hotmail.com)

5: University of Sousse, Faculty of Medicine, Sousse, Tunisia; [zarroukamira@gmail.com](mailto:zarroukamira@gmail.com)

6: University Tunis-El Manar, Laboratory of Neurophysiology, Cellular Physiopathology and Valorisation of BioMolecules, LR18ES03. Department of Biologie, Faculty of Sciences, 2092 Tunis, Tunisia, [mohamedksila44@gmail.com](mailto:mohamedksila44@gmail.com) ; [taoufik.ghraïri@fst.utm.tn](mailto:taoufik.ghraïri@fst.utm.tn)

7: Laboratory of Clinical Chemistry, Hospitals of Desio, ASST-Brianza and Department of Medicine and Surgery, University of Milano-Bicocca, 20900 Monza, Italy; [valerio.leoni@unimib.it](mailto:valerio.leoni@unimib.it)

8: Laboratory of Venoms and Therapeutic Molecules, Pasteur Institute of Tunis & University of Tunis El Manar, 1002 Tunis, Tunisia; [balkiss.bouhaouala@pasteur.rns.tn](mailto:balkiss.bouhaouala@pasteur.rns.tn)

9: Department of Physiology, School of Medicine, University College Cork, Cork, Ireland; [J.Mackrill@ucc.ie](mailto:J.Mackrill@ucc.ie)

10: LCPMC-A2, ICPM, Dept of Chemistry, Univ. Lorraine, Metz Technopôle, Metz, France; [mohammad.samadi@univ-lorraine.fr](mailto:mohammad.samadi@univ-lorraine.fr)

**Correspondence:** Dr Gérard LIZARD (PhD), Laboratoire Bio-peroxIL, Université de Bourgogne / Inserm, Faculté des Sciences Gabriel, 6 Bd Gabriel, 21000 Dijon, France ; E.mail : [gerard.lizard@u-bourgogne.fr](mailto:gerard.lizard@u-bourgogne.fr) ; phone: 33 3 80 39 62 56 / Fax: 33 3 80 39 62 50

**Abstract:** 7-ketocholesterol, which is one of the earliest cholesterol oxidization products identified, is essentially formed by the auto-oxidation of cholesterol. In the body, 7-ketocholesterol is both provided by food and produced endogenously. This pro-oxidant and pro-inflammatory molecule, which can activate apoptosis and autophagy at high concentrations, is an abundant component of oxidized Low Density Lipoproteins. 7-Ketocholesterol appears to significantly contribute to the development of age-related diseases (cardiovascular diseases, age-related macular degeneration, and Alzheimer's disease), chronic inflammatory bowel diseases and to certain cancers. Recent studies have also shown that 7-ketocholesterol has anti-viral activities, including on SARS-CoV-2, which are, however, lower than those of oxysterols resulting from the oxidation of cholesterol on the side chain. Furthermore, 7-ketocholesterol is increased in the serum of moderately and severely affected COVID-19 patients. In the case of COVID-19, it can be assumed that the antiviral activity of 7-ketocholesterol could be counterbalanced by its toxic effects, including pro-oxidant, pro-inflammatory and pro-coagulant activities that might promote the induction of cell death in alveolar cells. It is therefore suggested that this oxysterol might be involved in the pathophysiology of COVID-19 by contributing to the acute respiratory distress syndrome and promoting a deleterious, even fatal outcome. Thus, 7-ketocholesterol could possibly constitute a lipid biomarker of COVID-19 outcome and counteracting its toxic effects with adjuvant therapies might have beneficial effects in COVID-19 patients.

**Keywords:** adjuvant therapies, COVID-19, 7-ketocholesterol, oxysterols, SARS-CoV-2, viral diseases.

### Origin and metabolism of 7-ketocholesterol

Cholesterol ((3 $\beta$ )-cholest-5-en-3-ol; C<sub>27</sub>H<sub>46</sub>O; molecular weight: 386.65 g/mol) is a lipid which is both provided by the diet and formed endogenously, except in the brain where the cholesterol present is only produced by astrocytes ([1] [2]). After a meal, cholesterol transiting through enterocytes is taken up by the enteric capillaries as chylomicrons which will transport it to the liver, where it will be distributed to the peripheral organs via low density lipoproteins (LDLs); its return to the liver is then ensured by the high density lipoproteins (HDL) [3]. In all the cells of the body, acetyl-CoA is the first element involved in the biosynthesis of cholesterol. Endogenous cholesterol synthesis involves several enzymes which, after conversion of acetyl-CoA into mevalonate by the enzyme HMG-CoA reductase, will then be converted to squalene, which is subsequently metabolized to lanosterol. From lanosterol, which contains thirty carbon atoms (C<sub>30</sub>), the biosynthesis of cholesterol (C<sub>27</sub>) takes two routes: the Bloch pathway and the Kandutsch-Russell pathway. The Bloch pathway produces cholesterol precursors ranging from 14-demethyl-14-dehydrolanosterol (ff-MAS; C<sub>29</sub>) to desmosterol (C<sub>27</sub>) and includes zymosterol (C<sub>27</sub>); the Kandutsch-Russell pathway generates products spanning from 24,25-dihydrolanosterol (C<sub>30</sub>) to 7-dehydrocholesterol (C<sub>27</sub>) and includes zymostenol (C<sub>27</sub>) and lathosterol (C<sub>27</sub>) [4]. These two pathways can generate cholesterol via the enzymes 24-dehydrocholesterol reductase (DHCR24) and 7-dhydrocholesterol reductase (DHCR7): the enzyme DHCR24 generates cholesterol from desmosterol while the enzyme DHCR7 generates cholesterol from 7-dehydrocholesterol ([5] [6]). Whether endogenous or exogenous in nature, in a pro-oxidant environment cellular, cholesterol can then give oxidized derivatives in positions 4, 5, 6 and 7 ([7] [8] [9]). Most often radical attacks by reactive oxygen or nitrogen species (ROS or RNS) take place on carbon 7, because of the weak link between carbon and hydrogen at this position ([9] [10]). This local oxidation at carbon 7 then generates a peroxy radical (ROO•) which, by reacting with hydrogen, forms cholesterol hydroperoxide (7 $\alpha$ - or 7 $\beta$ -OOHC) [11]. As the hydroperoxide bond is very unstable, 7-ketocholesterol (7KC), also named 7-oxo-cholesterol ([12] [13]), is formed in the majority of cases, and 7 $\beta$ -hydroxycholesterol and 7 $\alpha$ -hydroxycholesterol in smaller quantities [11] (**Figure 1**). Under certain conditions, it has also been shown that 7KC can be formed via the enzyme sterol 7-hydroxylase (CYP7A1) from 7-dehydrocholesterol (7-DHC) present in large quantities in the plasma of patients with Smith Lemli Opitz (SLO) syndrome [4]. In addition, 11 $\beta$ -hydroxysteroid-dehydrogenase type 2 (11 $\beta$ -

HSD2), which converts cortisol to cortisone, is also responsible for the conversion of 7 $\beta$ -hydroxycholesterol to 7KC [14].

Overall, 7KC can be detected in the plasma at LDL level or bound to albumin as well as in the plasma membrane of the cells of different tissues [15]. It is also well established that 7KC can react with different molecules (sulfate, fatty acids) via the hydroxyl group (OH) localized on C3 in the ring A of the sterane nucleus, which leads to inhibition of its toxicity ([16] [17]). The enzyme sulfotransferase 2B1b (SULT 2B1b) is involved in the sulfonation of 7KC [18]. As for the esterification of 7KC, a combined action of cytosolic phospholipase A2 alpha (CPLA2 a) and sterol-O-acyltransferase (SOAT1) has been reported [19]. In addition, Acyl-coenzyme A transferase (ACAT, also abbreviated as SOATs) which converts cholesterol into cholesterol ester can use oxysterols as substrates: oxysterols are also substrates for SOAT1 and SOAT2 [20]. The lecithin cholesterol acyl-transferase (LCAT) also esterifies oxysterols in the plasma [15]. At the moment, it has been described in retinal pigment epithelium that 7KC can be metabolised by the enzyme 27-hydroxylase cytochrome P450 27A1 (CYP27A1) to form two metabolites, 3 $\beta$ ,27-dihydroxy-5-cholesten-7-one (7KC-27OHC) and 3 $\beta$ -hydroxy-5-cholesten-7-one-26-oic acid (7KC-27COOH), to reduce the toxicity of 7KC ([21] [22]). It is therefore important to know the biological activities of 7KC, and of its metabolites which are still not well known ([23] [24]), as well as their related signalling pathways, in order to identify therapeutic targets and develop drugs acting on them to treat diseases associated with high levels of 7KC in tissues and biological fluids.

### **Pathologies associated with 7-ketocholesterol**

Large amounts of oxysterols formed by auto-oxidation, particularly 7KC, have been initially detected in oxidised LDL (LDLox) and in atheromatous plaques. Since there is a positive correlation between the content of oxysterol-rich LDLox and induction of cell death, it has been suggested that 7KC may play a **crucial** role in atherosclerosis and cardiovascular diseases ([25] [26]). Moreover, analogies between atherosclerosis and the development of age-related macular degeneration (AMD) (significant presence of 7KC in lipid deposits called drüsens located between the Bruch membrane and the monolayer of retinal epithelial cells) also indicate

involvement of 7KC in the pathophysiology of this disease ([27] [28]). A potential involvement of 7KC in the pathophysiology of cataract, which affects one person in five from the age of 65, and more than 60% of people aged 85 and over, is also suspected [29]. Thus, the exposure of membranes isolated from transparent human lenses to 2,2'-azobis(2-amidinopropane) hydrochloride, a free radical generator, promotes the production of 7KC (74%) as the main cholesterol oxidation product [30]. In addition, cataract lenses contain quantifiable amounts of 7KC (4.2 +/- 0.32 mmol / mol of cholesterol), whereas clear lenses from cataract-free subjects do not contain detectable amounts [30]. Moreover, the presence of high levels of 7KC in the brain lesions of Alzheimer's disease patients suggests an involvement of 7KC in this prevalent neurodegenerative disease ([31] [32]). 7KC also appears to be involved in chronic inflammatory bowel diseases [33] as well as in some rare diseases of lipid metabolism, such as X-linked adrenoleukodystrophy (X-ALD) ([11] [34]). High levels of 7KC have also been described in non-infectious but severe inflammatory lung diseases such as silicosis ([35] [36]). In addition, air pollution which affects the respiratory system by promoting oxidative stress, increases the production of 7KC and promotes atherosclerosis by activating CD-36 positive macrophages [37]. Currently, several ozone-oxidized cholesterol products have been identified and can be considered as a new class of oxysterols ([10] [38]). The common denominators of 7KC-associated diseases are high levels of oxidative stress and inflammation which can in turn amplify the formation of 7KC, as well as organelle dysfunction (of mitochondria, peroxisomes, lysosomes, endoplasmic reticulum) [39] and subsequently contribute to the amplification of this stress [40]. Furthermore, organelle oxidative stress, mainly at the mitochondrial and peroxisomal level, could favour the formation of 7KC, which is itself strongly pro-oxidative and pro-inflammatory ([41] [42] [43]).

Beside age-related diseases, increases in 7KC have also been described in the context of viral infections. For example, in patients with type 2 diabetes who are co-infected with herpes virus type 8 (HHV8), significant increases in plasma levels of 7KC have been observed possibly amplifying diabetic complications [44]. It is hypothesized that the HHV8-infection may contribute to ROS overproduction which would trigger lipid peroxidation and cholesterol autoxidation, leading to 7KC formation [44]. Similarly, elevated plasma levels of 7KC have also been measured in patients infected with influenza A virus [45]. In patients infected by the SARS-CoV-2 coronavirus with severe forms of COVID-19 (COrona VIRus Disease - 2019), a

potentially fatal acute respiratory distress syndrome due to a bilateral pneumonia, elevated plasma levels of 7KC were observed whereas 27-hydroxycholesterol (also known for its strong anti-viral activity) ([46] [47] [48] [49]) was simultaneously significantly decreased compared to the control group, reaching a marked 50% reduction in severe COVID-19 cases [49]. SARS-CoV-2 is an enveloped RNA virus; its genome encodes for **fifteen genes** including a surface protein, the Spike protein, which allows it to enter and to infect the target cells at the level of vital organs [50]. Thus, this Spike protein binds to the angiotensin-converting enzyme 2 (ACE-2) receptor expressed in almost all tissues and abundant in the lungs, kidneys, brain stem, adipose tissue, heart, vasculature, stomach, liver, nasal and oral mucosa [51]. The innate immune cells (neutrophils, monocytes) and adaptive immune cells (T cells) are involved in the response to COVID-19 infection ([52] [53]). **In a retrospective study realized on 175 patients, it was noted that the highest co-morbidity has been observed in COVID-19 patients with cardiovascular diseases [54] in whom it is well known that the level of oxysterols formed by auto-oxidation, including 7KC, is already high [11].** An increased prevalence of fungal and *Pseudomonas aeruginosa* colonization has also been observed in patients with severe forms of COVID-19, suggesting that this association may be the result of a failure in the regulation of immune defenses against pathogens other than SARS-CoV-2, in the case of co-infection [55]. **From a clinical point of view, atherosclerosis, AMD, cataract, and Alzheimer's disease are age-related diseases ([42] [56]). At the beginning of the pandemic, COVID-19 could be considered as an emerging age-related disease. This concept is no longer relevant as it is well established that children, adolescents and young adults can also be infected ([57] [58]). However, severe forms of COVID-19 most often affect people over 65 years of age due to a combination of several parameters, including nutritional aspects but especially age-related immunological and metabolic alterations ([59] [60]).** About the cardiovascular diseases, it can be assumed that the presence of high concentrations of 7KC in the elderly could be a risk factor that may promote co-morbidity in the case of infection with the SARS-CoV-2. A possible involvement of the 7KC in the pathophysiology and in the fatal or non-fatal outcome of severe forms of COVID-19, would strengthen the interest for this oxysterol in the context of viral diseases.

### **Anti-viral activities of 7-ketocholesterol**

Currently, there is evidence that several oxysterols have antiviral effects against both enveloped and non-enveloped viruses. An in-depth review of the literature on this topic was carried out in 2016 by Lembo *et al.* [61]. In addition, several studies have shown the ability of certain cholesterol and oxysterol metabolites to regulate both intrinsic/innate immunity and adaptive immunity, each of which may be involved in bacterial and viral infections [62]. Among the oxysterols with strong anti-viral properties are mainly those derived from cholesterol oxidation on the aliphatic side chain such as 25-hydroxycholesterol and 27-hydroxycholesterol, which could act both by inhibiting viral replication and by activating the immune response ([63] [46] [64] [65]). 7 $\alpha$ , 25-hydroxycholesterol generated by cholesterol by the sequential action of cholesterol 25 hydroxylase (CH25H) and 7-alpha-hydroxylase (CYP7B1) also has anti-viral activities, and acts as chemo-attractant for cells expressing the cell surface receptor GPR183 present on dendritic and B cells, as well as type 3 innate lymphoid cells (ILC3) ([66] [67]). Less potent antiviral activities have been reported with 7 $\beta$ -hydroxycholesterol and 7KC [61]. However, 7KC shows anti-viral activities (inhibition of viral replication, viral inactivation) on three enveloped viruses (human papillomavirus-16 (HPV16), human rotavirus (HRoV), human rhinovirus (HRhV)) [46]. In addition, in cultures of green monkey kidney cells (Vero cells) and differentiated human nerve cells (hNP1 cells), 7KC reduces the viral titer of the Zika virus, which is an emerging African virus characterized by significant neurotropism, without affecting cell viability: 7KC reduces the number and infectivity of viral particles released into the culture medium [68]. In VeroE6/TMPRSS2 cells inoculated with SARS-CoV-2, weak anti-viral activities were demonstrated with 7KC, 22(R)-hydroxycholesterol and 22(S)-hydroxycholesterol; the latter are weak inhibitors of viral replication, in contrast to 27-hydroxycholesterol and 24(S)-hydroxycholesterol [69]. These observations led to the development of semi-synthetic oxysterols with anti-SARS-CoV-2 activity when administered orally to mice which are named **Oxy210** (3S,8S,9S,10R,13S,14S,17S)-17-((R)-2-hydroxy-4-(pyridin-3-yl)butan-2-yl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-ol) and **Oxy232** ((3S,5S,8R,9S,10S,13S,14S,17S)-17-((R)-3-hydroxy-1-(pyridin-3-yl)pentan-3-yl)-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-3-ol) [69]. **These different results suggest that 7KC has generally weak anti-viral activities; however, this oxysterol could nevertheless contribute to the immune response due to its pro-inflammatory properties including stimulation of IL-8 and IL-1 $\beta$  production ([70, 71]).**



### **Potential role of 7-ketocholesterol in the pathophysiology of COVID-19**

Although 7KC has weak anti-viral activities, it has strong cytotoxic activities. Indeed, there are several data that favour the pro-oxidative and pro-inflammatory activities of 7KC that may be associated with a type of death defined as oxiaoptophagy (OXIdative stress + APOPTOsis + autoPHAGY), due to the ability of 7KC (at high concentration) to induce oxidative stress and an apoptotic mode of cell death associated with autophagy criteria ([14] [72] [73]). In several cells from different types and from different species, i) oxidative stress is characterized by ROS overproduction including at the mitochondrial level, protein carbonylation, lipid peroxidation and decrease and/or enhancement of anti-oxidant enzymes (catalase, superoxide dismutases (SODs), glutathione peroxidase (GPx), ii) apoptosis is associated by an activation of the mitochondrial pathway leading to a loss of transmembrane mitochondrial potential ( $\Delta\Psi_m$ ), an externalization of phosphatidylserine, an increase of plasma membrane rigidity associated with an increased cell permeability contributing to an altered packing of the lipid bilayer, a cytoplasmic release of cytochrome c, a caspase cascade activation (caspases-3,-7,-8 and -9), poly-ADP ribose (PARP) fragmentation and an internucleosomal DNA fragmentation associated with condensation and/or fragmentation of the nuclei and iii) autophagy is characterized by the presence the presence of autophagosomes and autophagolysosomes revealed by transmission electron microscopy as well as activation of LC3-I into LC3-II [73] [74]. In addition, 7KC-induced oxiaoptophagy is associated with a production of inflammatory cytokines (IL-1 $\beta$ , IL-8) ([72] [14]). Noteworthy, 7KC, together with 7 $\beta$ -hydroxycholesterol (7 $\beta$ -OHC), was significantly increased in the plasma collected from COVID-19 patients compared to age matched healthy controls [49]. A progressive positive trend was found together with the severity of the disease: the highest 7KC and 7 $\beta$ -OHC concentrations were observed in severe COVID-19 patients. In these patients, it was also observed a progressive reduction of the plasma concentrations of lanosterol, lathosterol, and desmosterol, markers of cholesterol synthesis and of the 27-hydroxycholesterol, which has antiviral and immunomodulatory activities against SARS-CoV-2 ([49] [69]). The rise of 7KC and 7 $\beta$ -OHC was related to the increased oxidative stress caused by respiratory distress [49].

**In the case of COVID-19, high serum levels of 7KC were observed: In male/female control subjects of the same age range as COVID-19 patients (70  $\pm$  10 years old), the serum level of 7KC**

is about 20  $\mu\text{g/L}$  whereas the serum level of 7KC was increased by a factor of 2 to 3.5 and of 2 to 5 in moderate and severe COVID-19 patients, respectively [49]. Similarly, while the serum level of 7 $\beta$ -OHC is about 8  $\mu\text{g/L}$  in control subjects, this later was increased by a factor of 1.5 to 2.5 and of 1.5 to 5 in moderate and severe COVID-19 patients, respectively [49]. As patients with severe forms of COVID-19 may have the same 7KC and 7 $\beta$ -OHC levels as patients with moderate forms, this suggests that increased serum levels of these oxysterols are not indicative of disease severity. On the other hand, the very high serum levels of 7KC observed in some patients with severe COVID-19 - which may be a consequence of the infection and result from an exacerbated immune response - may contribute to the deterioration of the patients' condition. Thus, it can be assumed that the anti-viral activity of 7KC should be outweighed by its toxic effects, which could contribute to cytokine storm ([12] [75]) and also to the activation of coagulation in the capillaries of the pulmonary alveoli [76], and to the deterioration of alveolar epithelial cells [77]. It is assumed that the organism, with the contribution of 7KC, is overwhelmed by a storm of cytokines that have a deleterious impact on host cells, particularly alveolar epithelial cells, and is therefore unable to inhibit and destroy the SARS-CoV-2. Through its cytotoxic activities associated with oxiaoptophagy, it is hypothesized that 7KC could contribute to the progression and outcome of COVID-19 pathophysiology at different levels: a) by activating the TLR4 receptor which would contribute to increased secretion of pro-inflammatory cytokines (IL-1 $\beta$ , IL-8) [78]; b) by promoting the externalization of phosphatidylserine at the level of endothelial cells of the alveolar capillaries [79] and the eryptosis of red blood cells [80] as well as platelet aggregation [76] which would activate coagulation and the induction of thrombosis c) by altering the viability of the epithelial alveolar cells (oxidative stress, induction of cell death) [77] which would have the consequence of favouring acute respiratory distress syndrome. In addition, elevated plasma levels of 7KC could have systemic effects by altering mitochondrial function [81] with a consequent decrease in ATP production and general exhaustion of the patient. In addition, mitochondrial dysfunction could lead to peroxisomal alterations with subsequent amplification of oxidative stress, deregulation of non-cytokine-mediated inflammation due to the fact that leukotrienes, some of which are also pro-coagulants, are catabolized at the peroxisome level ([82] [83] [84] [85]). Furthermore, the activation of autophagy by 7KC could also have consequences on the infection. Indeed, autophagy initially considered as anti-viral also seems capable of promoting infection by acting at

the level of viral replication and viral cycle ([86] [87]). Given the toxic characteristics of 7KC, its high level in COVID-19 patients could contribute to cytokine storm, thrombosis and respiratory distress (**Figure 2**). To better understand the involvement of 7KC in the pathophysiology of COVID-19, rapid and easy to use analytical tests are a necessity. These could include ELISA tests using anti-7KC mouse monoclonal antibodies [88] or dried blood tests [89] associated with gas chromatography or liquid chromatography coupled with mass spectrometry allowing the simultaneous identification and quantification of several oxysterols and of their metabolites ([90] [24]).

### **7-ketocholesterol-modifying drugs in COVID-19**

Depending on the risk factors presented by SARS-CoV-2 infected patients, the infection may cause an acute respiratory distress syndrome, and multiple organ failure, which can be fatal. To prevent and/or cure COVID-19, the vaccine strategy seems the most promising approach whereas conventional drugs can provide potential alternative treatments as adjuvants [91] [92]. These alternative approaches using drugs acting on the viral cycle and/or on the cytopathic effects of SARS-CoV-2 should not be neglected. As cholesterol is essential for the assembly, replication and infectivity of enveloped viruses such as SARS-CoV-2, several cholesterol-modifying drugs could alter the SARS-CoV-2 life cycle [93]. In addition, as cholesterol is also a major component of immune cell membranes, excess cholesterol levels in patients with obesity and/or cardiovascular diseases could contribute to dysregulate acquire immunity and promote abnormal inflammatory responses [93]. Moreover, as cholesterol oxidation under the effect of ROS or RNS or through the intermediary of enzymes can lead to the formation of oxysterols [94], and in particular 7KC, which can impact the redox status, inflammation, coagulation, and cell viability, cholesterol-modifying drugs could be of interest to target the side effects of 7KC in the management of SARS-CoV-2 infection [95].

To have a more targeted activity against 7KC, natural and synthetic cytoprotective molecules have been identified and some of them could be used ([72] [73] [85]). Among the natural molecules, fatty acids that can be used as a basis for the design of synthetic analogues, making it possible to neutralise 7KC by esterifying it at the level of the hydroxyl residue on carbon 3 of the

A ring of the sterane nucleus, and/or by acting on its signalling pathways, are promising molecules ([96] [72]). Natural polyphenols ([72] [73]) and derivatives, such as aza- and azo-stilbenes, which are bio-isosteric analogues of resveratrol [97], could be also of interest, as well as tocopherols which have strong anti-oxidant properties and prevent the accumulation of 7KC in the lipid rafts to trigger several signalling pathways ([98] [99]). In addition, the phospholipid bis(monoacylglycero)phosphate (BMP) is a structural isomer of phosphatidylglycerol that exhibits an unusual sn1:sn1' stereoconfiguration based on the position of the phosphate moiety of its two glycerol units [100]. BMP prevents the formation of 7KC in murine macrophagic RAW264.7 cells [101]. Recently, it has been suggested that BMP could be useful in the prevention of SARS-CoV-2 infection [102]. It is hypothesised that BMP, which is present in the endosomes and is involved in the intracellular cholesterol trafficking, could act on the SARS-CoV-2 viral cycle and reduce virus production.

Among the synthetic molecules, dimethyl-fumarate (DMF), which is marketed under the name Tecfidera (Biogen) for the treatment of multiple sclerosis and psoriasis, exerts anti-inflammatory activities on T and B cells, as well as on dendritic cells via an inhibitory effect on nuclear factor kappa B (NF- $\kappa$ B) [103] and anti-oxidant activities by activating the erythroid 2-related factor 2 (Nrf2) signaling pathway. This induces the expression of several anti-oxidant enzymes (hemeoxygenase 1 (HO-1), catalase (CAT), superoxide dismutase (SOD), glutathione peroxidase (GPx)), the expression of phase II detoxifying enzymes, such as glutathione S-transferase (GST), and enzymes responsible for glutathione (GSH) synthesis, such as glutamine-cysteine ligase (GCL) and glutathione synthetase (GS) ([104] [105]). However, as the suppression of pyroptosis by DMF is independent of Nrf2, this supports that several signaling pathways can be activated by this molecule [106]. DMF also attenuates *in vitro* oxidative stress and cell death induction triggered by 7KC on 158N oligodendrocytes ([107] [108]). Noteworthy, it is currently reported that DMF could reduce lung alveolar cells damage in COVID-19 patients [109].

As 7KC accumulates in the lysosome, a strategy based on the use of bacterial enzymes targeting this organelle, defined as medical bioremediation, could also be used to inactivate 7KC and to prevent 7KC-induced cytotoxic activities ([110] [111]).

## Conclusion

Compared to oxysterols derived from oxidation on the alkyl chain of cholesterol, 7KC has weak anti-viral activities. In severe and often fatal forms of COVID-19, it is hypothesized that 7KC could be a predictive biomarker for assessing the severity of the disease and its progression to a fatal outcome. Indeed, while 7KC has slight antiviral activities, its cytotoxic activities can be considered dominant in many diseases associated with high levels of this oxysterol. These include age-related diseases, among which COVID-19 could be included. It can be assumed that decreasing the amount of 7KC by promoting its degradation and inhibiting or mitigating its toxicity could constitute adjuvant therapies that would if not eliminate, at least reduce, mortality and morbidity associated with COVID-19 infection. Due to the severity of the pandemic, it is reasonable to consider all hypotheses and explore all avenues to treat patients with COVID-19, especially those with potentially fatal outcomes. Therefore, given the biological activities of 7KC, adjuvant treatment with drugs that reduce 7KC toxicity could help to reduce the number of patients with severe forms of COVID-19 and thus reduce the number of patients on respiratory support in emergency departments.

**Acknowledgements:** The authors acknowledge the European Network for Oxysterols Research (ENOR: <http://oxysterols.net>) which permits discussion and collaborative research between the members. Khoulood Sassi (MS; Ph.D student), Amira Zarrouk (Ph.D), Thomas Nury (Ph.D), Valerio Leoni (M.D, Ph.D), John J. Mackrill (Ph.D), Mohammad Samadi (Ph.D), Anne Vejux (Ph.D) and Gérard Lizard (Ph.D, D.Sc) are all members of ENOR. The authors acknowledge the University Tunis El Manar for the financial support of Mrs Imen Ghzaïel (M.S, Ph.D student), Mrs Khoulood Sassi (M.S, Ph.D student) and Mr Mohamed Ksila (M.S, Ph.D student) as well as the “Association Bourguignonne pour les Applications des Sciences de l'Information en Médecine” (A.B.A.S.I.M).

**Conflict of interest:** The authors declare no conflict of interest.

## References

- [1] I. Björkhem, S. Meaney, Brain cholesterol: long secret life behind a barrier, *Arteriosclerosis, thrombosis, and vascular biology* 24(5) (2004) 806-815.
- [2] A. Zarrouk, M. Debbabi, M. Bezine, E.M. Karym, A. Badreddine, O. Rouaud, T. Moreau, M. Cherkaoui-Malki, M. El Ayeb, B. Nasser, M. Hammami, G. Lizard, Lipid Biomarkers in Alzheimer's Disease, *Current Alzheimer research* 15(4) (2018) 303-312.
- [3] B.C. Kwan, F. Kronenberg, S. Beddhu, A.K. Cheung, Lipoprotein metabolism and lipid management in chronic kidney disease, *Journal of the American Society of Nephrology : JASN* 18(4) (2007) 1246-1261.
- [4] K.A. Tallman, H.H. Kim, Z. Korade, T.C. Genaro-Mattos, P.A. Wages, W. Liu, N.A. Porter, Probes for protein adduction in cholesterol biosynthesis disorders: Alkynyl lanosterol as a viable sterol precursor, *Redox biology* 12 (2017) 182-190.
- [5] M.A. Mitsche, J.G. McDonald, H.H. Hobbs, J.C. Cohen, Flux analysis of cholesterol biosynthesis in vivo reveals multiple tissue and cell-type specific pathways, *eLife* 4 (2015) e07999.
- [6] Z. Hubler, R.M. Friedrich, J.L. Sax, D. Allimuthu, F. Gao, A.M. Rivera-León, M.J. Pleshinger, I. Bederman, D.J. Adams, Modulation of lanosterol synthase drives 24,25-epoxysterol synthesis and oligodendrocyte formation, *Cell chemical biology* (2021).
- [7] U. Diczfalusy, H. Nylén, P. Elander, L. Bertilsson, 4 $\beta$ -Hydroxycholesterol, an endogenous marker of CYP3A4/5 activity in humans, *British journal of clinical pharmacology* 71(2) (2011) 183-189.
- [8] T. Nury, M. Samadi, A. Zarrouk, J.M. Riedinger, G. Lizard, Improved synthesis and in vitro evaluation of the cytotoxic profile of oxysterols oxidized at C4 (4 $\alpha$ - and 4 $\beta$ -hydroxycholesterol) and C7 (7-ketocholesterol, 7 $\alpha$ - and 7 $\beta$ -hydroxycholesterol) on cells of the central nervous system, *European journal of medicinal chemistry* 70 (2013) 558-567.
- [9] L. Iuliano, Pathways of cholesterol oxidation via non-enzymatic mechanisms, *Chemistry and physics of lipids* 164(6) (2011) 457-468.
- [10] C. Zerbinati, L. Iuliano, Cholesterol and related sterols autoxidation, *Free radical biology & medicine* 111 (2017) 151-155.
- [11] A. Anderson, A. Campo, E. Fulton, A. Corwin, W.G. Jerome, 3rd, M.S. O'Connor, 7-Ketocholesterol in disease and aging, *Redox biology* 29 (2020) 101380.
- [12] B. Buttari, L. Segoni, E. Profumo, D. D'Arcangelo, S. Rossi, F. Facchiano, R. Businaro, L. Iuliano, R. Riganò, 7-Oxo-cholesterol potentiates pro-inflammatory signaling in human M1 and M2 macrophages, *Biochemical pharmacology* 86(1) (2013) 130-137.
- [13] W.J. Griffiths, E. Yutuc, J. Abdel-Khalik, P.J. Crick, T. Hearn, A. Dickson, B.W. Bigger, T. Hoi-Yee Wu, A. Goenka, A. Ghosh, S.A. Jones, D.F. Covey, D.S. Ory, Y. Wang, Metabolism of Non-Enzymatically Derived Oxysterols: Clues from sterol metabolic disorders, *Free radical biology & medicine* 144 (2019) 124-133.
- [14] A. Vejux, D. Abed-Vieillard, K. Hajji, A. Zarrouk, J.J. Mackrill, S. Ghosh, T. Nury, A. Yammine, M. Zaibi, W. Mihoubi, H. Bouchab, B. Nasser, Y. Grosjean, G. Lizard, 7-Ketocholesterol and 7 $\beta$ -hydroxycholesterol: In vitro and animal models used to characterize their activities and to identify molecules preventing their toxicity, *Biochemical pharmacology* 173 (2020) 113648.
- [15] A.J. Brown, W. Jessup, Oxysterols: Sources, cellular storage and metabolism, and new insights into their roles in cholesterol homeostasis, *Molecular aspects of medicine* 30(3) (2009) 111-122.
- [16] I. Tabas, Consequences of cellular cholesterol accumulation: basic concepts and physiological implications, *The Journal of clinical investigation* 110(7) (2002) 905-911.
- [17] S. Monier, M. Samadi, C. Prunet, M. Denance, A. Laubriet, A. Athias, A. Berthier, E. Steinmetz, G. Jürgens, A. Nègre-Salvayre, G. Bessède, S. Lemaire-Ewing, D. Néel, P. Gambert, G. Lizard, Impairment of

the cytotoxic and oxidative activities of 7 beta-hydroxycholesterol and 7-ketocholesterol by esterification with oleate, *Biochemical and biophysical research communications* 303(3) (2003) 814-824.

[18] H. Fuda, N.B. Javitt, K. Mitamura, S. Ikegawa, C.A. Strott, Oxysterols are substrates for cholesterol sulfotransferase, *Journal of lipid research* 48(6) (2007) 1343-1352.

[19] J.W. Lee, J.D. Huang, I.R. Rodriguez, Extra-hepatic metabolism of 7-ketocholesterol occurs by esterification to fatty acids via cPLA2 $\alpha$  and SOAT1 followed by selective efflux to HDL, *Biochimica et biophysica acta* 1851(5) (2015) 605-619.

[20] M.A. Rogers, J. Liu, B.L. Song, B.L. Li, C.C. Chang, T.Y. Chang, Acyl-CoA:cholesterol acyltransferases (ACATs/SOATs): Enzymes with multiple sterols as substrates and as activators, *The Journal of steroid biochemistry and molecular biology* 151 (2015) 102-107.

[21] J.W. Lee, H. Fuda, N.B. Javitt, C.A. Strott, I.R. Rodriguez, Expression and localization of sterol 27-hydroxylase (CYP27A1) in monkey retina, *Experimental eye research* 83(2) (2006) 465-469.

[22] G.Y. Heo, I. Bederman, N. Mast, W.L. Liao, I.V. Turko, I.A. Pikuleva, Conversion of 7-ketocholesterol to oxysterol metabolites by recombinant CYP27A1 and retinal pigment epithelial cells, *Journal of lipid research* 52(6) (2011) 1117-1127.

[23] Y. Wang, W.J. Griffiths, Unravelling new pathways of sterol metabolism: lessons learned from in-born errors and cancer, *Current opinion in clinical nutrition and metabolic care* 21(2) (2018) 90-96.

[24] W.J. Griffiths, Y. Wang, Oxysterols as lipid mediators: Their biosynthetic genes, enzymes and metabolites, *Prostaglandins & other lipid mediators* 147 (2020) 106381.

[25] H. Hughes, B. Mathews, M.L. Lenz, J.R. Guyton, Cytotoxicity of oxidized LDL to porcine aortic smooth muscle cells is associated with the oxysterols 7-ketocholesterol and 7-hydroxycholesterol, *Arteriosclerosis and thrombosis : a journal of vascular biology* 14(7) (1994) 1177-1185.

[26] S.M. Colles, J.M. Maxson, S.G. Carlson, G.M. Chisolm, Oxidized LDL-induced injury and apoptosis in atherosclerosis. Potential roles for oxysterols, *Trends in cardiovascular medicine* 11(3-4) (2001) 131-138.

[27] L. Malvitte, T. Montange, C. Joffre, A. Vejux, C. Maïza, A. Bron, C. Creuzot-Garcher, G. Lizard, [Analogies between atherosclerosis and age-related maculopathy: expected roles of oxysterols], *Journal francais d'ophtalmologie* 29(5) (2006) 570-578.

[28] A. Pariente, R. Peláez, Á. Pérez-Sala, I.M. Larráyoz, Inflammatory and cell death mechanisms induced by 7-ketocholesterol in the retina. Implications for age-related macular degeneration, *Experimental eye research* 187 (2019) 107746.

[29] A. Vejux, M. Samadi, G. Lizard, Contribution of cholesterol and oxysterols in the physiopathology of cataract: implication for the development of pharmacological treatments, *J Ophthalmol* 2011 (2011) 471947.

[30] H. Girão, M.C. Mota, J. Ramalho, P. Pereira, Cholesterol oxides accumulate in human cataracts, *Experimental eye research* 66(5) (1998) 645-652.

[31] G. Poli, F. Biasi, G. Leonarduzzi, Oxysterols in the pathogenesis of major chronic diseases, *Redox biology* 1(1) (2013) 125-130.

[32] G. Testa, E. Staurenghi, C. Zerbinati, S. Gargiulo, L. Iuliano, G. Giaccone, F. Fantò, G. Poli, G. Leonarduzzi, P. Gamba, Changes in brain oxysterols at different stages of Alzheimer's disease: Their involvement in neuroinflammation, *Redox biology* 10 (2016) 24-33.

[33] G. Testa, D. Rossin, G. Poli, F. Biasi, G. Leonarduzzi, Implication of oxysterols in chronic inflammatory human diseases, *Biochimie* 153 (2018) 220-231.

[34] T. Nury, A. Zarrouk, K. Ragot, M. Debbabi, J.M. Riedinger, A. Vejux, P. Aubourg, G. Lizard, 7-Ketocholesterol is increased in the plasma of X-ALD patients and induces peroxisomal modifications in microglial cells: Potential roles of 7-ketocholesterol in the pathophysiology of X-ALD, *The Journal of steroid biochemistry and molecular biology* 169 (2017) 123-136.

- [35] L. Iuliano, R. Monticolo, G. Straface, S. Zullo, F. Galli, M. Boaz, S. Quattrucci, Association of cholesterol oxidation and abnormalities in fatty acid metabolism in cystic fibrosis, *The American journal of clinical nutrition* 90(3) (2009) 477-484.
- [36] N. Aksu, A. Samadi, A. Yalçinkaya, T. Çetin, B. Eser, İ. Lay, T.N. Öziş, Y. Öztaş, S. Sabuncuoğlu, Evaluation of oxysterol levels of patients with silicosis by LC-MS/MS method, *Molecular and cellular biochemistry* 467(1-2) (2020) 117-125.
- [37] X. Rao, J. Zhong, A. Maiseyeu, B. Gopalakrishnan, F.A. Villamena, L.C. Chen, J.R. Harkema, Q. Sun, S. Rajagopalan, CD36-dependent 7-ketocholesterol accumulation in macrophages mediates progression of atherosclerosis in response to chronic air pollution exposure, *Circulation research* 115(9) (2014) 770-780.
- [38] A.C. Raghavamenon, X. Gao, D.S. Atkins-Ball, S. Varikuti, N.L. Parinandi, R.M. Uppu, 'Ozone-Specific' Oxysterols and Neuronal Cell Signaling, in: L.J. Berliner, N.L. Parinandi (Eds.), *Measuring Oxidants and Oxidative Stress in Biological Systems*, Springer. Copyright 2020, Springer Nature Switzerland AG., Cham (CH), 2020. pp. 109-122.
- [39] I.H. Dias, K. Borah, B. Amin, H.R. Griffiths, K. Sassi, G. Lizard, A. Iriondo, P. Martinez-Lage, Localisation of oxysterols at the sub-cellular level and in biological fluids, *The Journal of steroid biochemistry and molecular biology* 193 (2019) 105426.
- [40] F. Di Cara, P. Andreoletti, D. Trompier, A. Vejux, M.H. Bülow, J. Sellin, G. Lizard, M. Cherkaoui-Malki, S. Savary, Peroxisomes in Immune Response and Inflammation, *International journal of molecular sciences* 20(16) (2019).
- [41] A. Vejux, G. Lizard, Cytotoxic effects of oxysterols associated with human diseases: Induction of cell death (apoptosis and/or oncosis), oxidative and inflammatory activities, and phospholipidosis, *Molecular aspects of medicine* 30(3) (2009) 153-170.
- [42] A. Zarrouk, A. Vejux, J. Mackrill, Y. O'Callaghan, M. Hammami, N. O'Brien, G. Lizard, Involvement of oxysterols in age-related diseases and ageing processes, *Ageing research reviews* 18 (2014) 148-162.
- [43] D. Trompier, A. Vejux, A. Zarrouk, C. Gondcaille, F. Geillon, T. Nury, S. Savary, G. Lizard, Brain peroxisomes, *Biochimie* 98 (2014) 102-110.
- [44] A. Incani, L. Marras, G. Serreli, A. Ingianni, R. Pompei, M. Deiana, F. Angius, Human Herpesvirus 8 infection may contribute to oxidative stress in diabetes type 2 patients, *BMC research notes* 13(1) (2020) 75.
- [45] M.P. Ng, J.C. Lee, W.M. Loke, L.L. Yeo, A.M. Quek, E.C. Lim, B. Halliwell, R.C. Seet, Does influenza A infection increase oxidative damage?, *Antioxidants & redox signaling* 21(7) (2014) 1025-1031.
- [46] A. Civra, V. Cagno, M. Donalisio, F. Biasi, G. Leonarduzzi, G. Poli, D. Lembo, Inhibition of pathogenic non-enveloped viruses by 25-hydroxycholesterol and 27-hydroxycholesterol, *Scientific reports* 4 (2014) 7487.
- [47] V. Cagno, A. Civra, D. Rossin, S. Calfapietra, C. Caccia, V. Leoni, N. Dorma, F. Biasi, G. Poli, D. Lembo, Inhibition of herpes simplex-1 virus replication by 25-hydroxycholesterol and 27-hydroxycholesterol, *Redox biology* 12 (2017) 522-527.
- [48] A. Civra, R. Francese, P. Gamba, G. Testa, V. Cagno, G. Poli, D. Lembo, 25-Hydroxycholesterol and 27-hydroxycholesterol inhibit human rotavirus infection by sequestering viral particles into late endosomes, *Redox biology* 19 (2018) 318-330.
- [49] A. Marcello, A. Civra, R. Milan Bonotto, L. Nascimento Alves, S. Rajasekharan, C. Giacobone, C. Caccia, R. Cavalli, M. Adami, P. Brambilla, D. Lembo, G. Poli, V. Leoni, The cholesterol metabolite 27-hydroxycholesterol inhibits SARS-CoV-2 and is markedly decreased in COVID-19 patients, *Redox biology* 36 (2020) 101682.
- [50] R. Chilamakuri, S. Agarwal, COVID-19: Characteristics and Therapeutics, *Cells* 10(2) (2021).



- [51] H. Xu, L. Zhong, J. Deng, J. Peng, H. Dan, X. Zeng, T. Li, Q. Chen, High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa, *International journal of oral science* 12(1) (2020) 8.
- [52] W.J. Wiersinga, A. Rhodes, A.C. Cheng, S.J. Peacock, H.C. Prescott, Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): A Review, *Jama* 324(8) (2020) 782-793.
- [53] B. Hu, H. Guo, P. Zhou, Z.L. Shi, Characteristics of SARS-CoV-2 and COVID-19, *Nature reviews. Microbiology* 19(3) (2021) 141-154.
- [54] A. Sulejmani, E. Galimberti, C. Giacobone, A. Milano, E. Scopetta, J. Intra, R. Falbo, C. Sarto, V. Leoni, P. Brambilla, Baseline characteristics of COVID-19 Italian patients admitted to Desio Hospital, Lombardy: a retrospective study, *Scandinavian journal of clinical and laboratory investigation* 81(1) (2021) 18-23.
- [55] J. Intra, C. Sarto, E. Beck, N. Tiberti, V. Leoni, P. Brambilla, Bacterial and fungal colonization of the respiratory tract in COVID-19 patients should not be neglected, *American journal of infection control* 48(9) (2020) 1130-1131.
- [56] A. Samadi, S. Sabuncuoglu, M. Samadi, S.Y. Isikhan, S. Chirumbolo, M. Peana, I. Lay, A. Yalcinkaya, G. Bjørklund, A Comprehensive Review on Oxysterols and Related Diseases, *Current medicinal chemistry* 28(1) (2021) 110-136.
- [57] G. Costagliola, E. Spada, R. Consolini, Age-related differences in the immune response could contribute to determine the spectrum of severity of COVID-19, *Immunity, inflammation and disease* 9(2) (2021) 331-339.
- [58] A.P.S. Munro, S.N. Faust, COVID-19 in children: current evidence and key questions, *Current opinion in infectious diseases* 33(6) (2020) 540-547.
- [59] E. Rothenberg, Coronavirus Disease 19 from the Perspective of Ageing with Focus on Nutritional Status and Nutrition Management-A Narrative Review, *Nutrients* 13(4) (2021).
- [60] R. Channappanavar, S. Perlman, Age-related susceptibility to coronavirus infections: role of impaired and dysregulated host immunity, *The Journal of clinical investigation* 130(12) (2020) 6204-6213.
- [61] D. Lembo, V. Cagno, A. Civra, G. Poli, Oxysterols: An emerging class of broad spectrum antiviral effectors, *Molecular aspects of medicine* 49 (2016) 23-30.
- [62] S.Y. Bah, P. Dickinson, T. Forster, B. Kampmann, P. Ghazal, Immune oxysterols: Role in mycobacterial infection and inflammation, *The Journal of steroid biochemistry and molecular biology* 169 (2017) 152-163.
- [63] M. Blanc, W.Y. Hsieh, K.A. Robertson, K.A. Kropp, T. Forster, G. Shui, P. Lacaze, S. Watterson, S.J. Griffiths, N.J. Spann, A. Meljon, S. Talbot, K. Krishnan, D.F. Covey, M.R. Wenk, M. Craigon, Z. Ruzsics, J. Haas, A. Angulo, W.J. Griffiths, C.K. Glass, Y. Wang, P. Ghazal, The transcription factor STAT-1 couples macrophage synthesis of 25-hydroxycholesterol to the interferon antiviral response, *Immunity* 38(1) (2013) 106-118.
- [64] J.G. Cyster, E.V. Dang, A. Reboldi, T. Yi, 25-Hydroxycholesterols in innate and adaptive immunity, *Nature reviews. Immunology* 14(11) (2014) 731-743.
- [65] J. Zhao, J. Chen, M. Li, M. Chen, C. Sun, Multifaceted Functions of CH25H and 25HC to Modulate the Lipid Metabolism, Immune Responses, and Broadly Antiviral Activities, *Viruses* 12(7) (2020).
- [66] T. Willinger, Oxysterols in intestinal immunity and inflammation, *Journal of internal medicine* 285(4) (2019) 367-380.
- [67] C. Choi, D.K. Finlay, Diverse Immunoregulatory Roles of Oxysterols-The Oxidized Cholesterol Metabolites, *Metabolites* 10(10) (2020).
- [68] K.A. Willard, C.L. Elling, S.L. Stice, M.A. Brindley, The Oxysterol 7-Ketocholesterol Reduces Zika Virus Titers in Vero Cells and Human Neurons, *Viruses* 11(1) (2018).

- [69] H. Ohashi, F. Wang, F. Stappenbeck, K. Tsuchimoto, C. Kobayashi, W. Saso, M. Kataoka, M. Yamasaki, K. Kuramochi, M. Muramatsu, T. Suzuki, C. Sureau, M. Takeda, T. Wakita, F. Parhami, K. Watashi, Identification of Anti-Severe Acute Respiratory Syndrome-Related Coronavirus 2 (SARS-CoV-2) Oxysterol Derivatives In Vitro, *International journal of molecular sciences* 22(6) (2021).
- [70] S. Lemaire-Ewing, C. Prunet, T. Montange, A. Vejux, A. Berthier, G. Bessède, L. Corcos, P. Gambert, D. Néel, G. Lizard, Comparison of the cytotoxic, pro-oxidant and pro-inflammatory characteristics of different oxysterols, *Cell biology and toxicology* 21(2) (2005) 97-114.
- [71] C. Prunet, T. Montange, A. Vêjux, A. Laubriet, J.F. Rohmer, J.M. Riedinger, A. Athias, S. Lemaire-Ewing, D. Néel, J.M. Petit, E. Steinmetz, R. Brenot, P. Gambert, G. Lizard, Multiplexed flow cytometric analyses of pro- and anti-inflammatory cytokines in the culture media of oxysterol-treated human monocytic cells and in the sera of atherosclerotic patients, *Cytometry. Part A : the journal of the International Society for Analytical Cytology* 69(5) (2006) 359-373.
- [72] F. Brahmi, A. Vejux, R. Sghaier, A. Zarrouk, T. Nury, W. Meddeb, L. Rezig, A. Namsi, K. Sassi, A. Yammine, I. Badreddine, D. Vervandier-Fasseur, K. Madani, L. Boulekbache-Makhlouf, B. Nasser, G. Lizard, Prevention of 7-ketocholesterol-induced side effects by natural compounds, *Critical reviews in food science and nutrition* 59(19) (2019) 3179-3198.
- [73] T. Nury, A. Zarrouk, A. Yammine, J.J. Mackrill, A. Vejux, G. Lizard, Oxiaoptophagy: a type of cell death induced by some oxysterols, *British journal of pharmacology* (2020).
- [74] T. Nury, A. Yammine, I. Ghzaïel, K. Sassi, A. Zarrouk, F. Brahmi, M. Samadi, S. Rup-Jacques, D. Vervandier-Fasseur, J.P. Pais de Barros, V. Bergas, S. Ghosh, M. Majeed, A. Pande, A. Atanasov, S. Hammami, M. Hammami, J. Mackrill, B. Nasser, P. Andreoletti, M. Cherkaoui-Malki, A. Vejux, G. Lizard, Attenuation of 7-ketocholesterol- and 7 $\beta$ -hydroxycholesterol-induced oxiaoptophagy by nutrients, synthetic molecules and oils: potential for the prevention of age-related diseases, *Ageing research reviews* (2021) 101324.
- [75] I.M. Larrayoz, J.D. Huang, J.W. Lee, I. Pascual, I.R. Rodríguez, 7-ketocholesterol-induced inflammation: involvement of multiple kinase signaling pathways via NF $\kappa$ B but independently of reactive oxygen species formation, *Investigative ophthalmology & visual science* 51(10) (2010) 4942-4955.
- [76] M.L. Selley, J.A. McGuiness, N.G. Ardlie, The effect of cholesterol oxidation products on human platelet aggregation, *Thrombosis research* 83(6) (1996) 449-461.
- [77] B. Kosmider, J.E. Loader, R.C. Murphy, R.J. Mason, Apoptosis induced by ozone and oxysterols in human alveolar epithelial cells, *Free radical biology & medicine* 48(11) (2010) 1513-1524.
- [78] J.D. Huang, J. Amaral, J.W. Lee, I.R. Rodríguez, 7-Ketocholesterol-induced inflammation signals mostly through the TLR4 receptor both in vitro and in vivo, *PLoS one* 9(7) (2014) e100985.
- [79] E. Kahn, G. Lizard, F. Frouin, J.C. Bernengo, C. Souchier, G. Bessède, O. Clément, H. Siitari, P. Gambert, G. Frija, A. Todd-Pokropek, Confocal analysis of phosphatidylserine externalization with the use of biotinylated annexin V revealed with streptavidin-FITC, -europium, -phycoerythrin or -Texas Red in oxysterol-treated apoptotic cells, *Analytical and quantitative cytology and histology* 23(1) (2001) 47-55.
- [80] A. Attanzio, A. Frazzitta, A. Cilla, M.A. Livrea, L. Tesoriere, M. Allegra, 7-Keto-Cholesterol and Cholestan-3 $\beta$ , 5 $\alpha$ , 6 $\beta$ -Triol Induce Eryptosis through Distinct Pathways Leading to NADPH Oxidase and Nitric Oxide Synthase Activation, *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology* 53(6) (2019) 933-947.
- [81] V. Leoni, T. Nury, A. Vejux, A. Zarrouk, C. Caccia, M. Debbabi, A. Fromont, R. Sghaier, T. Moreau, G. Lizard, Mitochondrial dysfunctions in 7-ketocholesterol-treated 158N oligodendrocytes without or with  $\alpha$ -tocopherol: Impacts on the cellular profile of tricarboxylic cycle-associated organic acids, long chain saturated and unsaturated fatty acids, oxysterols, cholesterol and cholesterol precursors, *The Journal of steroid biochemistry and molecular biology* 169 (2017) 96-110.

- [82] E. Mayatepek, W.D. Lehmann, J. Fauler, D. Tsikas, J.C. Frölich, R.B. Schutgens, R.J. Wanders, D. Keppler, Impaired degradation of leukotrienes in patients with peroxisome deficiency disorders, *The Journal of clinical investigation* 91(3) (1993) 881-888.
- [83] C. Lismont, M. Nordgren, P.P. Van Veldhoven, M. Fransen, Redox interplay between mitochondria and peroxisomes, *Frontiers in cell and developmental biology* 3 (2015) 35.
- [84] T. Nury, R. Sghaier, A. Zarrouk, F. Ménétrier, T. Uzun, V. Leoni, C. Caccia, W. Meddeb, A. Namsi, K. Sassi, W. Mihoubi, J.M. Riedinger, M. Cherkaoui-Malki, T. Moreau, A. Vejux, G. Lizard, Induction of peroxisomal changes in oligodendrocytes treated with 7-ketocholesterol: Attenuation by  $\alpha$ -tocopherol, *Biochimie* 153 (2018) 181-202.
- [85] T. Nury, A. Yammine, F. Menetrier, A. Zarrouk, A. Vejux, G. Lizard, 7-Ketocholesterol- and 7 $\beta$ -Hydroxycholesterol-Induced Peroxisomal Disorders in Glial, Microglial and Neuronal Cells: Potential Role in Neurodegeneration : 7-ketocholesterol and 7 $\beta$ -hydroxycholesterol-Induced Peroxisomal Disorders and Neurodegeneration, *Advances in experimental medicine and biology* 1299 (2020) 31-41.
- [86] S.B. Kudchodkar, B. Levine, Viruses and autophagy, *Reviews in medical virology* 19(6) (2009) 359-378.
- [87] Y. Choi, J.W. Bowman, J.U. Jung, Autophagy during viral infection - a double-edged sword, *Nature reviews. Microbiology* 16(6) (2018) 341-354.
- [88] M. Indaram, W. Ma, L. Zhao, R.N. Fariss, I.R. Rodriguez, W.T. Wong, 7-Ketocholesterol increases retinal microglial migration, activation, and angiogenicity: a potential pathogenic mechanism underlying age-related macular degeneration, *Scientific reports* 5 (2015) 9144.
- [89] K. Gupta, R. Mahajan, Applications and Diagnostic Potential of Dried Blood Spots, *International journal of applied & basic medical research* 8(1) (2018) 1-2.
- [90] W.J. Griffiths, Y. Wang, Sterolomics in biology, biochemistry, medicine, *Trends in analytical chemistry* : TRAC 120 (2019) 115280.
- [91] H. Chugh, A. Awasthi, Y. Agarwal, R.K. Gaur, G. Dhawan, R. Chandra, A comprehensive review on potential therapeutics interventions for COVID-19, *European journal of pharmacology* 890 (2021) 173741.
- [92] P.K. Zarandi, M.R. Zinatizadeh, M. Zinatizadeh, M.H. Yousefi, N. Rezaei, SARS-CoV-2: From the pathogenesis to potential anti-viral treatments, *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 137 (2021) 111352.
- [93] N.M. Schmidt, P.A.C. Wing, J.A. McKeating, M.K. Maini, Cholesterol-modifying drugs in COVID-19, *Oxford open immunology* 1(1) (2020) iqaa001.
- [94] V. Mutemberezi, O. Guillemot-Legris, G.G. Muccioli, Oxysterols: From cholesterol metabolites to key mediators, *Progress in lipid research* 64 (2016) 152-169.
- [95] D.J. Drucker, Diabetes, obesity, metabolism, and SARS-CoV-2 infection: the end of the beginning, *Cell metabolism* 33(3) (2021) 479-498.
- [96] M. Debbabi, A. Zarrouk, M. Bezine, W. Meddeb, T. Nury, A. Badreddine, E.M. Karym, R. Sghaier, L. Bretillon, S. Guyot, M. Samadi, M. Cherkaoui-Malki, B. Nasser, M. Mejri, S. Ben-Hammou, M. Hammami, G. Lizard, Comparison of the effects of major fatty acids present in the Mediterranean diet (oleic acid, docosahexaenoic acid) and in hydrogenated oils (elaidic acid) on 7-ketocholesterol-induced oxiaoptophagy in microglial BV-2 cells, *Chemistry and physics of lipids* 207(Pt B) (2017) 151-170.
- [97] G. Lizard, N. Latruffe, D. Vervandier-Fasseur, Aza- and Azo-Stilbenes: Bio-Isosteric Analogs of Resveratrol, *Molecules (Basel, Switzerland)* 25(3) (2020).
- [98] M.C. Royer, S. Lemaire-Ewing, C. Desrumaux, S. Monier, J.P. Pais de Barros, A. Athias, D. Néel, L. Lagrost, 7-ketocholesterol incorporation into sphingolipid/cholesterol-enriched (lipid raft) domains is impaired by vitamin E: a specific role for alpha-tocopherol with consequences on cell death, *The Journal of biological chemistry* 284(23) (2009) 15826-15834.

- [99] K. Ragot, J.J. Mackrill, A. Zarrouk, T. Nury, V. Aires, A. Jacquin, A. Athias, J.P. Pais de Barros, A. Véjux, J.M. Riedinger, D. Delmas, G. Lizard, Absence of correlation between oxysterol accumulation in lipid raft microdomains, calcium increase, and apoptosis induction on 158N murine oligodendrocytes, *Biochemical pharmacology* 86(1) (2013) 67-79.
- [100] F. Hullin-Matsuda, C. Luquain-Costaz, J. Bouvier, I. Delton-Vandenbroucke, Bis(monoacylglycero)phosphate, a peculiar phospholipid to control the fate of cholesterol: Implications in pathology, Prostaglandins, leukotrienes, and essential fatty acids 81(5-6) (2009) 313-324.
- [101] M. Arnal-Levron, Y. Chen, P. Greimel, F. Calevro, K. Gaget, F. Riols, A. Batut, J. Bertrand-Michel, F. Hullin-Matsuda, V.M. Olkkonen, I. Delton, C. Luquain-Costaz, Bis(monoacylglycero)phosphate regulates oxysterol binding protein-related protein 11 dependent sterol trafficking, *Biochimica et biophysica acta. Molecular and cell biology of lipids* 1864(9) (2019) 1247-1257.
- [102] C. Luquain-Costaz, M. Rabia, F. Hullin-Matsuda, I. Delton, Bis(monoacylglycero)phosphate, an important actor in the host endocytic machinery hijacked by SARS-CoV-2 and related viruses, *Biochimie* 179 (2020) 247-256.
- [103] R. Gold, R.A. Linker, M. Stangel, Fumaric acid and its esters: an emerging treatment for multiple sclerosis with antioxidative mechanism of action, *Clinical immunology (Orlando, Fla.)* 142(1) (2012) 44-48.
- [104] F.V. Francisqueti-Ferron, A.J.T. Ferron, J.L. Garcia, C. Silva, M.R. Costa, C.S. Gregolin, F. Moreto, A.L.A. Ferreira, I.O. Minatel, C.R. Correa, Basic Concepts on the Role of Nuclear Factor Erythroid-Derived 2-Like 2 (Nrf2) in Age-Related Diseases, *International journal of molecular sciences* 20(13) (2019).
- [105] S.A. Scuderi, A. Ardizzone, I. Paterniti, E. Esposito, M. Campolo, Antioxidant and Anti-inflammatory Effect of Nrf2 Inducer Dimethyl Fumarate in Neurodegenerative Diseases, *Antioxidants (Basel, Switzerland)* 9(7) (2020).
- [106] F. Humphries, L. Shmuel-Galia, N. Ketelut-Carneiro, S. Li, B. Wang, V.V. Nemmara, R. Wilson, Z. Jiang, F. Khalighinejad, K. Muneeruddin, S.A. Shaffer, R. Dutta, C. Ionete, S. Pesiridis, S. Yang, P.R. Thompson, K.A. Fitzgerald, Succination inactivates gasdermin D and blocks pyroptosis, *Science (New York, N.Y.)* 369(6511) (2020) 1633-1637.
- [107] A. Zarrouk, T. Nury, E.M. Karym, A. Vejux, R. Sghaier, C. Gondcaille, P. Andreoletti, D. Trompier, S. Savary, M. Cherkaoui-Malki, M. Debbabi, A. Fromont, J.M. Riedinger, T. Moreau, G. Lizard, Attenuation of 7-ketocholesterol-induced overproduction of reactive oxygen species, apoptosis, and autophagy by dimethyl fumarate on 158N murine oligodendrocytes, *The Journal of steroid biochemistry and molecular biology* 169 (2017) 29-38.
- [108] R. Sghaier, T. Nury, V. Leoni, C. Caccia, J.P. Pais De Barros, A. Cherif, A. Vejux, T. Moreau, K. Limem, M. Samadi, J.J. Mackrill, A.S. Masmoudi, G. Lizard, A. Zarrouk, Dimethyl fumarate and monomethyl fumarate attenuate oxidative stress and mitochondrial alterations leading to oxiaoptophagy in 158N murine oligodendrocytes treated with 7 $\beta$ -hydroxycholesterol, *The Journal of steroid biochemistry and molecular biology* 194 (2019) 105432.
- [109] S.M. Hassan, M.J. Jawad, S.W. Ahjel, R.B. Singh, J. Singh, S.M. Awad, N.R. Hadi, The Nrf2 Activator (DMF) and Covid-19: Is there a Possible Role?, *Medical archives (Sarajevo, Bosnia and Herzegovina)* 74(2) (2020) 134-138.
- [110] J.M. Mathieu, J. Schloendorn, B.E. Rittmann, P.J. Alvarez, Medical bioremediation of age-related diseases, *Microbial cell factories* 8 (2009) 21.
- [111] S. Ghosh, S.K. Khare, Biodegradation of 7-Ketocholesterol by *Rhodococcus erythropolis* MTCC 3951: Process optimization and enzymatic insights, *Chemistry and physics of lipids* 207(Pt B) (2017) 253-259.
- [112] T. Nury, A. Yammine, I. Ghzaïel, K. Sassi, A. Zarrouk, F. Brahmi, M. Samadi, S. Rup-Jacques, D. Vervandier-Fasseur, J.P. Pais de Barros, V. Bergas, S. Ghosh, M. Majeed, A. Pande, A. Atanasov, S. Hammami, M. Hammami, J. Mackrill, B. Nasser, P. Andreoletti, M. Cherkaoui-Malki, A. Vejux, G. Lizard,

Attenuation of 7-ketocholesterol- and 7 $\beta$ -hydroxycholesterol-induced oxiaoptophagy by nutrients, synthetic molecules and oils: Potential for the prevention of age-related diseases, Ageing research reviews 68 (2021) 101324.

## Figures Legends

**Figure 1: Biosynthesis of 7-ketocholesterol.** The biosynthesis and the metabolism of 7-ketocholesterol (7KC) are described in the detail by Wang *et al.* [23], Brahmi *et al.* [72], Vejux *et al.* [14] and Griffiths *et al.* [24]. For detailed information on the biosynthesis and degradation of 7KC and 7 $\beta$ -hydroxycholesterol, please see the review from Nury T *et al.* [112].

**Figure 2: Potential contribution of 7-ketocholesterol in the progression and outcome of COVID-19.** Due to high serum levels of 7KC in COVID-19 patients, it is hypothesized that this oxysterol may be involved in the progression and outcome of the disease. The anti-viral effects of 7KC have been described, but these are often weak compared to those of oxysterols such as 24(S)-, 25- or 27-hydroxycholesterol ([61] [69]). On the other hand, 7KC at high concentrations is known for its toxicity (pro-oxidative and pro-inflammatory effects, induction of cell death) [73]. Because of these different activities, 7KC could contribute to the pathophysiology of COVID-19. It is suggested that the measurement of 7KC plasma levels could provide information on the evolution of patients with COVID-19 (use of 7KC as a prognostic biomarker) and that adjuvant therapies mitigating 7KC toxicity could benefit patients and help reduce the number of patients on ventilatory support in emergency departments.




Figure 1: Ghzaiel I et al.




Figure 2: Ghzaiei I et al.

# Hypothetic contribution of 7-ketocholesterol (7KC) to the outcome of COVID-19

