



**HAL**  
open science

# Optimisation et validation de CHEMFRANCE : un modèle régional de fugacité niveau III appliqué à la France

Sylvain Bintein

► **To cite this version:**

Sylvain Bintein. Optimisation et validation de CHEMFRANCE : un modèle régional de fugacité niveau III appliqué à la France. Toxicologie. Université Paul Verlaine - Metz, 1996. Français. NNT : 1996METZ030S . tel-01775906

**HAL Id: tel-01775906**

**<https://hal.univ-lorraine.fr/tel-01775906>**

Submitted on 24 Apr 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## AVERTISSEMENT

Ce document est le fruit d'un long travail approuvé par le jury de soutenance et mis à disposition de l'ensemble de la communauté universitaire élargie.

Il est soumis à la propriété intellectuelle de l'auteur. Ceci implique une obligation de citation et de référencement lors de l'utilisation de ce document.

D'autre part, toute contrefaçon, plagiat, reproduction illicite encourt une poursuite pénale.

Contact : [ddoc-theses-contact@univ-lorraine.fr](mailto:ddoc-theses-contact@univ-lorraine.fr)

## LIENS

Code de la Propriété Intellectuelle. articles L 122. 4

Code de la Propriété Intellectuelle. articles L 335.2- L 335.10

[http://www.cfcopies.com/V2/leg/leg\\_droi.php](http://www.cfcopies.com/V2/leg/leg_droi.php)

<http://www.culture.gouv.fr/culture/infos-pratiques/droits/protection.htm>

**THESE**

présentée à l'université de Metz  
pour l'obtention du Diplôme de Doctorat  
Spécialité Toxicologie de l'Environnement

Sylvain BINTEIN

**OPTIMISATION ET VALIDATION DE  
CHEMFRANCE: UN MODELE REGIONAL  
DE FUGACITE NIVEAU III APPLIQUE A LA  
FRANCE**

Soutenue le 31 Octobre 1996 devant la Commission d'Examen:

MM. J. DEVILLERS  
J.M. JOUANY  
W. KARCHER  
Mlle P. VASSEUR

Rapporteur  
Rapporteur

08 JAN. 1997

-1996-

| BIBLIOTHEQUE UNIVERSITAIRE<br>- METZ |            |
|--------------------------------------|------------|
| N° inv.                              | 19960605   |
| Cote                                 | S/Mz 96/30 |
| Loc                                  | Magasin    |

A Sandra

A mes parents

A mes frères et soeurs

Aux mineurs

Je remercie Monsieur J. Devillers de m'avoir accueilli dans son équipe et de m'avoir fourni un sujet de recherche aussi porteur et intéressant que celui-ci. Qu'il trouve ici l'expression de toute ma reconnaissance pour la compétence scientifique et le soutien quotidien qu'il a su m'apporter.

Je remercie Monsieur J.M. Jouany et Monsieur W. Karcher pour m'avoir fait l'honneur de juger cette thèse et de participer au jury. Je voudrais également exprimer ma reconnaissance à Monsieur W. Karcher pour avoir permis le financement par la Communauté Européenne de la plupart des études présentées dans cette thèse.

Je remercie Mademoiselle P. Vasseur de m'avoir permis d'intégrer cette formation doctorale et pour l'honneur qu'elle m'a fait de bien vouloir juger mon travail.

Enfin, je voudrais exprimer ma reconnaissance à toutes les personnes qui m'ont aidé à divers titres. Mes remerciements vont plus particulièrement:

- à mes collègues de laboratoire pour leur disponibilité et notamment à D. Domine pour son accueil chaleureux, son aide et ses produits régionaux,
- à tous les exilés du Nord et du Grand Nord pour leur amitié,
- aux "Bachelardiens" pour tous les moments de bonheur qu'ils m'ont procurés.

## SOMMAIRE

|  |              |
|--|--------------|
| <b>1. INTRODUCTION</b>   | <b>p. 6</b>  |
| <b>2. ESTIMATION DE LA BIOCONCENTRATION ET DE L'ADSORPTION</b>         | <b>p. 7</b>  |
| <b>3. CHEMFRANCE</b>   | <b>p. 10</b> |
| <b>4. "VALIDATION" DE CHEMFRANCE</b>                                   | <b>p. 11</b> |
| <b>5. ETUDE COMPARATIVE DES EQUATIONS ESTIMANT LA BIOCONCENTRATION</b> | <b>p. 13</b> |
| <b>6. CONCLUSION</b>   | <b>p. 15</b> |
| <b>7. REFERENCES</b>   | <b>p. 15</b> |

Cette thèse est constituée des articles suivants:

**ARTICLE I:** Bintein, S., Devillers, J. et Karcher, W. (1993). Nonlinear dependence of fish bioconcentration on *n*-octanol/water partition coefficient. *SAR QSAR Environ. Res.* **1**, 29-39. **p. 19**

**ARTICLE II:** Bintein, S. et Devillers, J. (1994). QSAR for organic chemical sorption in soils and sediments. *Chemosphere* **28**, 1171-1188. **p. 31**

**ARTICLE III:** Devillers, J., Bintein, S. et Karcher, W. (1995). CHEMFRANCE: A regional level III fugacity model applied to France. *Chemosphere* **30**, 457-476. **p. 50**

**ARTICLE IV:** Bintein, S. et Devillers, J. (1996). Evaluating the environmental fate of lindane in France. *Chemosphere* **32**, 2427-2440. **p. 71**

**ARTICLE V:** Bintein, S. et Devillers, J. (1996). Evaluating the environmental fate of atrazine in France. *Chemosphere* **32**, 2441-2456. **p. 86**

**ARTICLE VI:** Devillers, J., Bintein, S. et Domine, D. (1996). Modeling the environmental fate of atrazine. Présenté au 211<sup>ème</sup> Congrès de l'American Chemical Society (Nouvelles-Orléans, Louisiane, 24-28 Mars, 1996).

Soumis dans *Sulfur Triazine Herbicides: Risk Assessment* (L. Ballantine, J. McFarland et D. Hackett, Eds.). ACS Symposium Series, ACS Books, Washington, D.C. **p. 103**

**ARTICLE VII:** Devillers, J., Bintein, S. et Domine, D. (1996). Comparison of BCF models based on log P. *Chemosphere* (sous presse). **p. 117**

## 1. INTRODUCTION

Le comportement d'une substance chimique dans l'environnement est complexe. Il met en jeu de nombreux phénomènes liés à ses propriétés physico-chimiques et aux caractéristiques des écosystèmes. Pour les produits que l'on souhaite introduire sur le marché ou ceux déjà présents, l'estimation de leur devenir dans l'environnement est primordiale. Le nombre considérable de composés chimiques employés dans nos activités rendant inconcevable la réalisation exclusive d'études *in situ*, on fait de plus en plus appel aux techniques de modélisation. La souplesse de ces approches permet de concevoir des modèles globaux (e.g., Mackay *et al.*, 1992) ou restreints à un phénomène particulier (e.g., PRZM (Carsel *et al.*, 1985)). Ces outils mathématiques sont utilisés pour comparer le comportement de plusieurs molécules, rechercher des informations avant la réalisation d'essais en laboratoire ou sur le terrain, ou encore identifier des compartiments ou des secteurs à risques. Cependant, l'emploi de modèles dans les analyses de risques exige une grande sûreté dans les résultats obtenus et une connaissance parfaite de leurs limites d'utilisation. Dans ces conditions, l'objet de notre travail a été d'optimiser et de valider CHEMFRANCE, un modèle régional de fugacité niveau III appliqué à la France (Chancrogne, 1991). Nous devons élaborer de nouvelles relations structure-bioconcentration et structure-adsorption possédant un large domaine d'application, mais également confirmer les fondements théoriques et empiriques de CHEMFRANCE en mettant en parallèle les observations faites en laboratoire ou sur le terrain et les résultats des simulations effectuées sur ordinateur.



## 2. ESTIMATION DE LA BIOCONCENTRATION ET DE L'ADSORPTION (Articles I et II)

Les modèles de distribution sont pour la plupart fondés sur la notion de répartition compartimentale nécessitant l'utilisation de coefficients de partage. Chacun d'eux, défini pour deux phases, gouverne la distribution à l'équilibre du composé et donc de sa concentration finale dans les différents compartiments de l'environnement. Ainsi, pour estimer le potentiel d'accumulation des molécules dans les sols, les sédiments et les matières en suspension, on utilise le coefficient d'adsorption (i.e.,  $K_p$ ). De même, le facteur de bioconcentration (i.e., BCF) permet d'évaluer l'accumulation des substances chimiques chez les êtres vivants par des voies non-alimentaires. La plupart des études sur la bioconcentration sont réalisées sur le poisson, compte tenu de son importance économique et de la disponibilité de tests standardisés. De ce fait, cet organisme est fréquemment utilisé pour représenter le compartiment biotique dans les modèles de distribution. Le  $K_p$  et le BCF peuvent être déterminés expérimentalement ou par des relations de type structure-activité (i.e., QSAR). Depuis le début des années 80, l'importance considérable de ces deux paramètres a engendré la publication d'une multitude d'équations (Lyman *et al.*, 1990; ECETOC, 1995; Güsten et Sabljic, 1995). En dépit de ce fait, des problèmes de modélisation restaient en suspens. La première partie de notre travail a donc consisté à établir de nouvelles équations possédant un domaine d'application plus étendu. Pour estimer la bioconcentration, les modèles régressifs linéaires entre les transformations logarithmiques des BCF et des coefficients de partage *n*-octanol/eau (i.e., log

P) sont couramment utilisés. Cependant, pour les substances très hydrophobes ( $\log P > 6$ ), ils ne sont plus applicables (Banerjee et Baughman, 1991). Cette cassure dans la relation linéaire entre une activité biologique et le caractère hydrophobe des molécules a été soulignée dans diverses études (Devillers et Lipnick, 1990; Hansch et Leo, 1995). En pharmacologie et en toxicologie, ce problème de linéarité a été résolu grâce à l'utilisation de modèles paraboliques et bilinéaires (Kubinyi, 1993; Hansch et Leo, 1995). **L'article I** présente l'aptitude de ces techniques non-linéaires à modéliser la bioconcentration chez le poisson. Dans un premier temps, une analyse bibliographique a été réalisée pour obtenir de nombreuses valeurs de BCF et de  $\log P$ . Une banque de données contenant 154 molécules dont les valeurs de BCF suivaient des critères de sélection précis a été constituée. A partir de cet échantillon, trois équations (i.e., linéaire, parabolique, bilinéaire) reliant le BCF au  $\log P$  ont été développées. L'analyse des paramètres statistiques et des résidus correspondant à ces relations a mis en évidence, le faible pouvoir prédictif du modèle linéaire dans notre étendue de  $\log P$  (i.e., 1,12 à 8,60) et la supériorité du modèle bilinéaire face à l'équation parabolique. Un échantillon test constitué de 29 molécules a permis de confirmer ces conclusions et d'étendre la validité de la relation bilinéaire aux substances possédant une valeur de  $\log P$  comprise entre 0,39 et 9,50.

Les problèmes de modélisation du coefficient d'adsorption sont différents. En effet, la trop grande spécificité et l'absence de paramètres prenant en compte le degré d'ionisation des acides et des bases correspondent aux principaux défauts des équations existantes. La finalité de l'étude décrite dans **l'article II** était donc de pallier ces problèmes en élaborant un modèle général

intégrant le caractère hydrophobe et le potentiel d'ionisation des substances (i.e., log P et pKa, respectivement) ainsi que le pH et le pourcentage de carbone organique (i.e., %OC) pour décrire le substrat. Ce modèle a été construit à partir de 229 valeurs de Kp correspondant à 53 composés et a été testé sur 500 Kp mesurés pour 87 molécules. Ces données ont été extraites d'articles originaux et non de compilations. Trois étapes ont été nécessaires pour obtenir une équation applicable aux composés ionisés et non-ionisés. Dans un premier temps, une relation intégrant le log P et le %OC a confirmé la nécessité d'introduire des facteurs correctifs intégrant le degré de dissociation des molécules. En effet, il est admis que pour les substances capables de former des ions (i.e., acides, bases) des phénomènes de répulsion ou d'affinité avec les colloïdes du substrat chargés négativement induisent, respectivement, une baisse du coefficient d'adsorption pour les substances présentant une charge négative et une augmentation pour celles chargées positivement (Bailey *et al.*, 1968; Jafvert, 1990). Pour prendre en compte ces phénomènes, le pKa des molécules et le pH de la solution des sols ou des sédiments ont été incorporés dans une seconde équation. Leur intégration a provoqué un net accroissement de la qualité des résidus pour les composés ionisés, tout en conservant un haut pouvoir prédictif pour les autres substances. Cependant, une analyse précise des résidus a mis en évidence une sous-estimation générale des valeurs de Kp pour les bases. Cette tendance s'explique par le fait que l'adsorption de ces molécules dépend de l'acidité de surface inférieure de deux unités de pH à celle de la solution. L'intégration de cette observation empirique dans notre équation a permis d'obtenir une distribution aléatoire des résidus pour les bases. La supériorité de cette

dernière équation a été confirmée par les simulations réalisées sur notre échantillon test, qui ont également permis d'étendre son domaine d'application à de nouvelles classes de substances chimiques et à des conditions pédologiques très variées.

### 3. CHEMFRANCE (Article III)

Les modèles de distribution peuvent être ponctuels ou globaux. Les premiers permettent d'obtenir, en fonction de leur degré de complexité, des résultats précis et une bonne compréhension des phénomènes étudiés. Cependant, la masse d'information nécessaire à leur emploi limite leur domaine d'application. A l'inverse, la finalité des modèles globaux est d'estimer le comportement des molécules dans un environnement représenté par plusieurs compartiments. Pour cela, un compromis entre la réduction des informations nécessaires à leur fonctionnement et la pertinence des résultats fournis doit être trouvé. Ces outils de simulation sont fondés, en quasi-totalité, sur le concept de fugacité (Mackay, 1991; Cowan *et al.*, 1995). Parmi eux, on distingue les modèles régionaux et les modèles d'évaluation employant un "monde unité" représentant un environnement hypothétique. Les premiers tentent de s'approcher le plus possible d'une situation réelle afin d'obtenir des informations comparables aux données expérimentales. Les seconds permettent uniquement de comparer les molécules entre elles ou d'identifier les principaux processus mis en jeu. L'article III présente CHEMFRANCE, un modèle régional de fugacité niveau III appliqué à la France. Le niveau III

signifie que le système est à l'état stable (i.e., flux d'entrée = flux de sortie) mais que des transferts entre les compartiments sont possibles (e.g., volatilisation, ruissellement, lessivage). CHEMFRANCE comprend six compartiments globaux (i.e., air, eaux de surface, sol, sédiments, eaux côtières, eaux souterraines). Les quatre premiers sont constitués de sous-compartiments (e.g., poissons, matières en suspension). CHEMFRANCE permet de simuler, pour les quatre saisons, le comportement des molécules en France et dans 12 régions françaises définies en fonction de leurs caractéristiques pédologiques, climatiques et hydrologiques. Les paramètres environnementaux (e.g., pH du sol, volume des eaux de surface) indispensables au fonctionnement de CHEMFRANCE ont été déterminés pour chacune de ces régions. Ainsi, l'utilisateur doit fournir simplement les propriétés physico-chimiques et les quantités d'émissions de la molécule à analyser pour obtenir la concentration, la quantité absolue, les cinétiques d'élimination et d'échange pour chaque compartiment...

#### **4. "VALIDATION" DE CHEMFRANCE (Articles IV à VI)**

Afin d'obtenir un outil de simulation efficace et fiable, tout logiciel doit être vérifié et testé (Devillers *et al.*, 1992, 1995; Fredenslund *et al.*, 1995; Vincent *et al.*, 1996). Pour les modèles de distribution, la validation des informations produites présente le niveau de difficulté le plus élevé. La complexité de cette procédure est due à la rareté mais surtout à la variabilité spatiale et temporelle des données expérimentales utilisées comme éléments de

comparaison. Dans ces conditions, la validité d'un modèle ne pourra jamais être pleinement affirmée. On pourra simplement confirmer la pertinence des résultats obtenus en démontrant les concordances entre observations et prédictions. Dans ce contexte, le pouvoir de simulation de CHEMFRANCE a été évalué sur le lindane (Article IV) et l'atrazine (Articles V et VI). Du fait de l'utilisation massive de ces pesticides, de multiples études ont été entreprises afin de connaître avec précision tous les aspects de leur comportement dans l'environnement. Une synthèse de ces informations nous a permis d'établir un profil environnemental assez précis de ces molécules. Une fois cette base de travail établie, les données indispensables au fonctionnement du programme (i.e., propriétés physico-chimiques, tonnages émis) ont été déterminées le plus précisément possible pour faciliter la comparaison entre les valeurs expérimentales et celles calculées par CHEMFRANCE. Quatre scénarios intégrant les méthodes d'application de ces substances actives et leurs déversements dans les eaux de surface ont été simulés. Cette approche possède deux intérêts. D'une part, elle permet d'identifier les risques de contamination et les principaux processus mis en jeu en fonction des compartiments d'émission et d'autre part, d'obtenir une variabilité dans les résultats. De fortes similitudes entre les résultats extraits de ces simulations et ceux obtenus en laboratoire ou sur le terrain ont été observées. Ainsi, le lindane est présent principalement dans les sols, les sédiments et les éléments biotiques du fait de ses capacités d'adsorption et de bioconcentration. A l'inverse, l'atrazine se retrouve principalement dans les milieux aquatiques bien que la contamination atmosphérique ne puisse pas être écartée. Cette molécule est faiblement adsorbée par les sols induisant par conséquent un fort

potentiel de ruissellement et de lessivage. Enfin, elle est très peu bioaccumulée. Une analyse sensitive portant sur le temps de demi-vie de l'atrazine dans les sols est présentée dans l'article VI. Cette étude montre que la concentration dans les sols et les eaux souterraines et les processus de volatilisation au niveau du sol, de ruissellement et de lessivage sont les paramètres les plus sensibles aux variations de la vitesse de dégradation de l'atrazine dans les sols.

## **5. ETUDE COMPARATIVE DES EQUATIONS ESTIMANT LA BIOCONCENTRATION (Article VII)**

Les modèles de type QSAR peuvent s'avérer des outils précieux pour évaluer les risques écotoxicologiques s'ils sont performants et surtout utilisés à bon escient. Il est donc indispensable de connaître précisément leurs limites d'utilisation. Dans ce contexte, nous avons entrepris de comparer différents modèles permettant d'estimer la bioconcentration des molécules organiques. Tous les modèles testés utilisaient le log P comme descripteur moléculaire.

La sélection des équations a été motivée par une étude d'occurrences effectuée à partir de rapports élaborés pour des organisations officielles (e.g., OCDE, EPA, UE). L'intégration de ces équations dans les modèles de distribution a également influencé notre choix. La réalisation de nos études comparatives soulevait un certain nombre de problèmes méthodologiques. En effet, les données expérimentales sur la bioconcentration des molécules sont relativement peu nombreuses et ont donc largement été utilisées pour élaborer

les équations à comparer. Or, lorsque l'on veut estimer le pouvoir prédictif d'un modèle de type QSAR, il est indispensable d'utiliser un échantillon test constitué de molécules n'appartenant pas à l'échantillon d'apprentissage. La solution optimale aurait été de mesurer des valeurs de BCF pour de nouvelles substances. Dans le cadre de notre travail, pour des raisons de temps et de coût, une telle alternative ne pouvait être envisagée. Nous n'avons donc pas cherché à sélectionner une seule valeur par composé mais plutôt, dans la mesure du possible, à retenir un ensemble de résultats expérimentaux cohérents permettant de fournir un profil de bioconcentration pour une molécule donnée. Une étude bibliographique visant à récolter un grand nombre de valeurs expérimentales a été réalisée à partir de publications originales et non de compilations afin d'obtenir des informations sur les conditions expérimentales. L'article VII présente une étude comparative réalisée sur sept équations linéaires ou non-linéaires permettant d'estimer le BCF à partir du log P (Veith *et al.*, 1979, 1980; Mackay, 1982; Connell et Hawker, 1988; Isnard et Lambert, 1988; Nendza, 1991; Bintein *et al.*, 1993). Cette analyse a été effectuée à partir d'une banque de données constituée de 342 valeurs de BCF provenant d'essais dynamiques et 94 déterminées par essais statiques ou semi-statiques. Pour des valeurs de log P inférieures à 6, les différents modèles testés donnent des résultats équivalents. Par contre, pour les substances très hydrophobes (i.e., log P > 6), les relations linéaires ne sont plus applicables et sont nettement devancées par les équations non-linéaires. Parmi celles-ci, notre équation possède le meilleur pouvoir prédictif et le plus large domaine d'application.



## 6. CONCLUSION

En élaborant de nouvelles relations structure-bioconcentration et structure-adsorption nous avons accru le domaine d'application et le pouvoir de simulation de CHEMFRANCE. Les études réalisées sur le lindane et l'atrazine ont démontré le haut potentiel de simulation de CHEMFRANCE et ont permis d'effectuer une synthèse du comportement dans l'environnement de ces pesticides. Cependant, nous envisageons de poursuivre ce processus de validation avec d'autres molécules aux propriétés physico-chimiques variées pour définir d'une façon plus précise les limites d'utilisation de ce logiciel. Nous prévoyons également d'intégrer dans CHEMFRANCE la méthode SIRIS (Vaillant *et al.*, 1995) ainsi que des informations écotoxicologiques afin de mieux appréhender les risques (éco)toxicologiques et de créer un outil d'aide à la décision plus facile à utiliser.

## 7. REFERENCES

- Bailey, G.W., White, J.L. et Rothberg, T. (1968). Adsorption of organic herbicides by montmorillonite: Role of pH and chemical character of adsorbate. *Soil Sci. Soc. Amer. Proc.* **32**, 222-234.
- Banerjee, S. et Baughman, G.L. (1991). Bioconcentration factors and lipid solubility. *Environ. Sci. Technol.* **25**, 536-539.
- Bintein, S., Devillers, J. et Karcher, W. (1993). Nonlinear dependence of fish bioconcentration on *n*-octanol/water partition coefficient. *SAR QSAR Environ. Res.* **1**, 29-39.

Carsel, R.F., Mulkey, L.A., Lorber, M.N. et Baskin, L.B. (1985). The pesticide root zone model (PRZM): A procedure for evaluating pesticide leaching threats to groundwater. *Ecol. Modelling* **30**, 49-69.

Chancrogne, A. (1991). *CHEMFRANCE: Un modèle de fugacité niveau III appliqué à la France*. D.E.A. Toxicologie de l'Environnement. Centre des Sciences de l'Environnement, Metz, p. 43.

Connell, D.W. et Hawker, D.W. (1988). Use of polynomial expressions to describe the bioconcentration of hydrophobic chemicals by fish. *Ecotoxicol. Environ. Safety* **16**, 242-257.

Cowan, C.E., Mackay, D., Feijtel, T.C.J., van de Meent, D., Di Guardo, A., Davies, J. et Mackay, N. (1995). *The Multi-Media Fate Model: A Vital Tool for Predicting the Fate of Chemicals*. SETAC Press, Pensacola, p. 78.

Devillers, J. et Lipnick, R.L. (1990). Practical applications of regression analysis in environmental QSAR studies. Dans, *Practical Applications of Quantitative Structure-Activity Relationships (QSAR) in Environmental Chemistry and Toxicology* (W. Karcher et J. Devillers, Eds.). Kluwer Academic Publishers, Dordrecht, pp. 129-143.

Devillers, J., Bintein, S. et Domine, D. (1995). *Evaluation of USES 1.0. French Contribution to the Development of EU Risk Assessment Guidances for New and Existing Substances*. Ministère de l'Environnement (BSPC)/CTIS, p. 39.

Devillers, J., Domine, D. et Bintein, S. (1992). *Study of Four Exposure Models applied to Ten Chemicals Substances*. Ministère de l'Environnement (BSPC)/CTIS, p. 51.

ECETOC (1995). *The Role of Bioaccumulation in Environmental Risk Assessment: The Aquatic Environment and Related Food Webs*. Technical Report No. 67, p. 112.

Fredenslund, F.C., Severinsen, M. et Andersen, M.B. (1995). *Evaluation of the SimpleBox Model for Danish Conditions*. Environmental Project n° 307. Ministry of Environment and Energy, Danish Environmental Protection Agency, Copenhagen, p. 96.

Güsten, H. et Sabljic, A. (1995). QSARs for soil sorption. Dans, *Overview of*

*Structure-Activity Relationships for Environmental Endpoints. Part 1: General Outline and Procedure.* Report of the EU-DG-XII Project QSAR for Predicting Fate and Effects of Chemicals in the Environment. Contract # EV5V-CT92-0211. Chapter III.

Hansch, C. et Leo, A. (1995). *Exploring QSAR. Fundamentals and Applications in Chemistry and Biology.* ACS, Washington, p. 557.

Isnard, P. et Lambert, S. (1988). Estimating bioconcentration factors from octanol-water partition coefficient and aqueous solubility. *Chemosphere* **17**, 21-34.

Jafvert, C.T. (1990). Sorption of organic acid compounds to sediments: Initial model development. *Environ. Toxicol. Chem.* **9**, 1259-1268.

Kubinyi, H. (1993). *QSAR: Hansch Analysis and Related Approaches.* VCH, Weinheim, p. 240.

Lyman, W.J., Reehl, W.F. et Rosenblatt, D.H. (1990). *Handbook of Chemical Property Estimation Methods.* ACS, Washington.

Mackay, D. (1982). Correlation of bioconcentration factors. *Environ. Sci. Technol.* **16**, 274-278.

Mackay, D. (1991). *Multimedia Environmental Models. The Fugacity Approach.* Lewis Publishers, Chelsea, p. 257.

Mackay, D., Paterson, S. et Shiu, W.Y. (1992). Generic models for evaluating the regional fate of chemicals. *Chemosphere* **24**, 695-717.

Nendza, M. (1991). QSARs of bioconcentration: Validity assessment of log Pow/log BCF correlations. Dans, *Bioaccumulation in Aquatic Systems* (R. Nagel et R. Loskill, Eds.). VCH, Weinheim, pp. 43-66.

Vaillant, M., Jouany, J.M. et Devillers, J. (1995). A multicriteria estimation of the environmental risk of chemicals with SIRIS method. *Toxicol. Model.* **1**, 57-72.

Veith, G.D., DeFoe, D.L. et Bergstedt, B.V. (1979). Measuring and estimating the bioconcentration factor of chemicals in fish. *J. Fish. Res. Board Can.* **36**, 1040-1048.

Veith, G.D., Macek, K.J., Petrocelli, S.R. et Carroll, J. (1980). An evaluation of using partition coefficients and water solubility to estimate

bioconcentration factors for organic chemicals in fish. Dans, *Aquatic Toxicology*, ASTM STP 707 (J.G. Eaton, P.R. Parrish et A.C. Hendricks, Eds.). American Society for Testing and Materials, pp. 116-129.

Vincent, R., Devillers, J., Bintein, S., Sandino, J.P. et Karcher, W. (1996). *Evaluation of EASE 2.0. Assessment of Occupational Exposure to Chemicals*. INRS/CTIS/ECB, p. 48.

# ARTICLE I

# NONLINEAR DEPENDENCE OF FISH BIOCONCENTRATION ON *n*-OCTANOL/WATER PARTITION COEFFICIENT

S. BINTEIN and J. DEVILLERS\*

*CTIS, 21 rue de la Bannière, 69003 Lyon (France).*

W. KARCHER

*Commission of the European Communities, Joint Research Centre,  
ISPRA Establishment, I-21020 Ispra Varese (Italy).*

*(accepted July 28, 1992)*

The log-log relationship between the bioconcentration tendency of organic chemicals in fish and the *n*-octanol/water partition coefficients breaks down for very hydrophobic compounds. The use of parabolic and bilinear models allows this problem to be overcome. The QSAR equation  $\log \text{BCF} = 0.910 \log P - 1.975 \log (6.8 \cdot 10^{-7} P + 1) - 0.786$  ( $n = 154$ ;  $r = 0.950$ ;  $s = 0.347$ ;  $F = 463.51$ ) was found to be a good predictor of bioconcentration in fish.

KEY WORDS: Bioconcentration; BCF; fish; bilinear model; linear model; parabolic model.

## INTRODUCTION

Bioconcentration is the process of accumulation of chemicals by organisms through nondietary routes.<sup>1</sup> In aquatic ecosystems, the bioconcentration factor (BCF) of an organic chemical is defined as the ratio of its concentration in a target organism to that in water at steady state.<sup>2</sup> Typically, fish are the target organisms of BCF assessments due to their importance as a human food source and the availability of standardized testing protocols.<sup>1</sup> The most common method for estimating chemicals' BCFs consists of establishing correlations between BCFs and hydrophobicity of the chemicals. The majority of these are obtained from linear regression models between the log transformations of the BCF values and the *n*-octanol/water partition coefficients ( $\log P$ ) of the chemicals.<sup>3-6</sup> The regression equations have the following general form:

$$\log \text{BCF} = a \log P + b \quad (1)$$

\* Author to whom all correspondence should be addressed.

where  $a$  and  $b$  are constants. In most published BCF equations, the slope ( $a$ ) is  $< 1$  and the intercept ( $b$ ) is negative.<sup>6</sup> Equation 1 breaks down for strongly hydrophobic chemicals ( $\log P > 6$ ).<sup>7</sup> The break in the linear relationship between a biological activity (BA) and the hydrophobic character of the organic molecules has been stressed by different authors in numerous QSAR studies.<sup>8</sup> Hansch initiated the use of a simple parabolic model<sup>8</sup> in  $\log P$  (Eq. 2) to overcome this problem.

$$\log BA = a \log P + b (\log P)^2 + c \quad (2)$$

Kubinyi<sup>9-12</sup> subsequently proposed a better model (Eq. 3) that adequately accounted for the fact that the biological activity and partition coefficient initially vary in a linear fashion on a log-log scale, reaching an optimum value, followed by a second decreasing linear portion.

$$\log BA = a \log P - b \log (\beta P + 1) + c \quad (3)$$

This bilinear model (Eq. 3) has been successfully used in many drug design and environmental QSAR studies.<sup>13-18</sup>


The aim of this paper is to compare the performance of the parabolic and bilinear models to overcome the "cut-off" problem encountered in the estimation of BCF values from the *n*-octanol/water partition coefficient ( $\log P$ ).

## MATERIALS AND METHODS

### Data selection

BCF values for 154 chemicals (Figure 1) were obtained from an extensive compilation.<sup>19</sup> Data were selected according to the following criteria:

1) Only experimental data, measured in whole fish (wet weight), and obtained under flow-through conditions, were used.



**Figure 1** Data set constitution. (PCB = polychlorobiphenyls; PBB = polybromobiphenyls; PCDD = polychlorodibenzo-*p*-dioxins; PCDF = polychlorodibenzofurans; PAC = polyaromatic compounds).

2) Data were included only if a steady state was reached or when BCF values were obtained from the kinetic method.

3) BCF data were rejected if contamination by food and/or presence of adsorbents (e.g., suspended sediments) was suspected.

4) Only studies using freshwater fish were considered. Five families were represented in the selected data:

- Salmonidae: rainbow trout, whitefish,
- Cyprinidae: fathead minnow, goldfish, carps (from different geographical origins), zebrafish, bleak, and topmouth gudgeon
- Centrarchidae: bluegill sunfish,
- Poeciliidae: guppy,
- Cyprinodontidae: American flagfish and killifish.

Fish species distribution in the study is summarized in Figure 2. The log  $P$  values were retrieved from an extensive bibliographical review aimed at preferentially selecting data obtained by direct methods (e.g., shake-flask, generator column).

#### Model development

Description and calculation procedures for the bilinear model can be found in numerous comprehensive papers.<sup>9-12</sup> Briefly, the bilinear model is derived from the McFarland multicompartamental model which allows the estimation of the probability  $p$  of a molecule crossing aqueous-lipid interfaces and reaching its receptor site. Thus, if we consider a hypothetical biological system made up of alternating aqueous phases  $A$  and lipid (membrane) phases  $L$  (Figure 3), the probability of a





Figure 2 Number of BCF values per species.





**Figure 3** The McFarland hypothetical biological system ( $k_1$  and  $k_2$  are the rate constants of the distribution process).

molecule leaving  $A_0$  and entering into  $L_1$  is:

$$p_{0,1} = \frac{k_1}{k_1 + k_2} \quad (4)$$

In the same way, the probability that a molecule goes from  $L_1$  to  $A_2$  is:

$$p_{1,2} = \frac{k_2}{k_1 + k_2} \quad (5)$$

If both parts of the fractions in Eqs. 4 and 5 are divided by  $k_2$  and if  $k_1/k_2$  is replaced by  $P$  (the partition coefficient), we obtain:

$$p_{0,1} = \frac{P}{P + 1} \quad (6)$$

$$p_{1,2} = \frac{1}{P + 1} \quad (7)$$

Therefore, the probability that a molecule goes from  $A_0$  to  $A_2$  and more generally from  $A_0$  to  $A_n$  is given by Eqs. 8 and 9 respectively.

$$p_{0,2} = p_{0,1} \times p_{1,2} = \frac{P}{(P + 1)^2} \quad (8)$$

$$p_{0,n} = p_{0,1} \times p_{1,2} \times \cdots \times p_{n-1,n} = \frac{P^{n/2}}{(P + 1)^n} \quad (9)$$


If we consider first that the biological activity (BA) of a molecule is a function of its intrinsic activity and of its probability of reaching the receptor site, and second that the intrinsic activities of homologous series of chemicals are identical, we can write:

$$BA = \frac{\text{constant} \times P^n}{(P + 1)^{2n}} \quad (10)$$

The logarithmic transformation of Eq. 10 leads to Eq. 11.

$$\log BA = n \log P - 2n \log (P + 1) + c \quad (11)$$

Symmetrical curves with linear ascending and descending sides and a parabolic part within the range of optimum lipophilicity result from Eq. 11.



**Figure 4** Hypothetical biological system for the bilinear model ( $k_1$  and  $k_2$  are the rate constants of the distribution process).

The bilinear model is derived from the McFarland model (Eq. 11) by taking into account the different volumes of the aqueous and organic phases in a biological system comprising four phases (Figure 4). The general equation of the bilinear model (Eq. 3) applied to BCF and  $\log P$  is:

$$\log \text{BCF} = a \log P - b \log (\beta P + 1) + c \quad (12)$$

where  $\beta$  is the ratio volume between the lipid and the aqueous phases. For small  $P$  values,  $(\beta P + 1)$  is approximately equal to one, therefore  $\log (\beta P + 1)$  equals zero. For large  $P$  values  $(\beta P + 1)$  is nearly equal to  $\beta P$ , therefore  $\log (\beta P + 1)$  is proportional to  $\log P$ .

The parameters  $a$ ,  $b$  and  $c$  are linear terms, which can be calculated by linear multiple regression analysis,  $\beta$  is a nonlinear parameter, which is generally estimated by a Taylor series iteration method.<sup>20</sup>

Calculations were performed as described by Kubinyi and Kehrhan,<sup>10</sup> from software written in Turbo basic and running on an IBM PC compatible.

## RESULTS AND DISCUSSION

Figure 5 depicts the ability of the linear (Eq. 13), parabolic (Eq. 14) and bilinear (Eq. 15) models to describe the set of 154 BCF values from  $\log P$ .

$$\log \text{BCF} = 0.516 \log P + 0.576 \quad (13)$$

$$n = 154; \quad r = 0.772; \quad s = 0.702; \quad F = 224.32$$

$$\log \text{BCF} = -0.164 (\log P)^2 + 2.059 \log P - 2.592 \quad (14)$$

$$n = 154; \quad r = 0.914; \quad s = 0.450; \quad F = 382.09$$

$$\log \text{BCF} = 0.910 \log P - 1.975 \log (6.8 \cdot 10^{-7} P + 1) - 0.786 \quad (15)$$

$$n = 154; \quad r = 0.950; \quad s = 0.347; \quad F = 463.51$$

From the above and from Figure 5, it is obvious that Eq. 15 is the best regression model and that the linear solution given by Eq. 13 is not valid in our range of  $\log P$  values (i.e. 1.12–8.60). This is confirmed by inspection of the statistical parameters of Eq. 13 and the graphical analysis of the residuals in Figures 6 and 7. Furthermore, we can note the unsatisfactory behaviour of the residuals in Figure 6 which are displayed according to a "parabolic band". This confirms the possibility of using a




Figure 5 Relationships between  $\log BCF$  and  $\log P$ .




Figure 6 Distribution of the residual values obtained with the linear model (Eq. 13).




Figure 7 Residual analysis of the linear model (Eq. 13).

quadratic term in the BCF model in order to increase its statistical qualities and predictive power.<sup>20</sup> Thus, introduction of  $(\log P)^2$  in the model (Eq. 14) leads to a better distribution of the residuals (Figures 8 and 9). In the same way, the high quality of the bilinear model (Eq. 15) is confirmed by analysis of the residuals (Figures 10 and 11) which reveals that no chemicals can be considered as outliers. To estimate the predictive power of the bilinear model (Eq. 15) a testing set of 29 chemicals has been constituted (Table I). In Table I, experimental BCF values were obtained under flow-through or static conditions and generally did not satisfy all the other constraints defined for the selection of the training set. Comparison of the expected BCF values obtained from the different models (i.e., Eq. 13 to 15) shows that the bilinear model (Eq. 15) is the most suitable to describe the BCF behaviour of the chemicals listed in Table I.

While it is recognised that the bioconcentration of organic chemicals in fish is a complex phenomenon involving numerous biotic and abiotic factors, our results underline the ability of the bilinear model of Kubinyi<sup>9-12</sup> to simulate this process in




Figure 8 Distribution of the residual values obtained with the parabolic model (Eq. 14).




Figure 9 Residual analysis of the parabolic model (Eq. 14).




Figure 10 Distribution of the residual values obtained with the bilinear model (Eq. 15).

a rather simplistic way. If only a few data points are available or if the  $\log P$  values are not too high, the linear models<sup>3-6</sup> are the most suitable to describe the bioconcentration in fish due to their simplicity. However, if enough data points are present and if the  $\log P$  values vary over a wide range, the parabolic model and preferentially the bilinear model are more relevant. Thus, it is interesting to note that if we consider the range of  $\log P$  values in our training and testing sets, the bilinear model (Eq. 15) allows the prediction of the bioconcentration factor of organic chemicals having  $\log P$  values between 0.39 and 9.50.

Chemicals which are released to the environment as the result of a variety of human-related activities migrate through the ecosystems according to their physico-chemical properties (e.g., vapour pressure, hydrosolubility). Numerous models varying in complexity are available to evaluate such behaviour.<sup>32</sup> Among them, the fugacity models of Mackay<sup>33</sup> provide a convenient and accurate simulation tool for exposure and risk analysis. In the classical fugacity models,<sup>33</sup> the bioconcentration




Figure 11 Residual analysis of the bilinear model (Eq. 15).

**Table I** Comparison between experimental and calculated BCF values for 29 chemicals.

| No. | Chemical                                | $\log P \log BCF_{exp} (ref)$ |            | Calculated $\log BCF$ |        |        |
|-----|---|-------------------------------|------------|-----------------------|--------|--------|
|     |   |                               |            | Eq. 13                | Eq. 14 | Eq. 15 |
| 1   | 2-( <i>t</i> -Butoxy) ethanol           | 0.39 <sup>a</sup>             | -0.22 (21) | 0.78                  | -1.81  | -0.43  |
| 2   | <i>t</i> -Butyl isopropyl ether         | 2.14 <sup>a</sup>             | 0.76 (21)  | 1.68                  | 1.06   | 1.16   |
| 3   | Octachloronaphthalene                   | 8.40 <sup>b</sup>             | 2.52 (22)  | 4.91                  | 3.13   | 2.44   |
| 4   | Pigment, monoazo <sup>c</sup>           | 9.50                          | 0.70 (23)  | 5.48                  | 2.17   | 1.28   |
| 5   | Pigment, monoazo <sup>c</sup>           | 9.30                          | 0.50 (23)  | 5.37                  | 2.37   | 1.49   |
| 6   | Aniline                                 | 0.90                          | 0.41 (24)  | 1.04                  | -0.87  | 0.03   |
| 7   | <i>t</i> -Butylphenyldiphenyl phosphate | 5.12                          | 3.04 (25)  | 3.22                  | 3.65   | 3.80   |
| 8   | 2-Chloroaniline                         | 1.90                          | 1.18 (24)  | 1.56                  | 0.73   | 0.94   |
| 9   | 3-Chloroaniline                         | 1.88                          | 1.06 (24)  | 1.55                  | 0.70   | 0.92   |
| 10  | 4-Chloroaniline                         | 1.83                          | 0.91 (24)  | 1.52                  | 0.63   | 0.88   |
| 11  | 2-Chloronaphthalene                     | 4.19                          | 3.63 (26)  | 2.74                  | 3.16   | 3.02   |
| 12  | 2,4-Dichloroaniline                     | 2.79                          | 1.98 (24)  | 2.02                  | 1.88   | 1.75   |
| 13  | 3,4-Dichloroaniline                     | 2.79                          | 1.48 (24)  | 2.02                  | 1.88   | 1.75   |
| 14  | 1,4-Dichloronaphthalene                 | 4.88                          | 3.36 (26)  | 3.09                  | 3.55   | 3.61   |
| 15  | 1,8-Dichloronaphthalene                 | 4.41                          | 3.79 (26)  | 2.85                  | 3.30   | 3.21   |
| 16  | 2,3-Dichloronaphthalene                 | 4.71                          | 4.04 (26)  | 3.01                  | 3.47   | 3.47   |
| 17  | 2,7-Dichloronaphthalene                 | 4.81                          | 4.04 (26)  | 3.06                  | 3.52   | 3.55   |
| 18  | 2-Nitroaniline                          | 1.78                          | 0.91 (24)  | 1.49                  | 0.55   | 0.83   |
| 19  | 3-Nitroaniline                          | 1.31                          | 0.92 (24)  | 1.25                  | -0.18  | 0.41   |
| 20  | 4-Nitroaniline                          | 1.31                          | 0.64 (24)  | 1.25                  | -0.18  | 0.41   |
| 21  | 2,3,7,8-TCDD                            | 6.42 <sup>d</sup>             | 3.90 (27)  | 3.89                  | 3.87   | 4.18   |
| 22  | 3,3',4,4'-Tetrachlorobiphenyl           | 5.82                          | 4.59 (28)  | 3.58                  | 3.84   | 4.19   |
| 23  | 3,3',4,4'-Tetrachlorodiphenyl ether     | 5.78                          | 4.51 (28)  | 3.56                  | 3.83   | 4.18   |
| 24  | 2,4,5-Trichlorobiphenyl                 | 5.51                          | 4.26 (28)  | 3.42                  | 3.77   | 4.06   |
| 25  | 2,4,5-Trichlorodiphenyl ether           | 5.44                          | 4.18 (28)  | 3.38                  | 3.76   | 4.02   |
| 26  | 3,4,5-Trichloroveratrole                | 4.60                          | 3.50 (29)  | 2.95                  | 3.41   | 3.38   |
| 27  | Tetrachloroveratrole                    | 5.80                          | 4.40 (29)  | 3.57                  | 3.83   | 4.19   |
| 28  | 1,3,7-Trichloronaphthalene              | 5.59                          | 4.43 (26)  | 3.46                  | 3.79   | 4.10   |
| 29  | Triphenyl phosphate                     | 3.90 <sup>e</sup>             | 2.76 (25)  | 2.59                  | 2.94   | 2.76   |

<sup>a</sup> mean of two values.<sup>b</sup> Ref. 26.<sup>c</sup> Chemicals no. 45 and 46 in Anliker *et al.*,<sup>23</sup> (Table I [part B] p. 267).<sup>d</sup> Ref. 30.<sup>e</sup> Ref. 31.

in the biota (i.e., fish) is estimated from a simple linear regression equation (Eq. 1). This induces erroneous estimations for highly lipophilic chemicals. To overcome this problem, we propose to introduce in the fugacity models our bilinear model (Eq. 15) in order to stretch their domain of application.

## References

1. Barron, M. G. (1990). Bioconcentration. *Environ. Sci. Technol.* **24**, 1612-1618.
2. Hamelink, J. L. (1977). Current bioconcentration test methods and theory. In, *Aquatic Toxicology and Hazard Evaluation* (F. L. Mayer and J. L. Hamelink, Eds.). ASTM STP **634**, pp. 149-161.
3. Kanazawa, J. (1981). Measurement of the bioconcentration factors of pesticides by freshwater fish and their correlation with physicochemical properties or acute toxicities. *Pestic. Sci.* **12**, 417-424.
4. Schüürmann, G. and Klein, W. (1988). Advances in bioconcentration prediction. *Chemosphere* **17**, 1551-1574.

5. Isnard, P. and Lambert, S. (1988). Estimating bioconcentration factors from octanol-water partition coefficient and aqueous solubility. *Chemosphere* **17**, 21–34.
6. Samiullah, Y. (1990). *Prediction of the Environmental Fate of Chemicals*. Elsevier Science Publishers, London, p. 285.
7. Banerjee, S. and Baughman, G. L. (1991). Bioconcentration factors and lipid solubility. *Environ. Sci. Technol.* **25**, 536–539.
8. Devillers, J. and Lipnick, R. L. (1990). Practical applications of regression analysis in environmental QSAR studies. In, *Practical Applications of Quantitative Structure-Activity Relationships (QSAR) in Environmental Chemistry and Toxicology* (W. Karcher and J. Devillers, Eds.). Kluwer Academic Publishers, Dordrecht, pp. 129–143.
9. Kubinyi, H. (1976). Quantitative structure-activity relationships. IV. Non-linear dependence of biological activity on hydrophobic character: a new model. *Arzneim. Forsch. Drug Res.* **26**, 1991–1997.
10. Kubinyi, H. and Kehrhahn, O. H. (1978). Quantitative structure-activity relationships. VI. Non-linear dependence of biological activity on hydrophobic character: calculation procedures for the bilinear model. *Arzneim. Forsch. Drug Res.* **28**, 598–601.
11. Kubinyi, H. (1977). Quantitative structure-activity relationships. 7. The bilinear model, a new model for nonlinear dependence of biological activity on hydrophobic character. *J. Med. Chem.* **20**, 625–629.
12. Kubinyi, H. (1979). Nonlinear dependence of biological activity on hydrophobic character: the bilinear model. *Il Farmaco* **34**, 247–276.
13. Lipnick, R. L., Pritzker, C. S., and Bentley, D. L. (1985). A QSAR study of the rat LD<sub>50</sub> for alcohols. In, *QSAR and Strategies in the Design of Bioactive Compounds: Proceedings of the 5th European Symposium on Quantitative Structure-Activity Relationships* (J. K. Seydel, Ed.). VCH, Weinheim, pp. 420–423.
14. Lipnick, R. L., Pritzker, C. S., and Bentley, D. L. (1987). Application of QSAR to model the acute toxicity of industrial organic chemicals to mammals and fish. In, *QSAR in Drug Design and Toxicology* (D. Hadzi and B. Jerman-Blazic, Eds.). Elsevier Science Publishers, Amsterdam, p. 301–306.
15. Lipnick, R. L., Watson, K. R., and Strausz, A. K. (1987). A QSAR study of the acute toxicity of some industrial organic chemicals to goldfish. Narcosis, electrophile, and proelectrophile mechanisms. *Xenobiotica* **17**, 1011–1025.
16. Vaishnav, D. D., Boethling, R. S., and Babeu, L. (1987). Quantitative structure-biodegradability relationships for alcohols, ketones and alicyclic compounds. *Chemosphere* **16**, 695–703.
17. Veith, G. D., Call, D. J., and Brooke, L. T. (1983). Structure-toxicity relationships for the fathead minnow, *Pimephales promelas*: Narcotic industrial chemicals. *Can. J. Fish. Aquat. Sci.* **40**, 743–748.
18. Hansch, C., Kim, D., Leo, A. J., Novellino, E., Silipo, C., and Vittoria, A. (1989). Toward a quantitative comparative toxicology of organic compounds. *CRC Critic. Rev. Toxicol.* **19**, 185–226.
19. Bintein, S. and Devillers, J. (1992). Data compilation of BCF values for fish and aquatic invertebrates. CTIS, R19203, p. 42.
20. Draper, N. and Smith, H. (1981). *Applied Regression Analysis*, Second Edition. John Wiley & Sons, New York, p. 709.
21. Fujiwara, Y., Kinoshita, T., Sato, H., and Kojima, I. (1984). Biodegradation and bioconcentration of alkyl ethers. *Yukagaku* **33**, 111–114.
22. Oliver, B. G. and Niimi, A. J. (1985). Bioconcentration factors of some halogenated organics for rainbow trout: Limitations in their use for prediction of environmental residues. *Environ. Sci. Technol.* **19**, 842–849.
23. Anliker, R., Clarke, E. A., and Moser, P. (1981). Use of the partition coefficient as an indicator of bioaccumulation tendency of dyestuffs in fish. *Chemosphere* **10**, 263–274.
24. Zok, S., Gorge, G., Kalsch, W., and Nagel, R. (1991). Bioconcentration, metabolism and toxicity of substituted anilines in the zebrafish (*Brachydanio rerio*). *Sci. Total Environ.* **109/110**, 411–421.
25. Muir, D. C. G., Yarechewski, A. L., and Grift, N. P. (1983). Environmental dynamics of phosphate esters. III. Comparison of the bioconcentration of four triaryl phosphates by fish. *Chemosphere* **12**, 155–166.
26. Opperhuizen, A., Van der Velde, E. W., Gobas, F. A. P. C., Liem, D. A. K., and Van der Steen, J. M. D. (1985). Relationship between bioconcentration in fish and steric factors of hydrophobic chemicals. *Chemosphere* **14**, 1871–1896.
27. Adams, W. J., de Graeve, G. M., Sabourin, T. D., Cooney, J. D., and Mosher, G. M. (1986). Toxicity and bioconcentration of 2,3,7,8-TCDD to fathead minnows (*Pimephales promelas*). *Chemosphere* **15**, 1503–1511.
28. Opperhuizen, A. and Voors, P. I. (1987). Bioconcentration kinetics of 2,4,5-tri- and 3,3',4,4'-tetrachlorobiphenyl and 2,4,5-tri- and 3,3',4,4'-tetrachlorodiphenylether in fish. *Chemosphere* **16**, 2379–2388.

29. Neilson, A. H., Allard, A. S., Reiland, S., Remberger, M., Tärholm, A., Viktor, T., and Landner, L. (1984). Tri- and tetra-chloroveratrole, metabolites produced by bacterial *O*-methylation of tri- and tetra-chloroguaiacol: An assessment of their bioconcentration potential and their effects on fish reproduction. *Can. J. Fish. Aquat. Sci.* **41**, 1502-1512.
30. Sijm, D. T. H. M., Wever, H., de Vries, P. J., and Opperhuizen, A. (1989). Octan-1-ol/water partition coefficients of polychlorinated dibenzo-*p*-dioxins and dibenzofurans: Experimental values determined with a stirring method. *Chemosphere* **19**, 263-266.
31. Bengtsson, B. E., Tarkpea, M., Sletten, T., Carlberg, G. E., Kringstad, A., and Renberg, L. (1986). Bioaccumulation and effects of some technical triaryl phosphate products in fish and *Nitocra spinipes*. *Environ. Toxicol. Chem.* **5**, 853-861.
32. Cohen, Y. (1986). Organic pollutant transport. *Environ. Sci. Technol.* **20**, 538-544.
33. Mackay, D. (1991). *Multimedia Environmental Models. The Fugacity Approach*. Lewis Publishers, Chelsea, p. 257.



# ARTICLE II

0045-6535(94)E0050-4

## QSAR FOR ORGANIC CHEMICAL SORPTION IN SOILS AND SEDIMENTS

S. Bintein and J. Devillers\*

CTIS, 21 rue de la Bannière, 69003 Lyon, France

(Received in Germany 30 November 1993; accepted 18 January 1994)

### ABSTRACT

Sorption phenomena exert important influences on the environmental fate of anthropogenic organic substances. Under these conditions, the aim of this study was to propose a general QSAR model using the physicochemical properties of the molecules (i.e.  $\log K_{ow}$  and  $pK_a$ ) and some relevant properties of soils or sediments (i.e. pH and %OC) to estimate the sorption behavior of both ionized and non-ionized chemicals. The proposed model was elaborated from 229  $K_p$  values recorded for 53 chemicals. The model was then tested on 500 other  $K_p$  values obtained for 87 chemicals.

### INTRODUCTION

The fate of the organic chemicals introduced into the environment depends on a variety of physicochemical and biological processes.<sup>1-2</sup> Mathematical models which attempt to integrate these phenomena are widely used to predict the environmental transport and distribution of the organic pollutants between the different compartments of the biosphere.<sup>3-7</sup> Use of these models requires a variety of abiotic and biotic parameters as inputs. Among them, the soil or sediment sorption coefficient of chemicals ( $K_p$ ) is one of the key input parameters in models used to estimate the environmental mobility and fate of the pollutants.<sup>8</sup> Because its experimental determination is time-consuming and expensive, estimated values based on QSAR (Quantitative Structure-Activity Relationship) equations are now widely used. The distribution of organic chemicals between soils or sediments and water can be quantified

-----  
\* Author to whom all correspondence should be addressed.

through the use of the Freundlich isotherm:

$$S = K_p * C^N \quad (1)$$

where  $S$  is the sorbed concentration,  $C$  is the solution concentration, and  $K_p$  and  $N$  (traditionally expressed as  $1/n$ ) are empirical constants. The values of  $N$  commonly range between 0.7 and 1.2.<sup>9</sup>

The influence of the organic matter from soils and sediments on the sorption behavior of the organic chemicals has been discussed in many studies.<sup>10-12</sup> Consequently, the sorption partition coefficient  $K_p$  is generally related to the fraction of organic carbon ( $f_{oc}$ ) associated with the sorbent to yield an organic-carbon-partition coefficient,  $K_{oc}$ ,

$$K_{oc} = (K_p * 100)/\%OC \quad (2)$$

Sometimes, sorption coefficients are normalized to the organic matter content instead of the organic carbon content. This yields an organic-matter-partition coefficient,  $K_{om}$ ,

$$K_{om} = (K_p * 100)/\%OM \quad (3)$$

Several regression equations, developed with physicochemical properties such as the  $n$ -octanol/water partition coefficient ( $K_{ow}$ ) or aqueous solubility have been used to estimate  $K_{oc}$  or  $K_{om}$ .<sup>13-18</sup> Topological descriptors (e.g. molecular connectivity indices) have also been used to predict  $K_{oc}$  or  $K_{om}$ .<sup>1, 19-23</sup> However, many of these correlations were developed for specific classes of compounds (e.g. PAH<sup>13</sup>) and are not applicable to particular chemicals such as acids and bases. Furthermore, some of these models fail by using in the same data set  $K_{om}$  and  $K_{oc}$  values without correction factors. Indeed, the relationship  $K_{oc} = 1.724 * K_{om}$  must be used to convert  $K_{om}$  to  $K_{oc}$ .<sup>1, 15, 23</sup> Last, in ecotoxicology modeling, it is obvious that  $K_p$  is more interesting to model than  $K_{om}$  or  $K_{oc}$ . Under these conditions, the aim of this study was to present a  $K_p$  equation easily incorporable in the main environmental fate models and allowing the prediction of the sorption behavior of structurally diverse chemicals in various sediments and soils.

## MODEL DEVELOPMENT

$K_p$  values for 53 organic chemicals including aliphatics, aromatics, pesticides, PCB, PAH, and related compounds (Table 1) were retrieved from the literature.<sup>9, 13, 16, 24-34</sup> They were selected according to the following criteria:

- only experimental results obtained from Freundlich isotherms with  $N = 1$  were selected in order to standardize the units,<sup>35</sup>
- $K_p$  data recorded in soils or sediments having a  $\%OC < 0.1$  were rejected since for organic-poor sorbents, the interactions of the chemicals with the inorganic matrix of sorbent may become important,<sup>8, 28, 36, 37</sup>
- values corresponding to suspended sediments were rejected.

When necessary, the relationship  $\%OC = \%OM/1.724$  was used to convert  $\%OM$  to  $\%OC$ . The log  $K_{ow}$  and pKa values (Table 1) were obtained from the literature.<sup>13, 16, 24, 26-30, 32, 38-49</sup> Experimental values or data obtained from critical compilations were preferentially selected.

Table 1: Training set.

| Chemical                      | n#      | log K <sub>ow</sub> | pKa      | %OC         | pH          |
|-------------------------------|---------|---------------------|----------|-------------|-------------|
| Acetophenone                  | 12(24*) | 1.59(24)            |          | 0.15-2.38** |             |
| Acridine                      | 12(25)  | 3.40(38)            | 5.68(39) | 0.48-2.38   | 4.54-7.83** |
| Acrylonitrile                 | 2(26)   | 0.12(26)            |          | 0.66-1.49   |             |
| 2-Aminoanthracene             | 13(27)  | 4.13(27)            | 4.10(38) | 0.15-2.38   | 4.54-8.32   |
| 6-Aminochrysene               | 13(27)  | 4.98(27)            | 3.70(38) | 0.15-2.38   | 4.54-8.32   |
| Anisole                       | 1(16)   | 2.11(16)            |          | 1.10        |             |
| Anthracene-9-carboxylic acid  | 12(27)  | 4.30(38)            | 3.65(38) | 0.15-2.38   | 4.54-8.32   |
| Benzamide                     | 1(9)    | 0.64(38)            |          | 1.25        |             |
| Benzene                       | 2(26)   | 2.13(16)            |          | 0.66-1.49   |             |
|                               | 1(16)   | 2.13(16)            |          | 1.10        |             |
| <i>n</i> -Butylbenzene        | 1(28)   | 4.13(28)            |          | 0.15        |             |
| Carbon tetrachloride          | 2(26)   | 2.64(26)            |          | 0.66-1.49   |             |
| Chlorobenzene                 | 2(26)   | 2.84(16)            |          | 0.66-1.49   |             |
|                               | 1(16)   | 2.84(16)            |          | 1.10        |             |
|                               | 1(28)   | 2.84(16)            |          | 0.15        |             |
| 2-Chlorobiphenyl              | 1(16)   | 4.51(16)            |          | 1.10        |             |
| Chloroform                    | 2(26)   | 1.97(26)            |          | 0.66-1.49   |             |
| 3-Chlorophenol                | 1(29)   | 2.50(29)            | 8.85(40) | 2.80        | 8.00        |
| 1,2:5,6-Dibenzanthracene      | 14(30)  | 6.50(30)            |          | 0.11-2.38   |             |
| 1,2-Dichlorobenzene           | 2(26)   | 3.38(16)            |          | 0.66-1.49   |             |
|                               | 1(16)   | 3.38(16)            |          | 1.10        |             |
|                               | 1(31)   | 3.38(16)            |          | 0.93        |             |
| 1,3-Dichlorobenzene           | 1(16)   | 3.38(16)            |          | 1.10        |             |
| 1,4-Dichlorobenzene           | 1(16)   | 3.39(16)            |          | 1.10        |             |
|                               | 1(28)   | 3.39(16)            |          | 0.15        |             |
| 2,2'-Dichlorobiphenyl         | 1(16)   | 4.80(16)            |          | 1.10        |             |
| 2,4'-Dichlorobiphenyl         | 1(16)   | 5.10(16)            |          | 1.10        |             |
| 1,2-Dichloroethane            | 1(31)   | 1.45(41)            |          | 0.93        |             |
| 3,4-Dichlorophenol            | 2(29)   | 3.44(42)            | 7.39(40) | 2.80        | 6.00-8.00   |
| 7,12-Dimethylbenzanthracene   | 13(30)  | 5.98(30)            |          | 0.15-2.38   |             |
| 1,4-Dimethylbenzene           | 2(26)   | 3.15(28)            |          | 0.66-1.49   |             |
|                               | 1(28)   | 3.15(28)            |          | 0.15        |             |
| 2,4-Dinitro- <i>o</i> -cresol | 12(32)  | 2.85(43)            | 4.46(32) | 0.15-3.04   | 4.27-8.29   |
| Ethylbenzene                  | 1(16)   | 3.15(16)            |          | 1.10        |             |
| Furan                         | 2(26)   | 1.34(26)            |          | 0.66-1.49   |             |
| Hexachlorobenzene             | 1(26)   | 5.73(44)            |          | 0.66        |             |
| Hexanoic acid                 | 1(9)    | 1.90(45)            | 4.85(46) | 4.85        | 2.80        |
| Methoxychlor                  | 13(13)  | 5.08(13)            |          | 0.13-3.29   |             |

Table 1 (continued)

| Chemical                   | n#      | log K <sub>ow</sub> | pKa      | %OC         | pH          |
|----------------------------|---------|---------------------|----------|-------------|-------------|
| 3-Methylcholanthrene       | 12(30*) | 6.42(30)            |          | 0.11-2.38** |             |
| Nitrobenzene               | 2(26)   | 1.87(26)            |          | 0.66-1.49   |             |
| Pentachlorophenol          | 1(29)   | 5.04(42)            | 4.92(40) | 3.20        | 4.70        |
| Pyrene                     | 14(30)  | 5.09(30)            |          | 0.11-2.38   |             |
|                            | 12(13)  | 5.09(30)            |          | 0.13-3.29   |             |
| Silvex                     | 12(32)  | 3.41(47)            | 3.07(32) | 0.15-3.04   | 4.27-8.29** |
| 2,3,7,8-TCDD               | 1(33)   | 6.42(48)            |          | 0.66        |             |
| 1,2,3,4-Tetrachlorobenzene | 1(28)   | 4.64(44)            |          | 0.15        |             |
| 1,2,4,5-Tetrachlorobenzene | 1(28)   | 4.72(28)            |          | 0.15        |             |
| 1,1,2,2-Tetrachloroethane  | 1(31)   | 2.39(41)            |          | 0.93        |             |
| Tetrachloroethylene        | 1(28)   | 2.60(28)            |          | 0.15        |             |
| 2,3,4,6-Tetrachlorophenol  | 2(29)   | 4.42(42)            | 5.38(40) | 1.70-3.20   | 3.40-4.70   |
| Tetrahydrofuran            | 2(26)   | 0.46(26)            |          | 0.66-1.49   |             |
| 1,2,4,5-Tetramethylbenzene | 1(28)   | 4.05(28)            |          | 0.15        |             |
| Toluene                    | 2(26)   | 2.71(26)            |          | 0.66-1.49   |             |
|                            | 1(28)   | 2.71(26)            |          | 0.15        |             |
| 1,2,3-Trichlorobenzene     | 1(28)   | 4.14(44)            |          | 0.15        |             |
|                            | 1(34)   | 4.14(44)            |          | 4.70        |             |
| 1,2,4-Trichlorobenzene     | 1(16)   | 4.02(16)            |          | 1.10        |             |
|                            | 1(28)   | 4.02(16)            |          | 0.15        |             |
| 2,4,4'-Trichlorobiphenyl   | 1(16)   | 5.62(16)            |          | 1.10        |             |
| 1,1,1-Trichloroethane      | 1(31)   | 2.47(41)            |          | 0.93        |             |
| 2,4,5-Trichlorophenol      | 1(29)   | 3.72(49)            | 7.43(40) | 2.80        | 8.00        |
| 1,2,3-Trichloropropane     | 2(26)   | 2.01(26)            |          | 0.66-1.49   |             |
| 1,2,3-Trimethylbenzene     | 1(28)   | 3.60(28)            |          | 0.15        |             |
| 1,3,5-Trimethylbenzene     | 1(28)   | 3.60(28)            |          | 0.15        |             |

#number of experimental K<sub>p</sub> values; \*reference; \*\*range.

Regression analysis of the 229 K<sub>p</sub> values obtained from the 53 chemicals under study versus their log K<sub>ow</sub> and log f<sub>oc</sub> values (Table 1) yields Eq. 4.

$$\log K_p = 0.96 \log K_{ow} + 1.04 \log f_{oc} - 0.13 \quad (4)$$

$$n = 229 \quad s = 0.639 \quad r = 0.924 \quad F = 660.07 \quad p < 0.01\%$$

Graphical analysis of the residuals obtained with Eq. 4 shows (Fig. 1) that this model overestimates the K<sub>p</sub> values of the acids and underestimates those of the bases. At the

opposite, Fig. 1 reveals that the residuals for the other chemicals under study are well distributed.




Fig. 1. Graphical analysis of the residuals ( $\log K_p \text{ obs.} - \log K_p \text{ calc.}$ ) obtained with Eq. 4.

Our results are not surprising since numerous authors<sup>32, 36, 39, 50-53</sup> have underlined that it was dangerous to neglect the role of pH in the modeling of the adsorption of acids and bases in soils and sediments. Indeed, it is well admitted that most weakly acidic chemicals are in predominantly ionic (i.e. negatively charged) form at the pH of most natural soils, while most weakly basic chemicals are in molecular form. In either case, as natural sorbent pH decreases toward a value equal to the dissociation constant (pKa) of the chemical, sorption tends to increase because at low pH, the acidic chemical has more of the molecular species and the basic chemical has more of the protonated (positive) ionic species.<sup>8</sup> The colloidal surfaces of most natural soils are negatively charged and therefore have an affinity for positively charged molecules, but not much affinity for negatively charged molecules.<sup>8, 54</sup>

Therefore, to optimize the model, it could be interesting to introduce two different correction factors (one for acids and another for bases) allowing to quantify the variations of the concentration of ionic species in the range of soil pH values.

Thus, for acids, the evolution of the anionic species concentration in relation to pH values can be estimated by:<sup>55</sup>

$$CFa = \log \frac{1}{1 + 10^{\text{pH} - \text{pKa}}} \quad (5)$$

For bases, the relation between the protonated species concentration and the pH values can be expressed by:<sup>55</sup>

$$CFb = \log \frac{1}{1 + 10^{pKa - pH}} \quad (6)$$

Introduction of CFa and CFb in Eq. 4 allows to obtain the following model:

$$\log K_p = 0.92 \log K_{ow} + 1.09 \log f_{oc} + 0.33 CFa - 1.32 CFb + 0.30 \quad (7)$$

$$n = 229 \quad s = 0.453 \quad r = 0.963 \quad F = 713.57 \quad p < 0.01\%$$

Like Wauchope and coworkers,<sup>56</sup> we have defined acids as the sole chemicals whose neutral (molecular) forms are capable of forming negatively charged ions. Therefore, in Eq. 7, CFa must be taken equal to zero for bases and other compounds since this parameter is null when the concentration of anionic species is zero.<sup>55</sup> In the same way,<sup>56</sup> we have defined bases as the sole chemicals whose neutral forms are able to form positively charged ions. Therefore in Eq. 7, CFb must be taken equal to zero for acids and other compounds since this parameter is null when the concentration of cationic species is zero.<sup>55</sup>

Eq. 7 gives a better distribution of the residuals for the acids and bases (Fig. 2) than Eq. 4 (Fig. 1), but some  $K_p$  values for basic chemicals are still underestimated.




Fig. 2. Graphical analysis of the residuals ( $\log K_p \text{ obs.} - \log K_p \text{ calc.}$ ) obtained with Eq. 7.

The remaining difference between the adsorption behavior of bases and acids could be

explained by the fact that the adsorption of basic compounds in certain soils (e.g. montmorillonite clay systems) is principally dependent upon the surface acidity and not upon the pH of the bulk solution, while the converse is true for the adsorption of acidic compounds.<sup>50</sup> Bailey *et al.*<sup>50</sup> have estimated that the surface acidity of montmorillonite appears to be 3-4 pH units lower than the pH of bulk solution. Green and Karickhoff<sup>8</sup> have stressed that a reasonable estimate of surface acidity (pHs) is two pH units lower than the bulk suspension pH (i.e. pHs = pH - 2). Under these conditions, we have tried to introduce this correction in our model (Eq. 7) defining the new following variable:

$$CFb' = \log \frac{1}{1 + 10^{pKa - (pH - 2)}} \quad (8)$$

Introduction of CFb' in Eq. 7 instead of CFb yields Eq. 9.

$$\log K_p = 0.93 \log K_{ow} + 1.09 \log f_{oc} + 0.32 CFa - 0.55 CFb' + 0.25 \quad (9)$$

$$n = 229 \quad s = 0.433 \quad r = 0.966 \quad F = 786.07 \quad p < 0.01\%$$

Fig. 3 depicts the high quality of Eq. 9 to describe the 229  $K_p$  values of the training set (Table 1). Note that CFb' (like CFb) must be taken equal to zero for acids and other chemicals since this parameter is null when the concentration of cationic species is zero.




Fig. 3. Observed versus calculated (Eq. 9)  $K_p$  values for the training set.



Introduction of CFb' in the model leads to a better distribution of the residuals of basic compounds (Fig. 4).




Fig. 4. Graphical analysis of the residuals ( $\log K_p \text{ obs.} - \log K_p \text{ calc.}$ ) obtained with Eq. 9.

## MODEL VALIDATION

To estimate the predictive power of our final model (Eq. 9), a test set of 500 experimental  $K_p$  values for 87 chemicals has been constituted (Table 2). It is important to note that these values did not satisfy all the constraints defined for the selection of the original training set since they were generally obtained from Freundlich isotherms with  $0.8 < N < 1.2$  or by one initial concentration.<sup>9, 15, 29, 34, 39, 51, 57-76</sup>

The  $\log K_{ow}$  and  $pK_a$  values (Table 2) were obtained from the literature.<sup>4, 15, 16, 29, 38-46, 49, 52, 59, 60, 63, 70, 71, 75-85</sup> Experimental values or data obtained from critical compilations were preferentially selected.

Table 2: Test set.

| Chemical    | n#    | $\log K_{ow}$ | $pK_a$   | $N$        | %OC        | pH   |
|-------------|-------|---------------|----------|------------|------------|------|
| Acetanilide | 2(9*) | 1.16(38)      |          | 0.82-0.84† | 1.58-4.85† |      |
| Acridine    | 1(39) | 3.40(38)      | 5.68(39) | 1.02       | 0.58       | 7.64 |
| Alachlor    | 1(57) | 2.64(77)      |          | 1.01       | 1.17       |      |

Table 2 (continued)

| Chemical            | n#     | log K <sub>ow</sub> | pKa      | N          | %OC         | pH         |
|---------------------|--------|---------------------|----------|------------|-------------|------------|
| Aldicarb            | 1(15*) | 1.08(77)            |          | 1\$        | 2.05        |            |
| Aldrin              | 1(15)  | 5.66(4)             |          | 1\$        | 2.05        |            |
| Ametryne            | 34(58) | 2.58(78)            | 4.00(79) | 1\$        | 0.35-20.88† | 4.50-9.00† |
| Aniline             | 1(59)  | 0.90(38)            | 4.60(46) | 0.83       | 0.84§       | 5.40§      |
| Atrazine            | 4(60)  | 2.33(77)            | 1.68(79) | 1          | 0.40-7.60   | 4.40-8.00  |
|                     | 1(61)  | 2.33(77)            | 1.68(79) | 0.92       | 0.43        | 6.05       |
|                     | 25(62) | 2.33(77)            | 1.68(79) | 1\$        | 0.35-2.84   | 4.30-7.10  |
| Benzamide           | 1(9)   | 0.64(38)            |          | 0.88       | 4.85        |            |
| Benzoic acid (B.a.) | 1(9)   | 1.87(38)            | 4.20(46) | 0.90       | 4.85        | 2.80       |
| B.a. ethyl ester    | 3(9)   | 2.64(38)            |          | 0.81-0.88† | 1.25-4.85   |            |
| B.a. methyl ester   | 3(9)   | 2.20(38)            |          | 0.81-0.85  | 1.25-4.85   |            |
| B.a. phenyl ester   | 2(9)   | 3.59(38)            |          | 0.91-0.93  | 1.25-1.58   |            |
| 4-Bromophenol       | 1(15)  | 2.59(15)            | 9.34(46) | 1\$        | 1.46        | 6.70       |
| Captafol            | 1(15)  | 3.83(15)            |          | 1\$        | 2.05        |            |
| Captan              | 1(15)  | 2.54(15)            |          | 1\$        | 2.05        |            |
| Carbendazim         | 15(63) | 1.52(63)            | 4.48(63) | 1\$        | 0.63-3.47   | 4.14-7.54  |
| Carbaryl            | 1(15)  | 2.32(15)            |          | 1\$        | 2.05        |            |
| Carbofuran          | 2(64)  | 1.63(77)            |          | 0.90-0.94  | 1.60-2.50   |            |
| Chlorfenvinphos     | 1(15)  | 3.10(15)            |          | 1\$        | 2.05        |            |
| Chlorobenzene       | 1(65)  | 2.84(16)            |          | 0.87       | 4.70        |            |
| 2-Chlorophenol      | 1(66)  | 2.17(43)            | 8.49(40) | 0.80       | 2.96        | 5.70       |
| 3-Chlorophenol      | 5(29)  | 2.50(29)            | 8.85(40) | 0.80       | 1.70-3.20   | 3.40-6.00  |
|                     | 3(34)  | 2.50(29)            | 8.85(40) | 0.85-0.96  | 2.15-9.05   | 3.60-5.90  |
|                     | 1(66)  | 2.50(29)            | 8.85(40) | 0.83       | 2.96        | 5.70       |
| Chlorsulfuron       | 1(67)  | 2.20(80)            | 3.60(52) | 0.87       | 1.42        | 4.60       |
|                     | 23(68) | 2.20(80)            | 3.60(52) | 1\$        | 0.95-20.36  | 5.20-7.90  |
| 3-Cresol            | 1(66)  | 1.98(38)            | 10.0(46) | 0.87       | 2.96        | 5.70       |
| 2,4-D               | 3(9)   | 2.81(81)            | 2.64(46) | 0.89-1.06  | 1.25-4.85   | 2.80-7.10  |
|                     | 19(69) | 2.81(81)            | 2.64(46) | 0.82-1.16  | 0.28-2.73   | 6.30-8.30  |
| Diazinon            | 1(15)  | 3.11(15)            |          | 1\$        | 2.05        |            |
| Dibenzothiophene    | 11(70) | 4.38(70)            |          | 0.88-1.10  | 0.15-2.38   |            |
| 2,4-Dichloroaniline | 3(34)  | 2.78(82)            | 2.05(83) | 0.85-1.11  | 3.54-9.05   | 3.60-5.90  |
| 3,5-Dichloroaniline | 1(59)  | 2.69(59)            | 2.37(46) | 0.91       | 0.84§       | 5.40§      |
| 1,4-Dichlorobenzene | 1(65)  | 3.39(16)            |          | 0.96       | 2.15        |            |
| 2,4-Dichlorophenol  | 3(71)  | 3.21(42)            | 7.68(40) | 1\$        | 0.20-3.70   | 4.20-7.40  |
| 3,4-Dichlorophenol  | 5(29)  | 3.44(42)            | 7.39(40) | 0.80-0.90  | 0.90-3.20   | 3.40-7.50  |
|                     | 4(34)  | 3.44(42)            | 7.39(40) | 0.81-0.94  | 2.15-9.05   | 3.60-5.90  |
| Dieldrin            | 1(15)  | 4.60(4)             |          | 1\$        | 2.05        |            |

Table 2 (continued)

| Chemical                    | n#     | log K <sub>ow</sub> | pKa      | N          | %OC        | pH         |
|-----------------------------|--------|---------------------|----------|------------|------------|------------|
| Dimethoate                  | 1(15*) | 0.79(15)            |          | 1\$        | 2.05       |            |
| <i>N,N</i> -Dimethylaniline | 3(9)   | 2.31(38)            | 5.15(46) | 0.87-0.91† | 1.25-4.85† | 2.80-7.10† |
| Diuron                      | 34(58) | 2.60(77)            |          | 1\$        | 0.35-20.88 |            |
| Fenamiphos                  | 1(15)  | 3.18(15)            |          | 1\$        | 2.05       |            |
| Folpet                      | 1(15)  | 3.63(15)            |          | 1\$        | 2.05       |            |
| Hexachlorobenzene           | 1(15)  | 5.73(44)            |          | 1\$        | 1.53       |            |
| Hexanoic acid               | 2(9)   | 1.90(45)            | 4.85(46) | 1.01       | 1.25-1.58  | 6.70-7.10  |
| 3-Methoxyphenol             | 1(66)  | 1.58(49)            | 9.65(46) | 0.89       | 2.96       | 5.70       |
| <i>N</i> -Methylaniline     | 1(9)   | 1.66(38)            | 4.85(46) | 0.89       | 4.85       | 2.80       |
| 4-Methylaniline             | 1(9)   | 1.39(38)            | 5.08(38) | 0.96       | 4.85       | 2.80       |
| 4-Methylbenzoic acid        | 1(9)   | 2.34(38)            | 4.36(46) | 0.82       | 4.85       | 2.80       |
| Metsulfuron-methyl          | 23(68) | 2.20(80)            | 3.30(80) | 1\$        | 0.95-20.36 | 5.20-7.90  |
| 4-Nitroaniline              | 1(9)   | 1.39(45)            | 1.01(46) | 0.81       | 1.25       | 6.70       |
| 4-Nitrobenzoic acid         | 1(9)   | 1.89(45)            | 3.44(46) | 1.14       | 4.85       | 2.80       |
| 2-Nitrophenol               | 1(66)  | 1.79(45)            | 7.22(46) | 0.89       | 2.96       | 5.70       |
| Parathion                   | 1(15)  | 3.76(77)            |          | 1\$        | 2.05       |            |
|                             | 6(72)  | 3.76(77)            |          | 1.02-1.11  | 0.44-14.28 |            |
| 2,2',5,5'-PCB               | 2(73)  | 6.10(84)            |          | 1\$        | 0.16-1.87  |            |
|                             | 5(74)  | 6.10(84)            |          | 1\$        | 0.16-1.87  |            |
| 3,3',4,4'-PCB               | 5(74)  | 6.10(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',3,4,5'-PCB             | 2(73)  | 6.50(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',3,5',6-PCB             | 2(73)  | 6.40(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',3',4,5-PCB             | 2(73)  | 6.60(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',4,5,5'-PCB             | 2(73)  | 6.40(84)            |          | 1\$        | 0.16-1.87  |            |
|                             | 5(74)  | 6.40(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,3,3',4',6-PCB             | 2(73)  | 6.53(85)            |          | 1\$        | 0.16-1.87  |            |
| 2,3',4,4',5-PCB             | 2(73)  | 6.40(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',3,4,4',5-PCB           | 2(73)  | 7.00(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',3,4,5,5'-PCB           | 2(73)  | 7.59(85)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',3,4',5',6-PCB          | 2(73)  | 6.80(84)            |          | 1\$        | 0.16-1.87  |            |
| 2,2',4,4',5,5'-PCB          | 2(73)  | 6.90(84)            |          | 1\$        | 0.16-1.87  |            |
|                             | 5(74)  | 6.90(84)            |          | 1\$        | 0.16-1.87  |            |
| Pentachlorobenzene          | 2(73)  | 4.94(41)            |          | 1\$        | 0.16-1.87  |            |
| Pentachlorophenol           | 6(29)  | 5.04(42)            | 4.92(40) | 0.80-0.90  | 0.90-29.80 | 3.40-7.50  |
|                             | 4(34)  | 5.04(42)            | 4.92(40) | 0.82-0.92  | 2.15-9.05  | 3.60-5.90  |
|                             | 3(71)  | 5.04(42)            | 4.92(40) | 1\$        | 0.20-3.70  | 4.20-7.40  |
| Phenol                      | 1(15)  | 1.46(15)            | 9.99(46) | 1\$        | 1.46       | 6.70       |
| Phenylacetic acid           | 2(9)   | 1.41(38)            | 4.31(46) | 0.95-1.12  | 2.80-6.70  | 1.25-4.85  |

Table 2 (continued)

| Chemical                | n#     | log K <sub>ow</sub> | pKa      | N          | %OC        | pH         |
|-------------------------|--------|---------------------|----------|------------|------------|------------|
| Phorate                 | 1(15*) | 3.83(77)            |          | 1\$        | 2.05       |            |
| Prometone               | 2(61)  | 1.94(78)            | 4.28(79) | 0.92-0.99† | 0.33-0.43† | 6.05-6.30† |
|                         | 25(62) | 1.94(78)            | 4.28(79) | 1\$        | 0.35-2.84  | 4.30-7.10  |
| Prometryne              | 25(62) | 2.99(78)            | 4.05(79) | 1\$        | 0.35-2.84  | 4.30-7.10  |
| Propazine               | 25(62) | 3.09(78)            | 1.85(79) | 1\$        | 0.35-2.84  | 4.30-7.10  |
| Pyridine                | 1(39)  | 0.65(38)            | 5.23(39) | 1.04       | 0.58       | 7.64       |
| Quinoline               | 1(39)  | 2.03(38)            | 4.92(39) | 0.87       | 0.58       | 7.64       |
|                         | 2(51)  | 2.03(38)            | 4.92(39) | 0.84-0.91  | 0.35-0.58  | 7.46-7.64  |
| Simazine                | 1(15)  | 2.27(78)            | 1.65(79) | 1\$        | 2.05       | 6.10       |
|                         | 25(62) | 2.27(78)            | 1.65(79) | 1\$        | 0.35-2.84  | 4.30-7.10  |
| 2,4,5-T                 | 3(9)   | 3.36(81)            | 2.90(75) | 0.84-1.14  | 1.25-4.85  | 2.80-7.10  |
|                         | 2(75)  | 3.36(81)            | 2.90(75) | 0.81-0.85  | 0.46-1.74  | 5.90-7.70  |
| 1,2,3,4-TCB£            | 2(65)  | 4.64(44)            |          | 0.91-1.13  | 2.15-4.70  |            |
|                         | 2(73)  | 4.64(44)            |          | 1\$        | 0.16-1.87  |            |
| 1,2,3,5-TCB£            | 2(73)  | 4.46(41)            |          | 1\$        | 0.16-1.87  |            |
| Tetrachloroguaiacol     | 3(71)  | 4.45(42)            | 5.97(71) | 1\$        | 0.20-3.70  | 4.20-7.40  |
| 2,3,4,5-TCP&            | 2(34)  | 4.82(42)            | 6.96(40) | 0.81-0.82  | 2.15-3.54  | 4.80-5.60  |
| 2,3,4,6-TCP&            | 5(29)  | 4.42(42)            | 5.38(40) | 0.80-0.90  | 0.90-29.80 | 4.60-7.50  |
|                         | 3(71)  | 4.42(42)            | 5.38(40) | 1\$        | 0.20-3.70  | 4.20-7.40  |
| 3,4,5-Trichloroaniline  | 6(76)  | 3.49(76)            | 1.78(83) | 1          | 0.12-6.34  | 4.50-5.10  |
| 1,2,3-Trichlorobenzene  | 3(34)  | 4.14(44)            |          | 0.96-1.01  | 2.15-9.05  |            |
|                         | 2(73)  | 4.14(44)            |          | 1\$        | 0.16-1.87  |            |
| 1,2,4-Trichlorobenzene  | 2(73)  | 4.02(16)            |          | 1\$        | 0.16-1.87  |            |
| 1,3,5-Trichlorobenzene  | 2(73)  | 4.19(44)            |          | 1\$        | 0.16-1.87  |            |
| 4,5,6-Trichloroguaiacol | 3(71)  | 3.74(42)            | 7.40(71) | 1\$        | 0.20-3.70  | 4.20-7.40  |
| 2,4,5-Trichlorophenol   | 8(29)  | 3.72(49)            | 7.43(40) | 0.80-0.90  | 0.90-29.80 | 3.40-7.50  |
|                         | 4(34)  | 3.72(49)            | 7.43(40) | 0.81-0.98  | 2.15-9.05  | 3.60-5.90  |
| 2,4,6-Trichlorophenol   | 3(71)  | 3.75(42)            | 7.42(40) | 1\$        | 0.20-3.70  | 4.20-7.40  |
| Trifluralin             | 4(60)  | 5.07(60)            |          | 1          | 0.40-7.60  |            |

# number of experimental K<sub>p</sub> values

\*reference; †range; \$single concentration; §ref. 86; £Tetrachlorobenzene; &Tetrachlorophenol.

Comparison of Figs. 5 to 7 shows that Eq. 9 is the most suitable model to describe the sorption behavior of the 87 organic chemicals listed in Table 2.




Fig. 5. Observed versus calculated (Eq. 4)  $K_p$  values for the test set.




Fig. 6. Observed versus calculated (Eq. 7)  $K_p$  values for the test set.




Fig. 7. Observed versus calculated (Eq. 9)  $K_p$  values for the test set.

## CONCLUSION

Our study shows that it is possible to propose a unique general QSAR equation (Eq. 9) for estimating the sorption behavior of all the organic chemicals which can potentially contaminate the ecosystems. Introduction of Eq. 9 in the environmental fate models (e.g. fugacity models<sup>6, 87</sup>) could be useful to increase the accuracy of their outputs and stretch their domain of application. It could be also interesting to test the usefulness of our correction factors for ionization problems to improve the predictive power of the models based upon linear relationships between soil and/or sediment sorption coefficients and topological descriptors such as molecular connectivity indices.

## ACKNOWLEDGEMENTS

We acknowledge with appreciation the comments of Pr. M. Chastrette (Laboratoire de Chimie Organique Physique, University Lyon I, France) and Dr. T. E. McKone (Health and Ecological Assessments Division, Lawrence Livermore National Laboratory, California) which improved this paper.

## REFERENCES

1. Bahnick, D.A. and Doucette, W.J. (1988). Use of molecular connectivity indices to estimate soil sorption coefficients for organic chemicals. *Chemosphere* **17**, 1703-1715.
2. Domine, D., Devillers, J., Chastrette, M., and Karcher, W. (1992). Multivariate structure-property relationships (MSPR) of pesticides. *Pestic. Sci.* **35**, 73-82.
3. Devillers, J., Karcher, W., and Isnard, P. (1991). Graphical display on the fugacity model level I. *Chemosphere* **23**, 633-642.
4. Devillers, J., Thioulouse, J., Domine, D., Chastrette, M., and Karcher, W. (1991). Multivariate analysis of the input and output data in the fugacity model level I. In, *Applied Multivariate Analysis in SAR and Environmental Studies* (J. Devillers and W. Karcher, Eds.). Kluwer Academic Publishers, Dordrecht, pp. 281-345.
5. Jaffé, R. (1991). Fate of hydrophobic organic pollutants in the aquatic environment: a review. *Environ. Pollut.* **69**, 237-257.
6. Mackay, D. (1991). *Multimedia Environmental Models. The Fugacity Approach*. Lewis Publishers, Chelsea, p. 257.
7. McKone, T.E. (1993). The precision of QSAR methods for estimating intermedia transfer factors in exposure assessments. *SAR QSAR Environ. Res.* **1**, 41-51.
8. Green, R.E. and Karickhoff, S.W. (1990). Sorption estimates for modeling. In, *Pesticides in the Soil Environment: Processes, Impacts, and Modeling* (H.H. Cheng, Ed.). Soil Science Society of America, Madison, pp. 79-101.
9. von Oepen, B., Kördel, W., and Klein, W. (1991). Sorption of nonpolar and polar compounds to soils: processes, measurements and experience with the applicability of the modified OECD-Guideline 106. *Chemosphere* **22**, 285-304.
10. Lambert, S.M. (1968). Omega ( $\Omega$ ), a useful index of soil sorption equilibria. *J. Agric. Food Chem.* **16**, 340-343.
11. Grathwohl, P. (1990). Influence of organic matter from soils and sediments from various origins on the sorption of some chlorinated aliphatic hydrocarbons: implications on  $K_{oc}$  correlations. *Environ. Sci. Technol.* **24**, 1687-1693.
12. Rutherford, D.W., Chiou, C.T., and Kile, D.E. (1992). Influence of soil organic matter composition on the partition of organic compounds. *Environ. Sci. Technol.* **26**, 336-340.
13. Karickhoff, S.W., Brown, D.S., and Scott, T.A. (1979). Sorption of hydrophobic pollutants on natural sediments. *Wat. Res.* **13**, 241-248.
14. Kenaga, E.E. and Goring, C.A.I. (1980). Relationship between water solubility, soil sorption, octanol-water partitioning, and concentration of chemicals in biota. In, *Aquatic Toxicology* (J.G. Eaton, P.R. Parrish, and A.C. Hendricks, Eds.). ASTM STP 707, pp. 78-115.
15. Briggs, G.G. (1981). Theoretical and experimental relationships between soil adsorption, octanol-water partition coefficients, water solubilities, bioconcentration factors, and the parachor. *J. Agric. Food Chem.* **29**, 1050-1059.
16. Chiou, C.T., Porter, P.E., and Schmedding, D.W. (1983). Partition equilibria of nonionic organic compounds between soil organic matter and water. *Environ. Sci. Technol.* **17**, 227-

231.

17. Chin, Y.P., Peven, C.S., and Weber, W.J. (1988). Estimating soil/sediment partition coefficients for organic compounds by high performance reverse phase liquid chromatography. *Wat. Res.* **22**, 873-881.
18. Noegrohati, S. and Hammers, W.E. (1992). Regression models for some solute distribution equilibria in the terrestrial environment. *Toxicol. Environ. Chem.* **34**, 175-185.
19. Sabljic, A. and Protic, M. (1982). Relationship between molecular connectivity indices and soil sorption coefficients of polycyclic aromatic hydrocarbons. *Bull. Environ. Contam. Toxicol.* **28**, 162-165.
20. Koch, R. (1983). Molecular connectivity index for assessing ecotoxicological behaviour of organic compounds. *Toxicol. Environ. Chem.* **6**, 87-96.
21. Sabljic, A. (1984). Predictions of the nature and strength of soil sorption of organic pollutants by molecular topology. *J. Agric. Food Chem.* **32**, 243-246.
22. Sabljic, A. (1987). On the prediction of soil sorption coefficients of organic pollutants from molecular structure: application of molecular topology model. *Environ. Sci. Technol.* **21**, 358-366.
23. Meylan, W., Howard, P.H., and Boethling, R.S. (1992). Molecular topology/fragment contribution method for predicting soil sorption coefficients. *Environ. Sci. Technol.* **26**, 1560-1567.
24. Khan, A., Hassett, J.J., Banwart, W.L., Means, J.C., and Wood, S.G. (1979). Sorption of acetophenone by sediments and soils. *Soil Sci.* **128**, 297-302.
25. Banwart, W.L., Hassett, J.J., Wood, S.G., and Means, J.C. (1982). Sorption of nitrogen-heterocyclic compounds by soils and sediments. *Soil Sci.* **133**, 42-47.
26. Walton, B.T., Hendricks, M.S., Anderson, T.A., Griest, W.H., Merriweather, R., Beauchamp, J.J., and Francis, C.W. (1992). Soil sorption of volatile and semivolatile organic compounds in a mixture. *J. Environ. Qual.* **21**, 552-558.
27. Means, J.C., Wood, S.G., Hassett, J.J., and Banwart, W.L. (1982). Sorption of amino- and carboxy-substituted polynuclear aromatic hydrocarbons by sediments and soils. *Environ. Sci. Technol.* **16**, 93-98.
28. Schwarzenbach, R.P. and Westall, J. (1981). Transport of nonpolar organic compounds from surface water to groundwater. Laboratory sorption studies. *Environ. Sci. Technol.* **15**, 1360-1367.
29. Lagas, P. (1988). Sorption of chlorophenols in the soil. *Chemosphere* **17**, 205-216.
30. Means, J.C., Wood, S.G., Hassett, J.J., and Banwart, W.L. (1980). Sorption of polynuclear aromatic hydrocarbons by sediments and soils. *Environ. Sci. Technol.* **14**, 1524-1528.
31. Chiou, C.T., Peters, L.J., and Freed, V.H. (1979). A physical concept of soil-water equilibria for nonionic organic compounds. *Science* **206**, 831-832.
32. Jafvert, C.T. (1990). Sorption of organic acid compounds to sediments: initial model development. *Environ. Toxicol. Chem.* **9**, 1259-1268.
33. Walters, R.W., Ostazeski, S.A., and Guiseppi-Elie, A. (1989). Sorption of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin from water by surface soils. *Environ. Sci. Technol.* **23**, 480-484.



34. van Gestel, C.A.M. and Ma, W.C. (1990). An approach to quantitative structure-activity relationships (QSARs) in earthworm toxicity studies. *Chemosphere* **21**, 1023-1033.
35. Bowman, B.T. (1982). Conversion of Freundlich adsorption K values to the mole fraction format and the use of  $S_y$  values to express relative adsorption of pesticides. *Soil Sci. Soc. Am. J.* **46**, 740-743.
36. Schellenberg, K., Leuenberger, C., and Schwarzenbach, R.P. (1984). Sorption of chlorinated phenols by natural sediments and aquifer materials. *Environ. Sci. Technol.* **18**, 652-657.
37. Karickhoff, S.W. (1984). Organic pollutant sorption in aquatic systems. *J. Hydraul. Eng.* **110**, 707-735.
38. Sangster, J. (1989). Octanol-water partition coefficients of simple organic compounds. *J. Phys. Chem. Ref. Data* **18**, 1111-1229.
39. Zachara, J.M., Ainsworth, C.C., Cowan, C.E., and Thomas, B.L. (1987). Sorption of binary mixtures of aromatic nitrogen heterocyclic compounds on subsurface materials. *Environ. Sci. Technol.* **21**, 397-402.
40. Jones, P.A. (1981). *Chlorophenols and their Impurities in the Canadian Environment*. Environment Canada, Report SPE 3-EC-81-2F, p. 322.
41. Veith, G.D., Macek, K.J., Petrocelli, S.R., and Carroll, J. (1980). An evaluation of using partition coefficients and water solubility to estimate bioconcentration factors for organic chemicals in fish. In, *Aquatic Toxicology* (J.G. Eaton, P.R. Parrish, and A.C. Hendricks, Eds.). ASTM STP 707, pp. 116-129.
42. Xie, T.M., Hulthe, B., and Folestad, S. (1984). Determination of partition coefficients of chlorinated phenols, guaiacols, and catechols by shake-flask GC and HPLC. *Chemosphere* **13**, 445-459.
43. Westall, J.C., Leuenberger, C., and Schwarzenbach, R.P. (1985). Influence of pH and ionic strength on the aqueous-nonaqueous distribution of chlorinated phenols. *Environ. Sci. Technol.* **19**, 193-198.
44. de Bruijn, J., Busser, F., Seinen, W., and Hermens, J. (1989). Determination of octanol/water partition coefficients for hydrophobic organic chemicals with the "slow-stirring" method. *Environ. Toxicol. Chem.* **8**, 499-512.
45. El Tayar, N., Testa, B., and Carrupt, P.A. (1992). Polar intermolecular interactions encoded in partition coefficients: an indirect estimation of hydrogen-bond parameters of polyfunctional solutes. *J. Phys. Chem.* **96**, 1455-1459.
46. Dean, J.A. (1985). *Lange's Handbook of Chemistry*, thirteenth edition. McGraw-Hill Book Company, New York.
47. Howard, P.H. (1990). *Handbook of Environmental Fate and Exposure Data for Organic Chemicals III*. Lewis publishers, Chelsea, p. 684.
48. Sijm, D.T.H.M., Wever, H., de Vries, P.J., and Opperhuizen, A. (1989). Octan-1-ol/water partition coefficients of polychlorinated dibenzo-*p*-dioxins and dibenzofurans: experimental values determined with a stirring method. *Chemosphere* **19**, 263-266.
49. Leo, A., Hansch, C., and Elkins, D. (1971). Partition coefficients and their uses. *Chem. Rev.* **71**, 525-616.

50. Bailey, G.W., White, J.L., and Rothberg, T. (1968). Adsorption of organic herbicides by montmorillonite: role of pH and chemical character of adsorbate. *Soil Sci. Soc. Amer. Proc.* **32**, 222-234.
51. Zachara, J.M., Ainsworth, C.C., Felice, L.J., and Resch, C.T. (1986). Quinoline sorption to subsurface materials: role of pH and retention of the organic cation. *Environ. Sci. Technol.* **20**, 620-627.
52. Nicholls, P.H. and Evans, A.A. (1991). Sorption of ionisable organic compounds by field soils. Part 1: acids. *Pestic. Sci.* **33**, 319-330.
53. Nicholls, P.H. and Evans, A.A. (1991). Sorption of ionisable organic compounds by field soils. Part 2: cations, bases and zwitterions. *Pestic. Sci.* **33**, 331-345.
54. Green, R.E. (1974). Pesticide-clay-water interactions. In, *Pesticides in Soil and Water* (W.D. Guenzi, Ed.). Soil Science Society of America, Madison, pp. 3-37.
55. Scherrer, R.A. and Howard, S.M. (1977). Use of distribution coefficients in quantitative structure-activity relationships. *J. Med. Chem.* **20**, 53-58.
56. Wauchope, R.D., Buttler, T.M., Hornsby, A.G., Augustijn-Beckers, P.W.M., and Burt, J.P. (1992). The SCS/ARS/CES pesticide properties database for environmental decision-making. *Rev. Environ. Contam. Toxicol.* **123**, 1-155.
57. Sato, T., Kohnosu, S., and Hartwig, J.F. (1987). Adsorption of butachlor to soils. *J. Agric. Food Chem.* **35**, 397-402.
58. Liu, L.C., Cibes-Viadé, H., and Koo, F.K.S. (1970). Adsorption of ametryne and diuron by soils. *Weed Sci.* **18**, 470-474.
59. Dao, T.H., Bouchard, D., Mattice, J., and Lavy, T.L. (1986). Soil sorption of aniline and chloroanilines: direct and indirect concentration measurements of the adsorbed phase. *Soil Sci.* **141**, 26-30.
60. Francioso, O., Bak, E., Rossi, N., and Sequi, P. (1992). Sorption of atrazine and trifluralin in relation to the physico-chemical characteristics of selected soils. *Sci. Total Environ.* **123/124**, 503-512.
61. Singh, G., Spencer, W.F., Cliath, M.M., and van Genuchten, M.Th. (1990). Sorption behavior of *s*-triazine and thiocarbamate herbicides on soils. *J. Environ. Qual.* **19**, 520-525.
62. Talbert, R.E. and Fletchall, O.H. (1965). The adsorption of some *s*-triazines in soils. *Weeds* **13**, 46-52.
63. Austin, D.J. and Briggs, G.G. (1976). A new extraction method for benomyl residues in soil and its application in movement and persistence studies. *Pestic. Sci.* **7**, 201-210.
64. Achik, J., Schiavon, M., and Jamet, P. (1991). Study of carbofuran movement in soils part I-soil structure. *Environ. Int.* **17**, 73-79.
65. van Gestel, C.A.M., Ma, W.C., and Smit, C.E. (1991). Development of QSARs in terrestrial ecotoxicology: earthworm toxicity and soil sorption of chlorophenols, chlorobenzenes and dichloroaniline. *Sci. Total Environ.* **109/110**, 589-604.
66. Boyd, S.A. (1982). Adsorption of substituted phenols by soil. *Soil Sci.* **134**, 337-343.
67. Mersie, W. and Foy, C.L. (1986). Adsorption, desorption, and mobility of chlorsulfuron in soils. *J. Agric. Food Chem.* **34**, 89-92.
68. Walker, A., Cotterill, E.G., and Welch, S.J. (1989). Adsorption and degradation of

- chlorsulfuron and metsulfuron-methyl in soils from different depths. *Weed Res.* **29**, 281-287.
69. Hermosin, M.C. and Cornejo, J. (1991). Soil adsorption of 2,4-D as affected by the clay mineralogy. *Toxicol. Environ. Chem.* **31-32**, 69-77.
70. Hassett, J.J., Means, J.C., Banwart, W.L., Wood, S.G., Ali, S., and Khan, A. (1980). Sorption of dibenzothiophene by soils and sediments. *J. Environ. Qual.* **9**, 184-186.
71. Seip, H.M., Alstad, J., Carlberg, G.E., Martinsen, K., and Skaane, R. (1986). Measurement of mobility of organic compounds in soils. *Sci. Total Environ.* **50**, 87-101.
72. Wahid, P.A. and Sethunathan, N. (1978). Sorption-desorption of parathion in soils. *J. Agric. Food Chem.* **26**, 101-105.
73. Payá-Pérez, A.B., Riaz, M., and Larsen, B.R. (1991). Soil sorption of 20 PCB congeners and six chlorobenzenes. *Ecotoxicol. Env. Safety* **21**, 1-17.
74. Cortes, A., Riego, J., Payá-Pérez, A.B., and Larsen, B.R. (1991). Soil sorption of coplanar and non-planar PCBs. *Toxicol. Environ. Chem.* **31-32**, 79-86.
75. Koskinen, W.C, O'Connor, G.A., and Cheng, H.H. (1979). Characterization of hysteresis in the desorption of 2,4,5-T from soils. *Soil Sci. Soc. Am. J.* **43**, 871-874.
76. Payá-Pérez, A.B. and Pelusio, F. (1992). Sorption of 3,4,5-trichloroaniline in a soil profile. *Sci. Total Environ.* **123/124**, 519-529.
77. Suntio, L.R., Shiu, W.Y., Mackay, D., Seiber, J.N., and Glotfelty, D. (1988). Critical review of Henry's law constants for pesticides. *Rev. Environ. Contam. Toxicol.* **103**, 1-59.
78. Gerstl, Z. and Helling, C.S. (1987). Evaluation of molecular connectivity as a predictive method for the adsorption of pesticides by soils. *Environ. Sci. Health* **B22**, 55-69.
79. Weber, J.B. (1970). Mechanisms of adsorption of *s*-triazines by clay colloids and factors affecting plant availability. *Residue. Rev.* **32**, 93-130.
80. Grayson, B.T. and Kleier, D.A. (1990). Phloem mobility of xenobiotics. IV. Modelling of pesticide movement in plants. *Pestic. Sci.* **30**, 67-79.
81. Garten, C.T. and Trabalka J.R. (1983). Evaluation of models for predicting terrestrial food chain behavior of xenobiotics. *Environ. Sci. Technol.* **17**, 590-595.
82. Hulzebos, E.M., Adema, D.M.M., Dirven-van Breemen, E.M., Henzen, L., and van Gestel, C.A.M. (1991). QSARs in phytotoxicity. *Sci. Total Environ.* **109/110**, 493-497.
83. Perrin, D.D., Dempsey, B., and Serjeant, E.P. (1981). *pK<sub>a</sub> Prediction for Organic Acids and Bases*. Chapman and Hall, London, p. 146.
84. Shiu, W.Y. and Mackay, D. (1986). A critical review of aqueous solubilities, vapor pressures, Henry's law constants, and octanol-water partition coefficients of the polychlorinated biphenyls. *J. Phys. Chem. Ref. Data* **15**, 911-929.
85. Sklarew, D.S. and Girvin, D.C. (1987). Attenuation of polychlorinated biphenyls in soils. *Rev. Environ. Contam. Toxicol.* **98**, 1-41.
86. Dao, T.H., Marx, D.B., Lavy, T.L., and Dragun, J. (1982). Effect, and statistical evaluation, of soil sterilization on aniline and diuron adsorption isotherms. *Soil Sci. Soc. Am. J.* **46**, 963-969.
87. Devillers, J., Bintein, S., Domine, D., and Karcher, W. CHEMFRANCE: A regional fugacity level III model applied to France. (In preparation).

# ARTICLE III



## CHEMFRANCE: A REGIONAL LEVEL III FUGACITY MODEL APPLIED TO FRANCE

J. Devillers\* and S. Bintein  
CTIS, 21 rue de la Bannière, 69003 Lyon, France

W. Karcher  
Commission of the European Communities, Joint Research Centre, Ispra Establishment,  
I-21020 Ispra Varese, Italy

(Received in Germany 22 September 1994; accepted 31 October 1994)

### ABSTRACT

CHEMFRANCE, a computer model allowing to estimate the environmental fate of organic chemicals in France, is presented. This multimedia model is represented by six bulk compartments (i.e., air, surface water, soil, bottom sediment, groundwater, coastal water), and by ten subcompartments. The model employs the fugacity concept and requires information on chemical properties (i.e., molecular weight, aqueous solubility, vapor pressure, *n*-octanol/water partition coefficient, dissociation constant, melting point, reaction half-lives) and emission rates of the pollutants. The outputs of the model consist of estimated chemical distribution between environmental media, transport and transformation process rates, and steady state concentrations in one of the twelve defined regions of France, or France as a whole, at a chosen season.

The nature of the model and its underlying assumptions are described. An illustrative example dealing with the modeling of the environmental fate of isobutylene is presented. Aspects of model validation are discussed and recommendations are made for a proper use of CHEMFRANCE.

-----  
\* Author to whom all correspondence should be addressed.

## INTRODUCTION

Pollutants discharged into the environment are distributed across environmental media (e.g., air, water, soil) and among biota as the result of their physicochemical properties (e.g., aqueous solubility, vapor pressure, *n*-octanol/water partition coefficient (log *P*), dissociation constant) and the nature of the environment (e.g., climatic parameters).<sup>1, 2</sup> The human mind is not able to take into account simultaneously the above parameters in order to estimate the theoretical distribution of a chemical between the different compartments of the biosphere. Mathematical models, at different levels of complexity can be used to overcome this problem. These models are very attractive since they provide a rapid and inexpensive simulation tool to produce a comprehensive picture of the environmental fate of the organic chemicals.<sup>3, 4</sup> Among them, the level III fugacity models are particularly powerful to estimate the environmental distribution of pollutants in abiotic and biotic media.<sup>4-6</sup> The assessment is based on information relative to emission rates, environmental conditions, and physicochemical and reactive properties of the organic chemicals.

In this paper, we describe a level III computer model (CHEMFRANCE) which can be used to estimate the environmental fate of organic chemicals in France. The country has been divided into twelve regions where the environment is represented by a multimedia system constituted of six bulk compartments (i.e., air, surface water, soil, bottom sediment, groundwater, coastal water). The first four bulk compartments are considered as a combination of subcompartments of varying proportions of pure and particle phases. Equilibrium is assumed to apply within each bulk compartment, but is not assumed between compartments. Expressions for emissions, advective flows, degrading reactions, and interphase transports by diffusive and non-diffusive processes are included in CHEMFRANCE.

The aim of this paper is basically to present the theoretical foundations of CHEMFRANCE and underline the difficulties encountered to construct such a model. For illustrative purposes, the modeling of the environmental fate of isobutylene is presented.

## MODEL DESCRIPTION

### Regions of CHEMFRANCE

To develop CHEMFRANCE, it was necessary to divide France into a reasonable number of regions of similar hydrogeological, climatic, and ecologic characteristics. These regions were defined after a careful study of the factors susceptible to have an effect on the environmental fate of chemicals in France. Thus, it is obvious that all the economical or political parameters inducing artifactitious divisions (e.g., limits of department, high density population area) were excluded. The factors selected in priority were the drainage basins constituting the French hydrographical network, the climate (i.e., precipitations, temperatures, wind), the nature of the soil, and the different types of vegetation.

Under these conditions, France was divided into the twelve following regions (Figure 1):

Region 1- Bretagne, Pays de Loire

Region 2- Nord, Normandie

Region 3- Bassin de Paris

Region 4- Alsace, Lorraine

Region 5- Vallée du Rhône

Region 6- Massif central

Region 7- Alpes

Region 8- Bassin méditerranéen

Region 9- Aquitaine, Midi-Pyrénées

Region 10- Pyrénées

Region 11- Bordure atlantique

Region 12- Centre

Region 13 being France

Dividing a country into regions is ultimately a matter of judgement, and is almost certain to generate controversy. Therefore, the segmentation presented in Figure 1 has to be regarded as an optimal compromise which can be modified for particular simulation purposes. Indeed, in CHEMFRANCE an option allows a user to define its own regions without any kind of restriction.

## Environmental media

The choice of environmental compartment structures requires to identify the acceptable level of model complexity. The modeler should make a balance between the current perceptions of the system and the experimental and environmental data available from the system. In CHEMFRANCE, the environment is rationalized and simplified to a multimedia system constituted of six bulk compartments (i.e., air, surface water, soil, bottom sediment, groundwater, coastal water). The first four bulk compartments (i.e., principal compartments) consist of a combination of subcompartments of varying proportions of pure and particle phases. A brief description of these compartments is given below.

### 1- Air

The air compartment is divided into two subcompartments (i.e., pure air, air particles). In CHEMFRANCE, chemicals enter the air compartment by local emissions, advective inflow from adjacent regions, and by diffusion (i.e., volatilization) from soil and surface water. The environmental loss processes are advective outflow, reaction, transfer to higher altitude, wet and dry depositions, rain dissolution of gaseous chemicals, and diffusion (i.e., adsorption) to soil and water. An average value of 5 days was selected for the air residence time, in order to calculate the advective flows.<sup>5</sup> The total surface area for each region was calculated from the surface of departments and the fraction of departments included in each region.<sup>7</sup> The air height was reduced to 1000 m, since the depth of the troposphere, into which emissions generally occur and are constrained, is typically around 100 m during the night and 1000 m during the day.<sup>8, 9</sup>




Figure 1: Defined regions of CHEMFRANCE.



## 2- Surface water

The surface water compartment is composed of pure water, suspended sediment, and biota (only represented by fish). In CHEMFRANCE, pollutants move into this compartment by local emissions, advective inflow from adjacent regions, diffusion from air and sediment, soil and water runoffs, wet and dry depositions, rain dissolution of gaseous chemicals, and sediment resuspension. The removal processes are advective outflow, reaction, diffusion to air and sediment, and deposition. In order to calculate the advective flows, an average value of 500 days was selected for the surface water residence time.<sup>5</sup> This compartment is used to represent lakes, ponds, dams, rivers, and canals. The surface and volume of surface water were estimated from numerous bibliographical references.<sup>10-18</sup> When only the surface area was available, we assumed an average depth of 10 m, 2 m, and 15 m for lakes, ponds, and dams, respectively. The volume fraction of suspended sediment was also estimated for each region.<sup>19, 20</sup> In order to take into account the possible ionization of chemicals, an average pH value of suspended sediment equal to 7 was adopted for all the regions.<sup>21</sup>

## 3- Soil

The soil compartment consists of three subcompartments (i.e., pure soil, soil water, soil air). Pollutants move into this compartment by local emissions, diffusion from water (i.e., adsorption), wet and dry depositions, and by rain dissolution of gaseous chemicals. Chemicals are assumed to be removed by reaction, diffusion to air (i.e., volatilization), soil and water runoffs, and by leaching to groundwater. Soil was characterized by depth (15 cm), density, air and water content, fraction of organic carbon, and the pH allowing to take into account the ionization of acidic and basic compounds. In CHEMFRANCE, only the last two parameters were adapted to each region. The nature of the soil of the twelve regions was determined from a pedologic map,<sup>22</sup> which allowed to select the predominant soil. The soil was then characterized from the upper layer type (i.e., horizon A<sub>0</sub> or A<sub>1</sub> in pedology).<sup>23-25</sup>

## 4- Bottom sediment

This compartment is constituted of two subcompartments (i.e., pure sediment, sediment pore water). Chemicals enter this compartment by local emissions, deposition, and by diffusion from water. The environmental loss mechanisms are reaction, diffusion to water, resuspension, and sediment burial. As for the suspended sediment, an average pH value equal to 7 was adopted for all the regions,<sup>21</sup> in order to take into account the possible ionization of chemicals.

## 5- Groundwater

The groundwater compartment represents the saturated material of the lowest soil horizon. For each region, the volume of groundwater was estimated from the nature, the surface area, the thickness and the porosity of the substratum.<sup>26-35</sup>

## 6- Coastal water

The coastal water was assumed to extend 1 km offshore with a mean depth of 100 m. A flushing time of 100 h and an advective inflow rate of 500 days were selected for all the regions possessing coasts. Marine biota and sediments were ignored in our model.

### Fugacity and basic equations

CHEMFRANCE is a level III fugacity model. The concept of fugacity<sup>36</sup> and its usefulness in environmental modeling<sup>4-6, 37-46</sup> have been clearly described in the seminal book of Mackay.<sup>4</sup> Briefly, fugacity, which means escaping or fleeing tendency, has units of pressure (F in Pa) and can be viewed as the partial pressure which a chemical exerts as it attempts to escape from one phase and migrate to another.<sup>41</sup> Fugacity is linearly related to concentration (C in mol/m<sup>3</sup>) through the fugacity capacity (Z in mol/m<sup>3</sup> Pa):

$$C = Z F \quad (1)$$

The Z values depend on the physicochemical properties of the studied substance, the nature of the phase into which the chemical partitions, and the environmental temperature.

In CHEMFRANCE, the Z values are calculated according to Mackay and Paterson<sup>40</sup> (Tables I and II) except for those related to the bioconcentration of chemicals in fish (BCF) and to the sorption phenomena (Kp).

Table I. Definition of Z values (mol/m<sup>3</sup> Pa) for the subcompartments.

| Subcompartment (i,j)      | Z <sub>(i,j)</sub>  |
|---------------------------|---|
| pure air (1,1)            | 1/RT  |
| air particles (1,3)       | 6 · 10 <sup>6</sup> Z <sub>(1,1)</sub> /VPL                     |
| pure water (2,2)          | 1/H   |
| suspended sediment (2,3)  | Kp <sub>(2,3)</sub> ρ <sub>(2,3)</sub> Z <sub>(2,2)</sub> /1000 |
| biota (fish) (2,4)        | BCF ρ <sub>(2,4)</sub> Z <sub>(2,2)</sub> /1000                 |
| soil air (3,1)            | 1/RT  |
| soil water (3,2)          | 1/H   |
| pure soil (3,3)           | Kp <sub>(3,3)</sub> ρ <sub>(3,3)</sub> Z <sub>(2,2)</sub> /1000 |
| sediment pore water (4,2) | 1/H   |
| pure sediment (4,3)       | Kp <sub>(4,3)</sub> ρ <sub>(4,3)</sub> Z <sub>(2,2)</sub> /1000 |

R = 8.314 (Pa m<sup>3</sup>/mol K); T = environmental temperature (K); VPL\* = liquid vapor pressure (Pa); H = VP/S = Henry's law constant (Pa m<sup>3</sup>/mol); S<sup>£</sup> = solubility in water (mol/m<sup>3</sup>); VP<sup>£</sup> = vapor pressure (Pa); Kp = sorption coefficient; BCF = bioconcentration factor; ρ<sub>(i,j)</sub> = density of subcompartment (i,j) (kg/m<sup>3</sup>).

\* for solids VPL = VP/exp (6.79 (1-MP/T)) in which MP is the melting point (K) of the chemical.

£ temperature dependence is included.<sup>40</sup>

Table II. Definition of Z values (mol/m<sup>3</sup> Pa) for the bulk compartments.

| Compartment (i)     | Z <sub>(i)</sub>   |
|---------------------|--|
| air (1)             | Z <sub>(1,1)</sub> +Z <sub>(1,3)</sub> ϕ <sub>(1,3)</sub>  |
| surface water (2)   | Z <sub>(2,2)</sub> +Z <sub>(2,3)</sub> ϕ <sub>(2,3)</sub> +Z <sub>(2,4)</sub> ϕ <sub>(2,4)</sub>                     |
| soil (3)            | Z <sub>(3,3)</sub> ϕ <sub>(3,3)</sub> +Z <sub>(3,1)</sub> ϕ <sub>(3,1)</sub> + Z <sub>(3,2)</sub> ϕ <sub>(3,2)</sub> |
| bottom sediment (4) | Z <sub>(4,3)</sub> ϕ <sub>(4,3)</sub> +Z <sub>(4,2)</sub> ϕ <sub>(4,2)</sub>   |
| groundwater (5)     | Z <sub>(2,2)</sub> = 1/H   |
| coastal water (6)   | 0.8 Z <sub>(2,2)</sub> = 0.8/H   |

ϕ<sub>(i,j)</sub> = volume fraction of subcompartment (i,j).

Indeed, to estimate the BCF values in fish a bilinear model<sup>47</sup> (Eq. 2) particularly suitable for very hydrophobic chemicals is used.

$$\log \text{BCF} = 0.910 \log P - 1.975 \log (6.8 \cdot 10^{-7} P + 1) - 0.786 \quad (2)$$

$$n = 154 \quad s = 0.347 \quad r = 0.950 \quad F = 463.51$$

The K<sub>p</sub> values are estimated from an original model taking into account the possible ionization of basic and acidic chemicals.<sup>48</sup> This model<sup>48</sup> requires to know the log P and pK<sub>a</sub> values of the chemicals and also the fraction of organic carbon (foc) and pH of the sorbent (soil or sediment).

CHEMFRANCE describes a steady-state system with non-equilibrium distribution between the bulk compartments in which each phase may have a different fugacity.<sup>39</sup> Therefore, intermedia transfer coefficients D (in mol/h Pa) for the various diffusive and non-diffusive processes<sup>40</sup> have to be calculated as displayed in Tables III and IV.

Table III. Calculation of D parameters (mol/h Pa).

| Compartment Process | Individual D   |  |
|---------------------|----------------|--|
| (1)-(2)             | diffusion      | $D_V = 1/((1/\text{TK}_{(1,2)} \text{AR}_{(2)} Z_{(1,1)})+(1/\text{TK}_{(2,1)} \text{AR}_{(2)} Z_{(2,2)}))$                      |
|                     | rain           | $D_{QW} = \text{AR}_{(2)} Z_{(2,2)} U_Q$   |
|                     | wet deposition | $D_{DW} = \text{AR}_{(2)} Z_{(1,3)} \phi_{(1,3)} U_D$  |
|                     | dry deposition | $D_{PW} = \text{AR}_{(2)} Z_{(1,3)} \phi_{(1,3)} U_P$  |
| (2)-(1)             | diffusion      | $D_V = 1/((1/\text{TK}_{(1,2)} \text{AR}_{(2)} Z_{(1,1)})+(1/\text{TK}_{(2,1)} \text{AR}_{(2)} Z_{(2,2)}))$                      |
| (1)-(3)             | diffusion      | $D_S = 1/((1/\text{TK}_{(1,3)} \text{AR}_{(3)} Z_{(1,1)})+(Y_3/(\text{AR}_{(3)} (\text{BA}_3 Z_{(1,1)}+\text{BW}_3 Z_{(2,2)})))$ |
|                     | rain           | $D_{QS} = \text{AR}_{(3)} Z_{(2,2)} U_Q$   |
|                     | wet deposition | $D_{DS} = \text{AR}_{(3)} Z_{(1,3)} \phi_{(1,3)} U_D$  |
|                     | dry deposition | $D_{PS} = \text{AR}_{(3)} Z_{(1,3)} \phi_{(1,3)} U_P$  |
| (3)-(1)             | diffusion      | $D_S = 1/((1/\text{TK}_{(1,3)} \text{AR}_{(3)} Z_{(1,1)})+(Y_3/(\text{AR}_{(3)} (\text{BA}_3 Z_{(1,1)}+\text{BW}_3 Z_{(2,2)})))$ |
| (3)-(2)             | water runoff   | $D_{WW} = \text{AR}_{(3)} Z_{(2,2)} U_{WW}$  |
|                     | soil runoff    | $D_{SW} = \text{AR}_{(3)} Z_{(3,3)} U_{SW}$  |

Table III. (Continued)

| Compartment   | Process      | Individual D  |
|---|--------------|---|
| (2)-(4)   | diffusion    | $D_Y = 1/((1/TK_{(2,4)} AR_{(4)} Z_{(2,2)})+(Y_4/BW_4 AR_{(4)} Z_{(2,2)}))$ |
|   | deposition   | $D_{DX} = AR_{(4)} Z_{(2,3)} U_{DX}$  |
| (4)-(2)   | diffusion    | $D_Y = 1/((1/TK_{(2,4)} AR_{(4)} Z_{(2,2)})+(Y_4/BW_4 AR_{(4)} Z_{(2,2)}))$ |
|   | resuspension | $D_{RX} = AR_{(4)} Z_{(4,3)} U_{RX}$  |
| D parameters for advective, pseudo-advective and reaction processes |              |   |
| advection (i = 1 to 2)  |              | $D_{A(i)} = Z_{(i)} GA_{(i)} = Z_{(i)} V_{(i)}/\tau_{(i)}$                  |
| transfer to higher altitude   |              | $D_{ST} = Z_{(1)} U_{ST} AR_{(1)}$  |
| leaching from soil  |              | $D_L = AR_{(3)} Z_{(2,2)} U_L$  |
| advective flow from coastal water                                   |              | $D_{CW} = Z_{(6)} V_{(6)}/\tau_{CW}$  |
| advective flow to coastal water                                     |              | $D_{A(6)} = D_{A(2)}$   |
| sediment burial   |              | $D_{BX} = AR_{(4)} Z_{(4,3)} U_{BX}$  |
| reaction (i = 1 to 4)   |              | $D_{R(i)} = V_{(i)} Z_{(i)} k_{(i)}$  |

$AR_{(i)}$  = area of compartment i and  $V_{(i)}$  = volume of compartment i.

Table IV. Total D value ( $DT_{(i)}$ ) for the principal compartments.

$$DT_{(1)} = D_{(1,2)} + D_{(1,3)} + D_{A(1)} + D_{R(1)} + D_{ST}$$

$$DT_{(2)} = D_{(2,1)} + D_{(2,4)} + D_{A(2)} + D_{R(2)}$$

$$DT_{(3)} = D_{(3,1)} + D_{(3,2)} + D_{R(3)} + D_L$$

$$DT_{(4)} = D_{(4,2)} + D_{R(4)} + D_{BX}$$

$$D_{(i,j)} = \sum \text{individual } D_{(i,j)} \text{ of Table III.}$$

Equations for calculation of the D values (Tables III and IV) require to use the transport velocity parameters described in Table V.

Table V. Assumed transport parameters.

| Parameter                                     | Symbol       | Value                | Unit    |
|---|--------------|----------------------|---------|
| air-side mass transfer coefficient over water | $TK_{(1,2)}$ | 3                    | m/h     |
| water-side mass transfer coefficient          | $TK_{(2,1)}$ | 0.03                 | m/h     |
| rain rate                                     | $U_Q$        | $12 U_{QM}/(24+365)$ | m/h     |
| rain rate                                     | $U_{QM}$     | Region dependent     | m/month |
| wet deposition velocity                       | $U_D$        | $U_Q Q$              | m/h     |
| scavenging ratio                              | $Q$          | 200000               | no unit |
| dry deposition velocity                       | $U_P$        | 10.8                 | m/h     |
| air-side mass transfer coefficient over soil  | $TK_{(1,3)}$ | 1                    | m/h     |

Table V. (Continued)

| Parameter  | Symbol       | Value  | Unit    |
|--|--------------|--|---------|
| diffusion path length in soil                      | $Y_3$        | 0.05   | m       |
| effective diffusivity in soil air                  | $BA_3$       | $BMA (\phi_{(3,1)})^{3.33}/(1-\phi_{(3,3)})^2$     | $m^2/h$ |
| molecular diffusivity in air                       | BMA          | 0.04   | $m^2/h$ |
| effective diffusivity in soil water                | $BW_3$       | $BMW (\phi_{(3,2)})^{3.33}/(1-\phi_{(3,3)})^2$     | $m^2/h$ |
| molecular diffusivity in water                     | BMW          | $4 \cdot 10^{-6}$                                  | $m^2/h$ |
| water runoff rate from soil                        | $U_{WW}$     | $12 U_{WWM}/(24 \cdot 365)$                        | m/h     |
| velocity of water runoff from soil                 | $U_{WWM}$    | $0.4 U_{QM}$                                       | m/month |
| solids runoff rate from soil                       | $U_{SW}$     | $2 \cdot 10^{-4}/(24 \cdot 365)$                   | m/h     |
| water-side mass transfer coefficient over sediment | $TK_{(2,4)}$ | 0.01   | m/h     |
| diffusion path length in sediment                  | $Y_4$        | $5 \cdot 10^{-3}$                                  | m       |
| effective diffusivity in sediment pore water       | $BW_4$       | $BMW (\phi_{(4,2)})^{1.33}$                        | $m^2/h$ |
| sediment deposition rate                           | $U_{DX}$     | $4 \cdot 10^{-4}/(24 \cdot 365)$                   | m/h     |
| sediment resuspension rate                         | $U_{RX}$     | $1 \cdot 10^{-4}/(24 \cdot 365)$                   | m/h     |
| transfer to higher altitude                        | $U_{ST}$     | $90/(24 \cdot 365)$                                | m/h     |
| leaching rate from soil to groundwater             | $U_L$        | $0.4 U_Q$  | m/h     |
| flushing time for coastal water                    | $\tau_{CW}$  | 100  | h       |
| air residence time                                 | $\tau_{(1)}$ | 120  | h       |
| water residence time                               | $\tau_{(2)}$ | 12000  | h       |
| sediment burial rate                               | $U_{BX}$     | $U_{DX}-U_{RX} = (3 \cdot 10^{-4}/(24 \cdot 365))$ | m/h     |

For the four principal compartments, the previous calculations (Tables I to IV) allow to setup steady state mass balance equations (Table VI), which incorporate emissions, transfers between adjacent phases, reaction, and advection. The solutions for the four unknown fugacity values (Table VI) can be easily obtained by a matrix inversion technique such as the Gaussian elimination.

Table VI. Mass balance equations.

| Compartment     | Equation   |
|-----------------|--|
| air             | $E_{(1)} + G_{A(1)} CB_{(1)} + F_{(2)} D_{(2,1)} + F_{(3)} D_{(3,1)} = F_{(1)} DT_{(1)}$                     |
| surface water   | $E_{(2)} + G_{A(2)} CB_{(2)} + F_{(1)} D_{(1,2)} + F_{(3)} D_{(3,2)} + F_{(4)} D_{(4,2)} = F_{(2)} DT_{(2)}$ |
| soil            | $E_{(3)} + F_{(1)} D_{(1,3)} = F_{(3)} DT_{(3)}$   |
| bottom sediment | $E_{(4)} + F_{(2)} D_{(2,4)} = F_{(4)} DT_{(4)}$   |

$CB_{(i)}$  = background inflow concentration ( $mol/m^3$ ) of adjacent region in compartment  $i$  and  $E_{(i)}$  = emission ( $mol/m^3$ ) in compartment  $i$ .

In order to estimate the fugacity in the two remaining bulk phases (i.e., groundwater, coastal water), we use fugacity values relative to the principal phases:

$$\text{Groundwater} \quad F(5) = F(3) \quad (3)$$

$$\text{Coastal water} \quad F(6) = F(2) D_{A(6)}/D_{CW} \quad (4)$$

Eq. (3) assumes that the groundwater and the soil compartments are in equilibrium. This assumption is based on the relatively prolonged contact between the soil water and the surface soil during infiltration.<sup>40</sup> Eq. (4) allows to calculate the fugacity of the coastal water from a consideration of the advective inflow rate and the flushing time using a simple dilution calculation.<sup>40</sup>

Fugacity values of the subcompartments are obtained from those of the principal compartments, since equilibrium partitioning is assumed to apply within each of the four principal compartments (i.e.,  $F(i) = F(i,j)$ ). This implies that the time required to reach equilibrium within a bulk phase is short compared to the time required to reach equilibrium between bulk phases.<sup>40</sup>

CHEMFRANCE allows one to obtain a considerable amount of information which is summarized in Table VII.

Table VII. Some output information obtained with CHEMFRANCE.

| Output                                  | Equation  | Unit               |
|---|---|--------------------|
| concentration                           | $C(i) = F(i) Z(i)$  | mol/m <sup>3</sup> |
| amount                                  | $M(i) = V(i) C(i)$  | mol                |
| diffusive flux between two phases       | $N_{d(i,j)} = D_{(i,j)} (F(i) - F(j))$                                      | mol/h              |
| non-diffusive flux between phases       | $N_{nd(i)} = D_{(i)} F(i)$  | mol/h              |
| first-order reaction process in a phase | $N_{k(i)} = DR_{(i)} F(i)$  | mol/h              |
| net flux between two phases             | $N_{f(i,j)} = D_{(i,j)} F(i) - D_{(j,i)} F(j)$                              | mol/h              |
| persistence time                        | $TP = \frac{\sum M(i)}{\sum DR_{(i)} F(i)}$                                 | h                  |
| residence time                          | $TR = \frac{\sum M(i)}{\sum DA_{(i)} \frac{CB_{(i)}}{Z(i)} + \sum E_{(i)}}$ | h                  |

i and j represent compartments.

### Environmental parameters

The monthly mean precipitations and temperatures were determined from maps<sup>49</sup> and data given locality by locality.<sup>50</sup> Table VIII summarizes the environmental parameters for which no region dependence was included. The values of the remaining parameters (i.e., area and volume of the compartments, coastal length, monthly mean precipitation and temperature for the four seasons, fraction of organic carbon content and pH of soil, volume fraction of suspended sediment) were selected for each region.

Table VIII. Values of constant environmental parameters selected for the twelve regions of CHEMFRANCE\*.

| Parameter   | Selected value  |
|---|---|
| fraction of organic carbon content of suspended sediment          | $foc_{(2,3)} = 0.2$   |
| fraction of organic carbon content of pure sediment               | $foc_{(4,3)} = 0.04$  |
| pH of suspended sediment  | $pH_{(2,3)} = 7$  |
| pH of pure sediment   | $pH_{(4,3)} = 7$  |
| volume fraction of air particles                                  | $\phi_{(1,3)} = 2 \cdot 10^{-11}$   |
| volume fraction of biota  | $\phi_{(2,4)} = 1 \cdot 10^{-6}$  |
| volume fraction of soil air                                       | $\phi_{(3,1)} = 0.2$  |
| volume fraction of soil water                                     | $\phi_{(3,2)} = 0.3$  |
| volume fraction of sediment pore water                            | $\phi_{(4,2)} = 0.7$  |
| density of pure air and soil air                                  | $\rho_{(1,1)} = \rho_{(3,1)} = 1.2 \text{ kg/m}^3$                                |
| density of air particles and suspended sediment                   | $\rho_{(1,3)} = \rho_{(2,3)} = 1500 \text{ kg/m}^3$                               |
| density of pure soil and pure sediment                            | $\rho_{(3,3)} = \rho_{(4,3)} = 2400 \text{ kg/m}^3$                               |
| density of pure water, biota, soil water, and sediment pore water | $\rho_{(2,2)} = \rho_{(2,4)} = \rho_{(3,2)} = \rho_{(4,2)} = 1000 \text{ kg/m}^3$ |
| height of air   | $DEP_{(1)} = 1000 \text{ m}$  |
| depth of soil   | $DEP_{(3)} = 0.15 \text{ m}$  |
| depth of sediment   | $DEP_{(3)} = 0.03 \text{ m}$  |
| depth of coastal water  | $DEP_{(6)} = 100 \text{ m}$   |

\* all these values can be changed by the user, as well as all those selected for each region.

## CASE STUDY: ISOBUTYLENE

Isobutylene [CAS RN 115-11-7] has become an important starting material for the manufacture of polymers and chemicals. There are many patents describing the use of isobutylene or its derivatives to produce insecticides, antioxidants, elastomers, additives for lubricating oils, adhesives, sealants, and caulking compounds.<sup>51, 52</sup> The estimated Western Europe isobutylene consumption in 1984 was  $1640 \cdot 10^3 \text{ t/y}$ .<sup>53</sup> The main sources for the commercial production of isobutylene are catalytic or thermal cracking ( $730 \cdot 10^3 \text{ t/y}$  in Western Europe) and steam cracking ( $920 \cdot 10^3 \text{ t/y}$  in Western Europe). There are other processes available for the production of isobutylene but they are more site or manufacturer specific (e.g., dehydration of *tert*-butanol,  $50 \cdot 10^3 \text{ t/y}$  in Western Europe).<sup>51, 53</sup> Environmental contaminations of isobutylene principally occur during industrial processes of production and use as well as during fuel combustions (stationary and mobile sources).<sup>54-57</sup> To illustrate the usefulness of CHEMFRANCE as simulation tool, we have modeled the environmental fate of isobutylene in France. In our case study, the physicochemical properties required to run CHEMFRANCE were the following:

- Molecular weight:  $56.11 \text{ g/mol}$ ,

- Solubility in water: 263 mg/l at room temperature,<sup>58</sup>
- Vapor pressure: 257.4 kPa at 20°C,<sup>59</sup>
- *n*-Octanol/water partition coefficient (log P): 2.11,<sup>59</sup>
- Melting point: -140 °C,<sup>59</sup>
- Half-life in air: 4 h.<sup>59</sup>

In addition to the six above input data, it was necessary to make the following assumptions:

- due to the lack of information on the degradation processes of isobutylene in the water, soil, and sediment compartments, it was necessary to estimate the half-lives of this chemical in these compartments. As generally assumed, in that case, fictitiously large half-life values are supplied inducing that there is no reaction.<sup>40</sup> In this study, a half-life value of 10<sup>11</sup> h was arbitrarily selected.

- as no information was available on the amounts of isobutylene released to the environment in France, a scenario was designed from the estimation of isobutylene emissions in the California's South Coast Basin.<sup>55, 60</sup> Thus, considering that in the California's South Coast Basin, an atmospheric emission of 7 10<sup>3</sup> kg/d was measured for a surface area of 60000 km<sup>2</sup>, the emissions in the 12 regions of CHEMFRANCE were calculated using the ratio of these two above values and the surface area of the corresponding region. Only the results obtained with region n°8 (i.e., bassin méditerranéen) in summer are presented here.

Thus, the outputs of CHEMFRANCE (Tables IX to XII) clearly show that the concentrations of isobutylene are low in all the compartments. The highest amount (99.99%) is found in the air compartment (Table X). The high percentage found in air is due to the high vapor pressure of this chemical. It therefore exists as a gas at ambient temperature. This high vapor pressure value combined to low log P and high Henry's law constant values induces that the transfers from air to the other compartments are not important. It is noteworthy that the major part of the emissions reacts in air (i.e., 95.4%) and the remaining is advected (i.e., 4.6%) (Table XII). All other reactions and transfer processes are negligible. The individual intermedia transport coefficients (D) are small compared to the reaction (D<sub>R(1)</sub>) and advection (D<sub>A(1)</sub>) values (Table XI). Therefore isobutylene tends to remain in air where it is rapidly degraded or advected. Indeed, its residence and persistence time values are very low (5.51 h and 5.77 h, respectively). From a practical point of view, this means that the outflow to other regions is unlikely due to the rapid degradation of this chemical.

Table IX. Z and F values obtained in region n°8 (summer) for isobutylene.

| Compartment             | Z (mol/m <sup>3</sup> Pa)                | F (Pa)                                    |
|-------------------------|--|---|
| <b>Bulk compartment</b> |  |   |
| air                     | Z <sub>(1)</sub> = 4.08 10 <sup>-4</sup> | F <sub>(1)</sub> = 1.17 10 <sup>-6</sup>  |
| surface water           | Z <sub>(2)</sub> = 1.63 10 <sup>-5</sup> | F <sub>(2)</sub> = 1.19 10 <sup>-6</sup>  |
| soil                    | Z <sub>(3)</sub> = 2.71 10 <sup>-4</sup> | F <sub>(3)</sub> = 1.17 10 <sup>-6</sup>  |
| bottom sediment         | Z <sub>(4)</sub> = 8.80 10 <sup>-5</sup> | F <sub>(4)</sub> = 1.19 10 <sup>-6</sup>  |
| groundwater             | Z <sub>(5)</sub> = 1.63 10 <sup>-5</sup> | F <sub>(5)</sub> = 1.17 10 <sup>-6</sup>  |
| coastal water           | Z <sub>(6)</sub> = 1.30 10 <sup>-5</sup> | F <sub>(6)</sub> = 6.42 10 <sup>-10</sup> |



Table IX. (Continued)

| Compartment           | Z (mol/m <sup>3</sup> Pa)        | F (Pa)                           |
|-----------------------|----------------------------------|----------------------------------|
| <b>Subcompartment</b> |                                  |                                  |
| pure air              | $Z_{(1,1)} = 4.08 \cdot 10^{-4}$ | $F_{(1,1)} = 1.17 \cdot 10^{-6}$ |
| air particles         | $Z_{(1,3)} = 8.27 \cdot 10^{-3}$ | $F_{(1,3)} = 1.17 \cdot 10^{-6}$ |
| pure water            | $Z_{(2,2)} = 1.63 \cdot 10^{-5}$ | $F_{(2,2)} = 1.19 \cdot 10^{-6}$ |
| suspended sediment    | $Z_{(2,3)} = 7.98 \cdot 10^{-4}$ | $F_{(2,3)} = 1.19 \cdot 10^{-6}$ |
| biota (fish)          | $Z_{(2,4)} = 2.17 \cdot 10^{-4}$ | $F_{(2,4)} = 1.19 \cdot 10^{-6}$ |
| soil air              | $Z_{(3,1)} = 4.08 \cdot 10^{-4}$ | $F_{(3,1)} = 1.17 \cdot 10^{-6}$ |
| soil water            | $Z_{(3,2)} = 1.63 \cdot 10^{-5}$ | $F_{(3,2)} = 1.17 \cdot 10^{-6}$ |
| pure soil             | $Z_{(3,3)} = 3.70 \cdot 10^{-4}$ | $F_{(3,3)} = 1.17 \cdot 10^{-6}$ |
| sediment pore water   | $Z_{(4,2)} = 1.63 \cdot 10^{-5}$ | $F_{(4,2)} = 1.19 \cdot 10^{-6}$ |
| pure sediment         | $Z_{(4,3)} = 2.55 \cdot 10^{-4}$ | $F_{(4,3)} = 1.19 \cdot 10^{-6}$ |

Table X. Chemical fate of isobutylene.

| Compartment             | Percentage | Amount<br>moles      | Concentrations        |                       |                      |
|-------------------------|------------|----------------------|-----------------------|-----------------------|----------------------|
|                         |            |                      | mol/m <sup>3</sup>    | mg/kg                 | µg/m <sup>3</sup>    |
| <b>Bulk compartment</b> |            |                      |                       |                       |                      |
| air                     | 99.99      | $24.36 \cdot 10^3$   | $4.77 \cdot 10^{-10}$ | $2.23 \cdot 10^{-5}$  | $2.68 \cdot 10^{-2}$ |
| surface water           | 0.00       | $7.95 \cdot 10^{-2}$ | $1.94 \cdot 10^{-11}$ | $1.09 \cdot 10^{-9}$  | $1.09 \cdot 10^{-3}$ |
| soil                    | 0.01       | 2.38                 | $3.18 \cdot 10^{-10}$ | $1.19 \cdot 10^{-8}$  | $1.78 \cdot 10^{-2}$ |
| bottom sediment         | 0.00       | $3.58 \cdot 10^{-3}$ | $1.05 \cdot 10^{-10}$ | $4.15 \cdot 10^{-9}$  | $5.89 \cdot 10^{-3}$ |
| groundwater             | 0.00       | 0.88                 | $1.91 \cdot 10^{-11}$ | $1.07 \cdot 10^{-9}$  | $1.07 \cdot 10^{-3}$ |
| coastal water           | 0.00       | $6.62 \cdot 10^{-4}$ | $8.36 \cdot 10^{-15}$ | $4.69 \cdot 10^{-13}$ | $4.69 \cdot 10^{-7}$ |
| <b>Subcompartment</b>   |            |                      |                       |                       |                      |
| pure air                | 99.99      | $24.36 \cdot 10^3$   | $4.77 \cdot 10^{-10}$ | $2.23 \cdot 10^{-5}$  | $2.68 \cdot 10^{-2}$ |
| air particles           | 0.00       | $9.88 \cdot 10^{-6}$ | $9.68 \cdot 10^{-9}$  | $3.62 \cdot 10^{-7}$  | 0.54                 |
| pure water              | 0.00       | $7.95 \cdot 10^{-2}$ | $1.94 \cdot 10^{-11}$ | $1.09 \cdot 10^{-9}$  | $1.09 \cdot 10^{-3}$ |
| suspended sediment      | 0.00       | $3.89 \cdot 10^{-5}$ | $9.49 \cdot 10^{-10}$ | $3.55 \cdot 10^{-8}$  | $5.32 \cdot 10^{-2}$ |
| biota (fish)            | 0.00       | $1.06 \cdot 10^{-6}$ | $2.58 \cdot 10^{-10}$ | $1.45 \cdot 10^{-8}$  | $1.45 \cdot 10^{-2}$ |
| soil air                | 0.00       | 0.71                 | $4.77 \cdot 10^{-10}$ | $2.23 \cdot 10^{-5}$  | $2.68 \cdot 10^{-2}$ |
| soil water              | 0.00       | $4.29 \cdot 10^{-2}$ | $1.91 \cdot 10^{-11}$ | $1.07 \cdot 10^{-9}$  | $1.07 \cdot 10^{-3}$ |
| pure soil               | 0.01       | 1.62                 | $4.33 \cdot 10^{-10}$ | $1.01 \cdot 10^{-8}$  | $2.43 \cdot 10^{-2}$ |
| sediment pore water     | 0.00       | $4.64 \cdot 10^{-4}$ | $1.94 \cdot 10^{-11}$ | $1.09 \cdot 10^{-9}$  | $1.09 \cdot 10^{-3}$ |
| pure sediment           | 0.00       | $3.11 \cdot 10^{-3}$ | $3.05 \cdot 10^{-10}$ | $7.12 \cdot 10^{-9}$  | $1.71 \cdot 10^{-2}$ |

Table XI. D values and transfer rates.

| Compartment   | Process        | Individual D (mol/h Pa)           | Transfer rate (mol/h) |
|---|----------------|-----------------------------------|-----------------------|
| (1)-(2)   | diffusion      | $D_V = 55.483 \cdot 10^1$         | $6.50 \cdot 10^{-4}$  |
|   | rain           | $D_{QW} = 0.760$                  | $8.90 \cdot 10^{-7}$  |
|   | wet deposition | $D_{DW} = 1.543 \cdot 10^{-3}$    | $1.81 \cdot 10^{-9}$  |
|   | dry deposition | $D_{PW} = 2.028 \cdot 10^{-3}$    | $2.37 \cdot 10^{-9}$  |
| (2)-(1)   | diffusion      | $D_V = 55.483 \cdot 10^1$         | $6.60 \cdot 10^{-4}$  |
| (1)-(3)   | diffusion      | $D_S = 30.171 \cdot 10^4$         | 0.35                  |
|   | rain           | $D_{QS} = 33.431$                 | $3.91 \cdot 10^{-5}$  |
|   | wet deposition | $D_{DS} = 6.784 \cdot 10^{-2}$    | $7.94 \cdot 10^{-8}$  |
|   | dry deposition | $D_{PS} = 8.915 \cdot 10^{-2}$    | $1.04 \cdot 10^{-7}$  |
| (3)-(1)   | diffusion      | $D_S = 30.171 \cdot 10^4$         | 0.35                  |
| (3)-(2)   | water runoff   | $D_{WW} = 13.372$                 | $1.57 \cdot 10^{-5}$  |
|   | soil runoff    | $D_{SW} = 0.422$                  | $4.94 \cdot 10^{-7}$  |
| (2)-(4)   | diffusion      | $D_Y = 8.774$                     | $1.04 \cdot 10^{-5}$  |
|   | deposition     | $D_{DX} = 4.136 \cdot 10^{-2}$    | $4.92 \cdot 10^{-8}$  |
| (4)-(2)   | diffusion      | $D_Y = 8.774$                     | $1.05 \cdot 10^{-5}$  |
|   | resuspension   | $D_{RX} = 3.309 \cdot 10^{-3}$    | $3.95 \cdot 10^{-9}$  |
| <b>Advective, pseudo-advective and reaction processes</b> |                |                                   |                       |
| advection   |                | $D_{A(1)} = 17.36 \cdot 10^7$     | $20.32 \cdot 10^1$    |
|   |                | $D_{A(2)} = 5.57$                 | $6.62 \cdot 10^{-6}$  |
| transfer to higher altitude                               |                | $D_{ST} = 21.38 \cdot 10^4$       | 0.25                  |
| leaching from soil  |                | $D_L = 13.37$                     | $1.57 \cdot 10^{-5}$  |
| advective flow from coastal water                         |                | $D_{CW} = 10.32 \cdot 10^3$       |                       |
| sediment burial   |                | $D_{BX} = 9.93 \cdot 10^{-3}$     | $1.18 \cdot 10^{-8}$  |
| reaction  |                | $D_{R(1)} = 36.0518 \cdot 10^8$   | $42.20 \cdot 10^2$    |
|   |                | $D_{R(2)} = 4.6317 \cdot 10^{-7}$ | $5.51 \cdot 10^{-13}$ |
|   |                | $D_{R(3)} = 1.4089 \cdot 10^{-5}$ | $1.65 \cdot 10^{-11}$ |
|   |                | $D_{R(4)} = 2.0772 \cdot 10^{-8}$ | $2.48 \cdot 10^{-14}$ |

Table XII. Summary of 4 principal compartment mass balance (mol/h).

| Compartment     | Emission           | Inflow | Reaction              | Outflow              | Net flux out to other compts. |
|-----------------|--------------------|--------|-----------------------|----------------------|-------------------------------|
| air             | $44.24 \cdot 10^2$ | 0.0    | $42.20 \cdot 10^2$    | $20.32 \cdot 10^1$   | $2.23 \cdot 10^{-5}$          |
| surface water   | 0.0                | 0.0    | $5.51 \cdot 10^{-13}$ | $6.62 \cdot 10^{-6}$ | $-6.62 \cdot 10^{-6}$         |
| soil            | 0.0                | 0.0    | $1.65 \cdot 10^{-11}$ | 0.0                  | $-1.56 \cdot 10^{-5}$         |
| bottom sediment | 0.0                | 0.0    | $2.48 \cdot 10^{-14}$ | 0.0                  | $-1.18 \cdot 10^{-8}$         |
| total           | $44.24 \cdot 10^2$ | 0.0    | $42.20 \cdot 10^2$    | $20.32 \cdot 10^1$   |                               |

From the simulation results of the environmental fate of isobutylene with CHEMFRANCE, it appears that this chemical is of minor concern regarding its potential hazards to the environment since it is rapidly degraded in air and it is not accumulated in other media.

In order to test the predictive power of CHEMFRANCE, we have tried to compare the results obtained with CHEMFRANCE to concentration data actually found in the environment. Only concentrations in air were retrieved from the literature. There are listed in Table XIII.

Table XIII. Recorded concentrations of isobutylene in air.

| Location, date  | Concentration                  |
|---|--------------------------------|
| Slottskogen Park (Sweden, 25/07/90) <sup>61</sup>   | 0.09 $\mu\text{g}/\text{m}^3$  |
| Avenue of Göteborg (Sweden, 25/07/90, 20°C) <sup>61</sup>   | 0.7 $\mu\text{g}/\text{m}^3$   |
| Avenue of Göteborg near intersection with traffic lights (Sweden, 30/11/89, 0°C) <sup>61</sup>    | 7.7 $\mu\text{g}/\text{m}^3$   |
| Avenue of Göteborg near intersection with traffic lights (Sweden, 7/05/90, 20°C) <sup>61</sup>    | 3.2 $\mu\text{g}/\text{m}^3$   |
| Bridge 5 m above motorway approach to Göteborg (Sweden, 17/11/89, 0°C) <sup>61</sup>              | 1.5 $\mu\text{g}/\text{m}^3$   |
| Motorway 10 km south of Göteborg (Sweden, 9/05/90, 10°C) <sup>61</sup>                            | 0.7 $\mu\text{g}/\text{m}^3$   |
| Inside parking garage near cars leaving in the morning (Sweden, 23/05/90, 15°C) <sup>61</sup>     | 7.5 $\mu\text{g}/\text{m}^3$   |
| Service station during refueling. No petrol vapor recovery (Sweden, 7/06/88, 22°C) <sup>62</sup>  | 1300 $\mu\text{g}/\text{m}^3$  |
| Service station during refueling. No petrol vapor recovery (Sweden, 22/06/88, 15°C) <sup>62</sup> | 700 $\mu\text{g}/\text{m}^3$   |
| Service station during refueling. Petrol vapor recovery (Sweden, 10/06/88, 21°C) <sup>62</sup>    | 10 $\mu\text{g}/\text{m}^3$    |
| Service station. Ambient air. Petrol vapor recovery (Sweden, 15/06/88, 22°C) <sup>62</sup>        | 9 $\mu\text{g}/\text{m}^3$     |
| Fos-Berre basin* (France, 17/09/87) <sup>63</sup>   | 0.058 $\mu\text{g}/\text{m}^3$ |

\* Located in the region n°8 of CHEMFRANCE.

Table XIII shows that if we consider locations relatively far from sources of emission, the concentrations recorded in air (Slottskogen park: 0.09  $\mu\text{g}/\text{m}^3$ ; Fos-Berre basin: 0.058  $\mu\text{g}/\text{m}^3$ ) are in accordance with our results (0.0268  $\mu\text{g}/\text{m}^3$  in Table X). However, note that the concentrations recorded near the emission sources are significantly higher (Table XIII).

From the simulations performed on isobutylene and those carried out on *o*-cresol and atrazine (to be published), it appears that CHEMFRANCE is particularly suitable to estimate the environmental fate of organic chemicals in France. However, it is obvious that to be validated, CHEMFRANCE has to be tested more thoroughly against other substances of widely differing physicochemical properties, reactivity and transport characteristics. The principal difficulties in accomplishing this task are the lack or the variability of emission data, environmental concentration data, reaction rate constants, and physicochemical properties.<sup>5, 64</sup>

CHEMFRANCE is basically a simulation tool for predicting the environmental fate behavior of new or existing chemicals for which there is insufficient environmental experience. To reach this goal, the predefined options of CHEMFRANCE easily allow to test different

scenarios. In addition, as CHEMFRANCE is totally open and flexible, the users can modify numerous parameters (e.g., compartment volumes and areas, densities, climatic parameters) in order to more easily interpret the different outputs or adapt CHEMFRANCE to local and global situations.

## ACKNOWLEDGEMENTS

This research which was initiated by A. Chancrogne in the frame of her MSc (1991) constitutes a part of the PhD of S. Bintein. It was sponsored by the European Community and by the French Ministry of the Environment. The authors wish to acknowledge the critical and constructive review by Dr. T. E. McKone (Health and Ecological Assessments Division, Lawrence Livermore National Laboratory, California). The elaboration of CHEMFRANCE would not have been possible without all the published and unpublished information provided by Pr. D. Mackay (Department of Chemical Engineering and Applied Chemistry, University of Toronto, Canada) on his fugacity models and more specifically on CHEMCAN which has constituted an invaluable methodological framework for our research. His comments on our manuscript were also helpful.

A first version of CHEMFRANCE was presented in the OECD Workshop on the Application of Simple Models for Environmental Exposure Assessment (Berlin, 11th-13th December 1991). As the version presented in this paper is radically different from that actually distributed via the different OECD Secretariats, it is preferable to contact CTIS to obtain this up-to-date version of CHEMFRANCE.

## REFERENCES

1. Cohen, Y. (1986). Organic pollutant transport. *Environ. Sci. Technol.* **20**, 538-544.
2. Domine, D., Devillers, J., Chastrette, M., and Karcher, W. (1992). Multivariate structure-property relationships (MSPR) of pesticides. *Pestic. Sci.* **35**, 73-82.
3. McKone, T.E. and Layton, D.W. (1986). Screening the potential risks of toxic substances using a multimedia compartment model: estimation of human exposure. *Regul. Toxicol. Pharmacol.* **6**, 359-380.
4. Mackay, D. (1991). *Multimedia Environmental Models. The Fugacity Approach*. Lewis Publishers, Chelsea, p. 257.
5. Mackay, D. and Paterson, S. (1991). Evaluating the multimedia fate of organic chemicals: a level III fugacity model. *Environ. Sci. Technol.* **25**, 427-436.
6. Mackay, D., Paterson, S., and Shiu, W.Y. (1992). Generic models for evaluating the regional fate of chemicals. *Chemosphere* **24**, 695-717.
7. Anonymous (1990). *La France et ses Régions*. INSEE.
8. Harrison, R.M. (1990). Chemistry of the troposphere. In: *Pollution: Causes, Effects, and Control* (R.M. Harrison, Ed.). Royal Society of Chemistry, Cambridge, pp. 157-179.

9. Williams, M.L. (1990). Atmospheric dispersal of pollutants and the modelling of air pollution. In: *Pollution: Causes, Effects, and Control* (R.M. Harrison, Ed.). Royal Society of Chemistry, Cambridge, pp. 201-219.
10. Anonymous. *Lacs, Etangs, Retenues - Carte de Situation - Notice Descriptive*. Ministère de la Qualité de la Vie, Secrétariat d'Etat à l'Environnement, p. 72.
11. Anonymous (1990). *Répertoire et Fichier Technique des Plans d'Eau - Doubs 25 - Présentation et Carte*. Agence de l'Eau Rhône Méditerranée Corse.
12. Anonymous (1990). *Répertoire et Fichier Technique des Plans d'Eau - Doubs 25 - Recueil des Données*. Agence de l'Eau Rhône Méditerranée Corse.
13. Anonymous (1991). *Répertoire et Fichier Technique des Plans d'Eau - Haute-Saône 70, Territoire de Belfort 90 - Présentation et Carte*. Agence de l'Eau Rhône Méditerranée Corse.
14. Anonymous (1991). *Répertoire et Fichier Technique des Plans d'Eau - Haute-Saône 70, Territoire de Belfort 90 - Recueil des Données*. Agence de l'Eau Rhône Méditerranée Corse.
15. Anonymous (1990). *Répertoire et Fichier Technique des Plans d'Eau - Jura 39 - Présentation et Carte*. Agence de l'Eau Rhône Méditerranée Corse.
16. Anonymous (1990). *Répertoire et Fichier Technique des Plans d'Eau - Jura 39 - Recueil des Données*. Agence de l'Eau Rhône Méditerranée Corse.
17. Anonymous (1988). *Plans d'Eau - De l'Autre Côté du Miroir*. Agence de Bassin Rhône Méditerranée Corse, Institut International de Gestion et de Génie de l'Environnement, p. 127.
18. Anonymous (1982). *Enquête Utilisation du Territoire. Année 1982. Répartition du Sol Départemental Selon les Utilisations Physiques*. Ministère de l'Agriculture - Service Central des Enquêtes et des Etudes Statistiques.
19. Anonymous (1974). *Analyse à l'Echelle de la France des Ressources en Eau et de leur Qualité ainsi que des Besoins Futurs et des Problèmes d'Approvisionnement à Moyen et Long Terme*. Arlab.
20. Loup, J. (1974). *Les Eaux Terrestres - Hydrologie Continentale*. Masson, Paris, p. 171.
21. OECD Environment Monographs n°69 (1993). *Report of the OECD Workshop on the Application of Simple Models for Environmental Exposure Assessment*, p. 143.
22. Carte pédologique de la France 1/1000000 (1966). INRA.
23. Chamayou, H. and Legros, J.P. (1989). *Les Bases Physiques, Chimiques et Minéralogiques de la Science du Sol*. Presse Universitaire de France, Paris, p. 593.
24. Duchaufour, P. (1970). *Précis de Pédologie*. Masson, Paris, p. 481.
25. Duchaufour, P. (1976). *Atlas Ecologique des Sols du Monde*. Masson, Paris, p. 178.
26. Anonymous (1970). *Atlas des Eaux Souterraines de la France*. Edition BRGM, Paris.
27. Anonymous (1979). *Bilan des Ressources en Eau Souterraine de la France*. Arlab, p. 269.
28. Anonymous (1986). *Synthèse Hydrogéologique de la Région Bourgogne Versant Méditerranéen avec Extension aux Régions Champagne-Ardenne et Lorraine pour le même Versant*. Bureau de Recherches Géologiques et Minières.
29. de Bellegarde, B. and Lafosse, J. (1977). La nappe du miocène du nord de la Drôme. In: *Les Eaux Souterraines et l'Approvisionnement en Eau de la France, Tome I*. National symposium, Nice, France, pp. 115-130.
30. Eisenlohr, B. and Lene, J.P. (1977). Utilisation des nappes alluviales côtières du département du Var. Exemples du fond de golfe de St Tropez et du bas Argens. In: *Les Eaux*

- Souterraines et l'Approvisionnement en Eau de la France, Tome I.* National symposium, Nice, France, pp. 267-279.
31. Erhard-Cassegrain, A. and Margat, J. (1983). *Introduction à l'Economie Générale de l'Eau.* Masson, Paris, p. 361.
32. Marchal, J.P., Paloc, H., and Viala, J.P. (1977). Ressources en eau souterraine et état actuel de leur exploitation dans le département des Pyrénées Orientales. In: *Les Eaux Souterraines et l'Approvisionnement en Eau de la France, Tome II.* National symposium, Nice, France, pp. 365-370.
33. Margat, J. (1987). Quand les eaux nous sont comptées. L'eau et les comptes du patrimoine naturel. *Annales des Mines Oct.*, 84-86.
34. Megnien, C. (1977). Les ressources en eau souterraine du centre du bassin de Paris. In: *Les Eaux Souterraines et l'Approvisionnement en Eau de la France, Tome II.* National symposium, Nice, France, pp. 447-455.
35. Ramon, S. (1977). Bilan d'exploitation des eaux souterraines dans le bassin Rhin-Meuse. In: *Les Eaux Souterraines et l'Approvisionnement en Eau de la France, Tome II.* National symposium, Nice, France, pp. 561-568.
36. Lewis, G.N. (1901). The law of physico-chemical change. *Proc. Am. Acad. Sci.* **37**, 49.
37. Mackay, D. and Diamond, M. (1989). Application of the QWASI (Quantitative Water Air Sediment Interaction) fugacity model to the dynamics of organic and inorganic chemicals in lakes. *Chemosphere* **18**, 1343-1365.
38. Mackay, D. and Paterson, S. (1981). Calculating fugacity. *Environ. Sci. Technol.* **15**, 1006-1014.
39. Mackay, D. and Paterson, S. (1982). Fugacity revisited. *Environ. Sci. Technol.* **16**, 654-660.
40. Mackay, D. and Paterson, S. (1990). Development of a fugacity model for assessing chemical fate in Canada. Report prepared for Health and Welfare Canada.
41. Mackay, D. and Paterson, S. (1990). Fugacity models. In: *Practical Applications of Quantitative Structure-Activity Relationships (QSAR) in Environmental Chemistry and Toxicology* (W. Karcher and J. Devillers, Eds.). Kluwer Academic Publishers, Dordrecht, pp. 433-460.
42. Mackay, D., Paterson, S., Cheung, B., and Neely, W.B. (1985). Evaluating the environmental behavior of chemicals with a level III fugacity model. *Chemosphere* **14**, 335-374.
43. Mackay, D., Paterson, S., and Joy, M. (1983). Application of fugacity models to the estimation of chemical distribution and persistence in the environment. In: *Fate of Chemicals in the Environment. Compartmental and Multimedia Models for Predictions* (R.L. Swann and A. Eschenroeder, Eds.). *ACS Symp. Ser.* **225**, 175-196.
44. Paterson, S. and Mackay, D. (1985). The fugacity concept in environmental modelling. In: *The Handbook of Environmental Chemistry, Vol. 2/Part C* (O. Hutzinger, Ed.). Springer-Verlag, Heidelberg, pp. 121-140.
45. Paterson, S. and Mackay, D. (1989). A model illustrating the environmental fate, exposure and human uptake of persistent organic chemicals. *Ecol. Model.* **47**, 85-114.

46. Mackay, D. and Paterson, S. (1983). Fugacity models of indoor exposure to volatile chemicals. *Chemosphere* **12**, 143-154.
47. Bintein, S., Devillers, J., and Karcher, W. (1993). Nonlinear dependence of fish bioconcentration on *n*-octanol/water partition coefficient. *SAR QSAR Environ. Res.* **1**, 29-39.
48. Bintein, S. and Devillers, J. (1994). QSAR for organic chemical sorption in soils and sediments. *Chemosphere* **28**, 1171-1188.
49. Arlery, R. (1979). *Le Climat de la France*. Ministère des transports, Direction de la Météorologie, Paris, p. 131.
50. Kessler, J. and Chambraud, A. (1990). *Météo de la France*. J.C. Lattès, Paris, p. 312.
51. *Kirk-Othmer Encyclopedia of Chemical Technology, Fourth Edition, Vol. 4* (1992). John Wiley & Sons, New York, p. 1117.
52. Perrin, R. and Scharff, J.P. (1993). *Chimie Industrielle. Vols. 1 and 2*. Masson, Paris, p. 1136.
53. *Ullmann's Encyclopedia of Industrial Chemistry, Fifth Edition, Vol. A4* (1985). VCH, Weinheim, p. 584.
54. Lonneman, W.A., Seila, R.L., and Meeks, S.A. (1986). Non-methane organic composition in the Lincoln tunnel. *Environ. Sci. Technol.* **20**, 790-796.
55. Russell, A.G. and Cass, G.R. (1986). Verification of a mathematical model for aerosol nitrate and nitric acid formation and its use for control measure evaluation. *Atmos. Environ.* **20**, 2011-2025.
56. Harley, R.A., Hannigan, M.P., and Cass, G.R. (1992). Respeciation of organic gas emissions and the detection of excess unburned gasoline in the atmosphere. *Environ. Sci. Technol.* **26**, 2395-2408.
57. Kleindienst, T.E., Smith, D.F., Hudgens, E.E., Snow, R.F., Perry, E., Claxton, L.D., Bufalini, J.J., Black, F.M., and Cupitt, L.T. (1992). The photo-oxidation of automobile emissions: measurements of the transformation products and their mutagenic activity. *Atmos. Environ.* **26A**, 3039-3053.
58. McAuliffe, C. (1966). Solubility in water of paraffin, cycloparaffin, olefin, acetylene, cycloolefin, and aromatic hydrocarbons. *J. Phys. Chem.* **70**, 1267-1275.
59. Bintein, S., Domine, D., and Devillers, J. (1994). Modeling the environmental fate of isobutylene and *o*-cresol in France with CHEMFRANCE. Rapport Ministère de l'Environnement, p. 38.
60. Grosjean, D. (1989). Organic acids in Southern California air: ambient concentrations, mobile source emissions, in situ formation and removal processes. *Environ. Sci. Technol.* **23**, 1506-1514.
61. Löfgren, L. and Petersson, G. (1992). Butenes and butadiene in urban air. *Sci. Total Environ.* **116**, 195-201.
62. Berglund, P.M. and Petersson, G. (1990). Hazardous petrol hydrocarbons from refuelling with and without vapour recovery. *Sci. Total Environ.* **91**, 49-57.
63. Coddeville, P., Guillermo, R., Dechaux, J.C., and Zimmermann, V. (1992). Photochemical air pollution and Lagrangian modeling. Validation of the EKMA model at the photochemical smog observed in the Fos-Berre basin on September, 1987. *Pollut. Atmos. Jul./Sept.*, 50-65.

64. Devillers, J., Thioulouse, J., Domine, D., Chastrette, M., and Karcher, W. (1991). Multivariate analysis of the input and output data in the fugacity model level I. In: *Applied Multivariate Analysis in SAR and Environmental Studies* (J. Devillers and W. Karcher, Eds.). Kluwer Academic Publishers, Dordrecht, pp. 281-345.



# ARTICLE IV



## EVALUATING THE ENVIRONMENTAL FATE OF LINDANE IN FRANCE

S. Bintein and J. Devillers\*

CTIS, 21 rue de la Bannière, 69003 Lyon, France

(Received in Germany 31 January 1996; accepted 18 March 1996)

### ABSTRACT

Lindane, a highly persistent and lipophilic pesticide, is still used in large quantities. This chemical can be found with appreciable concentrations in biota, atmosphere, and in other environmental compartments in the adsorbed form. This information provided a basis for assessing the simulation performances of CHEMFRANCE, a regional level III fugacity model allowing to generate environmental behavior profile of organic chemicals in France. The comparison between the estimated environmental fate and field and laboratory observations suggests that this fugacity model can be used to determine the processes that control the environmental fate of lindane in France. CHEMFRANCE also provides accurate estimates of environmental compartment contaminations. Copyright © 1996 Elsevier Science Ltd

### INTRODUCTION

Lindane, the gamma isomer of hexachlorocyclohexane ( $\gamma$ -HCH) is an organochlorine insecticide. Like DDT, lindane was at first praised as almost a perfect insecticide, being relatively cheap to manufacture, strikingly effective against a wide range of insects and some animal ectoparasites and yet apparently safe to man and warm-blooded animals.<sup>1</sup> However, lindane is notoriously persistent in the environment and possesses a lipophilic character which strongly favors its bioaccumulation. In addition, its use has led to the indiscriminate killing of beneficial as well as harmful insects and, increasingly, to the development of insect resistance.<sup>2,3</sup> Despite restrictions and banishments in several countries, large amounts of lindane are still applied each year for crop protection.<sup>4</sup> Nowadays, the world and French consumptions of lindane are estimated to be 720 000 t/year<sup>4</sup> and 1 600 t/year,<sup>5</sup> respectively.

---

\* Author to whom all correspondence should be addressed.

The huge use of lindane and its long persistence pose serious environmental problems. Significant concentrations of  $\gamma$ -HCH have been found in surface waters, soils, plants, animals, and food.<sup>6-13</sup> Moreover, due to its vapor pressure,  $\gamma$ -HCH is released to the atmosphere through spray applications<sup>4</sup> and volatilization from contaminated surfaces and then disseminated to adjacent or more distant regions.<sup>14-18</sup> Due to its economic and environmental concerns, numerous publications dealing with the environmental fate and hazards of lindane are available. This information was used for deriving the environmental fate profile of this chemical and then comparing experimental results with those obtained with CHEMFRANCE, a regional level III fugacity model allowing to simulate the environmental distribution of organic chemicals in France.<sup>19</sup>

## LITERATURE DATA ANALYSIS

### PHYSICOCHEMICAL PROPERTIES

An analysis of the literature allowed the selection of the following values:

- Molecular weight: 290.85 g/mol,
- Solubility in water: 7.52 mg/l at 25°C,<sup>20</sup>
- Vapor pressure:  $4.1 \cdot 10^{-3}$  Pa at 25°C,<sup>20</sup>
- *n*-Octanol/water partition coefficient (log P): 3.85,<sup>21</sup>
- Melting point: 112.5°C.<sup>22</sup>

### ENVIRONMENTAL BEHAVIOR

#### Degradation

The structure of lindane constituted entirely of C-C, C-H, and C-Cl bonds induces a high chemical stability in normal environmental conditions. Its degradation products are principally volatile chemicals. In marine systems and soils, one of its main metabolites,  $\gamma$ -pentachlorocyclohex-1-ene ( $\gamma$ -PCCH), is readily transferred to the atmosphere.<sup>23-25</sup> However, there are some divergences in the literature on the part of  $\gamma$ -PCCH in the volatilization process of lindane. Samuel and Pillai<sup>24</sup> reported a percentage slightly superior to ten. Conversely, Cliath and Spencer<sup>26</sup> found that two thirds to three fourths of the volatilized material was lost as  $\gamma$ -PCCH from a calcareous soil treated with lindane. Another principal metabolic product is  $\gamma$ -3,4,5,6-tetrachlorocyclohex-1-ene ( $\gamma$ -TCCH) identified in treated soils<sup>24,27</sup> and in aquatic sediments.<sup>23</sup> Mineralization producing CO<sub>2</sub><sup>24,28</sup> and bioisomerization of  $\gamma$ -HCH to  $\alpha$ ,  $\beta$ , and  $\delta$ -HCH in soils<sup>29</sup> are other processes of degradation of lindane. Some aquatic microorganisms can also convert lindane to  $\alpha$ -HCH.<sup>23,25</sup>

The degradation profile of lindane can be depicted by the following reaction half-lives:

- Half-life in soil: 2888 h (mean value calculated from data of Isnard and Lambert<sup>5</sup>),
- Half-life in sediment: 1188 h (mean value calculated from data of Isnard and Lambert<sup>5</sup>),
- Half-life in water: 1848 h (mean value calculated from data of Isnard and Lambert<sup>5</sup>),
- Half-life in air: 17000 h (in temperate climatic zones).<sup>30</sup>

Levels of contamination in the different compartments of the environment

#### *Soil contamination*

The soil concentrations in lindane depend on the method of application (e.g.; spray application, seed treatment), soil composition, and climatic conditions.<sup>24</sup> Five to six months after its soil application, the amount of lindane present in the top surface soil was estimated to be around 25% of the initial amount.<sup>29</sup> Another field study carried out at Delhi (India) revealed that the lindane concentration dropped to ~15% of the initial concentration one year after treatment.<sup>31</sup> Studies of the partitioning of  $\gamma$ -HCH in soils revealed low contamination of soil water.<sup>24,28</sup> The disappearance of lindane in soils depends principally on volatilization, plant adsorption, and degradation.<sup>24,28,29</sup>

#### *Surface water contamination*

Lindane has been detected at relatively low concentrations (i.e.; generally  $\ll 10^3$  ng/l) in surface waters in numerous countries.<sup>7,12,13,17,32,33</sup> In accordance with its low solubility, significant sorption in sediment and suspended sediment has been demonstrated.<sup>12,13,32</sup> Investigations on the lindane contamination of estuarine and sea waters in Europe showed lower contaminations than those reported in freshwater. In French estuaries, concentrations between 5 and 50 ng/l have been reported.<sup>7</sup> In the Scheldt estuaries located in southwest Netherlands and northwest Belgium, lindane was detected in June 1986 with dissolved concentrations ranging from 17 to 125 ng/l (median concentration of 82 ng/l) and with particulate concentrations ranging from 0.3 to 6.0 ng/g (median concentration of 2.5 ng/g).<sup>34</sup> A survey of the lindane contamination in the Baltic Sea and the North Sea reported concentrations in water samples varying between 1.0 to 7.7 ng/l.<sup>25</sup>

#### *Groundwater contamination*

Despite the persistent character of lindane, the risk of groundwater contamination is reduced due to its significant sorption in soils.<sup>31</sup> In U.S. surveys concerning groundwater contamination, lindane was seldom detected.<sup>22</sup> In France, between 1977 and 1990, lindane was detected in 11.6% of the 1314 groundwater samples from Calvados department with concentrations greater than 0.02  $\mu\text{g/l}$ , but only in 1% with concentrations above 0.05  $\mu\text{g/l}$  and 0.3% in concentrations superior to 0.1  $\mu\text{g/l}$ .<sup>35</sup>

### *Atmospheric contamination*

Numerous studies have reported the presence of lindane in air and rain samples. The presence of lindane in the atmospheric environment can be explained by spray droplet drift-off during its application and its volatilization from sprayed plants and soils.<sup>24,26,28</sup> Volatilization of incorporated lindane in soil contributes to a smaller extent to the atmospheric contamination.<sup>4</sup> The mean atmospheric concentration of lindane in Paris (France) was  $1.5 \cdot 10^{-3}$  ng/l from July 1989 to July 1990.<sup>18</sup> In Paris and adjacent areas, mean concentrations in rain samples in the range of 100-150 ng/l were recorded during 1988.<sup>36</sup> In Bavaria (Germany), rainwater samples had average concentrations between 11 and 17 ng/l in suburban stations and between 12 and 22 ng/l in rural stations in winter and summer, respectively.<sup>15</sup> In Jacksonville and Springfield (U.S.), mean air exposure estimates were 224 and 38 ng/day, respectively.<sup>37</sup>  $\gamma$ -HCH was detected in river and estuarine air from eastern and southern Asia and Oceania at concentrations ranging from 5.5 ng/l to  $4.0 \cdot 10^6$  ng/l.<sup>13</sup> Moreover, the presence of lindane in atmosphere and precipitations from sites remote from industrial and human activities indicates that long-range atmospheric transport of this pesticide occurs.<sup>14,38</sup> For illustrative purposes, at Enewetak Atoll in the North Pacific Ocean, lindane was found in concentrations ranging from  $0.006 \cdot 10^{-3}$  to  $0.021 \cdot 10^{-3}$  ng/l and from 0.34 to 1.59 ng/l in the atmosphere and bulk precipitation, respectively.<sup>14</sup>

### *Food and biota contamination*

The air and dietary exposures of lindane were considered to be the dominant public exposure pathways.<sup>22,37,39</sup> Due to its high persistence and lipophilic character, lindane enters into the food chain and accumulates in animals and vegetals. Numerous studies reported high levels of  $\gamma$ -HCH in plant biomass,<sup>9,10,40</sup> invertebrates,<sup>7,9,11,41</sup> vertebrates,<sup>11,12,41-44</sup> and food.<sup>8</sup> However, despite this contamination, neither the limit of maximum residues, nor the acceptable daily intake were exceeded in recent years in France.<sup>8,44</sup>

## MODELING STRATEGY

Besides the physicochemical properties and degradation half-life values listed in the previous section, information dealing with emissions was also necessary to run CHEMFRANCE.<sup>45</sup> In France, lindane is essentially used in corn crops.<sup>46</sup> The amounts of lindane applied on these crops in each of the 12 regions of CHEMFRANCE were estimated from the surface of corn crops treated (i.e.; 27% with lindane alone, 3.5% in association with chlorpyrifos), the area of this cultivation in the 12 regions, and the applied doses (i.e.; 1.35 kg of active matter/ha for lindane alone, 0.79 kg of active matter/ha when in association with chlorpyrifos).<sup>35</sup> With the above information, the quantities of lindane applied on corn crops in the 12 defined regions of CHEMFRANCE in 1990 could be estimated. However, in order to take into account the application of lindane on other tillages (e.g.; sugar-beet, colza, seed-treatment),

and as veterinary treatment, we assumed that these uses corresponded to 30% of the lindane applied on corn crops. These calculations yielded global releases summarized in Table I.

Table I. Estimated total emission rates of lindane for the regions of CHEMFRANCE.

| Region                       | Emission (kg/y) |
|------------------------------|-----------------|
| 1 (Bretagne, Pays de Loire)  | 355530          |
| 2 (Nord, Normandie)          | 224380          |
| 3 (Bassin de Paris)          | 135380          |
| 4 (Alsace, Lorraine)         | 116370          |
| 5 (Vallée du Rhône)          | 71530           |
| 6 (Massif central)           | 75340           |
| 7 (Alpes)                    | 49450           |
| 8 (Bassin méditerranéen)     | 11600           |
| 9 (Aquitaine, Midi-Pyrénées) | 161030          |
| 10 (Pyrénées)                | 57360           |
| 11 (Bordure atlantique)      | 240260          |
| 12 (Centre)                  | 147670          |
| France                       | 1645900         |

Spring was the season chosen to run CHEMFRANCE, since it is the period of year during which the agricultural applications of lindane occur.

The emission processes were simulated from four different scenarios.<sup>45</sup> The first considered a total emission of lindane into the soil compartment (scenario A), which can be related to a direct incorporation of lindane in soil from seed-treatment. Scenario B took into account accidental pollutions occurring in surface waters and the release via effluents at manufacturing sites (90% in soil and 10% in water). Due to the relative volatility of lindane, significant quantities may be released into the atmosphere through spray applications.<sup>4</sup> Therefore, a scenario C was designed to consider in addition to a soil emission (87%) a loss of 13%<sup>47</sup> into the air compartment. Last, scenario D summarized all the previous assumptions (i.e.; 77% in soil, 10% in surface water, 13% in air).

## SIMULATION RESULTS AND COMPARISON WITH LABORATORY AND FIELD OBSERVATIONS

Table II gives the average percentages corresponding to the removal processes obtained for the twelve defined regions of CHEMFRANCE. In all cases, lindane disappearance is principally governed by reaction in soil. Moreover, due to its long persistence time determined by CHEMFRANCE (i.e.; ~180 days), it can be concluded that the greater part of lindane applied on soil remains in this compartment, in which it biodegrades slowly. This is in accordance with experimental studies.<sup>28</sup> When only seed-treatment applications have been assumed (i.e.; scenarios A and B), the dissipation of lindane by advective processes for the air

compartment is low. However, in case of spray applications (i.e.; scenarios C and D), ~9% of the applied lindane is transported to adjacent regions. Furthermore, as lindane is relatively stable in air, the risk of contamination by atmospheric transports must not be neglected. To a lower degree, the advective process in water must be taken into account when emission of lindane in water occurs (i.e.; scenarios B and D). The above conclusions can be linked to the presence of lindane with significant atmospheric concentrations (i.e.; ~10 to 100  $\mu\text{g}/\text{m}^3$ ) in non-treated areas (e.g.; Arctic and Antarctic zones).<sup>14,48,49</sup>

Field studies clearly demonstrated the low groundwater contamination by lindane.<sup>22</sup> Thus, leaching to groundwater represents only about 0.35% of the emissions (Table II). This percentage is close to the results obtained in a field experiment (i.e.; 1% within 15 days of lindane application<sup>27</sup>).

Table II. Average percentages of each removal process obtained from the four scenarios.

| Removal process             | Scenario A        | Scenario B        | Scenario C        | Scenario D |
|-----------------------------|-------------------|-------------------|-------------------|------------|
| air advection               | 0.05              | 0.31              | 8.79              | 9.04       |
| water advection             | 0.07              | 1.40              | 0.08              | 1.40       |
| reaction in air             | $2 \cdot 10^{-4}$ | $2 \cdot 10^{-3}$ | 0.04              | 0.04       |
| reaction in water           | 0.31              | 6.28              | 0.33              | 6.31       |
| reaction in soil            | 99.07             | 89.29             | 90.28             | 80.50      |
| reaction in sediment        | 0.11              | 2.35              | 0.12              | 2.36       |
| transfer to higher altitude | $6 \cdot 10^{-5}$ | $4 \cdot 10^{-4}$ | 0.01              | 0.01       |
| leaching to groundwater     | 0.39              | 0.35              | 0.35              | 0.32       |
| sediment burial             | $7 \cdot 10^{-4}$ | 0.02              | $8 \cdot 10^{-4}$ | 0.02       |

Granier<sup>36</sup> reported a mean runoff rate equal to  $48 \mu\text{g}/\text{m}^2/\text{y}$  in the region of Paris. This rate is ten-fold superior to those calculated by CHEMFRANCE (i.e.; 4.9 to  $6.1 \mu\text{g}/\text{m}^2/\text{year}$ ). Nevertheless, the calculated percentages of loss by runoff of the applied lindane (i.e.; 0.23 to 0.28%) are in accordance with the mean percentage (i.e.; <0.2%) obtained in a field survey.<sup>46</sup>

In scenarios C and D, more than 20% of the atmospheric emissions of lindane are deposited by rain and in lower proportions by wet and dry depositions (93%, 4%, and 3% of the total atmospheric fallout, respectively). These results agree with the percentages observed during a study performed near Paris in 1988.<sup>36</sup> Indeed, Granier<sup>36</sup> found 94%, 5%, and 1% for rain, wet, and dry depositions, respectively. In the same study, a mean flux of total atmospheric fallout equal to  $77 \mu\text{g}/\text{m}^2/\text{y}$  was reported. This value is similar to those obtained in scenarios C and D (i.e.; 65 and  $67 \mu\text{g}/\text{m}^2/\text{y}$ , respectively). In the same way, Granier<sup>36</sup> observed a dry deposition rate of  $0.92 \mu\text{g}/\text{m}^2/\text{y}$ . This is in agreement with our results derived from scenarios C and D (i.e.; 2.01 and  $2.08 \mu\text{g}/\text{m}^2/\text{y}$ , respectively).

There are some discrepancies in the literature on the part of the volatilization process in the environmental fate of lindane. Firstly, the percentages of  $\gamma$ -HCH detected in the volatilized organics in field studies varied between 0% (i.e.; not detected) and >85%.<sup>24,28</sup> This large variation can be attributed to the fact that it is difficult to know whether it is the lindane itself

that volatilized or its degradation products which possess higher vapor pressures. Secondly, the variations of experimental conditions (e.g.; temperature) induce large fluctuations in the percentage of volatilized lindane.<sup>24,28,50,51</sup> In this context, it is very difficult to compare the transfer of lindane in the air compartment obtained with CHEMFRANCE (i.e.; ~0.1% of the applied lindane) to field and laboratory observations.

The transfer rates of lindane between the sediment and water compartments are dominated by the sorption process and by sediment deposition. This leads to the accumulation of lindane in the bottom sediment. The other transfer rates (i.e.; sediment resuspension, diffusion from air to soil and water) between the four principal compartments (i.e.; air, surface water, soil, bottom sediment) are negligible in all the scenarios.

Figure 1 summarizes the concentrations of lindane found in the six bulk compartments for the different regions of CHEMFRANCE. Observed concentrations in surface water, groundwater, air, rain, and fish as well as those derived from the four different scenarios are reported in Tables III and IV.

The surface water concentrations obtained in scenarios A and C are generally in accordance with the maximum actual concentrations (Table III). However, the results of scenarios B and D greatly exceed field observations. This discrepancy could be due to an overestimation of the percentage of lindane emitted in the water, which could be closer to 1% than to 10% of the total emission of lindane.

The groundwater concentrations measured in regions 9 and 10 are superior to the simulated concentrations (Table III). Nevertheless, a study performed between 1977 and 1990 in the department of Calvados (region 2) revealed that from 1314 groundwater samples, only 1.3% presented a concentration larger than 50 ng/l.<sup>35</sup> This maximum is in accordance with our results for this region (i.e.; 54-67 ng/l).

Very few data are available on lindane contamination in French coastal and sea waters. Marchand<sup>7</sup> reported concentrations in coastal waters ranging between <1 to 5 ng/l, which are slightly superior to the maximum concentration obtained with CHEMFRANCE (i.e.; 0.85 ng/l).

The atmospheric concentrations derived from scenarios C and D assuming emissions in the air compartment are relatively near to those observed in the region of Paris (Table III).

By means of an equation published by Mackay and coworkers,<sup>20</sup> the concentrations in rain for region 3 were calculated from the air concentrations obtained with CHEMFRANCE. For scenarios A and B, the calculated rain concentrations are lower than those observed in Paris.<sup>36</sup> Conversely, scenarios C and D generate values close to the actual data (Table III).

The fish concentrations obtained in scenarios A and C are in accordance with the actual concentrations (Table IV). However, the results obtained with scenarios B and D greatly exceed field observations (Table IV). This overestimation confirms the assumption that the percentage of lindane emitted in the water compartment is probably closer to 1%.

The concentration of residues of lindane detected in a soil historically treated<sup>29</sup> is in accordance with the soil concentrations derived by CHEMFRANCE (i.e.;  $\sim 6 \cdot 10^{-3}$  mg/kg).






Figure 1. Lindane concentrations (ng/l) for the bulk compartments calculated with CHEMFRANCE for the twelve defined regions and the four scenarios (see text).

Table III. Comparison between field and predicted (scenarios A-D) concentrations (ng/l).

| Date, region of CHEMFRANCE                       | Actual concentration           |           | Reference | Predicted concentration |                       |                       |                       |
|--|--------------------------------|-----------|-----------|-------------------------|-----------------------|-----------------------|-----------------------|
|  | mean&                          | maximum   |           | A                       | B                     | C                     | D                     |
| <b>Surface water compartment</b>                 |                                |           |           |                         |                       |                       |                       |
| 7 rivers (05/1980 to 07/1981), region 1          | 78.0 (n = 141)                 | 450       | 35        | 390.1                   | 7200                  | 387.3                 | 7197                  |
| 4 rivers (17/04/1990 to 24/07/1990), region 1    | 95.2 (n = 39)                  | 400       | 35        | 390.1                   | 7200                  | 387.3                 | 7197                  |
| 4 rivers (17/04/1991 to 24/07/1991), region 1    | 56.1 (n = 27)                  | 400       | 35        | 390.1                   | 7200                  | 387.3                 | 7197                  |
| 12 rivers (04/1991 to 11/1991), region 1         | 16.6 (n = 75)                  | 70        | 35        | 390.1                   | 7200                  | 387.3                 | 7197                  |
| 1 river (17/04/1990 to 24/07/1990), region 2     | 48 (n = 10)                    | 103       | 35        | 270.8                   | 4999                  | 260.7                 | 4989                  |
| 1 river (17/04/1991 to 24/07/1991), region 2     | 41.3 (n = 7)                   | 153       | 35        | 270.8                   | 4999                  | 260.7                 | 4989                  |
| 2 rivers (04/1991 to 11/1991), region 2          | 73.8 (n = 8)                   | 200       | 35        | 270.8                   | 4999                  | 260.7                 | 4989                  |
| Yonne river (01/1988 to 03/1991), region 3       | 22 (n = 3)                     | 36        | 43        | 53.35                   | 1908                  | 56.48                 | 1911                  |
| Seine barrage (26/03/90 to 19/11/90), region 3   | 13.6 (n = 46)                  | 29        | 35        | 53.35                   | 1908                  | 56.48                 | 1911                  |
| 29 rivers (summer 1990), region 9                | 37.2 (n = 46)                  | 550       | 35        | 137.0                   | 3707                  | 143.6                 | 3713                  |
| 17 rivers (summer 1990), region 11               | 63.2 (n = 24)                  | 550       | 35        | 127.3                   | 3444                  | 142.5                 | 3459                  |
| <b>Groundwater compartment</b>                   |                                |           |           |                         |                       |                       |                       |
| (05/1978 to 12/1987), region 1                   | 9.5 (n = 16)                   | 31        | 35        | 122.0                   | 110.0                 | 110.5                 | 98.46                 |
| (summer 1990), region 9                          | 183.8 (n = 14)                 | 1125      | 35        | 28.60                   | 25.78                 | 26.07                 | 23.25                 |
| (summer 1990), region 10                         | 92 (n = 6)                     | 213       | 35        | 38.77                   | 34.97                 | 35.85                 | 32.05                 |
| <b>Air compartment</b>                           |                                |           |           |                         |                       |                       |                       |
| air (05/1988 to 07/1989), Creteil, region 3      | 2.1 10 <sup>-3</sup> *         | not given | 18        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (05/1988 to 07/1989), Paris, region 3        | 1.7 10 <sup>-3</sup> *         | not given | 18        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (1986), Paris, region 3                      | 1.4 10 <sup>-3</sup> \$        | not given | 16        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (1987), Paris, region 3                      | 1.4 10 <sup>-3</sup> \$        | not given | 16        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (07/1989 to 07/1990), Paris, region 3        | 1.5 10 <sup>-3</sup> \$        | not given | 18        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (1986 to 1990), in spring at Paris, region 3 | 2.7 10 <sup>-3</sup> £ (n = 7) | not given | 18        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (1986 to 1990), in summer at Paris, region 3 | 1.4 10 <sup>-3</sup> £ (n = 6) | not given | 18        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (1986 to 1990), in fall at Paris, region 3   | 1.3 10 <sup>-3</sup> £ (n = 7) | not given | 18        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| air (1986 to 1990), in winter at Paris, region 3 | 0.7 10 <sup>-3</sup> £ (n = 5) | not given | 18        | 8.96 10 <sup>-6</sup>   | 97.1 10 <sup>-6</sup> | 2.82 10 <sup>-3</sup> | 2.91 10 <sup>-3</sup> |
| rain (1988), region of Paris, region 3           | 130 (n = 6)                    | 150       | 36        | 0.34                    | 3.71                  | 107.7                 | 111.2                 |
| rain (1986), region of Paris, region 3           | 39 (n = ?)                     | 90        | 36        | 0.34                    | 3.71                  | 107.7                 | 111.2                 |
| rain (1986), Paris, region 3                     | 29 (n = ?)                     | 40        | 36        | 0.34                    | 3.71                  | 107.7                 | 111.2                 |
| rain (1986), region of Paris, region 3           | 28 (n = ?)                     | 70        | 36        | 0.34                    | 3.71                  | 107.7                 | 111.2                 |

& n = number of values; \* mean monthly concentration; \$ mean annual concentration; £ mean seasonal concentration

Table IV. Comparison between field and predicted (scenarios A-D) fish concentrations in region 3 ( $\mu\text{g}/\text{kg}$  wet weight).

| Species, date, location                  | Actual concentration |           | Ref. | Predicted concentration |     |      |     |
|--|----------------------|-----------|------|-------------------------|-----|------|-----|
|  | mean&                | maximum   |      | A                       | B   | C    | D   |
| Roach, (04/1991 to 05/1991), Seine river | 29.25 (n = 7)        | 97.5      | 42   | 26.2                    | 940 | 27.8 | 940 |
| Roach, (04/1991 to 05/1991), Yonne river | 11.25 (n = ?)        | not given | 42   | 26.2                    | 940 | 27.8 | 940 |
| Roach, (04/1991 to 05/1991), Marne river | 12.25 (n = ?)        | not given | 42   | 26.2                    | 940 | 27.8 | 940 |
| Roach, (10/1991), Seine river            | 13.5 (n = ?)         | not given | 43   | 26.2                    | 940 | 27.8 | 940 |
| Perch, (10/1991), Seine river            | 15.0 (n = ?)         | not given | 43   | 26.2                    | 940 | 27.8 | 940 |

& n = number of values

The field bioconcentration factors of lindane in fish<sup>46</sup> ranging between 435 and 758 are on the same order of magnitude as that derived by the QSAR equation<sup>52</sup> (i.e.; ~500) included in CHEMFRANCE.

In the air compartment, lindane is principally present in pure air (i.e.; 99.3%), and only a low quantity is bound to the air particles (i.e.; 0.7%). This percentage of lindane associated with air particles is in agreement with field measurements in the Lake Ontario area indicating low particulate-associated fractions of lindane (i.e.; generally <1%).<sup>53</sup>

The great tendency of lindane to be sorbed<sup>31</sup> explains the high percentage (i.e.; >99%) of lindane bound to solid components in the soil and sediment compartments of CHEMFRANCE. This sorption capacity also induces an accumulation in the suspended sediments,<sup>32</sup> as depicted by the sorption coefficient of lindane in this subcompartment (i.e.; ~2000) obtained by the QSAR model<sup>54</sup> used in CHEMFRANCE.

## CONCLUSION

The literature data analysis on the environmental fate and hazards of lindane reveals its high potential for accumulation in biota and its capacity to be bound to the solid components of abiotic media. In addition to the risk of human contamination from food, inhalation exposure cannot be neglected due to potential atmospheric contamination through volatilization and spray applications. Surface waters, as well as the marine compartment, are only weakly contaminated by lindane. Indeed, runoff, water advection, and atmospheric deposition processes are not considered as major fate pathways and therefore lead to relatively low amounts of lindane being present in these compartments. The weak groundwater concentrations are linked to the slow leaching process. The similarity between the actual environmental fate profile of lindane and that obtained from CHEMFRANCE underlines the relevance of this fugacity model in simulating the environmental fate of this pesticide. From these results and those recently obtained,<sup>45</sup> it appears that CHEMFRANCE is a useful tool for estimating the environmental fate of organic chemicals in France. Information on the availability of this computer program can be obtained from the authors.

## ACKNOWLEDGMENTS

The authors are grateful to Dr. D.M. Kane (Health and Welfare, Ottawa, Canada) for his comments. This study was granted by the French Ministry of the Environment.

## REFERENCES

1. Worthing, C.R. (1987). *The Pesticide Manual, Eighth Edition*. The British Crop Protection Council, Thornton Heath, p. 1081.
2. Taylor, R.N. (1982). Insecticide resistance in houseflies from the Middle East and North Africa with notes on the use of various bioassay techniques. *Pestic. Sci.* **13**, 415-425.
3. Pap, L. and Farkas, R. (1994). Monitoring of resistance of insecticides in house fly (*Musca domestica*) populations in Hungary. *Pestic. Sci.* **40**, 245-258.
4. Voldner, E.C. and Li, Y.F. (1995). Global usage of selected persistent organochlorines. *Sci. Total Environ.* **160/161**, 201-210.
5. Isnard, P. and Lambert, S. (1987). Modélisation du devenir des pesticides dans l'environnement: application au lindane. *ANPP* **5**, 199-207.
6. Cole, R.H., Frederick, R.E., Healy, R.P., and Rolan, R.G. (1984). Preliminary findings of the priority pollutant monitoring project of the nationwide urban runoff program. *J. WPCF* **56**, 898-908.
7. Marchand, M. (1989). Les produits phytosanitaires agricoles et la qualité des eaux marines littorales. *T.S.M. L'Eau* **10**, 511-521.
8. Venant, A., Borrel, S., Mallet, J., and Gillon, E. (1989). Contamination in food by organochlorine compound residues from 1980 to 1987. Regulatory considerations. *Sci. Aliments* **9**, 473-489.
9. Hernandez, L.M., Fernandez, M.A., and Gonzalez, M.J. (1991). Lindane pollution near an industrial source in northeast Spain. *Bull. Environ. Contam. Toxicol.* **46**, 9-13.
10. Granier, L. and Chevreuil, M. (1992). On the use of tree leaves as bioindicators of the contamination of air by organochlorines in France. *Wat. Air Soil Pollut.* **64**, 575-584.
11. Nair, A. and Pillai, M.K.K. (1992). Trends in ambient levels of DDT and HCH residues in humans and the environment of Delhi, India. *Sci. Total Environ.* **121**, 145-157.
12. Kucklick, J.R., Bidleman, T.F., McConnell, L.L., Walla, M.D., and Ivanov, G.P. (1994). Organochlorines in the water and biota of lake Baikal, Siberia. *Environ. Sci. Technol.* **28**, 31-37.
13. Iwata, H., Tanabe, S., Sakai, N., Nishimura, A., and Tatsukawa, R. (1994). Geographical distribution of persistent organochlorines in air, water and sediments from Asia and Oceania, and their implications for global redistribution from lower latitudes. *Environ. Pollut.* **85**, 15-33.
14. Atlas, E. and Giam, C.S. (1981). Global transport of organic pollutants: ambient concentrations in the remote marine atmosphere. *Science* **211**, 163-165.

15. Thomas, W. (1986). Principal component analysis of trace substance concentrations in rainwater samples. *Atmos. Environ.* **20**, 995-1000.
16. Chevreuil, M., Chesterikoff, A., Létolle, R., and Granier, L. (1989). Atmospheric pollution and fallout by PCBS and organochlorine pesticides (Ile-de-France). *Wat. Air Soil Pollut.* **43**, 73-83.
17. Gomme, J.W., Shurvell, S., Hennings, S.M., and Clark, L. (1991). Hydrology of pesticides in a chalk catchment: surface waters. *J. IWEM* **5**, 546-552.
18. Granier, L. and Chevreuil, M. (1992). Changes in organochlorine levels in the Paris atmosphere: 1986-1990. *C. R. Acad. Sci. Paris* **314**, 1457-1462.
19. Devillers, J., Bintein, S., and Karcher, W. (1995). CHEMFRANCE: a regional level III fugacity model applied to France. *Chemosphere* **30**, 457-476.
20. Mackay, D., Paterson, S., and Schroeder, W.H. (1986). Model describing the rates of transfer processes of organic chemicals between atmosphere and water. *Environ. Sci. Technol.* **20**, 810-816.
21. Isnard, P. and Lambert, S. (1989). Aqueous solubility and *n*-octanol/water partition coefficient correlations. *Chemosphere* **18**, 1837-1855.
22. Howard, P.H. (1991). *Handbook of Environmental Fate and Exposure Data for Organic Chemicals, Vol. III*. Lewis Publishers, Chelsea, p. 684.
23. Benezet, J.H. and Matsumara, F. (1973). Isomerization of  $\gamma$ -BHC to  $\alpha$ -BHC in the environment. *Nature* **243**, 480-481.
24. Samuel, T. and Pillai, M.K.K. (1990). Effect of temperature and sunlight exposure on the fate of soil-applied [ $^{14}$ C]-gamma-hexachlorocyclohexane. *Arch. Environ. Contam. Toxicol.* **19**, 214-220.
25. Hühnerfuss, H., Faller, J., König, W.A., and Ludwig, P. (1992). Gas chromatographic separation of the enantiomers of marine pollutants. 4. Fate of hexachlorocyclohexane isomers in the Baltic and North Sea. *Environ. Sci. Technol.* **26**, 2127-2133.
26. Cliath, M.M. and Spencer, W.F. (1972). Dissipation of pesticides from soil by volatilization of degradation products. I. Lindane and DDT. *Environ. Sci. Technol.* **6**, 910-914.
27. Adhya, T.K., Rao, V.R., Dani, R.C., Panda, D., and Sethunathan, N. (1995). Leaching of hexachlorocyclohexane isomers in a flooded field unplanted or planted to rice. *Wat. Air Soil Pollut.* **81**, 305-313.
28. Drego, J., Murthy, N.B.K., and Raghu, K. (1990). [ $^{14}$ C]- $\gamma$ -hexachlorocyclohexane in a flooded soil with green manuring. *J. Agric. Food Chem.* **38**, 266-268.
29. Waliszewski, S.M. (1993). Residues of lindane, HCH isomers and HCB in the soil after lindane application. *Environ. Pollut.* **82**, 289-293.
30. Wania, F. and Mackay, D. (1995). A global distribution model for persistent organic chemicals. *Sci. Total Environ.* **160/161**, 211-232.
31. Samuel, T., Agarwal, H.C., and Pillai, M.K.K. (1988). Persistence and binding of DDT and gamma-HCH in a sandy loam soil under field conditions in Delhi, India. *Pestic. Sci.* **22**, 1-15.

32. Nerin, C., Martinez, M., and Cacho, J. (1992). Distribution of hexachlorocyclohexane isomers from several sources. *Toxicol. Environ. Chem.* **35**, 125-135.
33. Nwankwoala, A.U. and Osibanjo, O. (1992). Baseline levels of selected organochlorine pesticide residues in surface waters in Ibadan (Nigeria) by electron capture gas chromatography. *Sci. Total Environ.* **119**, 179-190.
34. van Zoest, R. and van Eck, G.T.M. (1991). Occurrence and behaviour of several groups of organic micropollutants in the Scheldt estuary. *Sci. Total Environ.* **103**, 57-71.
35. Devillers, J., Bintein, S., and Domine, D. (1992). Lindane in France. Report CTIS #92012, p. 48.
36. Granier, L. (1991). *Apports Atmosphériques en Organochlorés à l'Hydrosphère. Comparaison avec les Métaux Lourds*. Thesis, University of Paris VI, p. 160.
37. Whitmore, R.W., Immerman, F.W., Camann, D.E., Bond, A.E., Lewis, R.G., and Schaum, J.L. (1994). Non-occupational exposures to pesticides for residents of two U.S. cities. *Arch. Environ. Contam. Toxicol.* **26**, 47-59.
38. Kurtz, D.A. and Atlas, E.L. (1990). Distribution of hexachlorocyclohexanes in the Pacific Ocean basin, air and water. In, *Long Range Transport of Pesticides* (D.A. Kurtz, Ed.). Lewis Publishers, Chelsea, pp. 143-160.
39. Kastenbergh, W.E. and Yeh, H.C. (1993). Assessing public exposure to pesticide-contaminated ground water. *Ground Wat.* **31**, 746-752.
40. Calamari, D., Bacci, E., Focardi, S., Gaggi, C., Morosini, M., and Vighi, M. (1991). Role of plant biomass in the global environmental partitioning of chlorinated hydrocarbons. *Environ. Sci. Technol.* **25**, 1489-1495.
41. Duursma, E.K., Nieuwenhuize, J., van Liere, J.M., and Hillebrand, M.T.J. (1986). Partitioning of organochlorines between water, particulate matter and some organisms in estuarine and marine systems of the Netherlands. *Neth. J. Sea Res.* **20**, 239-251.
42. Chevreuil, M., Carru, A.M., Chesterikoff, A., Boët, P., Tales, E., and Allardi, J. (1995). Contamination of fish from different areas of the river Seine (France) by organic (PCB and pesticides) and metallic (Cd, Cr, Cu, Fe, Mn, Pb and Zn) micropollutants. *Sci. Total Environ.* **162**, 31-42.
43. Chevreuil, M., Granier, L., and Carru, A.M. (1995). Relationship between biological parameters and bioaccumulation of some organochlorines (pesticides, PCB) by fishes in the river Seine (France). *Wat. Air Soil Pollut.* **81**, 107-120.
44. Klein, D., Dillon, J.C., Jirou-Najou, J.L., Gagey, M.J., and Debry, G. (1986). Cinétique d'élimination des composés organochlorés au cours de la première semaine d'allaitement maternel. *Fd Chem. Toxic.* **24**, 869-873.
45. Bintein, S. and Devillers, J. (1996). Evaluating the environmental fate of atrazine in France. *Chemosphere* **32**, 2441-2456.
46. Seux, R., Clément, M., Quinio, J., and Sauvaget, G. (1987). *Transfert du Lindane des Zones d'Épandage vers les Eaux Superficielles du Département de la Mayenne*. Ecole Nationale de la Santé Publique, Rennes, p. 42.

47. Haque, A. and Schuphan, I. (1989). Field - outdoor lysimeter - closed cultivation chamber: comparative studies with lindane after spray application on potato plants. *J. Plant Diseases Protection* **96**, 173-186.
48. Weber, R.R. and Montone, R.C. (1990). Distribution of organochlorines in the atmosphere of the south Atlantic and Antarctic Oceans. In, *Long Range Transport of Pesticides* (D.A. Kurtz, Ed.). Lewis Publisher, Chelsea, pp. 185-197.
49. Cotham, W.E. and Bidleman, T.F. (1991). Estimating the atmospheric deposition of organochlorine contaminants to the Arctic. *Chemosphere* **22**, 165-188.
50. Spencer, W.F., Cliath, M.M., Jury, W.A., and Zhang, L.Z. (1988). Volatilization of organic chemicals from soil as related to their Henry's law constant. *J. Environ. Qual.* **17**, 504-509.
51. Boehncke, A., Siebers, J., and Nolting, H.G. (1990). Investigations of the evaporation of selected pesticides from natural and model surfaces in field and laboratory. *Chemosphere* **21**, 1109-1124.
52. Bintein, S., Devillers, J., and Karcher, W. (1993). Nonlinear dependence of fish bioconcentration on *n*-octanol/water partition coefficient. *SAR QSAR Environ. Res.* **1**, 29-39.
53. Johnson, N.D., Lane, D.A., Schroeder, W.H., and Strachan, W.M. (1990). Measurements of selected organochlorine compounds in air near Ontario Lakes: gas-particles relationships. In, *Long Range Transport of Pesticides* (D.A. Kurtz, Ed.). Lewis Publishers, Chelsea, pp. 105-114.
54. Bintein, S. and Devillers, J. (1994). QSAR for organic chemical sorption in soils and sediments. *Chemosphere* **28**, 1171-1188.

# ARTICLE V





## EVALUATING THE ENVIRONMENTAL FATE OF ATRAZINE IN FRANCE

S. Bintein and J. Devillers\*

CTIS, 21 rue de la Bannière, 69003 Lyon, France

(Received in Germany 31 January 1996; accepted 18 March 1996)

### ABSTRACT

Atrazine is used in large quantities in U.S. and European countries as a weed-control agent. As a result, numerous data on its environmental fate and hazards have been published. Analysis of the literature shows that this herbicide can be found with appreciable concentrations in groundwaters, rivers, lakes, and estuaries. This contamination principally results from leaching and runoff processes. Atrazine can also pollute fog and rain due to its release into the atmosphere through spray applications. This large amount of information constitutes a very attractive basis for assessing the simulation performances of environmental fate models. In this context, CHEMFRANCE, a regional fugacity model level III which calculates the environmental distribution of organic chemicals in twelve defined regions of France has been used to estimate the environmental fate of atrazine. The calculated values are comparable with field and laboratory results. Therefore, CHEMFRANCE can be considered as a useful tool for simulating the environmental fate of this agrochemical. Copyright © 1996 Elsevier Science Ltd

### INTRODUCTION

Atrazine is a selective pre- and post-emergent weed-control agent for corn and millet cultivations. It is also used in asparagus, forestry, grassland, grass crops, sugarcane, sorghum, and pineapple.<sup>1</sup> In addition, atrazine is employed at higher rates as a nonselective herbicide for general weed control (e.g.; along roads and railroads).<sup>2</sup> The world consumption of atrazine is estimated to be ~70000 t/year (90% for corn cultivations).<sup>3</sup> In France, large amounts of atrazine are applied each year (e.g.; 5000 t in 1986 used in pre-emergence on  $3 \cdot 10^6$  ha of corn crops).<sup>4</sup>

-----  
\* Author to whom all correspondence should be addressed.

Due to the agricultural and non-agricultural applications, subsequent surface runoff,<sup>5-7</sup> and leaching to groundwater,<sup>8</sup> the environment is widely contaminated by this chemical. Atrazine causes considerable pollution in the groundwater and surface water in France and all over the world.<sup>9-14</sup> Furthermore, atrazine can also pollute fog and rain due to its release into the atmosphere through spray applications.<sup>10,14-18</sup> Due to its economic and environmental concerns, numerous publications dealing with the environmental fate and hazards of atrazine are available. This bibliography provides a comprehensive understanding of the processes governing the fate and hazards of this chemical.

In contrast, it is well recognized that for estimating the environmental fate of chemicals, models at different levels of complexity represent very attractive tools when experimental results are scarce. Among them, the level III fugacity models are particularly powerful for estimating the environmental distribution of pollutants in biotic and abiotic media.<sup>19</sup> In this context, CHEMFRANCE, a level III fugacity computer model was developed to simulate the environmental fate of organic chemicals in France.<sup>20</sup> The assessment is based on information in connection with environmental parameters, emission rates, and physicochemical and reactive properties of the organic chemicals. However, the main problem with these types of models is to determine if the simulation results are realistic. Under these conditions, the aim of this paper was first to review the large amount of bibliography available on atrazine in order to derive the environmental fate profile of this agrochemical and then to compare field and laboratory results with those obtained with CHEMFRANCE. This exercise is required in order to underline the usefulness and limitations of CHEMFRANCE in the estimation of the environmental behavior of the organic chemicals in France.

## LITERATURE DATA ANALYSIS

### *PHYSICOCHEMICAL PROPERTIES*

The estimation of the environmental distribution of a pesticide highly depends on its basic physicochemical properties.<sup>21</sup> For atrazine (2-chloro-4-ethylamino-6-isopropylamino-*s*-triazine, CAS RN 1912-24-9), the physicochemical data found in the literature are numerous and an appreciable variability is observed. Therefore, great care must be taken in their selection. Indeed, it was shown that the physicochemical properties of the molecules significantly influence the results obtained with the fugacity models.<sup>22,23</sup> An extensive bibliographical survey permitted the selection of the following physicochemical properties:

- Molecular weight: 215.69 g/mol,
- Solubility in water: 30 mg/l at 20°C,<sup>24</sup>
- Vapor pressure:  $4 \cdot 10^{-5}$  Pa at 20°C,<sup>24</sup>
- *n*-Octanol/water partition coefficient (log P): 2.34,<sup>24</sup>
- Melting point: 174°C,<sup>25</sup>
- pKa: 1.70.<sup>24</sup>

## ENVIRONMENTAL BEHAVIOR

### Degradation

In its pure state, atrazine is relatively stable and may be stored for a period of several years.<sup>2</sup> In the environment, its transformation begins immediately after application<sup>26</sup> and follows three main pathways. The first deals with hydrolysis of the chlorine-carbon bond inducing the formation of hydroxyatrazine which is one of the main degradation products in both the soil and the aquatic systems.<sup>8,27-30</sup> Note that this hydrolysis yielding hydroxyatrazine is considered as a major step in the deactivation process of the herbicide.<sup>29,31</sup> The second is a *N*-dealkylation of carbon no. 4 and/or no. 6 which gives rise to deethylated atrazine, deisopropylated atrazine, and 2-chloro-4,6-diaminotriazine.<sup>8,28-30,32</sup> Both deethylated atrazine and deisopropylated atrazine are phytotoxic degradation products,<sup>32</sup> but 2-chloro-4,6-diaminotriazine has been shown to be nonphytotoxic in oat bioassays.<sup>29</sup> Nevertheless, atrazine is significantly more toxic than its degradation products towards *Chlorella pyrenoidosa*, *Scenedesmus quadricauda*, *Anabaena inaequalis*, *A. cylindrica*, and *A. variabilis*.<sup>33</sup> The third degradation process corresponds to the splitting of the triazine ring usually caused by microorganisms.<sup>2</sup>

The reported rates of decomposition in the various media of the environment are extremely variable.<sup>2</sup> In soil, the persistence of atrazine is influenced by a large number of physical, chemical, and biological factors.<sup>26,27,34,36</sup> In aquatic systems, the rate of hydrolytic and photolytic decomposition is affected by the chemical composition of the water.<sup>2,26</sup> When released into the atmosphere, reactions with photochemically produced hydroxyl radicals may be important.<sup>37</sup> Therefore, as stated above, the data concerning the environmental degradation of atrazine are numerous and variable. The degradation behavior of atrazine can be summarized by means of the following values of reaction half-lives:

- Half-life in soil: 1440 h,<sup>38</sup>
- Half-life in sediment: 420 h,<sup>39</sup>
- Half-life in water: 180 h,<sup>39</sup>
- Half-life in air: 19 h.<sup>39</sup>

The above values were selected from a perusal of the literature. However, they must only be regarded as a compromise to summarize the degradation profile of this chemical. Undoubtedly, there is considerable variation in these half-lives from place-to-place and from time-to-time, for example with temperature.

### Levels of contamination in the different compartments of the environment

#### Soil contamination

The atrazine soil concentrations depend on agricultural and non-agricultural practices, soil composition, and climatic conditions.<sup>14</sup> The recommended application level of atrazine in corn crops in the European community (EC) (i.e.; 1.5 kg of active ingredient per hectare)

corresponds to a concentration of 0.5  $\mu\text{g/g}$  of dry soil.<sup>14</sup> Pesticides added to soils are subject to sorption and to several chemical and biological degradation mechanisms that promote reduction of pesticide concentrations in the soil. One year after its soil application, the amount of atrazine present in the top surface soil is estimated to be around 8 to 20% of the initial level solution.<sup>14</sup>

### *Surface water contamination*

Numerous reports and publications have been written on the level of atrazine contamination of surface waters in various European countries and in North America.<sup>9-14,16,40</sup> The concentration of atrazine in water strongly depends upon the seasons, with maximum concentrations during the period when crops are planted. Table I gives the upper peak values in surface waters encountered during studies carried out between 1989 and 1991 in European countries.<sup>3,12</sup> However, the mean level in each case is far lower than the maximum concentration. In U.S. rivers, atrazine concentrations between 0 and 87  $\mu\text{g/l}$  have been found.<sup>2</sup> In aquatic systems, atrazine is taken up by nearly all members of the aquatic biocenoses, but in many cases, it is also quickly eliminated.<sup>2</sup> In France, a study performed in four rivers of Brittany highly contaminated by atrazine revealed no detectable concentration of atrazine in fish.<sup>41</sup> Only few investigations on the atrazine contamination of estuarine and sea waters in Europe have been carried out.<sup>40,42-44</sup> It is worth noting that these studies reported significant concentrations of atrazine in estuarine Mediterranean waters (e.g.; 17  $10^{-3}$  to 0.386  $\mu\text{g/l}$  in the Rhone Delta),<sup>40</sup> in the Baltic and North Sea (e.g.; 1  $10^{-3}$  to 0.11  $\mu\text{g/l}$  in German Bight),<sup>43</sup> and in British estuarine rivers.<sup>42</sup>

Table I. Maximum atrazine concentrations in surface waters (1989-1991).<sup>3,12</sup>

| Country         | Concentration        |
|-----------------|----------------------|
| Belgium         | 3.7 $\mu\text{g/l}$  |
| Denmark         | 0.42 $\mu\text{g/l}$ |
| France          | 7.5 $\mu\text{g/l}$  |
| Germany         | 0.42 $\mu\text{g/l}$ |
| Italy           | 2 $\mu\text{g/l}$    |
| Luxembourg      | 0.6 $\mu\text{g/l}$  |
| The Netherlands | 9.4 $\mu\text{g/l}$  |
| UK              |                      |
| - England       | 10 $\mu\text{g/l}$   |
| - Scotland      | 4.23 $\mu\text{g/l}$ |
| - Wales         | 0.17 $\mu\text{g/l}$ |

### *Groundwater contamination*

When applied on soils, pesticides migrate through the other environmental compartments by several processes such as volatilization, leaching, and runoff. The risk of groundwater

contaminations from vertical transports is ultimately determined by the balance between the rates of chemical sorption and degradation in soil profile.<sup>45</sup> Atrazine is only moderately sorbed in soil systems and is relatively persistent.<sup>24,36,46</sup> Indeed, one year after its application, atrazine residues are still found in soil.<sup>14,32</sup> Therefore, this herbicide can potentially contaminate groundwaters.<sup>47</sup> Thus, in the numerous sampling programs concerning groundwater contamination of various western European countries, atrazine was one of the most frequently detected pesticides.<sup>12,14</sup> Nevertheless, the concentrations recorded were generally lower than those found in surface water samples. In some cases the measured concentrations were greater than the EC limit (i.e.; 0.1 µg/l). This is clearly illustrated in Table II which shows that more than 30% of the French population is exposed to concentrations above the admissible limit. For comparison purposes, in the United States, during a national survey on the pesticides in drinking water wells, recorded atrazine concentrations ranged between 0.18 and 1.04 µg/l.<sup>2</sup>

Table II. Distribution of the French population as a function of maximum and mean atrazine concentrations (µg/l) in the water supplies (before any treatment) in 1990.<sup>12</sup>

|                   |                |                |                |                |                |                |
|-------------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>max. conc.</b> | <b>&lt;0.1</b> | <b>0.1-0.2</b> | <b>0.2-0.5</b> | <b>0.5-1.0</b> | <b>1.0-2.0</b> | <b>&gt;2.0</b> |
| % of population   | 66.3           | 15.5           | 14.2           | 2.4            | 0.9            | 0.7            |
| <b>mean conc.</b> | <b>&lt;0.1</b> | <b>0.1-0.2</b> | <b>0.2-0.5</b> | <b>0.5-1.0</b> | <b>1.0-2.0</b> | <b>&gt;2.0</b> |
| % of population   | 77.8           | 11.6           | 7.5            | 2.3            | 0.5            | 0.3            |

#### *Atmospheric contamination*

Despite the low rates of atrazine volatilization,<sup>48,49</sup> several studies have reported the presence of this herbicide in air, rain, snow, and fog samples. The presence of atrazine in the atmospheric environment could be due to spray droplet drift-off during its application or wind erosion of soil particles and wettable powder formulations from treated areas.<sup>14</sup> Atrazine volatilization, favored by the increasing moisture of soil surface, contributes to a smaller extent to the atmospheric contamination.<sup>14,50</sup>

Thus, for example, in the United States, concentrations in fog from 270 to 820 ng/l were recorded.<sup>15</sup> In air samples, concentrations between  $<0.2 \cdot 10^{-3}$  and  $3.4 \cdot 10^{-3}$  ng/l were measured.<sup>15</sup> In rain, the 1.0 µg/l level is frequently exceeded.<sup>16,17</sup> In Maryland, the mean concentration of atrazine in air samples was  $3.7 \cdot 10^{-3}$  ng/l in summer and  $0.026 \cdot 10^{-3}$  ng/l in winter 1982.<sup>16</sup> In France, concentrations of atrazine between <5 ng/l and 140 ng/l were found in rain water samples collected in Paris in 1991.<sup>18</sup> In the same study, this herbicide was only detected in the vapor phase between April and June at concentrations of ca.  $0.05 \cdot 10^{-3}$  ng/l.<sup>18</sup> In eastern France, concentrations in rain samples from 2300 to 3170 ng/l were recorded.<sup>14</sup> In Switzerland, the analysis of rain and snow samples revealed concentrations between 0 and 600 ng/l.<sup>10</sup> It is interesting to note that most of these studies showed that the concentrations of atrazine were strongly dependent upon the seasons.

**CHEMFRANCE***DESCRIPTION*

The complete description of CHEMFRANCE with all equations, definition of the regions, and so on can be found in a previous publication.<sup>20</sup> Briefly, in CHEMFRANCE, France has been divided into twelve regions of homogeneous hydrogeological, climatic, and ecologic characteristics. The environment is represented by a multimedia system constituted of six bulk compartments (i.e.; air, surface water, soil, bottom sediment, groundwater, coastal water). The first four bulk compartments are considered to be a combination of subcompartments of varying proportions of pure and particle phases. CHEMFRANCE is based on a level III fugacity model for which equilibrium within each bulk compartment is considered, but not between compartments. Expressions for emissions, advective flows, degrading reactions, and interphase transports by diffusive and non-diffusive processes are included. As input, the model requires information on the physicochemical properties (i.e.; molecular weight, aqueous solubility, vapor pressure, *n*-octanol/water partition coefficient, dissociation constant, melting point), reaction half-lives, and emission rates of the studied chemical. The output of the model principally consists of the estimated chemical distribution between environmental media, transport and transformation process rates, and steady-state concentrations in France or in a particular French region (Figure 1) at a defined season.




Figure 1. Defined regions of CHEMFRANCE.

*MODELING STRATEGY*

With the physicochemical properties and degradation half-life values listed in the previous section, almost all information necessary to run CHEMFRANCE is available. Indeed, only the

estimations of the amounts applied in each of the twelve defined regions of CHEMFRANCE (Figure 1) are missing as well as the definition of the media in which the release occurs. For obtaining the former information, some assumptions have to be made. In addition, numerous data dealing with the surface of treated crops, the applied dose, the number of treatments per year, and the percentage of active ingredient included in the formulations have to be collected. Thus, in France, atrazine is essentially used for pre- and post-emergence weed control in corn crops.<sup>4</sup> In 1991, corn crops represented 19.1% (i.e.; 3444082 ha) of French cultivated lands.<sup>51</sup> This indicates that an estimation of the total release of atrazine in France can be obtained principally from its application in corn crops. In addition, this shows that the release of atrazine into the environment principally occurs in spring and this was the season chosen to run CHEMFRANCE. A sample survey performed by the BVA institute on the use of weed control on corn crops in France in 1990 resulted in the following information on atrazine:

- Mean % of ha of corn crops treated: 99%,
- Mean number of treatment per year: 1.87,
- Mean % of ha of corn crops treated with atrazine alone: 36.9%,
- Mean % of ha of corn crops treated with atrazine associated with simazine: 2.6%,
- Mean % of ha of corn crops treated with atrazine associated with other active ingredients (e.g.; pendimethalin, cyanazine, metolachlor, alachlor, pyridate): 13.5%.

The survey also gave the mean dose applied for each formulation. The percentages of active ingredient of atrazine contained in these formulations were retrieved from a phytosanitary index<sup>52</sup> and the estimations of corn crop surfaces corresponding to the defined regions of CHEMFRANCE were estimated from the data given department by department.<sup>53</sup> With the above information, the amounts of atrazine applied on corn crops in the twelve defined regions of CHEMFRANCE in 1990 were estimated. In order to take into account the application of atrazine on other crops, and also on non-agricultural areas, we assumed that the use of atrazine as weed control in corn crops represented 90% of its total consumption.<sup>3</sup> These calculations yielded the global release in 1990 for the twelve regions of CHEMFRANCE (Table III). It is interesting to compare this estimation with the data related to the importations of atrazine in France between 1973 and 1991 (Figure 2).

The emission processes were simulated from four different scenarios. The first considered a total emission of atrazine into the soil compartment (scenario A). Scenario B took into account accidental pollutions occurring in surface waters (e.g.; cleaning tank near river) and the release via effluents at manufacturing sites. Therefore, in this scenario, emissions of 90% into the soil and 10% into the surface water compartment were assumed. The commercial formulations of atrazine usually involve spray applications on agricultural or non-agricultural areas which allow the dissemination of fine droplets providing a uniform distribution of a small volume over a large area. Through spray applications, significant quantities of atrazine may be released into the atmosphere.<sup>14,54</sup> Therefore, the third scenario (scenario C) was designed to also consider a loss of 1% into the air compartment of the atrazine applied on soil. The last scenario (scenario D) summarized all the previous assumptions (i.e.; 89% in soil, 10% in surface water, 1% in air).

Table III. Estimated total emission rates of atrazine for the twelve defined regions of CHEMFRANCE.

| Region* | Emission (kg/y) |
|---------|-----------------|
| 1       | 913130          |
| 2       | 576290          |
| 3       | 347700          |
| 4       | 298870          |
| 5       | 183720          |
| 6       | 193490          |
| 7       | 127010          |
| 8       | 29790           |
| 9       | 413590          |
| 10      | 147320          |
| 11      | 617080          |
| 12      | 510220          |
| France  | 4358210         |

\* For location, see Figure 1.




Figure 2. Imports and exports of atrazine between 1973 and 1991 in France.

## SIMULATION RESULTS AND COMPARISON WITH LABORATORY AND FIELD OBSERVATIONS

Table IV gives the average percentages corresponding to the removal processes obtained for the twelve defined regions of CHEMFRANCE. In all cases, atrazine is principally removed from the system by reaction in soil. This is in accordance with experimental studies.<sup>54,55</sup> As expected, the reaction in water significantly increases when emissions of atrazine in water compartment are simulated (i.e.; scenarios B and D). The dissipation of atrazine by advective processes for the air and water compartments is low. Furthermore, as atrazine is relatively rapidly degraded in air and water, the risk of contamination of adjacent regions is minor. However, this transport must not be neglected since it was suggested that this type of contamination was the major reason for the presence of atrazine in non-treated areas (e.g.; mountain lake).<sup>10</sup>

Field and laboratory studies clearly demonstrated the risk of groundwater contaminations by atrazine from vertical transports. Our results reveal that about 5% of the emissions reach to groundwater (Table IV). This percentage is close to the results obtained in a laboratory experiment in which 5.64% of the emissions are subjected to the leaching process.<sup>8</sup> However, it must be emphasized that the pollution of groundwater is principally due to the chlorinated metabolites of atrazine and more specifically deethylated atrazine, rather than atrazine. Indeed, a lysimeter study showed that atrazine represented only 10% of the leached compounds.<sup>56</sup>



Table IV. Average percentages of each removal process obtained from the four scenarios.

| Removal process             | Scenario A        | Scenario B        | Scenario C        | Scenario D        |
|-----------------------------|-------------------|-------------------|-------------------|-------------------|
| air advection               | $6 \cdot 10^{-5}$ | $5 \cdot 10^{-5}$ | $6 \cdot 10^{-3}$ | $6 \cdot 10^{-3}$ |
| water advection             | 0.12              | 0.31              | 0.12              | 0.31              |
| reaction in air             | $2 \cdot 10^{-4}$ | $4 \cdot 10^{-5}$ | 0.03              | 0.03              |
| reaction in water           | 5.34              | 14.45             | 5.35              | 14.46             |
| reaction in soil            | 88.98             | 80.09             | 88.94             | 80.05             |
| reaction in sediment        | 0.08              | 0.22              | 0.08              | 0.22              |
| transfer to higher altitude | $7 \cdot 10^{-8}$ | $6 \cdot 10^{-8}$ | $7 \cdot 10^{-6}$ | $5 \cdot 10^{-7}$ |
| leaching to groundwater     | 5.48              | 4.93              | 5.48              | 4.93              |
| sediment burial             | $2 \cdot 10^{-4}$ | $4 \cdot 10^{-4}$ | $2 \cdot 10^{-4}$ | $4 \cdot 10^{-4}$ |

The calculated percentage of loss of atrazine by runoff (i.e.; 5.54%) is in accordance with the mean percentage (i.e.; 3.19%) obtained in field analyses.<sup>7</sup> This shows the importance of horizontal transports in the contamination of the surface waters by atrazine. Despite the importance of the runoff process, the rain deposition of dissolved atrazine leading to low contamination levels cannot be neglected. In addition, from the results obtained with scenarios C and D, we can observe that more than 90% of the atmospheric emissions of atrazine are deposited by rain on the soil and water compartments. The wet and dry deposition processes are negligible. In the same way, the transfers of atrazine from soil and water to the air compartment are very weak. Schiavon and coworkers<sup>56</sup> reported volatilization data ranging between 0.017% and 60% of applied dose and selected the values of 0.05% and 0.08%.<sup>49</sup> These values agree with the average percentage of volatilized atrazine obtained in our four scenarios (i.e.; 0.02%). This confirms the minor influence of this process in the environmental fate of atrazine. The transfer rates of atrazine between the sediment and water compartments are dominated by the sorption process. This induces the accumulation of atrazine in the bottom sediment. The other transfer rates (i.e.; sediment resuspension, diffusion from air to soil and water) between the four principal compartments (i.e.; air, surface water, soil, bottom sediment) are negligible in all the scenarios.

Figure 3 summarizes the concentrations in the six bulk compartments for the twelve defined regions of CHEMFRANCE. It shows the relatively low differences between the four scenarios, except for the air compartment, in which the contamination significantly increases for scenarios C and D. Observed concentrations in air, surface water, and groundwater as well as those derived from the different simulation scenarios are reported in Table V.

The high surface water and groundwater concentrations obtained in all the simulations (Figure 3) are in accordance with atrazine field data (Table V). Nevertheless, the results obtained for the groundwater compartment are of lesser quality. No data are available for the French sea water contamination by this herbicide. Therefore, our results are not directly comparable with field data. Nevertheless, it is interesting to note that the study dealing with the contamination of the German Bight (North Sea) reported concentrations of atrazine between  $<<1$  to 110 ng/l,<sup>43</sup> which greatly exceed our results corresponding to the same area (i.e.;  $6.18 \cdot 10^{-2}$  to 0.15 ng/l).




Figure 3. Atrazine concentrations (ng/l) for the bulk compartments calculated with CHEMFRANCE for the twelve defined regions and the four scenarios (see text).

Table V. Comparison between field and predicted (scenarios A-D) concentrations (ng/l).

| Date, region of CHEMFRANCE                    | Actual concentration       |           | Reference | Predicted concentration |                       |                       |                       |
|---|----------------------------|-----------|-----------|-------------------------|-----------------------|-----------------------|-----------------------|
|   | mean <sup>s</sup>          | maximum   |           | A                       | B                     | C                     | D                     |
| <b>Surface water compartment</b>              |                            |           |           |                         |                       |                       |                       |
| 4 rivers (17/04/1990 to 24/07/1990), region 1 | 2149 (n = 40)              | 7460      | 41        | 1949                    | 4711                  | 1950                  | 4712                  |
| 4 rivers (17/04/1991 to 24/07/1991), region 1 | 984 (n = 26)               | 3050      | 41        | 1949                    | 4711                  | 1950                  | 4712                  |
| 12 rivers (04/1991 to 11/1991), region 1      | 210 (n = 75)               | 750       | 41        | 1949                    | 4711                  | 1950                  | 4712                  |
| 1 river (17/04/1990 to 24/07/1990), region 2  | 1747 (n = 10)              | 6030      | 41        | 1236                    | 2989                  | 1237                  | 2989                  |
| 1 river (17/04/1991 to 24/07/1991), region 2  | 788 (n = 7)                | 2500      | 41        | 1236                    | 2989                  | 1237                  | 2989                  |
| 2 rivers (04/1991 to 11/1991), region 2       | 1206 (n = 8)               | 2400      | 41        | 1236                    | 2989                  | 1237                  | 2989                  |
| Seine barrage, region 3                       | 98 (n = 46)                | 260       | 41        | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| 9 rivers (04/07/1989 to 28/08/1989), region 3 | 360 (n = 106)              | 1550      | 41        | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| Seine river (July 1987), region 3             | 810 (n = 1)                |           | 13        | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| Marne river (July 1987), region 3             | 60 (n = 1)                 |           | 13        | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| 1 river (18/03/1991 to 16/03/1992), region 4  | 159 (n = 51)               | 340       | 41        | 375.8                   | 1308                  | 376.4                 | 1309                  |
| 29 rivers (summer 1990), region 9             | 165 (n = 47)               | 710       | 41        | 593.5                   | 2066                  | 594.4                 | 2067                  |
| Rhone delta (10/1990 to 03/1994), region 9    | 17-386 <sup>£</sup>        |           | 40        | 593.5                   | 2066                  | 594.4                 | 2067                  |
| Rhone river, 7 stations (10/1990), region 9   | 12-33 <sup>£</sup>         |           | 44        | 593.5                   | 2066                  | 594.4                 | 2067                  |
| 15 rivers (summer 1990), region 11            | 277 (n = 24)               | 1000      | 41        | 460.6                   | 1856                  | 462.0                 | 1858                  |
| <b>Groundwater compartment</b>                |                            |           |           |                         |                       |                       |                       |
| (July 1987), region 1                         | 60 (n = 1)                 |           | 13        | 4614                    | 4153                  | 4612                  | 4151                  |
| (July 1987), region 3                         | 95 (n = 1)                 |           | 13        | 565.7                   | 509.1                 | 565.4                 | 508.8                 |
| (November 1987), region 3                     | 225 (n = 1)                |           | 13        | 565.7                   | 509.1                 | 565.4                 | 508.8                 |
| (07/1990 to 09/1990), region 4                | 182 (n = 39)               | 856       | 41        | 774.4                   | 697.0                 | 774.2                 | 696.7                 |
| (November 1987), region 4                     | 90 (n = 1)                 |           | 13        | 774.4                   | 697.0                 | 774.2                 | 696.7                 |
| (summer 1990), region 9                       | 174 (n = 18)               | 630       | 41        | 1093                    | 984.0                 | 1093                  | 983.6                 |
| (summer 1990), region 10                      | 238 (n = 6)                | 430       | 41        | 1408                    | 1267                  | 1407                  | 1266                  |
| (summer 1990), region 11                      | 86.6 (n = 3)               | 130       | 41        | 1276                    | 1148                  | 1275                  | 1148                  |
| (July 1987), region 11                        | 110 (n = 1)                |           | 13        | 1276                    | 1148                  | 1275                  | 1148                  |
| (April 1989); region 12                       | 200 (n = 1)                |           | 13        | 1937                    | 1743                  | 1936                  | 1742                  |
| <b>Air compartment</b>                        |                            |           |           |                         |                       |                       |                       |
| air (04/1991 to 06/1991), region 3            | 5 10 <sup>-5</sup> (n = ?) | not given | 18        | 3.66 10 <sup>-8</sup>   | 3.40 10 <sup>-8</sup> | 6.38 10 <sup>-6</sup> | 5.74 10 <sup>-6</sup> |
| rain (03/1991 to 12/1991), region 3           | <5-140 <sup>£</sup>        |           | 18        | 0.08                    | 0.07                  | 13.98                 | 12.56                 |

<sup>s</sup> n = number of values; <sup>£</sup> range

The calculated concentrations in the air compartment are relatively lower than those observed in the environment (Figure 3 and Table V). By means of an equation published by Mackay and coworkers<sup>57</sup>, the concentrations in rain for region 3 were calculated from the air concentrations obtained with CHEMFRANCE. For scenarios A and B, the calculated rain concentrations are lower than those observed in Paris between March and December 1991 (Table V).<sup>18</sup> At the opposite, for scenarios C and D, the generated values are relatively closer to the actual data (Table V).

The calculated soil concentrations are lower than those expected after an application (i.e.; 0.5 µg/g) at the rate recommended by the EC. However, these results agree with those observed in a soil historically treated (i.e.; 0.025 µg/g) but before any further treatment.<sup>30</sup>

The mass distribution obtained in the soil compartment (i.e.; 96.7% in the solid components and 3.3% in the water and air contained in the pore of the soil) is near this reported by Tasli.<sup>58</sup> Indeed, in her study, 96% of the amount included in the soil is sorbed in the soil solids and 4% is dissolved in the soil water. In the same way, the results obtained for the sediment compartment indicate that 89% of the atrazine is sorbed and 11% dissolved. Huckins *et al.*,<sup>48</sup> in a microcosm study, found a slightly smaller percentage (i.e.; 86.1%) although the organic carbon content of their studied sediment (i.e.; 7.28%) was higher than this considered in CHEMFRANCE (i.e.; 4%). In the water compartment, ca. 99.9% of the total atrazine content of this compartment is dissolved in pure water, 0.07% is sorbed in the suspended sediment, and 0.002% is accumulated in biota (i.e.; fish). Field studies gave similar results (i.e.; 99.5% and 0.5% of atrazine in the dissolved and particulate river matter, respectively).<sup>11</sup> The mass distributions obtained for the air compartment cannot be compared with experimental results. In the air compartment, atrazine is principally present in pure air (i.e.; 89.2%), and only a low quantity is bound to the air particles (i.e.; 10.8%).

## CONCLUSION

From the survey of literature data available on the environmental fate and hazards of atrazine, it appears that this chemical presents a high contamination potential for water systems, since leaching and runoff processes have been identified as major fate pathways. The results derived with CHEMFRANCE are in accordance with those obtained in laboratory or field experiments. Therefore, CHEMFRANCE can be considered as a useful tool for simulating the environmental fate of this agrochemical in France. In addition, CHEMFRANCE provides a concise environmental fate profile of atrazine while literature data on this herbicide are numerous and variable, and therefore, difficult to interpret. This underlines the usefulness of environmental fate models for confirming results of fields and laboratory studies but also to provide a comprehensive summary of the processes involved in the environmental fate of chemicals.

An interesting characteristic of models is that they allow the user to quickly test several hypotheses through different scenarios. Thus, in our study, four scenarios assuming different emission profiles were studied. By taking into account the accidental pollutions occurring in surface water and the releases into the atmosphere through spray applications (scenario D) gave

the most satisfactory results. However, many other scenarios integrating variations in seasons, quantities, and so on could have been tested. In the same way, it should be noted that models can be coupled with sensitivity and uncertainty analyses in order to test the influence of all input parameters and assumptions on the outputs.

## ACKNOWLEDGMENTS

The authors are grateful to Pr. D. Mackay (Department of Chemical Engineering and Applied Chemistry, University of Toronto, Canada) for his thoughtful comments. This study was granted by the French Ministry of the Environment.

## REFERENCES

1. Worthing, C.R. (1987). *The Pesticide Manual, Eighth Edition*. The British Crop Protection Council, Thornton Heath, p. 1081.
2. Huber, W. (1993). Ecotoxicological relevance of atrazine in aquatic systems. *Environ. Toxicol. Chem.* **12**, 1865-1881.
3. Persoone, G., Janssen, C.R., and van Steertegem, M. (1991). *Extensive and Updated Literature Study of the Toxicity and Ecotoxicity of Atrazine*. Commission of the European Communities.
4. Montiel, A. (1989). *Etude bibliographique. Utilisation des Herbicides et des Triazines en Particulier*. Ministère de la Santé, DDRASS Ile de France, p. 273.
5. Hall, J.K., Pawlus, M., and Higgins, E.R. (1972). Losses of atrazine in runoff water and soil sediment. *J. Environ. Qual.* **1**, 172-176.
6. Hall, J.K. (1974). Erosional losses of *s*-triazine herbicides. *J. Environ. Qual.* **3**, 174-180.
7. Wauchope, R.D. (1978). The pesticide content of surface water draining from agricultural fields-A review. *J. Environ. Qual.* **7**, 459-472.
8. Schiavon, M. (1988). Studies of the leaching of atrazine, of its chlorinated derivatives, and of hydroxyatrazine from soil using <sup>14</sup>C ring-labeled compounds under outdoor conditions. *Ecotoxicol. Environ. Safety* **15**, 46-54.
9. Frank, R. and Sirons, G.J. (1979). Atrazine: its use in corn production and its loss to stream waters in southern Ontario, 1975-1977. *Sci. Total Environ.* **12**, 223-239.
10. Buser, H.R. (1990). Atrazine and other *s*-triazine herbicides in lakes and in rain in Switzerland. *Environ. Sci. Technol.* **24**, 1049-1058.
11. Pereira, W.E. and Rostad, C.E. (1990). Occurrence, distributions, and transport of herbicides and their degradation products in the lower Mississippi River and its tributaries. *Environ. Sci. Technol.* **24**, 1400-1406.
12. ECPA (1992). *Monitoring Pesticides in Water and Definition of the Uniform Principles: Volume 2 - Case Studies of Atrazine and Simazine*. European Crop Protection Association.
13. Legrand, M.F., Costentin, E., and Bruchet, A. (1992). Occurrence of 38 pesticides in various French surface and ground waters. *Water Supply* **10**, 51-61.

14. Mirgain, I., Schenck, C., and Monteil, H. (1993). Atrazine contamination of groundwaters in eastern France in relation to the hydrogeological properties of the agricultural land. *Environ. Technol.* **14**, 741-750.
15. Glotfelty, D.E., Seiber, J.N., and Liljedahl, L.A. (1987). Pesticides in fog. *Nature* **325**, 602-605.
16. Glotfelty, D.E., Williams, G.H., Freeman, H.P., and Leech, M.M. (1990). Regional atmospheric transport and deposition of pesticides in Maryland. In, *Long Range Transport of Pesticides* (D.A. Kurtz, Ed.). Lewis Publishers, Chelsea, pp. 199-221.
17. Richards, R.P., Kramer, J.W., Baker, D.B., and Krieger, K.A. (1987). Pesticides in rainwater in the northeastern United States. *Nature* **327**, 129-131.
18. Chevreuril, M. and Garmouma, M. (1993). Occurrence of triazines in the atmospheric fallout on the catchment basin of the river Marne (France). *Chemosphere* **27**, 1605-1608.
19. Mackay, D. (1991). *Multimedia Environmental Models. The Fugacity Approach*. Lewis Publishers, Chelsea, p. 257.
20. Devillers, J., Bintein, S., and Karcher, W. (1995). CHEMFRANCE: a regional level III fugacity model applied to France. *Chemosphere* **30**, 457-476.
21. Domine, D., Devillers, J., Chastrette, M., and Karcher, W. (1992). Multivariate structure-property relationships (MSPR) of pesticides. *Pestic. Sci.* **35**, 73-82.
22. Devillers, J., Karcher, W., and Isnard, P. (1991). Graphical display of the fugacity model level I. *Chemosphere* **23**, 633-642.
23. Devillers, J., Thioulouse, J., Domine, D., Chastrette, M., and Karcher, W. (1991). Multivariate analysis of the input and output data in the fugacity model level I. In, *Applied Multivariate Analysis in SAR and Environmental Studies* (J. Devillers and W. Karcher, Eds.). Kluwer Academic Publishers, Dordrecht, pp. 281-345.
24. Francioso, O., Bak, E., Rossi, N., and Sequi, P. (1992). Sorption of atrazine and trifluralin in relation to the physico-chemical characteristics of selected soils. *Sci. Total Environ.* **123/124**, 503-512.
25. Bacci, E., Renzoni, A., Gaggi, C., Calamari, D., Franchi, A., Vighi, M., and Severi, A. (1989). Models, field studies, laboratory experiments: an integrated approach to evaluate the environmental fate of atrazine (*s*-triazine herbicide). *Agric. Ecosys. Environ.* **27**, 513-522.
26. Burkhard, N. and Guth, J.A. (1981). Chemical hydrolysis of 2-chloro-4,6-bis(alkylamino)-1,3,5-triazine herbicides and their breakdown in soil under the influence of adsorption. *Pestic. Sci.* **12**, 45-52.
27. Reinhardt, C.F., Ehlers, J.G., and Nel, P.C. (1990). Persistence of atrazine as affected by selected soil properties. *S.-Afr. Tydskr. Plant Grond* **7**, 182-187.
28. Winkelmann, D.A. and Klaine, S.J. (1991). Degradation and bound residue formation of atrazine in a western Tennessee soil. *Environ. Toxicol. Chem.* **10**, 335-345.
29. Winkelmann, D.A. and Klaine, S.J. (1991). Degradation and bound residue formation of four atrazine metabolites, deethylatrazine, deisopropylatrazine, dealkylatrazine and hydroxyatrazine, in a western Tennessee soil. *Environ. Toxicol. Chem.* **10**, 347-354.
30. Huart, B., Pailler, F.M., Trique, B., and Ricordel, I. (1993). Protocole d'étude de la diffusion d'un herbicide dans l'écosystème. *Ann. Fals. Exp. Chim.* **86**, 75-86
31. Malan, C., Visser, J.H., and van de Venter, H.A. (1987). Atrazine metabolism in field-grown maize and the effect of atrazine phytotoxicity on yield. *S. Afr. J. Plant Soil* **4**, 7-11.

32. Sirons, G.J., Frank, R., and Sawyer, T. (1973). Residues of atrazine, cyanazine, and their phytotoxic metabolites in a clay loam soil. *J. Agric. Food Chem.* **21**, 1016-1020.
33. Stratton, G.W. (1984). Effects of the herbicide atrazine and its degradation products, alone and in combination, on phototrophic microorganisms. *Arch. Environ. Contam. Toxicol.* **13**, 35-42.
34. Hiltbold, A.E. and Buchanan, G.A. (1977). Influence of soil pH on persistence of atrazine in the field. *Weed Sci.* **25**, 515-520.
35. Helling, C.S., Zhuang, W., Gish, T.J., Coffman, C.B., Isensee, A.R., Kearney, P.C., Hoagland, D.R., and Woodward, M.D. (1988). Persistence and leaching of atrazine, alachlor, and cyanazine under no-tillage practices. *Chemosphere* **17**, 175-187.
36. Singh, G., Spencer, W.F., Cliath, M.M., and van Genuchten, M.T. (1990). Dissipation of s-triazines and thiocarbamates from soil as related to soil moisture content. *Environ. Pollut.* **66**, 253-262.
37. Atkinson, R., Tuazon, E.C., Wallington, T.J., Aschmann, S.M., Arey, J., Winer, A.M., and Pitts, J.N. (1987). Atmospheric chemistry of aniline, *N,N*-dimethylaniline, pyridine, 1,3,5-triazine, and nitrobenzene. *Environ. Sci. Technol.* **21**, 64-72.
38. Wauchope, R.D., Buttler, T.M., Hornsby, A.G., Augustijn-Beckers, P.W.M., and Burt, J.P. (1992). The SCS/ARS/CES pesticide properties database for environmental decision-making. *Rev. Environ. Contam. Toxicol.* **123**, 1-164.
39. Mackay, D., Paterson, S., Cheung, B., and Neely, W.B. (1985). Evaluating the environmental behavior of chemicals with a level III fugacity model. *Chemosphere* **14**, 335-374.
40. Readman, J.W., Albanis, T.A., Barcelo, D., Galassi, S., Tronczynski, J., and Gabrielides, G.P. (1993). Herbicide contamination of Mediterranean estuarine waters: results from a MED POL pilot survey. *Marine Poll. Bull.* **26**, 613-619.
41. Devillers, J., Bintein, S., and Domine, D. (1992). Atrazine in France. Report CTIS #92011, p. 42.
42. Ahel, M., Evans, K.M., Fileman, T.W., and Mantoura, R.F.C. (1992). Determination of atrazine and simazine in estuarine samples by high-resolution gas chromatography and nitrogen selective detection. *Anal. Chim. Acta* **268**, 195-204.
43. Bester, K. and Hühnerfuss, H. (1993). Triazines in the Baltic and North Sea. *Marine Poll. Bull.* **26**, 423-427.
44. Tronczynski, J., Munsch, C., Durand, G., and Barcelo, D. (1993). Monitoring of trace-levels of herbicides and their degradation products in the river Rhône, France, by gas chromatography-mass spectrometry. *Sci. Total Environ.* **132**, 327-337.
45. Gamberdinger, A.P., Lemley, A.T., and Wagenet, R.J. (1991). Nonequilibrium sorption and degradation of three 2-chloro-s-triazine herbicides in soil-water systems. *J. Environ. Qual.* **20**, 815-822.
46. Talbert, R.E. and Fletchall, O.H. (1965). The adsorption of some s-triazines in soils. *Weeds* **13**, 46-52.
47. Bowman, B.T. (1990). Mobility and persistence of alachlor, atrazine and metolachlor in plainfield sand, and atrazine and isazofos in honeywood silt loam, using field lysimeters. *Environ. Toxicol. Chem.* **9**, 453-461.

48. Huckins, J.N., Petty, J.D., and England, D.C. (1986). Distribution and impact of trifluralin, atrazine, and fonofos residues in microcosms simulating a northern prairie wetland. *Chemosphere* 15, 563-588.
49. Scheunert, I., Qiao, Z., and Korte, F. (1986). Comparative studies of the fate of atrazine-<sup>14</sup>C and pentachlorophenol-<sup>14</sup>C in various laboratory and outdoor soil-plant systems. *J. Environ. Sci. Health B21*, 457-485.
50. Glotfelty, D.E., Leech, M.M., Jersey, J., and Taylor, A.W. (1989). Volatilization and wind erosion of soil surface applied atrazine, simazine, alachlor, and toxaphene. *J. Agric. Food Chem.* 37, 546-551.
51. AGRESTE (1992). *Statistique Agricole Annuelle. Résultats Provisoires 1991*. Ministère de l'Agriculture et de la Forêt, France, p. 123.
52. ACTA (1992). *Index Phytosanitaire*. Association de Coordination Technique Agricole, Paris, France, p. 519.
53. AGRESTE (1991). *Statistique Agricole Annuelle. Résultats 1990*. Ministère de l'Agriculture et de la Forêt, France, p. 125.
54. Paterson, K.G. and Schnoor, J.L. (1992). Fate of alachlor and atrazine in a riparian zone field site. *Water Environ. Res.* 64, 274-283.
55. Di Muccio, A., Chirico, M., Dommarco, R., Funari, E., Musmeci, L., Santilio, A., Vergori, F., Zapponi, G., Giuliano, G., and Sparacino, A.C. (1990). Vertical mobility of soil contaminants: preliminary results of a survey on the herbicide atrazine. *Intern. J. Environ. Anal. Chem.* 38, 211-220.
56. Schiavon, M., Portal, J.M., and Andreux, F. (1992). Données actuelles sur les transferts d'atrazine dans l'environnement. *Agronomie* 12, 129-139.
57. Mackay, D., Paterson, S., and Schroeder, W.H. (1986). Model describing the rates of transfer processes of organic chemicals between atmosphere and water. *Environ. Sci. Technol.* 20, 810-816.
58. Tasli, S. (1992). *Distribution de l'Atrazine entre le Sol, l'Eau et les Plantes en Culture de Maïs*. DEA d'Agrochimie, University of Grenoble, p. 23.



# ARTICLE VI

## Modeling the Environmental Fate of Atrazine

J. Devillers, S. Bintein, and D. Domine

CTIS, 21 rue de la Bannière, 69003 Lyon, France

Mathematical simulation models of fate and transport of chemicals have been identified by researchers and regulators as potentially valuable tools for improving the understanding of the environmental behavior of chemicals which may be released to the environment as a consequence of routine (i.e.; normal manufacturing, use, disposal) and non-routine (e.g.; accidental spill) events. In this context, CHEMFRANCE, a regional fugacity model level III, which calculates the environmental distribution of organic chemicals in 12 defined regions of France, or France as a whole, has been designed. The aim of this study is to show that CHEMFRANCE provides interesting simulation results for understanding the environmental fate behavior of atrazine.

Organic chemicals may be released into the compartments of the biosphere as a result of their normal manufacture, formulation, use, and disposal or as a result of their accidental spill. Therefore, it is essential to develop both a qualitative and quantitative understanding of their environmental fate behavior (1). These insights, along with information on their possible adverse effects towards organisms occupying different trophic levels in the ecosystems, enable the risk to human health and the environment to be assessed. To achieve this understanding, the physicochemical properties of the organic compounds and emission patterns must be combined with the characteristics of the environment into which they are released in order to predict how advective and inter-compartment transport and transformation processes will affect their environmental distribution. However, because the environmental fate behavior of a chemical is always the result of numerous complex and interacting processes, the use of models is highly recommended. Among them, the Mackay-type multimedia models have shown their efficiency for estimating the environmental fate profile of numerous organic chemicals at a global or regional scale (1-4). In this context, CHEMFRANCE, a level III fugacity computer model, has been designed for simulating the environmental fate of organic chemicals in France (5). In this paper, we show the usefulness of CHEMFRANCE for simulating the environmental fate behavior of atrazine. Indeed, in France this herbicide is of environmental concern since it is widely used for control of certain annual broadleaf and grass weeds in corn, and to a lesser extent, other crops and landscaping (6).

## Model Description

In CHEMFRANCE, France has been divided into twelve regions of homogeneous hydrogeological, climatic, and ecologic characteristics (Figure 1). The environment is represented by a multimedia system constituted of six bulk compartments (i.e.; air, surface water, soil, bottom sediment, groundwater, coastal water) presenting various characteristics summarized in Table I. The first four bulk compartments (Table I) are considered to be a combination of subcompartments of varying proportions of pure and particle phases. CHEMFRANCE is based on a level III fugacity model for which equilibrium within each bulk compartment is considered, but not between compartments. Expressions for emissions, advective flows, degrading reactions, and interphase transports by diffusive and non-diffusive processes are included. As input, CHEMFRANCE requires the molecular weight, aqueous solubility, vapor pressure, log P, pKa, melting point, reaction half-lives, and emission rates of the studied chemical. The outputs of CHEMFRANCE principally consist of the estimated chemical distribution between environmental media, transport and transformation process rates, and steady-state concentrations in France or in a particular French region (Figure 1) at a defined season. A more complete description of CHEMFRANCE can be found in a recent publication (5).

## Data on Atrazine and Simulation Scenarios

The physicochemical data and reaction half-lives selected for running CHEMFRANCE were the following:

- molecular weight: 215.69 g/mol,
- solubility in water: 30 mg/l at 20°C (7),
- vapor pressure:  $4 \cdot 10^{-5}$  Pa at 20°C (7),
- *n*-octanol/water partition coefficient (log P): 2.34 (7),
- melting point: 174°C (8),
- pKa: 1.70 (7),
- half-life in soil: 1440 h (9),
- half-life in sediment: 420 h (2),
- half-life in water: 180 h (2),
- half-life in air: 19 h (2).

With the above physicochemical properties and degradation half-life values, almost all information necessary to run CHEMFRANCE is available. Indeed, only the estimations of the amounts applied in each of the twelve defined regions of CHEMFRANCE (Figure 1) are missing as well as the definition of the compartment in which the release occurs. For obtaining the former information, some assumptions have to be made. In addition, numerous data dealing with the surface of treated crops, the applied dose, the number of treatments per year, and the percentage of active ingredient included in the formulations have to be collected. Thus, in France, atrazine is essentially used for pre- and post-emergence weed control in corn crops (6). In 1991, corn crops represented 19.1% (i.e.; 3444082 ha) of French cultivated lands (10). This indicates that an estimation of the total release of atrazine in France can be obtained principally from its application in corn crops. In addition, this shows that the release of atrazine into the environment principally occurs in spring and this was the season chosen to run CHEMFRANCE. A sample survey performed by the BVA institute on the use of weed control on corn crops in France in 1990 resulted in the following information on atrazine:

- mean % of ha of corn crops treated: 99%,
- mean number of treatments per year: 1.87,
- mean % of ha of corn crops treated with atrazine alone: 36.9%,
- mean % of ha of corn crops treated with atrazine associated with simazine: 2.6%,
- mean % of ha of corn crops treated with atrazine associated with other active ingredients (e.g.; pendimethalin, cyanazine, metolachlor, alachlor, pyridate): 13.5%.




Figure 1. Defined regions of CHEMFRANCE

Table I. Principal characteristics of the bulk compartments in CHEMFRANCE

| Compartment   | Region-dependent characteristics*      | Fixed model characteristics  |
|---------------|--|--|
| Air           | Total surface area                     | Residence time (5 d); Height (1000 m); Vf** of aerosols ( $2 \cdot 10^{-11}$ ); Pure air ( $1.2 \text{ kg/m}^3$ ) and aerosol density ( $1500 \text{ kg/m}^3$ )                                    |
| Water         | Surface; Volume; Vf of Ss <sup>‡</sup> | Residence time (500 d); OC <sup>§</sup> of Ss (20%); pH of Ss (7); Vf of biota ( $1 \cdot 10^{-6}$ ); Pure water and biota density ( $1000 \text{ kg/m}^3$ ); Ss density ( $1500 \text{ kg/m}^3$ ) |
| Soil          | Surface; OC; pH                        | Depth (15 cm); Vf of air (0.2) and water (0.3); Pure soil density ( $2400 \text{ kg/m}^3$ )  |
| Sediment      |  | Depth (3 cm); OC (4%); pH (7); Vf of water (0.7); Pure sediment density ( $2400 \text{ kg/m}^3$ )  |
| Groundwater   | Volume                                 |  |
| Coastal water | Volume; Length                         | Depth (100 m)  |

\*Monthly mean precipitations and temperatures for the four seasons are also region-dependent.

\*\*Volume fraction; <sup>‡</sup>Suspended sediment; <sup>§</sup>% of organic carbon content.

The BVA survey also gave the mean dose applied for each formulation. The percentages of active ingredient of atrazine contained in these formulations and the estimations of corn crop surfaces corresponding to the 12 regions of CHEMFRANCE were retrieved from literature (11, 12). With the above information, the amounts of atrazine applied on corn crops in the 12 regions of CHEMFRANCE in 1990 were estimated. In order to take into account the application of atrazine on other crops, and also on non-agricultural areas, we assumed that the use of atrazine as weed control in corn crops represented 90% of its total consumption (13). These calculations yielded the global release in 1990 for the 12 regions of CHEMFRANCE (Table II).

Table II. Estimated total emission rates of atrazine for the 12 defined regions of CHEMFRANCE

| Region* | Emission (kg/y) |
|---------|-----------------|
| 1       | 913130          |
| 2       | 576290          |
| 3       | 347700          |
| 4       | 298870          |
| 5       | 183720          |
| 6       | 193490          |
| 7       | 127010          |
| 8       | 29790           |
| 9       | 413590          |
| 10      | 147320          |
| 11      | 617080          |
| 12      | 510220          |
| France  | 4358210         |

\*For location see Figure 1.

The emission processes were simulated from four different scenarios. The first considered a total emission of atrazine into the soil compartment (scenario A). Scenario B took into account accidental pollutions occurring in surface waters (e.g.; cleaning tank near river) and the release via effluents at manufacturing sites. Therefore, in this scenario, emissions of 90% into the soil and 10% into the surface water compartment were assumed. The commercial formulations of atrazine usually involve spray applications inducing the release of significant quantities into the atmosphere (14, 15). Therefore, the third scenario (scenario C) was designed to also consider a loss of 1% into the air compartment of the atrazine applied on soil. The last scenario (scenario D) summarized all the previous assumptions (i.e.; 89% in soil, 10% in surface water, 1% in air).

### Simulation Results and Comparison with Laboratory and Field Observations

Table III gives the average percentages corresponding to the removal processes obtained for the 12 defined regions of CHEMFRANCE. In all cases, atrazine is principally removed from the system by reaction in soil. This is in accordance with experimental studies (15, 16). As expected, the reaction in water significantly increases when emissions of atrazine in water compartment are simulated (i.e.; scenarios B and D). The dissipation of atrazine by advective processes for the air and water compartments is low. Furthermore, as atrazine is relatively rapidly degraded in air and water, the risk of contamination of adjacent regions is minor. However, this transport must not be neglected since it was suggested that this type of contamination was the major reason for the presence of atrazine in non-treated areas (e.g.; mountain lake) (17).

Table III. Average percentages of each removal process obtained from the four scenarios

| Removal process             | Scenario A | Scenario B | Scenario C | Scenario D |
|-----------------------------|------------|------------|------------|------------|
| Air advection               | <<0.01     | <<0.01     | <<0.01     | <<0.01     |
| Water advection             | 0.12       | 0.31       | 0.12       | 0.31       |
| Reaction in air             | <<0.01     | <<0.01     | 0.03       | 0.03       |
| Reaction in water           | 5.34       | 14.45      | 5.35       | 14.46      |
| Reaction in soil            | 88.98      | 80.09      | 88.94      | 80.05      |
| Reaction in sediment        | 0.08       | 0.22       | 0.08       | 0.22       |
| Transfer to higher altitude | <<0.01     | <<0.01     | <<0.01     | <<0.01     |
| Leaching to groundwater     | 5.48       | 4.93       | 5.48       | 4.93       |
| Sediment burial             | <<0.01     | <<0.01     | <<0.01     | <<0.01     |

Table III shows that about 5% of the emissions leach to groundwater. This percentage is close to the results obtained in a laboratory experiment in which 5.64% of the emissions were subjected to the leaching process (18). However, it must be emphasized that the pollution of groundwater is principally due to the chlorinated metabolites of atrazine and more specifically deethylated atrazine, rather than atrazine. Indeed, a lysimeter study showed that atrazine represented only 10% of the leached compounds (19).

The calculated percentage of loss of atrazine by runoff (i.e.; 5.54%) is in accordance with the mean percentage (i.e.; 3.19%) obtained in field analyses (20). This shows the importance of horizontal transports in the contamination of the surface waters by atrazine. Despite the importance of the runoff process, the rain deposition of dissolved atrazine leading to low contamination levels cannot be neglected. In addition, from the results obtained with scenarios C and D, we can observe that more than 90% of the atmospheric emissions of atrazine are deposited by rain on the soil and water compartments. The wet and dry deposition processes are negligible. In the same way, the transfers of atrazine from soil and water to the air compartment are very weak. Schiavon and coworkers (19) reported volatilization data ranging between 0.017% and 60% of applied dose and selected the values of 0.05% and 0.08% (21). These values agree with the average percentage of volatilized atrazine obtained in our four scenarios (i.e.; 0.02%). This confirms the minor influence of this process in the environmental fate of atrazine. The transfer rates of atrazine between the sediment and water compartments are dominated by the sorption process. This induces the accumulation of atrazine in the bottom sediment. The other transfer rates (i.e.; sediment resuspension, diffusion from air to soil and water) between the four principal compartments (i.e.; air, surface water, soil, bottom sediment) are negligible in all the scenarios.

When the concentrations in atrazine in the six bulk compartments for the 12 defined regions of CHEMFRANCE are calculated, relatively low differences between the four scenarios are observed, except for the air compartment, in which the contamination significantly increases for scenarios C and D. Only the results obtained for the least and the most contaminated regions are given in Tables IV and V, respectively. Observed concentrations in air, surface water, and groundwater as well as those derived from the different simulation scenarios are reported in Table VI. The high surface water and groundwater concentrations obtained in all the simulations are generally in accordance with atrazine field data (Table VI). Nevertheless, the results obtained for the groundwater compartment are of lesser quality. No data are available for the French sea water contamination by atrazine. Therefore, our results are not directly comparable with field data. Nevertheless, it is interesting to note that the study dealing with the contamination of the German Bight (North Sea) reported concentrations of atrazine between <<1 and 110 ng/l (22), which greatly exceed our results corresponding to the same area (i.e.;  $6.18 \cdot 10^{-2}$  to 0.15 ng/l).

Table IV. Concentrations (ng/l) obtained with the four scenarios for region 8

| Compartment   | Scenario A            | Scenario B            | Scenario C            | Scenario D            |
|---------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Air           | 4.21 10 <sup>-9</sup> | 3.89 10 <sup>-9</sup> | 6.04 10 <sup>-7</sup> | 5.44 10 <sup>-7</sup> |
| Water         | 7.14                  | 27.24                 | 7.18                  | 27.28                 |
| Soil          | 879.4                 | 791.4                 | 878.8                 | 791.2                 |
| Sediment      | 27.67                 | 105.6                 | 27.83                 | 105.8                 |
| Groundwater   | 59.88                 | 53.89                 | 59.84                 | 53.87                 |
| Coastal water | 3.08 10 <sup>-3</sup> | 1.18 10 <sup>-2</sup> | 3.10 10 <sup>-3</sup> | 1.18 10 <sup>-2</sup> |

Table V. Concentrations (ng/l) obtained with the four scenarios for region 1

| Compartment   | Scenario A            | Scenario B            | Scenario C            | Scenario D            |
|---------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Air           | 2.79 10 <sup>-7</sup> | 2.55 10 <sup>-7</sup> | 2.08 10 <sup>-5</sup> | 1.88 10 <sup>-5</sup> |
| Water         | 1949                  | 4711                  | 1950                  | 4712                  |
| Soil          | 27920                 | 25130                 | 27910                 | 25120                 |
| Sediment      | 7554                  | 18260                 | 7559                  | 18270                 |
| Groundwater   | 4614                  | 4153                  | 4612                  | 4151                  |
| Coastal water | 0.16                  | 0.39                  | 0.16                  | 0.39                  |

The calculated concentrations in the air compartment are relatively lower than those observed in the environment (Table VI). By means of an equation published by Mackay and coworkers (23), the concentrations in rain for region 3 were calculated from the air concentrations obtained with CHEMFRANCE. For scenarios A and B, the calculated rain concentrations are lower than those observed in Paris between March and December 1991 (Table VI) (24). At the opposite, for scenarios C and D, the simulated values are relatively closer to the actual data (Table VI). The calculated soil concentrations are lower than those expected after an application (i.e.; 0.5 µg/g) at the rate recommended by the EC. However, these results agree with this observed in a soil historically treated (i.e.; 0.025 µg/g) but before any further treatment (25).

The mass distribution obtained in the soil compartment (i.e.; 96.7% in the solid components and 3.3% in the water and air contained in the pore of the soil) is near this reported by Tasli (26). Indeed, in her study, 96% of the amount included in the soil is sorbed in the soil solids and 4% is dissolved in the soil water. In the same way, the results obtained for the sediment compartment indicate that 89% of the atrazine is sorbed and 11% dissolved. Huckins and coworkers (27) in a microcosm study, found a slightly smaller percentage (i.e.; 86.1%) although the organic carbon content of their studied sediment (i.e.; 7.28%) was higher than this considered in CHEMFRANCE (i.e.; 4%). In the water compartment, ca. 99.9% of the total atrazine content of this compartment is dissolved in pure water, 0.07% is sorbed in the suspended sediment, and 0.002% is accumulated in biota (i.e.; fish). Field studies gave similar results (i.e.; 99.5% and 0.5% of atrazine in the dissolved and particulate river matter, respectively) (28). The mass distributions obtained for the air compartment cannot be compared with experimental results. In the air compartment, atrazine is principally present in pure air (i.e.; 89.2%), and only a low quantity is bound to the air particles (i.e.; 10.8%).

Table VI. Comparison between field and predicted (scenarios A-D) concentrations (ng/l)

| Date, region of CHEMFRANCE                    | Actual concentration       |           | Ref. | Predicted concentration |                       |                       |                       |
|---|----------------------------|-----------|------|-------------------------|-----------------------|-----------------------|-----------------------|
|   | mean <sup>s</sup>          | maximum   |      | A                       | B                     | C                     | D                     |
| <b>Surface water compartment</b>              |                            |           |      |                         |                       |                       |                       |
| 4 rivers (17/04/1990 to 24/07/1990), region 1 | 2149 (n = 40)              | 7460      | 29   | 1949                    | 4711                  | 1950                  | 4712                  |
| 4 rivers (17/04/1991 to 24/07/1991), region 1 | 984 (n = 26)               | 3050      | 29   | 1949                    | 4711                  | 1950                  | 4712                  |
| 12 rivers (04/1991 to 11/1991), region 1      | 210 (n = 75)               | 750       | 29   | 1949                    | 4711                  | 1950                  | 4712                  |
| 1 river (17/04/1990 to 24/07/1990), region 2  | 1747 (n = 10)              | 6030      | 29   | 1236                    | 2989                  | 1237                  | 2989                  |
| 1 river (17/04/1991 to 24/07/1991), region 2  | 788 (n = 7)                | 2500      | 29   | 1236                    | 2989                  | 1237                  | 2989                  |
| 2 rivers (04/1991 to 11/1991), region 2       | 1206 (n = 8)               | 2400      | 29   | 1236                    | 2989                  | 1237                  | 2989                  |
| Seine barrage, region 3                       | 98 (n = 46)                | 260       | 29   | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| 9 rivers (04/07/1989 to 28/08/1989), region 3 | 360 (n = 106)              | 1550      | 29   | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| Seine river (July 1987), region 3             | 810 (n = 1)                |           | 30   | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| Marne river (July 1987), region 3             | 60 (n = 1)                 |           | 30   | 195.1                   | 951.2                 | 195.5                 | 951.6                 |
| 1 river (18/03/1991 to 16/03/1992), region 4  | 159 (n = 51)               | 340       | 29   | 375.8                   | 1308                  | 376.4                 | 1309                  |
| 29 rivers (summer 1990), region 9             | 165 (n = 47)               | 710       | 29   | 593.5                   | 2066                  | 594.4                 | 2067                  |
| Rhone delta (10/1990 to 03/1994), region 9    | 17-386 <sup>£</sup>        |           | 31   | 593.5                   | 2066                  | 594.4                 | 2067                  |
| Rhone river, 7 stations (10/1990), region 9   | 12-33 <sup>£</sup>         |           | 32   | 593.5                   | 2066                  | 594.4                 | 2067                  |
| 15 rivers (summer 1990), region 11            | 277 (n = 24)               | 1000      | 29   | 460.6                   | 1856                  | 462.0                 | 1858                  |
| <b>Groundwater compartment</b>                |                            |           |      |                         |                       |                       |                       |
| (July 1987), region 1                         | 60 (n = 1)                 |           | 30   | 4614                    | 4153                  | 4612                  | 4151                  |
| (July 1987), region 3                         | 95 (n = 1)                 |           | 30   | 565.7                   | 509.1                 | 565.4                 | 508.8                 |
| (November 1987), region 3                     | 225 (n = 1)                |           | 30   | 565.7                   | 509.1                 | 565.4                 | 508.8                 |
| (07/1990 to 09/1990), region 4                | 182 (n = 39)               | 856       | 29   | 774.4                   | 697.0                 | 774.2                 | 696.7                 |
| (November 1987), region 4                     | 90 (n = 1)                 |           | 30   | 774.4                   | 697.0                 | 774.2                 | 696.7                 |
| (summer 1990), region 9                       | 174 (n = 18)               | 630       | 29   | 1093                    | 984.0                 | 1093                  | 983.6                 |
| (summer 1990), region 10                      | 238 (n = 6)                | 430       | 29   | 1408                    | 1267                  | 1407                  | 1266                  |
| (summer 1990), region 11                      | 86.6 (n = 3)               | 130       | 29   | 1276                    | 1148                  | 1275                  | 1148                  |
| (July 1987), region 11                        | 110 (n = 1)                |           | 30   | 1276                    | 1148                  | 1275                  | 1148                  |
| (April 1989), region 12                       | 200 (n = 1)                |           | 30   | 1937                    | 1743                  | 1936                  | 1742                  |
| <b>Air compartment</b>                        |                            |           |      |                         |                       |                       |                       |
| air (04/1991 to 06/1991), region 3            | 5 10 <sup>-5</sup> (n = ?) | not given | 24   | 3.66 10 <sup>-8</sup>   | 3.40 10 <sup>-8</sup> | 6.38 10 <sup>-6</sup> | 5.74 10 <sup>-6</sup> |
| rain (03/1991 to 12/1991), region 3           | <5-140 <sup>£</sup>        |           | 24   | 0.08                    | 0.07                  | 13.98                 | 12.56                 |

<sup>s</sup>n = number of values; <sup>£</sup> range



The correct use of a model requires to perform a sensitivity analysis showing how variations in the different model input parameters affect the outputs. In our study, a survey of the literature showed that the half-life of atrazine in soil widely varied according to the experimental and environmental conditions (Table VII). Similar observations were reported by Solomon and coworkers (33) for the half-lives of atrazine in water and sediments.

Table VII. Soil atrazine half-life ( $T_{1/2}$ ) values obtained in literature

| Experimental and environmental conditions  | Ref. | $T_{1/2}$ (d) |
|--|------|---------------|
| <b>Laboratory</b>  |      |               |
| Loam, clay-sandy <sup>*</sup> ; T°: 5; SMC: 7%                                     | 34   | 240.5         |
| Loam, clay-sandy <sup>*</sup> ; T°: 15; SMC: 7%                                    | 34   | 122.6         |
| Loam, clay-sandy <sup>*</sup> ; T°: 25; SMC: 7%                                    | 34   | 62.3          |
| Loam, clay-sandy <sup>*</sup> ; T°: 32; SMC: 7%                                    | 34   | 51.0          |
| Loam, clay-sandy <sup>*</sup> ; T°: 25; SMC: 4%                                    | 34   | 99.5          |
| Loam, clay-sandy <sup>*</sup> ; T°: 25; SMC: 10%                                   | 34   | 58.5          |
| Loam, clay-sandy <sup>*</sup> ; T°: 25; SMC: 13%                                   | 34   | 57.2          |
| Loam-sandy <sup>§</sup> ; T°: 5; SMC: 7%   | 34   | 271.0         |
| Loam-sandy <sup>§</sup> ; T°: 15; SMC: 7%  | 34   | 111.8         |
| Loam-sandy <sup>§</sup> ; T°: 25; SMC: 7%  | 34   | 65.7          |
| Loam-sandy <sup>§</sup> ; T°: 32; SMC: 7%  | 34   | 49.6          |
| Loam-sandy <sup>§</sup> ; T°: 25; SMC: 4%  | 34   | 78.5          |
| Loam-sandy <sup>§</sup> ; T°: 25; SMC: 10%   | 34   | 56.9          |
| Loamy <sup>£</sup> ; T°: 4; SMC: 17%   | 34   | 215.8         |
| Loamy <sup>£</sup> ; T°: 15; SMC: 17%  | 34   | 81.7          |
| Loamy <sup>£</sup> ; T°: 24; SMC: 17%  | 34   | 49.8          |
| Loamy <sup>£</sup> ; T°: 34; SMC: 17%  | 34   | 28.8          |
| Loamy <sup>£</sup> ; T°: 24; SMC: 5%   | 34   | 265.2         |
| Loamy <sup>£</sup> ; T°: 24; SMC: 10%  | 34   | 96.5          |
| Loamy <sup>£</sup> ; T°: 24; SMC: 15%  | 34   | 60.9          |
| Loamy <sup>£</sup> ; T°: 24; SMC: 20%  | 34   | 55.3          |
| pH: 4.8; OM: 1.8; Cy: 10.5; St: 12.4; Sd: 77.1; T°: 22; H: 40; C <sub>0</sub> : 10 | 35   | 53            |
| pH: 6.5; OM: 3.8; Cy: 8.9; St: 7.3; Sd: 83.8; T°: 22; H: 40; C <sub>0</sub> : 10   | 35   | 113           |
| Fine sandy medium sand; pH: 7.7; OC: 0.1; H: 30; C <sub>0</sub> : 3                | 8    | 90            |
| Very coarse sand-fine sand; pH: 7.3; OC: 4.3; H: 42; C <sub>0</sub> : 3            | 8    | 43            |
| Very coarse sandy fine sand; pH: 7.4; OC: 1.5; H: 39; C <sub>0</sub> : 3           | 8    | 50            |
| Fine sand-medium sand; pH: 7.7; OC: 0.6; H: 38; C <sub>0</sub> : 3                 | 8    | 47            |
| Silt loam <sup>+</sup> ; H: 80; AR: 2.20   | 36   | 20/21         |
| Sandy clay loam <sup>§</sup> ; pH: 5.7; OM: 2.7; Cy: 22; St: 26; Sd: 52; W: 0.224  | 37   | 41            |
| Loam <sup>§</sup> ; pH: 6.2; OM: 2.7; Cy: 26; St: 34; Sd: 40; W: 0.293             | 37   | 69            |
| Sandy clay loam <sup>§</sup> ; pH: 6.2; OM: 1.7; Cy: 28; St: 26; Sd: 46; W: 0.269  | 37   | 69            |
| Sandy clay loam <sup>§</sup> ; pH: 6.2; OM: 1.6; Cy: 26; St: 28; Sd: 46; W: 0.245  | 37   | 231           |
| Sandy clay loam <sup>§</sup> ; pH: 6.2; OM: 1.6; Cy: 26; St: 28; Sd: 46; W: 0.545  | 37   | 87            |
| Fine loam; pH: 5.9; OM: 1.1; Cy: 6; SMC: 15.8%; T°: 21; H: 80; C <sub>0</sub> : 5  | 38   | 59.6          |
| Sandy loam; pH: 5.5; OC: 0.91; Cy: 17; St: 14; Sd: 69; H: 60; W: 0.17              | 39   | 330           |
| Silty loam; pH: 6.4; OC: 0.91; Cy: 32; St: 52; Sd: 16; H: 100; W: 0.5              | 39   | 385           |
| Sandy loam <sup>£</sup> ; SMC: 3%; C <sub>0</sub> : 10                             | 40   | 36.7          |
| Sandy loam <sup>£</sup> ; SMC: 6%; C <sub>0</sub> : 10                             | 40   | 34.6          |
| Sandy loam <sup>£</sup> ; SMC: 12%; C <sub>0</sub> : 10                            | 40   | 26.6          |
| Sandy loam <sup>£</sup> ; SMC: 6%; C <sub>0</sub> : 1                              | 40   | 64.0          |
| Loamy-sand <sup>£</sup> ; SMC: 1%; C <sub>0</sub> : 10                             | 40   | 32.1          |
| Loamy-sand <sup>£</sup> ; SMC: 2%; C <sub>0</sub> : 10                             | 40   | 26.5          |
| Loamy-sand <sup>£</sup> ; SMC: 4%; C <sub>0</sub> : 10                             | 40   | 22.9          |
| Loamy-sand <sup>£</sup> ; SMC: 2%; C <sub>0</sub> : 1                              | 40   | 51.9          |

Table VII (continued)

| Experimental and environmental conditions              | Ref. | T <sub>1/2</sub> (d) |
|--|------|----------------------|
| <b>Field</b>   |      |                      |
| Silt loam; pH: 5.1-5.5; AR: 2.8; No-till corn crop     | 41   | 60/72                |
| Riparian zone; AR: 2.2; Depth: 90 cm; Poplar tree plot | 15   | 16                   |
| Silt loam <sup>g</sup> ; AR: 3.2; Depth: 3 cm          | 36   | 14                   |
| Fine sandy loam <sup>hh</sup> ; pH: 5.0; AR: 1.12      | 42   | 9/15                 |
| Fine sandy loam <sup>hh</sup> ; pH: 6.0; AR: 1.12      | 42   | 12/19                |
| Fine sandy loam <sup>hh</sup> ; pH: 7.0; AR: 1.12      | 42   | 15/23                |
| Fine sandy loam <sup>hh</sup> ; pH: 5.0; AR: 2.24      | 42   | 16/20                |
| Fine sandy loam <sup>hh</sup> ; pH: 6.0; AR: 2.24      | 42   | 18/23                |
| Fine sandy loam <sup>hh</sup> ; pH: 7.0; AR: 2.24      | 42   | 20/26                |
| Fine sandy loam <sup>hh</sup> ; pH: 5.0; AR: 3.36      | 42   | 17/22                |
| Fine sandy loam <sup>hh</sup> ; pH: 6.0; AR: 3.36      | 42   | 19/25                |
| Fine sandy loam <sup>hh</sup> ; pH: 7.0; AR: 3.36      | 42   | 21/28                |
| Silt loam <sup>kk</sup> ; pH: 5.0; AR: 1.12            | 42   | 18/19                |
| Silt loam <sup>kk</sup> ; pH: 6.0; AR: 1.12            | 42   | 27/28                |
| Silt loam <sup>kk</sup> ; pH: 7.0; AR: 1.12            | 42   | 36/37                |
| Silt loam <sup>kk</sup> ; pH: 5.0; AR: 2.24            | 42   | 22/27                |
| Silt loam <sup>kk</sup> ; pH: 6.0; AR: 2.24            | 42   | 28/34                |
| Silt loam <sup>kk</sup> ; pH: 7.0; AR: 2.24            | 42   | 36/41                |
| Silt loam <sup>kk</sup> ; pH: 5.0; AR: 3.36            | 42   | 25/27                |
| Silt loam <sup>kk</sup> ; pH: 6.0; AR: 3.36            | 42   | 31/33                |
| Silt loam <sup>kk</sup> ; pH: 7.0; AR: 3.36            | 42   | 37/39                |
| Sandy loam <sup>ll</sup> ; pH: 5.0; AR: 1.12           | 42   | 6                    |
| Sandy loam <sup>ll</sup> ; pH: 6.0; AR: 1.12           | 42   | 8                    |
| Sandy loam <sup>ll</sup> ; pH: 7.0; AR: 1.12           | 42   | 10                   |
| Sandy loam <sup>ll</sup> ; pH: 5.0; AR: 2.24           | 42   | 7/10                 |
| Sandy loam <sup>ll</sup> ; pH: 6.0; AR: 2.24           | 42   | 9/12                 |
| Sandy loam <sup>ll</sup> ; pH: 7.0; AR: 2.24           | 42   | 10/14                |
| Sandy loam <sup>ll</sup> ; pH: 5.0; AR: 3.36           | 42   | 8/10                 |
| Sandy loam <sup>ll</sup> ; pH: 6.0; AR: 3.36           | 42   | 9/12                 |
| Sandy loam <sup>ll</sup> ; pH: 7.0; AR: 3.36           | 42   | 11/13                |
| Riparian zone; AR: 1; Depth: 2 cm                      | 8    | 48                   |

<sup>\*</sup>pH: 8.6; OM: 1.22; Field capacity at -5 kPa: 17.4%; Bulk density: 1.3 g/cm<sup>3</sup>

<sup>g</sup>pH: 8.5; OM: 1.23; Field capacity at -5 kPa: 10.6%; Bulk density: 1.2 g/cm<sup>3</sup>

<sup>h</sup>pH: 7.8; OM: 0.93; Field capacity at -5 kPa: 20.4%; Bulk density: 1.4 g/cm<sup>3</sup>

<sup>i</sup>pH: 5.3-5.6; OC: 0.29-0.50; Cy: 13.8-15.7; St: 76.2-80.7; Sd: 5.5-8.2; T°: 25

<sup>j</sup>T°: 25; AR: 1.86; C<sub>0</sub>: 5

<sup>k</sup>pH: 6.05; OC: 0.43; CEC: 5.95 cmol/kg; Cy: 7.05; St: 25.84; Sd: 67.11; T°: 25

<sup>l</sup>pH: 6.30; OC: 0.33; CEC: 0.45 cmol/kg; Cy: 4.47; St: 13.52; Sd: 82.01; T°: 25

<sup>m</sup>pH: 5.3-5.5; OC: 0.43-0.50; Cy: 14.8-15.6; St: 78.4-80.7; Sd: 3.7-6.5

<sup>hh</sup>OM: 0.9; Cy: 6; St: 31; Sd: 63; CEC: 3.8 meq/100 g; No-till corn plot; Depth: 9 cm

<sup>kk</sup>OM: 1.6; Cy: 24; St: 58; Sd: 18; CEC: 8.3 meq/100 g; No-till corn plot; Depth: 9 cm

<sup>ll</sup>OM: 1.0; Cy: 12; St: 24; Sd: 64; CEC: 5.4 meq/100 g; No-till corn plot; Depth: 9 cm

T°: Temperature (°C); SMC: Soil moisture content; OM: % Organic matter; Cy: % Clay; St: % Silt; Sd: % Sand; H: % of water holding capacity; C<sub>0</sub>: Nominal initial concentration (mg/kg); OC: % Organic carbon; AR: application rate (kg/ha); W: Water content (g/g); CEC: Cation exchange capacity

Simulation experiments were performed in order to test the influence of soil half-life on the outputs calculated by CHEMFRANCE. Region 1 was selected due to its agricultural interest and scenario D was adopted since it represents a summary of the other scenarios. Five runs were performed with a starting value of 6 d and an increment of 95 d. The simulation results are displayed in Figure 2 and Tables VIII and IX. Figure 2 shows that the half-life in soil principally influences water and soil reactions and the leaching and runoff processes.




Figure 2. Influence of soil half-life on CHEMFRANCE outputs

It is obvious that changes in soil half-life values induce other variations than those displayed in Figure 2. Thus, Table VIII clearly shows that soil and water runoff, leaching, and soil volatilization are the most subject to variations. Water runoff and leaching are the most significant due to their magnitude. They induce an increase of the groundwater and surface water concentrations in atrazine (Table IX).

Our study shows that the results obtained with CHEMFRANCE for atrazine are in accordance with those obtained in laboratory or field experiments. This underlines the usefulness of the environmental fate models for confirming results of field and laboratory studies but also to provide a comprehensive summary of the processes involved in the environmental fate of chemicals. This is particularly interesting for atrazine since this herbicide is of environmental concern in France.

An interesting characteristic of CHEMFRANCE is that it allows the user to quickly test several hypotheses through different scenarios. Sensitivity and uncertainty analyses can also be performed in order to test the influence of all input parameters and assumptions on the outputs.

Table VIII. Influence of the soil half-life on the removal and transfer processes

| Multiplicative factor* | Process                     | Range (mol/h) <sup>§</sup>                   |
|------------------------|-----------------------------|--|
| 0.51                   | Soil reaction               | 4.283 10 <sup>2</sup> -2.204 10 <sup>2</sup> |
| 1.04                   | Rain deposition on water    | 3.41 10 <sup>-2</sup> -3.56 10 <sup>-2</sup> |
| 1.04                   | Rain deposition on soil     | 4.60-4.80                                    |
| 1.04                   | Dry deposition on soil      | 5.01 10 <sup>-3</sup> -5.23 10 <sup>-3</sup> |
| 1.04                   | Wet deposition on water     | 5.66 10 <sup>-5</sup> -5.91 10 <sup>-5</sup> |
| 1.04                   | Wet deposition on soil      | 7.63 10 <sup>-3</sup> -7.97 10 <sup>-3</sup> |
| 1.04                   | Air advection               | 3.6 10 <sup>-2</sup> -3.76 10 <sup>-2</sup>  |
| 1.04                   | Diffusion air/water         | 8.52 10 <sup>-5</sup> -8.90 10 <sup>-5</sup> |
| 1.04                   | Diffusion air/soil          | 3.78 10 <sup>-3</sup> -3.95 10 <sup>-3</sup> |
| 1.05                   | Dry deposition on water     | 3.72 10 <sup>-5</sup> -3.89 10 <sup>-5</sup> |
| 1.05                   | Transfer to higher altitude | 4.44 10 <sup>-5</sup> -4.64 10 <sup>-5</sup> |
| 3.01                   | Sediment reaction           | 0.94-2.83                                    |
| 3.02                   | Sediment burial             | 1.93 10 <sup>-3</sup> -5.82 10 <sup>-3</sup> |
| 3.02                   | Water volatilization        | 1 10 <sup>-3</sup> -3.02 10 <sup>-3</sup>    |
| 3.02                   | Sediment desorption         | 1.39-4.20                                    |
| 3.02                   | Sediment resuspension       | 6.42 10 <sup>-4</sup> -1.94 10 <sup>-3</sup> |
| 3.02                   | Water reaction              | 49.6-149.9                                   |
| 3.03                   | Sediment adsorption         | 2.31-6.99                                    |
| 3.03                   | Water advection             | 1.07-3.24                                    |
| 3.03                   | Sediment deposition         | 1.54 10 <sup>-2</sup> -4.67 10 <sup>-2</sup> |
| 33.07                  | Soil runoff                 | 2.57 10 <sup>-2</sup> -0.85                  |
| 33.14                  | Leaching                    | 3.22-106.7                                   |
| 33.14                  | Water runoff                | 3.22-106.7                                   |
| 33.18                  | Soil volatilization         | 6.63 10 <sup>-3</sup> -0.22                  |

\*Ratio between the rate of the process at the highest and this at the lowest half-life value. Only the processes with a ratio different from one have been included.

§Obtained with the lowest and the highest half-life, respectively.

Table IX. Influence of the soil half-life on the concentration in atrazine

| Compartment   | Multiplicative factor* | Concentration range (ng/l) <sup>§</sup>        |
|---------------|------------------------|--|
| Air           | 1.04                   | 2.060 10 <sup>-5</sup> -2.150 10 <sup>-5</sup> |
| Water         | 3.02                   | 3.158 10 <sup>3</sup> -9.540 10 <sup>3</sup>   |
| Soil          | 33.11                  | 2.848 10 <sup>3</sup> -9.431 10 <sup>4</sup>   |
| Sediment      | 3.02                   | 1.224 10 <sup>4</sup> -3.697 10 <sup>4</sup>   |
| Groundwater   | 33.10                  | 4.707 10 <sup>2</sup> -1.558 10 <sup>4</sup>   |
| Coastal water | 3.04                   | 0.26-0.79                                      |

\*Ratio between the concentration at the highest and this at the lowest half-life value.

§Concentration obtained with the lowest and the highest half-life, respectively.

### Acknowledgment

This study was granted by the French Ministry of the Environment.

## Literature Cited

1. *The Multi-Media Fate Model: A Vital Tool for Predicting the Fate of Chemicals*; Cowan, C.E.; Mackay, D.; Feijtel, T.C.J.; van de Meent, D.; Di Guardo, A.; Davies, J.; Mackay, N., Eds.; SETAC Press, Pensacola, FL, 1995.
2. Mackay, D.; Paterson, S.; Cheung, B.; Neely, W.B. *Chemosphere* 1985, 14, 335-374.
3. Mackay, D. *Multimedia Environmental Models. The Fugacity Approach*. Lewis Publishers, Chelsea, 1991, p. 257.
4. Mackay, D.; Paterson, S.; Shiu, W.Y. *Chemosphere* 1992, 24, 695-717.
5. Devillers, J.; Bintein, S.; Karcher, W. *Chemosphere* 1995, 30, 457-476.
6. Montiel, A. *Etude bibliographique. Utilisation des Herbicides et des Triazines en Particulier*. Ministère de la Santé, DDRASS Ile de France, 1989, p. 273.
7. Francioso, O.; Bak, E.; Rossi, N.; Sequi, P. *Sci. Total Environ.* 1992, 123/124, 503-512.
8. Bacci, E.; Renzoni, A.; Gaggi, C.; Calamari, D.; Franchi, A.; Vighi, M.; Severi, A. *Agric. Ecosys. Environ.* 1989, 27, 513-522.
9. Wauchope, R.D.; Buttler, T.M.; Hornsby, A.G.; Augustijn-Beckers, P.W.M.; Burt, J.P. *Rev. Environ. Contam. Toxicol.* 1992, 123, 1-164.
10. *Statistique Agricole Annuelle. Résultats Provisoires 1991*. Ministère de l'Agriculture et de la Forêt, France, 1992.
11. *Index Phytosanitaire*. Association de Coordination Technique Agricole, Paris, France, 1992.
12. *Statistique Agricole Annuelle. Résultats 1990*. Ministère de l'Agriculture et de la Forêt, France, 1991.
13. Persoone, G.; Janssen, C.R.; van Steertegem, M. *Extensive and Updated Literature Study of the Toxicity and Ecotoxicity of Atrazine*. Commission of the European Communities, 1991.
14. Mirgain, I.; Schenck, C.; Monteil, H. *Environ. Technol.* 1993, 14, 741-750.
15. Paterson, K.G.; Schnoor, J.L. *Water Environ. Res.* 1992, 64, 274-283.
16. Di Muccio, A.; Chirico, M.; Dommarco, R.; Funari, E.; Musmeci, L.; Santilio, A.; Vergori, F.; Zapponi, G.; Giuliano, G.; Sparacino, A.C. *Intern. J. Environ. Anal. Chem.* 1990, 38, 211-220.
17. Buser, H.R. *Environ. Sci. Technol.* 1990, 24, 1049-1058.
18. Schiavon, M. *Ecotoxicol. Environ. Safety* 1988, 15, 46-54.
19. Schiavon, M.; Portal, J.M.; Andreux, F. *Agronomie* 1992, 12, 129-139.
20. Wauchope, R.D. *J. Environ. Qual.* 1978, 7, 459-472.
21. Scheunert, I.; Qiao, Z.; Korte, F. *J. Environ. Sci. Health* 1986, B21, 457-485.
22. Bester, K.; Hühnerfuss, H. *Marine Poll. Bull.* 1993, 26, 423-427.
23. Mackay, D.; Paterson, S.; Schroeder, W.H. *Environ. Sci. Technol.* 1986, 20, 810-816.
24. Chevreuil, M.; Garmouma, M. *Chemosphere* 1993, 27, 1605-1608.
25. Huart, B.; Pailler, F.M.; Trique, B.; Ricordel, I. *Ann. Fals. Exp. Chim.* 1993, 86, 75-86
26. Tasli, S. *Distribution de l'Atrazine entre le Sol, l'Eau et les Plantes en Culture de Maïs*. DEA d'Agrochimie, University of Grenoble, 1992, p. 23.
27. Huckins, J.N.; Petty, J.D.; England, D.C. *Chemosphere* 1986, 15, 563-588.
28. Pereira, W.E.; Rostad, C.E. *Environ. Sci. Technol.* 1990, 24, 1400-1406.
29. Devillers, J.; Bintein, S.; Domine, D. *Atrazine in France*. Report CTIS #92011, 1992, p. 42.
30. Legrand, M.F.; Costentin, E.; Bruchet, A. *Water Supply* 1992, 10, 51-61.
31. Readman, J.W.; Albanis, T.A.; Barcelo, D.; Galassi, S.; Tronczynski, J.; Gabrielides, G.P. *Marine Poll. Bull.* 1993, 26, 613-619.
32. Tronczynski, J.; Munsch, C.; Durand, G.; Barcelo, D. *Sci. Total Environ.* 1993, 132, 327-337.

33. Solomon, K.R.; Baker, D.B.; Richards, R.P.; Dixon, K.R.; Klaine, S.J.; La Point, T.W.; Kendall, R.J.; Giddings, J.M.; Giesy, J.P.; Hall, L.W.; Williams, W.M. *Environ. Sci. Technol.* **1996**, *15*, 31-76.
34. Obrador, A.; Lechon, Y.; Tadeo, J.L. *Pestic. Sci.* **1993**, *37*, 301-308.
35. Burkhard, N.; Guth, J.A. *Pestic. Sci.* **1981**, *12*, 45-52.
36. Winkelmann, D.A.; Klaine, S.J. *Environ. Toxicol. Chem.* **1991**, *10*, 335-345.
37. Kruger, E.L.; Somasundaram, L.; Kanwar, R.S.; Coats, J.R. *Environ. Toxicol. Chem.* **1993**, *12*, 1959-1967.
38. Blumhorst, M.R.; Weber, J.B. *J. Agric. Food Chem.* **1992**, *40*, 894-897.
39. Jones, T.W.; Kemp, W.M.; Stevenson, J.C.; Means, J.C. *J. Environ. Qual.* **1982**, *11*, 632-638.
40. Singh, G.; Spencer, W.F.; Cliath, M.M.; van Genuchten, M.T. *Environ. Pollut.* **1990**, *66*, 253-262.
41. Helling, C.S.; Zhuang, W.; Gish, T.J.; Coffman, C.B.; Isensee, A.R.; Kearney, P.C.; Hoagland, D.R.; Woodward, M.D. *Chemosphere* **1988**, *17*, 175-187.
42. Hiltbold, A.E.; Buchanan, G.A. *Weed Sci.* **1977**, *25*, 515-520.

# ARTICLE

# VII

## COMPARISON OF BCF MODELS BASED ON LOG P

J. Devillers,\* S. Bintein, and D. Domine

CTIS, 21 rue de la Bannière, 69003 Lyon (France)

### ABSTRACT

Seven linear and nonlinear BCF models based on log P have been compared in order to estimate their accuracy, predictive power, and domain of application. This comparison was performed from a data set constituted of 436 experimental BCF values recorded for 227 chemicals. We showed that for chemicals with  $\log P < 6$ , the different models yielded equivalent results. At the opposite, for highly hydrophobic chemicals ( $\log P > 6$ ), the bilinear model  $\log \text{BCF} = 0.91 \log P - 1.975 \log (6.8 \cdot 10^{-7} P + 1) - 0.786$  ( $n = 154$ ;  $r = 0.95$ ;  $s = 0.347$ ;  $F = 463.5$ ) was superior to the other studied models.

### INTRODUCTION

Numerous attempts have been made for modeling the accumulation of organic chemicals by fish since the experimental determination of this important biological activity is a time-consuming and costly exercise. The most widely used method consists in correlating laboratory bioconcentration factors (BCFs) determined in fish and *n*-octanol/water partition coefficients ( $\log P$ ). Bioconcentration can be defined as the result of the uptake, distribution, and elimination of a chemical in a fish due to water-borne exposure (Barron, 1990), whereas bioaccumulation includes all routes of exposure, and biomagnification deals with accumulation via food chains. Numerous test guidelines for the experimental determination of BCF in fish are available (e.g., OECD, ASTM). BCF is determined as the concentration of the chemical in whole fish at steady state divided by the concentration of the chemical in water during the exposure period and/or as the ratio between the uptake ( $k_1$ ) and depuration ( $k_2$ ) rate constants assuming first order kinetics. The accumulation of an organic compound in a fish basically depends on its hydrophobicity. Therefore,  $\log P$  is the physicochemical descriptor of choice for modeling bioconcentration. As a result, numerous BCF QSAR equations derived from  $\log P$  are available in the literature and there is a need for the evaluation of their domain of application. Under these conditions, the aim of this study was to compare different BCF/ $\log P$  models in order to estimate their accuracy, predictive power, and domain of application.

---

\* Author to whom all correspondence should be addressed.



## MODEL SELECTION

The selection of the BCF/log P models was principally based on their occurrence of citation in the scientific literature and official reports (e.g., ECETOC, 1995; EEC, 1995). Thus, seven equations were selected. Their main characteristics are given below.

Equation no. 1 (Veith *et al.*, 1979)

$$\log \text{BCF} = 0.85 \log P - 0.70$$

n = 55      r = 0.95      s = not given      F = not given

Fathead minnows (*Pimephales promelas*) were used as test species.

Equation no. 2 (Veith *et al.*, 1980)

$$\log \text{BCF} = 0.76 \log P - 0.23$$

n = 84      r (?) = 0.95      s = not given      F = not given

Bioconcentration data were measured on *Pimephales promelas* and *Lepomis macrochirus*.

Equation no. 3 (Mackay, 1982)

$$\log \text{BCF} = 1.00 \log P - 1.32$$

n = 44 (?)      r = 0.97      s = 0.25      F = not given

The model was designed from bioconcentration data relating to several fish species.

Equation no. 4 (Isnard and Lambert, 1988)

$$\log \text{BCF} = 0.80 \log P - 0.52$$

n = 107      r = 0.904      s = 0.51      F = not given

The model was obtained from bioconcentration data relating to several fish species.

Equation no. 5 (Nendza, 1991)

$$\log \text{BCF} = 0.99 \log P - 1.47 \log (4.97 \cdot 10^{-8} P + 1) + 0.0135$$

n = 132

The above empirical model (without statistical parameters) was designed from bioconcentration data relating to several fish species and expressed as percentage of lipid content.

Equation no. 6 (Bintein *et al.*, 1993)

$$\log \text{BCF} = 0.91 \log P - 1.975 \log (6.8 \cdot 10^{-7} P + 1) - 0.786$$

n = 154      r = 0.95      s = 0.347      F = 463.51

The equation was derived from bioconcentration data measured on five families of freshwater fish (i.e., Salmonidae, Cyprinidae, Centrarchidae, Poeciliidae, Cyprinodontidae).

Equation no. 7 (Connell and Hawker, 1988)

$$\log \text{BCF} = 6.9 \cdot 10^{-3} (\log P)^4 - 0.185 (\log P)^3 + 1.55 (\log P)^2 - 4.18 \log P + 4.79$$

The model was designed from bioconcentration data relating to several fish species. No statistical parameters were given.

## BCF AND LOG P VALUES DATABASE CONSTITUTION

### *Experimental BCF Data*

To properly compare the accuracy, the predictive power, and the domain of application of the seven selected models, it would have been necessary to use new experimental BCF data specifically measured for this comparative study. However, for reasons of time and cost, this solution could not obviously be envisioned and only data retrieved from the literature were employed. This induced methodological problems. Indeed, there are relatively few experimental bioconcentration data towards fish and they have widely been used to derive QSAR models including those compared in this study. Under these conditions, in the present study, we can consider that a bias was systematically introduced when one of the BCF values used for the comparison exercise was included in the training set(s) of the studied models. However, this bias became lesser if this value was used in the derivation of all the seven models. In the same way, it is obvious that the models designed from a small number of compounds were penalized compared with those derived from large numbers of molecules. Indeed, the values selected in our study were more likely to be in the training sets of models derived from a large number of molecules than in those for which the training set was of a reduced size. To overcome these problems, rather than selecting a sole BCF value per molecule, when possible, a set of coherent experimental BCF data was compiled in order to provide bioconcentration profiles for a given molecule. This allowed to partly solve the problems stressed above and also to take into account the influence of the experimental protocols.

The experimental BCF values used in our comparative study are listed in Tables I and II. They only concern freshwater fish. They do not originate from data compilations but were all retrieved from the original publications. Only data obtained after a steady-state was reached (i.e., ratio between concentration in fish and concentration in water) or by the kinetic method (i.e., ratio between first-order uptake and elimination rate constants) were retained. Data for which contamination by food and/or presence of adsorbent was observed were not selected. The apparent BCF values were selected for the experiments using radiolabelled compounds. Data expressed as lipid content are indicated in Tables I and II. Table I only contains the experimental results obtained from tests performed under flow-through conditions (342 values) while Table II lists those obtained under static and semi-static conditions (94 values). The lipid content values were selected from the values given or recommended by the authors, otherwise the generally accepted estimate of 5 % was adopted (e.g., Connell and Hawker, 1988). In many instances, confirmation of the lipid content was obtained directly from the authors. When this compilation was performed, the BCF data being obviously out-of-line with those generally found in the literature were removed. For the sake of homogeneity, if one or more values appeared out-of-line in a publication, all the data contained in this paper were removed.

### *Compilation of log P Values*

In order to minimize the influence of the *n*-octanol/water partition coefficient values on the simulation results, the selection of log P data was based on the following criteria. First of all, direct methods involving normalized protocols were privileged. In addition, when several data were available, the selection was performed in order to obtain a coherent database for all the studied molecules. Indeed, a coherency between the selected log P values and the chemical structures of the molecules was maintained. Last, as far as possible, the log P selections made by the different authors who derived the models to be compared were also taken into account. The selected log P values for the 227 studied chemicals are shown in Tables I and II.

## RESULTS AND DISCUSSION

The obtained results are listed in Tables I to III and displayed in Figures 1 to 14. Thus, in the last column of Tables I and II, the seven models have been classified according to their decreasing predictive power estimated on the basis of the residuals. The first cited model (on the left-hand side) is the one which yields the lowest residual value (absolute value). At the opposite, the last model cited (on the right-hand side) presents the highest residual value. Thus, for example, for acenaphthene (i.e., first chemical of Table I), our comparative study shows that the most accurate model is that of Nendza (1991), followed by that designed by Mackay (1982), that of Isnard and Lambert (1988), Veith *et al.* (1979), Connell and Hawker (1988), Veith *et al.* (1980) and finally, the least accurate model to estimate the BCF of acenaphthene is Eq. (6) (Bintein *et al.*, 1993). The tendency of each model to under- or overestimate the BCF of the studied molecules is a critical feature to inspect. Thus, in Tables I and II, for each entry, the model numbers for which the predicted value is superior to the observed BCF are underlined. The analysis of the classifications obtained for the 227 studied chemicals (Tables I and II) shows that for chemicals with log P values inferior to 6, the seven models are broadly equivalent. At the opposite, for molecules presenting a log P value superior to 6, the nonlinear equations (6) and (7) bear better predictive performances. Among them, the bilinear model of Bintein *et al.* (1993) is the most often classified first. For some chemicals (e.g., permethrin), the use of different experimental protocols does not change the classification of the model performances while for some others (e.g., fenitrothion, fenthion), the classification can be completely modified. However, it is noteworthy that these classifications must be interpreted with care since the different residual values can be relatively weak or very high. Thus, for example, the residual values (i.e.,  $\log \text{BCF}_{\text{obs.}} - \log \text{BCF}_{\text{calc.}}$ ) which give the classification 5, 3, 4, 1, 7, 2, 6 obtained for acenaphthene (Table I) are the following: 0.00, -0.01, -0.03, -0.04, -0.12, -0.16, and -0.19. At the opposite, the residual values yielding the classification 6, 7, 2, 4, 1, 5, 3 for octachlorodibenzo-*p*-dioxin (Table I, last value for this chemical) are the following: 0.08, -1.00, -2.18, -2.19, -2.39, -2.47, and -2.91. In the former case, it can be advanced that all the models are able to produce satisfactory estimations for the BCF of acenaphthene. Therefore, the information provided by the classification is useless. However, for octachlorodibenzo-*p*-dioxin, the classification is far more significant. Indeed, it is obvious that Eq. (6) (Bintein *et al.*, 1993) is the most suitable for predicting the bioconcentration of this chemical in fish. As inspection of Tables I and II does not provide information on the magnitude of the differences observed between the studied models, the classification was completed by a graphical study of the residuals obtained with each of the seven studied models. For this purpose, the residuals were plotted versus the log P values (Figures 1 to 14). Inspection of these figures confirms the equivalence of the different models for chemicals having log P values inferior to 6. The comparison of these different figures also confirms that for highly hydrophobic chemicals, the nonlinear equations (6) and (7) provide better simulations. In addition, Figures 1 to 14 allow a better estimation of the effect of the different experimental protocols on the results of the BCF simulations since for a same chemical (i.e., same log P value), the residuals corresponding to different protocols appear on a same vertical line. Thus, for example, in Figures 1, 3, 5, and 7, it is possible to stress that the BCF values calculated by the linear models for octachlorodibenzo-*p*-dioxin are always largely superior to those experimentally observed. The same remark can be made for the bilinear equation (5) (Figure 9). At the opposite, for the nonlinear equations (6) and (7) (Figures 11 and 13), the results show a higher variability. It is particularly interesting to note that according to the selected experimental BCF values, Eq. (6) (Bintein *et al.*, 1993) can have a tendency to underestimate or at the opposite to overestimate the bioconcentration of

**Table I** *n*-octanol/water partition coefficients (log P), bioconcentration data (log BCF) obtained under flow-through conditions and classification of the seven models.

| Chemical                        | log P | log BCF            | Species                    | Reference                    | Classification <sup>o</sup> |
|---------------------------------|-------|--------------------|----------------------------|------------------------------|-----------------------------|
| Acenaphthene                    | 3.92  | 2.59               | <i>Lepomis macrochirus</i> | Veith et al. (1980)          | 5, 3, 4, 1, 7, 2, 6         |
| Acridine                        | 3.45  | 2.10               | <i>Pimephales promelas</i> | Southworth et al. (1979)     | 3-5, 7, 1, 4, 6, 2          |
| Acrolein                        | 0.90  | 2.54               | <i>Lepomis macrochirus</i> | Veith et al. (1980)          | 7, 2, 4, 1, 6, 5, 3         |
| Benzo[a]pyrene                  | 5.97  | 3.51               | <i>Lepomis macrochirus</i> | Jimenez et al. (1987)        | 6, 4, 2, 1, 7, 5, 3         |
| Bis(2-chloroethyl)ether         | 1.12  | 1.04               | <i>Lepomis macrochirus</i> | Veith et al. (1980)          | 2, 4, 7, 1, 6, 5, 3         |
| BPMC                            | 2.78  | 1.41               | <i>Pseudorasbora parva</i> | Kanazawa (1981)              | 3-5, 7, 1, 4, 6, 2          |
| Bromacil                        | 2.11  | 0.51               | <i>Pimephales promelas</i> | Call et al. (1987)           | 3, 5, 1, 6, 4, 7, 2         |
| 5-Bromoindole                   | 3.00  | 1.15               | <i>Pimephales promelas</i> | Veith et al. (1979)          | 3-5, 7, 1, 4, 6, 2          |
| 4-Bromophenol                   | 2.59  | 1.56               | <i>Brachydanio rerio</i>   | Butte et al. (1987)          | 4-6, 1, 7, 2, 3, 5          |
| Bromophos                       | 4.88  | 4.60 <sup>18</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991) | 3, 2, 6, 1, 4, 7, 5         |
|                                 | 4.88  | 4.65 <sup>1</sup>  | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991) | 6, 3, 2, 7, 1, 5, 4         |
| 2- <i>t</i> -Butoxy ethanol     | 0.39  | -0.22              | <i>Cyprinus carpio</i>     | Fujiwara et al. (1984)       | 4, 1, 6, 2, 5, 3, 7         |
| Butyl benzyl phthalate          | 4.05  | 2.89               | <i>Lepomis macrochirus</i> | Veith et al. (1980)          | 6, 2-7, 1, 3, 4-5           |
| <i>t</i> -Butyl isopropyl ether | 2.14  | 0.76               | <i>Cyprinus carpio</i>     | Fujiwara et al. (1984)       | 3, 5, 1, 6, 4, 7, 2         |
| <i>t</i> -Butyl methyl ether    | 1.24  | 0.18               | <i>Cyprinus carpio</i>     | Fujiwara et al. (1984)       | 6, 1, 5, 3, 4, 2, 7         |
| 4- <i>t</i> -Butyl phenol       | 3.31  | 1.86               | <i>Brachydanio rerio</i>   | Butte et al. (1987)          | 5, 3, 7, 1, 4, 6, 2         |
| Carbaryl                        | 2.31  | 0.95               | <i>Pseudorasbora parva</i> | Kanazawa (1981)              | 3, 5, 1, 6-7, 4, 2          |
| Carbon tetrachloride            | 2.73  | 1.48               | <i>Lepomis macrochirus</i> | Veith et al. (1980)          | 5, 3-7, 1, 4, 6, 2          |
| Chlordane                       | 6.00  | 4.58               | <i>Pimephales promelas</i> | Veith et al. (1979)          | 5, 7, 3, 1, 2, 4, 6         |
| $\alpha$ -Chlordane             | 6.00  | 4.45               | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1985)      | 7, 1, 2, 4, 6, 3, 5         |
| $\gamma$ -Chlordane             | 6.00  | 4.20               | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1985)      | 6, 4, 2, 1, 7, 3, 5         |
| Chlornitrofen                   | 4.70  | 3.04               | <i>Pseudorasbora parva</i> | Kanazawa (1981)              | 4, 1, 2, 5, 3, 6, 7         |
| 2-Chloroaniline                 | 1.93  | 0.30               | <i>Cyprinus carpio</i>     | Tsuda et al. (1993)          | 3, 5, 1, 6, 4, 2, 7         |
|                                 | 1.93  | 0.57               | <i>Cyprinus carpio</i>     | Tsuda et al. (1993)          | 3, 5, 1, 6, 4, 2, 7         |
| 3-Chloroaniline                 | 1.91  | -0.10              | <i>Cyprinus carpio</i>     | Tsuda et al. (1993)          | 3, 5, 1, 6, 4, 2, 7         |
|                                 | 1.91  | 0.34               | <i>Cyprinus carpio</i>     | Tsuda et al. (1993)          | 3, 5, 1, 6, 4, 2, 7         |
| 4-Chloroaniline                 | 1.88  | -0.10              | <i>Cyprinus carpio</i>     | Tsuda et al. (1993)          | 3, 5, 1, 6, 4, 2, 7         |
|                                 | 1.88  | 0.23               | <i>Cyprinus carpio</i>     | Tsuda et al. (1993)          | 3, 5, 1, 6, 4, 2, 7         |
| <i>p</i> -Chlorobiphenyl        | 4.61  | 3.54 <sup>8</sup>  | <i>Oryzias latipes</i>     | Sugiura et al. (1984)        | 7, 6, 3, 5, 2, 1, 4         |
| Chloroform                      | 1.90  | 0.78               | <i>Lepomis macrochirus</i> | Veith et al. (1980)          | 1, 6, 5, 3, 4, 2, 7         |
| 2-Chloronitrobenzene            | 2.52  | 2.10               | <i>Oncorhynchus mykiss</i> | Niimi et al. (1989)          | 2, 6, 4, 1, 5, 7, 3         |
| 3-Chloronitrobenzene            | 2.50  | 1.89               | <i>Oncorhynchus mykiss</i> | Niimi et al. (1989)          | 2, 6, 4, 1, 5-7, 3          |
| 4-Chloronitrobenzene            | 2.39  | 2.00               | <i>Oncorhynchus mykiss</i> | Niimi et al. (1989)          | 2, 4-6, 7, 1, 5, 3          |
| 2-Chlorophenol                  | 2.15  | 2.33               | <i>Lepomis macrochirus</i> | Veith et al. (1980)          | 2, 7, 4, 6, 1, 5, 3         |
| 3-Chlorophenol                  | 2.50  | 1.25               | <i>Brachydanio rerio</i>   | Butte et al. (1987)          | 3, 5-7, 1, 4, 6, 2          |
| Chlorothion                     | 3.63  | 2.46 <sup>18</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991) | 3, 4, 1-7, 6, 2, 5          |
|                                 | 3.63  | 2.61 <sup>1</sup>  | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991) | 3, 4, 1-7, 6, 2, 5          |
| Cresyldiphenyl phosphate        | 4.51  | 2.00               | <i>Alburnus alburnus</i>   | Bengtsson et al. (1986)      | 4, 1, 5, 3, 2, 6, 7         |
|                                 | 4.51  | 2.34               | <i>Alburnus alburnus</i>   | Bengtsson et al. (1986)      | 4, 1, 5, 3, 2, 6, 7         |
| 4-Cyanophenol                   | 1.60  | 0.91               | <i>Brachydanio rerio</i>   | Butte et al. (1987)          | 2, 4, 6, 1, 7, 3, 5         |
| Cyanophos                       | 2.71  | 2.49 <sup>18</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991) | 3, 7, 1, 5, 4, 6, 2         |
|                                 | 2.71  | 2.62 <sup>1</sup>  | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991) | 7, 1, 5, 4, 6, 3, 2         |
| Cypermethrin                    | 6.05  | 2.89               | <i>Oncorhynchus mykiss</i> | Muir et al. (1994)           | 6, 4, 2, 1, 7, 5, 3         |
|                                 | 6.05  | 2.92               | <i>Oncorhynchus mykiss</i> | Muir et al. (1994)           | 6, 4, 2, 1, 7, 5, 3         |
| 2,7-DCDD*                       | 6.38  | 2.56               | <i>Poecilia reticulata</i> | Gobas and Schrap (1990)      | 6, 4, 7, 2, 1, 3, 5         |
| <i>p,p'</i> -DDE                | 5.83  | 4.71               | <i>Pimephales promelas</i> | Veith et al. (1979)          | 3, 5, 7, 1, 2, 6, 4         |
|                                 | 5.83  | 4.91               | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1985)      | 5, 3, 7, 1, 2, 6, 4         |
| <i>o,p'</i> -DDT                | 6.00  | 4.57               | <i>Pimephales promelas</i> | Veith et al. (1979)          | 5, 7, 3, 1, 2, 4, 6         |
| <i>p,p'</i> -DDT                | 6.00  | 4.47               | <i>Pimephales promelas</i> | Veith et al. (1979)          | 7, 1, 2, 5, 4, 3, 6         |
|                                 | 6.00  | 4.81               | <i>Oncorhynchus mykiss</i> | Muir et al. (1994)           | 3, 5, 7, 1, 2, 4, 6         |
|                                 | 6.00  | 4.86               | <i>Oncorhynchus mykiss</i> | Muir et al. (1994)           | 3, 5, 7, 1, 2, 4, 6         |
|                                 | 6.00  | 4.95               | <i>Oncorhynchus mykiss</i> | Muir et al. (1994)           | 3, 5, 7, 1, 2, 4, 6         |
|                                 | 6.00  | 4.97               | <i>Oncorhynchus mykiss</i> | Muir et al. (1985)           | 3, 5, 7, 1, 2, 4, 6         |
|                                 | 6.00  | 4.99               | <i>Oncorhynchus mykiss</i> | Muir et al. (1994)           | 3, 5, 7, 1, 2, 4, 6         |
| Decachlorobiphenyl              | 8.27  | 3.92 <sup>8</sup>  | <i>Poecilia reticulata</i> | Gobas et al. (1989)          | 7, 6, 5, 2, 4, 1, 3         |
|                                 | 8.27  | 4.00 <sup>8</sup>  | <i>Poecilia reticulata</i> | Bruggeman et al. (1984)      | 7, 5, 6, 2, 4, 1, 3         |

Table I (continued)

| Chemical                  | log P | log BCF            | Species                           | Reference                      | Classification <sup>o</sup> |
|---------------------------|-------|--------------------|-----------------------------------|--------------------------------|-----------------------------|
| Deltamethrin              | 6.20  | 2.62               | <i>Oncorhynchus mykiss</i>        | Muir <i>et al.</i> (1994)      | 6, 4, 2, 7, 1, 5, 3         |
|                           | 6.20  | 2.70               | <i>Oncorhynchus mykiss</i>        | Muir <i>et al.</i> (1994)      | 6, 4, 2, 7, 1, 5, 3         |
| Diazinon                  | 3.81  | 1.24               | <i>Poecilia reticulata</i>        | Kanazawa (1978)                | 5, 3, 4, 1, 7, 2, 6         |
|                           | 3.81  | 1.34               | <i>Oryzias latipes</i>            | Tsuda <i>et al.</i> (1995)     | 5, 3, 4, 1, 7, 2, 6         |
|                           | 3.81  | 1.41               | <i>Misgurnus anguillicaudatus</i> | Seguchi and Asaka (1981)       | 5, 3, 4, 1, 7, 2, 6         |
|                           | 3.81  | 1.45               | <i>Oryzias latipes</i>            | Tsuda <i>et al.</i> (1995)     | 5, 3, 4, 1, 7, 2, 6         |
|                           | 3.81  | 1.56               | <i>Cyprinus auratus</i>           | Kanazawa (1978)                | 5, 3, 4, 1, 7, 2, 6         |
|                           | 3.81  | 1.80               | <i>Oncorhynchus mykiss</i>        | Seguchi and Asaka (1981)       | 3, 4, 1, 7, 2, 6, 5         |
|                           | 3.81  | 1.81               | <i>Cyprinus carpio</i>            | Kanazawa (1978)                | 5, 3, 4, 1, 7, 2, 6         |
|                           | 3.81  | 2.08               | <i>Cyprinus carpio</i>            | Seguchi and Asaka (1981)       | 3, 4, 1, 7, 2, 6, 5         |
|                           | 3.81  | 2.18               | <i>Pseudorasbora parva</i>        | Kanazawa (1981)                | 5, 3, 4, 1, 7, 2, 6         |
|                           | 3.81  | 2.18               | <i>Pseudorasbora parva</i>        | Kanazawa (1978)                | 5, 3, 4, 1, 7, 2, 6         |
| 1,3-Dibromobenzene        | 3.75  | 2.82               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1984)        | 5, 2-6, 7, 1, 4, 3          |
| 1,4-Dibromobenzene        | 3.79  | 1.96 <sup>§</sup>  | <i>Poecilia reticulata</i>        | Gobas <i>et al.</i> (1989)     | 3, 4, 1, 7, 5, 2, 6         |
| 4,4'-Dibromobiphenyl      | 5.72  | 4.24 <sup>§</sup>  | <i>Poecilia reticulata</i>        | Gobas <i>et al.</i> (1989)     | 1-6, 2-7, 3, 4, 5           |
| 2,6-Dibromo-4-cyanophenol | 2.61  | 1.67               | <i>Brachydanio rerio</i>          | Butte <i>et al.</i> (1987)     | 2-6, 4, 1, 7, 3, 5          |
| Dicaphon                  | 3.58  | 2.95 <sup>†</sup>  | <i>Poecilia reticulata</i>        | de Bruijn and Hermens (1991)   | 3, 1-4-7, 6, 2, 5           |
|                           | 3.58  | 3.09 <sup>†§</sup> | <i>Poecilia reticulata</i>        | de Bruijn and Hermens (1991)   | 3, 1-4-7, 6, 2, 5           |
| 1,2-Dichlorobenzene       | 3.43  | 1.95               | <i>Lepomis macrochirus</i>        | Veith <i>et al.</i> (1980)     | 3-5, 7, 1-4, 6, 2           |
|                           | 3.43  | 2.43               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1983)        | 2, 5, 6, 1-4, 7, 3          |
|                           | 3.43  | 2.75               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1983)        | 2, 5, 6, 1-4, 7, 3          |
| 1,3-Dichlorobenzene       | 3.53  | 1.82               | <i>Lepomis macrochirus</i>        | Veith <i>et al.</i> (1980)     | 3-5, 7, 1-4, 6, 2           |
|                           | 3.53  | 2.62               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1983)        | 2, 5, 6, 1-4, 7, 3          |
|                           | 3.53  | 2.87               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1983)        | 2, 5, 6, 1-4, 7, 3          |
| 1,4-Dichlorobenzene       | 3.44  | 1.78               | <i>Lepomis macrochirus</i>        | Veith <i>et al.</i> (1980)     | 3-5, 7, 1, 4, 6, 2          |
|                           | 3.44  | 2.47               | <i>Jordanella floridae</i>        | Smith <i>et al.</i> (1990)     | 2, 5, 6, 4, 1, 7, 3         |
|                           | 3.44  | 2.57               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1983)        | 2, 5, 6, 4, 1, 7, 3         |
|                           | 3.44  | 2.71               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1985)        | 2, 6, 5, 4, 1, 7, 3         |
|                           | 3.44  | 2.86               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1983)        | 2, 5, 6, 4, 1, 7, 3         |
|                           | 3.44  | 2.95               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1985)        | 2, 6, 5, 4, 1, 7, 3         |
| 2,5-Dichlorobiphenyl      | 5.16  | 4.00               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1985)        | 7, 5, 3, 6, 1-2, 4          |
|                           | 5.16  | 4.14 <sup>§</sup>  | <i>Carassius auratus</i>          | Bruggeman <i>et al.</i> (1981) | 7, 3, 6, 1-2, 5, 4          |
|                           | 5.16  | 4.24 <sup>§</sup>  | <i>Poecilia reticulata</i>        | Bruggeman <i>et al.</i> (1984) | 7, 3, 6, 1-2, 5, 4          |
|                           | 5.16  | 4.53               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1985)        | 7, 5, 3, 6, 1-2, 4          |
| 3,5-Dichlorobiphenyl      | 5.37  | 3.79               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1985)        | 4, 2, 1, 6, 3, 7, 5         |
|                           | 5.37  | 3.82               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1985)        | 2, 1-4, 6, 3, 7, 5          |
| 1,2-Dichloroethane        | 1.45  | 0.30               | <i>Lepomis macrochirus</i>        | Veith <i>et al.</i> (1980)     | 5, 3, 1-6, 4, 2, 7          |
| 4,5-Dichloroguaiacol      | 3.18  | 2.03               | <i>Oncorhynchus mykiss</i>        | Niimi <i>et al.</i> (1990)     | 4, 1-5, 6, 7, 2, 3          |
| 1,4-Dichloronaphthalene   | 4.66  | 3.75               | <i>Oncorhynchus mykiss</i>        | Oliver and Niimi (1984)        | 5, 7, 6, 3, 2, 1, 4         |
| 2,3-Dichloronitrobenzene  | 3.05  | 2.16               | <i>Oncorhynchus mykiss</i>        | Niimi <i>et al.</i> (1989)     | 2, 6, 5, 4, 1, 7, 3         |
| 2,4-Dichloronitrobenzene  | 3.05  | 2.07               | <i>Oncorhynchus mykiss</i>        | Niimi <i>et al.</i> (1989)     | 2, 6, 5, 4, 1, 7, 3         |
| 2,5-Dichloronitrobenzene  | 3.03  | 2.05               | <i>Oncorhynchus mykiss</i>        | Niimi <i>et al.</i> (1989)     | 2, 6, 5, 4, 1, 7, 3         |
| 3,4-Dichloronitrobenzene  | 3.04  | 2.07               | <i>Oncorhynchus mykiss</i>        | Niimi <i>et al.</i> (1989)     | 2, 6, 5, 4, 1, 7, 3         |
| 3,5-Dichloronitrobenzene  | 3.09  | 2.23               | <i>Oncorhynchus mykiss</i>        | Niimi <i>et al.</i> (1989)     | 2, 6, 5, 4, 1, 7, 3         |
| Dichlorvos                | 1.47  | -0.10              | <i>Gnathopogon aereulesens</i>    | Tsuda <i>et al.</i> (1992)     | 3, 5, 1-6, 4, 2, 7          |
| Dieldrin                  | 5.40  | 3.65               | <i>Pseudorasbora parva</i>        | Kanazawa (1981)                | 4, 2, 1, 6, 5, 3, 7         |
| Di(2-ethylhexyl)phthalate | 7.45  | 2.19               | <i>Pimephales promelas</i>        | Mayer (1976)                   | 6, 7, 2, 4, 5, 1, 3         |
|                           | 7.45  | 2.95               | <i>Pimephales promelas</i>        | Mayer (1976)                   | 6, 7, 2, 4, 5, 1, 3         |
| Diethyl phthalate         | 2.47  | 2.07               | <i>Lepomis macrochirus</i>        | Veith <i>et al.</i> (1980)     | 2, 4-6, 1, 7, 5, 3          |
| 2,4-Dimethylphenol        | 2.42  | 2.18               | <i>Lepomis macrochirus</i>        | Veith <i>et al.</i> (1980)     | 2, 4-6, 7, 1, 5, 3          |
| Dimethyl phthalate        | 1.56  | 1.76               | <i>Lepomis macrochirus</i>        | Veith <i>et al.</i> (1980)     | 7, 2, 4, 1-6, 5, 3          |
| Diphenylamine             | 3.42  | 1.48               | <i>Pimephales promelas</i>        | Veith <i>et al.</i> (1979)     | 3-5, 7, 1, 4, 6, 2          |
| Diuron                    | 2.68  | 2.16               | <i>Pimephales promelas</i>        | Call <i>et al.</i> (1987)      | 2, 6, 4, 1, 7, 5, 3         |
|                           | 2.68  | 2.20               | <i>Pimephales promelas</i>        | Call <i>et al.</i> (1987)      | 2, 6, 4, 1, 7, 5, 3         |
| EPN                       | 3.85  | 3.37               | <i>Pseudorasbora parva</i>        | Kanazawa (1981)                | 6, 2, 7, 1, 4, 3, 5         |
| Fenitrothion              | 3.47  | 1.65               | <i>Oryzias latipes</i>            | Tsuda <i>et al.</i> (1995)     | 3-5, 7, 1, 4, 6, 2          |
|                           | 3.47  | 1.68               | <i>Oryzias latipes</i>            | Tsuda <i>et al.</i> (1995)     | 3-5, 7, 1, 4, 6, 2          |
|                           | 3.47  | 3.36 <sup>†</sup>  | <i>Poecilia reticulata</i>        | de Bruijn and Hermens (1991)   | 4, 1, 6, 7, 5, 2, 3         |
|                           | 3.47  | 2.39               | <i>Pseudorasbora parva</i>        | Kanazawa (1981)                | 2-6, 4, 1, 7, 3-5           |

Table I (continued)

| Chemical                             | log P                             | log BCF Species    | Reference                    | Classification <sup>o</sup>    |                           |                     |
|--------------------------------------|-----------------------------------|--------------------|------------------------------|--------------------------------|---------------------------|---------------------|
| Fenitrothion                         | 3.47                              | 2.48               | <i>Oryzias latipes</i>       | Takimoto <i>et al.</i> (1987)  | 2, 6, 4, 1, 7, 3, 5       |                     |
|                                      | 3.47                              | 3.54 <sup>18</sup> | <i>Poecilia reticulata</i>   | de Bruijn and Hermens (1991)   | 2, 5, 6, 4, 1, 7, 3       |                     |
|                                      | 3.47                              | 2.74               | <i>Oryzias latipes</i>       | Takimoto <i>et al.</i> (1984)  | 5, 2, 6, 4, 1, 7, 3       |                     |
|                                      | 3.47                              | 2.75               | <i>Oryzias latipes</i>       | Takimoto <i>et al.</i> (1984)  | 5, 2, 6, 4, 1, 7, 3       |                     |
| Fenthion                             | 4.17                              | 1.96               | <i>Oryzias latipes</i>       | Tsuda <i>et al.</i> (1995)     | 4, 1-5, 3, 2, 7, 6        |                     |
|                                      | 4.17                              | 2.02               | <i>Oryzias latipes</i>       | Tsuda <i>et al.</i> (1995)     | 4, 1-5, 3, 2, 7, 6        |                     |
|                                      | 4.17                              | 2.68               | <i>Gnathopogon taenulens</i> | Tsuda <i>et al.</i> (1992)     | 4, 1-5, 3, 2, 7, 6        |                     |
|                                      | 4.17                              | 4.17 <sup>18</sup> | <i>Poecilia reticulata</i>   | de Bruijn and Hermens (1991)   | 5, 6, 7, 2, 3, 1, 4       |                     |
| Fenthion-S2145                       | 4.17                              | 4.22 <sup>†</sup>  | <i>Poecilia reticulata</i>   | de Bruijn and Hermens (1991)   | 5, 6, 7, 2, 3, 1, 4       |                     |
|                                      | 3.74                              | 3.65 <sup>18</sup> | <i>Poecilia reticulata</i>   | de Bruijn and Hermens (1991)   | 2-6, 5, 7, 1, 4, 3        |                     |
|                                      | 3.74                              | 3.84 <sup>†</sup>  | <i>Poecilia reticulata</i>   | de Bruijn and Hermens (1991)   | 5, 2-6, 7, 1, 4, 3        |                     |
| Fenvalerate                          | 6.20                              | 2.61               | <i>Oncorhynchus mykiss</i>   | Muir <i>et al.</i> (1994)      | 6, 4, 2, 7, 1, 5, 3       |                     |
|                                      | 6.20                              | 2.96               | <i>Oncorhynchus mykiss</i>   | Muir <i>et al.</i> (1994)      | 6, 4, 2, 7, 1, 5, 3       |                     |
| 1,2,3,4,7,8-HCDD <sup>s</sup>        | 7.79                              | 3.29               | <i>Oncorhynchus mykiss</i>   | Servos <i>et al.</i> (1989)    | 6, 7, 5, 2, 4, 1, 3       |                     |
|                                      | 7.79                              | 3.36               | <i>Oncorhynchus mykiss</i>   | Muir <i>et al.</i> (1985)      | 6, 7, 5, 2, 4, 1, 3       |                     |
|                                      | 7.79                              | 3.63               | <i>Pimephales promelas</i>   | Muir <i>et al.</i> (1985)      | 6, 7, 5, 2, 4, 1, 3       |                     |
|                                      | 7.79                              | 4.13 <sup>§</sup>  | <i>Poecilia reticulata</i>   | Loonen <i>et al.</i> (1994)    | 7, 6, 2, 4, 5, 1, 3       |                     |
| 1,2,3,4,6,7,8-HCDD <sup>s</sup>      | 8.20                              | 2.71               | <i>Pimephales promelas</i>   | Muir <i>et al.</i> (1985)      | 6, 7, 5, 2, 4, 1, 3       |                     |
|                                      | 8.20                              | 2.97               | <i>Oncorhynchus mykiss</i>   | Servos <i>et al.</i> (1989)    | 6, 7, 5, 2, 4, 1, 3       |                     |
|                                      | 8.20                              | 3.15               | <i>Oncorhynchus mykiss</i>   | Muir <i>et al.</i> (1985)      | 6, 7, 5, 2, 4, 1, 3       |                     |
|                                      | 8.20                              | 3.16               | <i>Oncorhynchus mykiss</i>   | Servos <i>et al.</i> (1989)    | 6, 7, 5, 2, 4, 1, 3       |                     |
|                                      | 8.20                              | 3.75 <sup>§</sup>  | <i>Poecilia reticulata</i>   | Loonen <i>et al.</i> (1994)    | 7, 6, 5, 2, 4, 1, 3       |                     |
|                                      | 8.20                              | 3.62 <sup>§</sup>  | <i>Poecilia reticulata</i>   | Loonen <i>et al.</i> (1994)    | 7, 6, 2, 5, 4, 1, 3       |                     |
| 1,2,3,4,6,7,8-HCDF <sup>f</sup>      | 7.92                              | 3.62 <sup>§</sup>  | <i>Poecilia reticulata</i>   | Loonen <i>et al.</i> (1994)    | 7, 6, 2, 5, 4, 1, 3       |                     |
| Heptachlor                           | 5.44                              | 3.98               | <i>Pimephales promelas</i>   | Veith <i>et al.</i> (1979)     | 6, 1, 2, 5, 3, 4, 7       |                     |
| Heptachlor epoxide                   | 5.40                              | 4.16               | <i>Pimephales promelas</i>   | Veith <i>et al.</i> (1979)     | 7, 3, 5, 6, 1, 2, 4       |                     |
| Heptachloronorbornene                | 5.28                              | 4.05               | <i>Pimephales promelas</i>   | Veith <i>et al.</i> (1979)     | 7, 3, 5, 6, 1, 2, 4       |                     |
| Hexabromobenzene                     | 6.07                              | 3.04               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1985)        | 6, 4, 2, 1, 7, 3, 5       |                     |
| Hexabromobiphenyl**                  | 6.39                              | 4.26               | <i>Pimephales promelas</i>   | Veith <i>et al.</i> (1979)     | 6, 4, 7, 2, 1, 5, 3       |                     |
| 2,2',4,4',6,6'-Hexabromobiphenyl     | 7.20                              | 4.66 <sup>§</sup>  | <i>Poecilia reticulata</i>   | Gobas <i>et al.</i> (1989)     | 7, 2-4, 1, 5, 6, 3        |                     |
| Hexabromocyclododecane               | 5.81                              | 4.26               | <i>Pimephales promelas</i>   | Veith <i>et al.</i> (1979)     | 1, 2-6, 4, 7, 5, 3        |                     |
| Hexachlorobenzene                    | 5.73                              | 3.74               | <i>Oncorhynchus mykiss</i>   | Veith <i>et al.</i> (1979)     | 4, 2, 6, 1, 7, 5, 3       |                     |
|                                      | 5.73                              | 4.16 <sup>§</sup>  | <i>Oryzias latipes</i>       | Sugiura <i>et al.</i> (1984)   | 6, 1, 2, 4, 7, 5, 3       |                     |
|                                      | 5.73                              | 4.21               | <i>Pimephales promelas</i>   | Veith <i>et al.</i> (1979)     | 1, 6, 2, 4-7, 5, 3        |                     |
|                                      | 5.73                              | 4.30               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1983)        | 7, 3, 1, 6, 2, 4, 5       |                     |
|                                      | 5.73                              | 4.34               | <i>Lepomis cyanellus</i>     | Veith <i>et al.</i> (1979)     | 7, 5, 3, 1, 6, 2, 4       |                     |
|                                      | 5.73                              | 4.43               | <i>Poecilia reticulata</i>   | Schrap and Opperhuizen (1990)  | 3, 5, 7, 1, 6, 2, 4       |                     |
|                                      | 2,2',4,4',5,5'-Hexachlorobiphenyl | 6.90               | 4.84                         | <i>Oncorhynchus mykiss</i>     | Muir <i>et al.</i> (1985) | 4, 2, 7, 1, 5, 3, 6 |
|                                      | 6.90                              | 5.30 <sup>§</sup>  | <i>Poecilia reticulata</i>   | Bruggeman <i>et al.</i> (1984) | 1, 5, 3, 2, 4, 7, 6       |                     |
| 1,1,2,3,4,4-Hexachloro-1,3-butadiene | 6.90                              | 5.65               | <i>Poecilia reticulata</i>   | Opperhuizen and Schrap (1987)  | 3, 5, 1, 2, 4, 7, 6       |                     |
|                                      | 4.78                              | 3.76               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1983)        | 5, 7, 6, 3, 2, 1, 4       |                     |
|                                      | 4.78                              | 4.23               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1983)        | 5, 7, 6, 3, 2, 1, 4       |                     |
| $\alpha$ -Hexachlorocyclohexane      | 3.80                              | 2.33               | <i>Oncorhynchus mykiss</i>   | Vigano <i>et al.</i> (1992)    | 3, 4, 1, 5-7, 2, 6        |                     |
|                                      | 3.80                              | 3.04               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 6, 2, 5, 7, 1, 4, 3       |                     |
|                                      | 3.80                              | 3.04               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 6, 2, 5, 7, 1, 4, 3       |                     |
|                                      | 3.80                              | 3.20               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1985)        | 6, 2, 5, 7, 1, 4, 3       |                     |
|                                      | 3.80                              | 3.38               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1985)        | 6, 2, 5, 7, 1, 4, 3       |                     |
| $\beta$ -Hexachlorocyclohexane       | 3.78                              | 3.16               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 6, 2, 5, 7, 1, 4, 3       |                     |
|                                      | 3.78                              | 3.18               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 6, 2, 5, 7, 1, 4, 3       |                     |
| $\delta$ -Hexachlorocyclohexane      | 4.14                              | 2.45               | <i>Oncorhynchus mykiss</i>   | Vigano <i>et al.</i> (1992)    | 5, 4, 1-3, 2, 7, 6        |                     |
|                                      | 4.14                              | 3.21               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 6, 7, 5, 2, 1-3, 4        |                     |
|                                      | 4.14                              | 3.25               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 6, 7, 5, 2, 1-3, 4        |                     |
|                                      | 3.72                              | 2.16               | <i>Oncorhynchus mykiss</i>   | Vigano <i>et al.</i> (1992)    | 5, 3, 1-4, 7, 2-6         |                     |
| $\gamma$ -Hexachlorocyclohexane      | 3.72                              | 2.26               | <i>Pimephales promelas</i>   | Veith <i>et al.</i> (1979)     | 3-5, 1-4, 7, 2-6          |                     |
|                                      | 3.72                              | 2.93               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 2-6, 5, 7, 1-4, 3         |                     |
|                                      | 3.72                              | 2.96               | <i>Brachydanio rerio</i>     | Butte <i>et al.</i> (1991)     | 2-6, 5, 7, 1-4, 3         |                     |
|                                      | 3.72                              | 3.08               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1985)        | 2-6, 5, 7, 1-4, 3         |                     |
|                                      | 3.72                              | 3.10               | <i>Pseudorasbora parva</i>   | Kanazawa (1981)                | 2-6, 7, 1-4, 3-5          |                     |
|                                      | 3.72                              | 3.30               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1985)        | 2-6, 5, 7, 1-4, 3         |                     |
|                                      | 3.72                              | 3.32               | <i>Oncorhynchus mykiss</i>   | Oliver and Niimi (1985)        | 2-6, 5, 7, 1-4, 3         |                     |

Table I (continued)

| Chemical                               | log P | log BCF Species   | Reference                  | Classification <sup>o</sup>    |                     |
|--|-------|-------------------|----------------------------|--------------------------------|---------------------|
| Hexachloroethane                       | 3.93  | 2.14              | <i>Lepomis macrochirus</i> | Veith <i>et al.</i> (1980)     | 5, 3, 4, 1, 7, 2, 6 |
|  | 3.93  | 2.71              | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1983)        | 7, 2, 1, 6, 4, 3, 5 |
|  | 3.93  | 3.08              | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1983)        | 5, 6, 2, 7, 1, 4, 3 |
| Hexachloronorbomadiene                 | 5.28  | 3.81              | <i>Pimephales promelas</i> | Veith <i>et al.</i> (1979)     | 1, 2, 6, 4, 5, 3, 7 |
| IBP                                    | 3.21  | 0.60              | <i>Pseudorasbora parva</i> | Kanazawa (1981)                | 3-5, 7, 1, 4, 6, 2  |
| Iodofenphos                            | 5.16  | 4.30 <sup>†</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991)   | 4, 1-2, 6, 3, 7, 5  |
|  | 5.16  | 4.68 <sup>†</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991)   | 4, 1-2, 6, 3, 7, 5  |
| Isophorone                             | 1.67  | 0.85              | <i>Lepomis macrochirus</i> | Veith <i>et al.</i> (1980)     | 4, 6, 1, 2, 7, 5, 3 |
| Leptophos                              | 5.88  | 3.78              | <i>Pseudorasbora parva</i> | Kanazawa (1981)                | 4, 6, 2, 1, 7, 5, 3 |
| Methidation                            | 2.42  | 1.26              | <i>Gnathopogon aeneus</i>  | Tsuda <i>et al.</i> (1992)     | 1, 7, 5, 3-4-6, 2   |
| Methoxychlor                           | 5.08  | 3.92              | <i>Pimephales promelas</i> | Veith <i>et al.</i> (1979)     | 7, 6, 3, 5, 2, 1, 4 |
| 2-Methyl-4,6-dinitrophenol             | 2.13  | 0.16              | <i>Brachydanio rerio</i>   | Butte <i>et al.</i> (1987)     | 5, 3, 1, 6, 4, 7, 2 |
| Methylisocyanothion                    | 3.58  | 3.39 <sup>†</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991)   | 1-4-7, 3, 6, 2, 5   |
|  | 3.58  | 3.53 <sup>†</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991)   | 2-6, 5, 1-4-7, 3    |
| Methylparathion                        | 2.94  | 2.98 <sup>†</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991)   | 6, 5, 2, 4, 1, 7, 3 |
|  | 2.94  | 3.04 <sup>†</sup> | <i>Poecilia reticulata</i> | de Bruijn and Hermens (1991)   | 2, 6, 5, 4, 1, 7, 3 |
| 2-Methylphenol                         | 1.95  | 1.03              | <i>Brachydanio rerio</i>   | Butte <i>et al.</i> (1987)     | 4, 6, 1, 2, 7, 3, 5 |
| Mirex                                  | 6.89  | 4.26              | <i>Pimephales promelas</i> | Veith <i>et al.</i> (1979)     | 6-7, 4, 2, 1, 5, 3  |
|  | 6.89  | 4.31 <sup>§</sup> | <i>Poecilia reticulata</i> | Gobas <i>et al.</i> (1989)     | 7, 6, 4, 2, 1, 5, 3 |
| Molinate                               | 3.21  | 1.41              | <i>Pseudorasbora parva</i> | Kanazawa (1981)                | 3-5, 7, 1, 4, 6, 2  |
| Naphthalene                            | 3.36  | 2.49              | <i>Lepomis macrochirus</i> | McCarthy and Jimenez (1985)    | 2, 6, 4, 1, 7, 3-5  |
|  | 3.36  | 2.51              | <i>Lepomis macrochirus</i> | McCarthy and Jimenez (1985)    | 2, 6, 4, 1, 7, 3-5  |
| 3-Nitrophenol                          | 2.00  | 1.40              | <i>Brachydanio rerio</i>   | Butte <i>et al.</i> (1987)     | 2, 7, 4, 6, 1, 3, 5 |
| 2,2',3,3',4,4',5,5'-Octachlorobiphenyl | 7.67  | 4.33 <sup>§</sup> | <i>Poecilia reticulata</i> | Bruggeman <i>et al.</i> (1984) | 7, 5, 6, 2, 4, 1, 3 |
| Octachlorodibenzo- <i>p</i> -dioxin    | 7.59  | 1.93              | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1985)      | 6, 7, 2-5, 4, 1, 3  |
|  | 7.59  | 2.59              | <i>Oncorhynchus mykiss</i> | Servos <i>et al.</i> (1989)    | 6, 7, 2-5, 4, 1, 3  |
|  | 7.59  | 2.67              | <i>Oncorhynchus mykiss</i> | Servos <i>et al.</i> (1989)    | 6, 7, 2-5, 4, 1, 3  |
|  | 7.59  | 2.85              | <i>Poecilia reticulata</i> | Gobas and Schrap (1990)        | 6, 7, 2, 4, 5, 1, 3 |
|  | 7.59  | 3.35              | <i>Pimephales promelas</i> | Muir <i>et al.</i> (1985)      | 6, 7, 2-5, 4, 1, 3  |
|  | 7.59  | 3.36 <sup>§</sup> | <i>Poecilia reticulata</i> | Loonen <i>et al.</i> (1994)    | 6, 7, 2, 4, 1, 5, 3 |
|  | 7.97  | 2.77              | <i>Poecilia reticulata</i> | Gobas and Schrap (1990)        | 6, 7, 5, 2, 4, 1, 3 |
|  | 7.97  | 3.10 <sup>§</sup> | <i>Poecilia reticulata</i> | Loonen <i>et al.</i> (1994)    | 6, 7, 5, 2, 4, 1, 3 |
| Octachloronaphthalene                  | 6.42  | 2.52              | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1985)        | 6, 7, 4, 2, 1, 3, 5 |
| Octachlorostyrene                      | 6.29  | 4.52              | <i>Pimephales promelas</i> | Veith <i>et al.</i> (1979)     | 4, 2, 7, 1, 6, 5, 3 |
| 1,2,3,4,7-PCDD <sup>SS</sup>           | 7.44  | 2.91              | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1985)      | 6, 7, 2, 4, 5, 1, 3 |
|  | 7.44  | 3.16              | <i>Pimephales promelas</i> | Muir <i>et al.</i> (1985)      | 6, 7, 2, 4, 5, 1, 3 |
| 1,2,3,7,8-PCDD <sup>SS</sup>           | 6.64  | 4.50 <sup>§</sup> | <i>Poecilia reticulata</i> | Loonen <i>et al.</i> (1994)    | 7, 4, 2, 6, 1, 3, 5 |
| 2,3,4,7,8-PCDF <sup>EE</sup>           | 6.92  | 4.36 <sup>§</sup> | <i>Poecilia reticulata</i> | Loonen <i>et al.</i> (1994)    | 7, 6, 4, 2, 1, 3, 5 |
| Pentachloroaniline                     | 5.08  | 2.56              | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1994)   | 4, 1, 2, 3, 6, 7, 5 |
| Pentachlorobenzene                     | 5.18  | 2.94 <sup>§</sup> | <i>Poecilia reticulata</i> | Bruggeman <i>et al.</i> (1984) | 4, 5, 1, 2, 6, 3, 7 |
|  | 5.18  | 3.53              | <i>Lepomis macrochirus</i> | Veith <i>et al.</i> (1980)     | 4, 1, 2, 5-6, 3, 7  |
|  | 5.18  | 4.11              | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1983)        | 5, 7, 3, 6, 2, 1, 4 |
|  | 5.18  | 4.30              | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1983)        | 5, 7, 3, 6, 2, 1, 4 |
|  | 5.18  | 4.36              | <i>Poecilia reticulata</i> | Schrap and Opperhuizen (1990)  | 7, 3, 5-6, 2, 1, 4  |
|  | 2.89  | 1.83              | <i>Lepomis macrochirus</i> | Veith <i>et al.</i> (1980)     | 6, 4, 1, 2, 7, 3-5  |
|  | 4.77  | 2.23              | <i>Oncorhynchus mykiss</i> | Niimi <i>et al.</i> (1989)     | 4, 1, 2, 3, 6, 7, 5 |
| Pentachloronitrobenzene                | 4.77  | 2.38              | <i>Pseudorasbora parva</i> | Kanazawa (1981)                | 4, 1, 2, 5, 3, 6, 7 |
|  | 4.77  | 2.41              | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1985)        | 4, 1, 2, 3, 6, 5, 7 |
|  | 4.77  | 2.77              | <i>Oncorhynchus mykiss</i> | Oliver and Niimi (1985)        | 4, 1, 2, 3, 6, 5, 7 |
|  | 5.01  | 2.33              | <i>Jordanella floridae</i> | Smith <i>et al.</i> (1990)     | 4, 1, 2, 3, 6, 7, 5 |
|  | 5.01  | 2.58 <sup>§</sup> | <i>Oryzias latipes</i>     | Sugiura <i>et al.</i> (1984)   | 4, 1, 2, 5, 3, 6, 7 |
| Pentachlorophenol                      | 5.01  | 2.89              | <i>Pimephales promelas</i> | Veith <i>et al.</i> (1979)     | 4, 1, 2, 5, 3, 6, 7 |
|  | 5.01  | 2.99              | <i>Brachydanio rerio</i>   | Butte <i>et al.</i> (1987)     | 4, 1, 5, 2, 3, 6, 7 |
|  | 5.01  | 3.23              | <i>Oryzias latipes</i>     | Tachikawa <i>et al.</i> (1991) | 4, 1, 2, 5, 3, 6, 7 |
|  | 6.50  | 3.23              | <i>Pimephales promelas</i> | Spehar <i>et al.</i> (1983)    | 6, 7, 4, 2, 1, 5, 3 |
|  | 6.50  | 3.29              | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1994)      | 6, 7, 4, 2, 1, 5, 3 |
| Permethrin                             | 6.50  | 3.39              | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1994)      | 6, 7, 4, 2, 1, 5, 3 |

Table I (continued)

| Chemical                       | log P                     | log BCF Species    | Reference                      | Classification <sup>o</sup>   |                     |
|--------------------------------|---------------------------|--------------------|--------------------------------|-------------------------------|---------------------|
| Permethrin                     | 6.50                      | 3.49               | <i>Pimephales promelas</i>     | Spehar et al. (1983)          | 6, 7, 4, 2, 1, 5, 3 |
|                                | 6.50                      | 3.52               | <i>Pimephales promelas</i>     | Spehar et al. (1983)          | 6, 7, 4, 2, 1, 5, 3 |
| Phenol                         | 1.46                      | 1.24               | <i>Brachydanio rerio</i>       | Butte et al. (1987)           | 7, 2, 4, 1-6, 3, 5  |
| Phenthoate                     | 3.69                      | 1.56               | <i>Pseudorasbora parva</i>     | Kanazawa (1981)               | 3-5, 4, 1, 7, 2, 6  |
| 2-Phenyldodecane               | 8.19                      | 2.65               | <i>Oncorhynchus mykiss</i>     | Burke et al. (1991)           | 6, 7, 5, 2, 4, 1, 3 |
| N-Phenyl-2-naphthylamine       | 4.38                      | 2.17               | <i>Pimephales promelas</i>     | Veith et al. (1979)           | 4, 1, 5, 3, 2, 6, 7 |
| Phosmet                        | 2.78                      | 1.56               | <i>Gnathopogon aenulescens</i> | Tsuda et al. (1992)           | 7, 1-3-5, 4, 6, 2   |
| Ronnell                        | 4.81                      | 4.55 <sup>18</sup> | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 3, 2-6, 1, 4-7, 5   |
|                                | 4.81                      | 4.64 <sup>1</sup>  | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 6, 7, 3, 5, 2, 1, 4 |
| Salithion                      | 2.67                      | 1.88               | <i>Gnathopogon aenulescens</i> | Tsuda et al. (1992)           | 2, 6, 4, 1, 7, 5, 3 |
| Simetryne                      | 2.54                      | 0.31               | <i>Gnathopogon aenulescens</i> | Tsuda et al. (1988)           | 3, 5, 7, 1, 4, 6, 2 |
| SV 5                           | 3.00                      | 2.95 <sup>1</sup>  | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 4, 1-5, 6, 7, 2, 3  |
|                                | 3.00                      | 3.22 <sup>18</sup> | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 2, 6, 5, 4, 1, 7, 3 |
| 1,2,3,4-TCDD <sup>a</sup>      | 6.20                      | 2.90               | <i>Poecilia reticulata</i>     | Gobas and Schrap (1990)       | 6, 4, 2, 7, 1, 3, 5 |
| 1,3,6,8-TCDD <sup>a</sup>      | 6.29                      | 3.20               | <i>Oncorhynchus mykiss</i>     | Servos et al. (1989)          | 6, 4, 2, 7, 1, 5, 3 |
|                                | 6.29                      | 3.32               | <i>Oncorhynchus mykiss</i>     | Muir et al. (1985)            | 6, 4, 2, 7, 1, 5, 3 |
|                                | 6.29                      | 3.32               | <i>Oncorhynchus mykiss</i>     | Servos et al. (1989)          | 6, 4, 2, 7, 1, 5, 3 |
|                                | 6.29                      | 3.39               | <i>Oncorhynchus mykiss</i>     | Servos et al. (1989)          | 6, 4, 2, 7, 1, 5, 3 |
|                                | 6.29                      | 3.57               | <i>Oncorhynchus mykiss</i>     | Servos et al. (1989)          | 6, 4, 2, 7, 1, 5, 3 |
|                                | 6.29                      | 3.76               | <i>Pimephales promelas</i>     | Muir et al. (1985)            | 6, 4, 2, 7, 1, 5, 3 |
|                                | 6.42                      | 4.01 <sup>8</sup>  | <i>Poecilia reticulata</i>     | Loonen et al. (1994)          | 6, 7, 4, 2, 1, 3, 5 |
|                                | 6.42                      | 4.56               | <i>Oncorhynchus mykiss</i>     | Mehrle et al. (1988)          | 7, 4, 2, 1, 6, 5, 3 |
|                                | 6.42                      | 4.57               | <i>Oncorhynchus mykiss</i>     | Mehrle et al. (1988)          | 7, 4, 2, 1, 6, 5, 3 |
|                                | 6.42                      | 4.59               | <i>Oncorhynchus mykiss</i>     | Mehrle et al. (1988)          | 7, 4, 2, 1, 5, 6, 3 |
| 2,3,7,8-TCDD <sup>a</sup>      | 6.42                      | 4.93               | <i>Oncorhynchus mykiss</i>     | Mehrle et al. (1988)          | 5, 1-3, 2, 4, 7, 6  |
|                                | 6.53                      | 3.31 <sup>8</sup>  | <i>Poecilia reticulata</i>     | Loonen et al. (1994)          | 6, 7, 4, 2, 1, 3, 5 |
|                                | 6.53                      | 3.42               | <i>Oncorhynchus mykiss</i>     | Mehrle et al. (1988)          | 6, 7, 4, 2, 1, 5, 3 |
|                                | 6.53                      | 3.65               | <i>Oncorhynchus mykiss</i>     | Mehrle et al. (1988)          | 6, 7, 4, 2, 1, 5, 3 |
|                                | 5.13                      | 3.57               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)       | 4, 1, 2, 3-6, 5, 7  |
| 1,2,4,5-Tetrabromobenzene      | 5.13                      | 3.81               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)       | 3-6, 5, 7, 2, 1, 4  |
|                                | 6.50                      | 4.97 <sup>8</sup>  | <i>Poecilia reticulata</i>     | Gobas et al. (1989)           | 1, 5, 3, 2, 4, 7, 6 |
| 2,2',5,5'-Tetrabromobiphenyl   | 4.57                      | 2.10               | <i>Poecilia reticulata</i>     | de Wolf et al. (1994)         | 4, 1, 2, 3, 6, 7, 5 |
| 2,3,4,5-Tetrachloroaniline     | 4.46                      | 2.46               | <i>Poecilia reticulata</i>     | de Wolf et al. (1994)         | 4, 1, 3, 2, 6, 7, 5 |
| 1,2,3,4-Tetrachlorobenzene     | 4.64                      | 3.72               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)       | 5, 7, 6, 3, 2, 1, 4 |
|                                | 4.64                      | 3.80               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)       | 7, 5, 6, 3, 2, 1, 4 |
|                                | 4.64                      | 3.82               | <i>Poecilia reticulata</i>     | Schrap and Opperhuizen (1990) | 7, 6, 3, 2-5, 1, 4  |
|                                | 4.64                      | 3.91               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)       | 7, 5, 6, 3, 2, 1, 4 |
|                                | 4.64                      | 4.08               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)       | 5, 7, 6, 3, 2, 1, 4 |
|                                | 4.66                      | 4.40 <sup>1</sup>  | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 3, 2, 1, 6, 4, 7, 5 |
|                                | 4.66                      | 4.61 <sup>1</sup>  | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 5, 7, 6, 3, 2, 1, 4 |
|                                | 4.66                      | 3.64               | <i>Poecilia reticulata</i>     | de Wolf et al. (1994)         | 7, 5, 6, 3, 2, 1, 4 |
| 1,2,4,5-Tetrachlorobenzene     | 4.66                      | 4.81 <sup>18</sup> | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 5, 7, 6, 3, 2, 1, 4 |
|                                | 4.66                      | 4.89 <sup>18</sup> | <i>Poecilia reticulata</i>     | de Bruijn and Hermens (1991)  | 5, 7, 6, 3, 2, 1, 4 |
|                                | 4.60                      | 3.61               | <i>Jordanella floridae</i>     | Smith et al. (1990)           | 5, 7, 6, 3, 2, 1, 4 |
|                                | 4.60                      | 3.72               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)       | 5, 7, 6, 3, 2, 1, 4 |
|                                | 4.60                      | 4.11               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)       | 5, 7, 6, 3, 2, 1, 4 |
|                                | 7.80                      | 3.36               | <i>Brachydanio rerio</i>       | Bouraly and Millischer (1989) | 6, 7, 5, 2, 4, 1, 3 |
| 2,2',3,3'-Tetrachlorobiphenyl  | 6.18                      | 4.69               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)       | 7, 1, 3, 2, 5, 4, 6 |
| 2,2',5,5'-Tetrachlorobiphenyl  | 6.09                      | 4.63               | <i>Poecilia reticulata</i>     | Opperhuizen and Schrap (1987) | 5, 7, 3, 1, 2, 4, 6 |
|                                | 6.09                      | 4.69 <sup>8</sup>  | <i>Carassius auratus</i>       | Bruggeman et al. (1981)       | 3, 7, 5, 1, 2, 4, 6 |
|                                | 6.09                      | 4.84 <sup>8</sup>  | <i>Poecilia reticulata</i>     | Gobas et al. (1989)           | 5, 3, 7, 1, 2, 4, 6 |
|                                | 6.09                      | 4.90 <sup>8</sup>  | <i>Poecilia reticulata</i>     | Bruggeman et al. (1984)       | 3, 7, 5, 1, 2, 4, 6 |
|                                | 6.09                      | 5.30               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)       | 5, 3, 7, 1, 2, 4, 6 |
|                                | 6.23                      | 4.62 <sup>8</sup>  | <i>Carassius auratus</i>       | Bruggeman et al. (1981)       | 1, 5, 7, 2, 4, 3, 6 |
|                                | 1,1,2,2-Tetrachloroethane | 2.39               | 0.90                           | <i>Lepomis macrochirus</i>    | Veith et al. (1980) |
| Tetrachloroethylene            | 2.53                      | 1.69               | <i>Lepomis macrochirus</i>     | Veith et al. (1980)           | 2, 6, 4, 1, 7, 5, 3 |
| Tetrachloroguaiacol            | 4.45                      | 2.26               | <i>Oncorhynchus mykiss</i>     | Niimi et al. (1990)           | 4, 1, 3, 2, 6, 7, 5 |
| 1,2,3,4-Tetrachloronaphthalene | 5.75                      | 3.71               | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)       | 4, 2, 6, 1, 7, 3, 5 |



Table I (continued)

| Chemical                        | log P | log BCF Species   | Reference                      | Classification <sup>o</sup>    |                     |
|---------------------------------|-------|-------------------|--------------------------------|--------------------------------|---------------------|
| 2,3,4,5-Tetrachloronitrobenzene | 3.93  | 1.87              | <i>Oncorhynchus mykiss</i>     | Niimi <i>et al.</i> (1989)     | 3, 4, 1, 7, 2, 6, 5 |
|                                 | 3.93  | 1.90              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 3, 4, 1, 7, 5, 2, 6 |
| 2,3,5,6-Tetrachloronitrobenzene | 4.38  | 3.13              | <i>Oncorhynchus mykiss</i>     | Niimi <i>et al.</i> (1989)     | 2, 6, 3, 7, 1, 5, 4 |
|                                 | 4.38  | 3.20              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 5-6-7, 2, 3, 1, 4   |
|                                 | 4.38  | 3.34              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 7, 5-6, 2, 3, 1, 4  |
|                                 | 4.39  | 2.15              | <i>Jordanella floridae</i>     | Smith <i>et al.</i> (1990)     | 4, 1, 3, 2, 6, 7, 5 |
| Thiobencarb                     | 3.40  | 1.82              | <i>Gnathopogon aenulescers</i> | Tsuda <i>et al.</i> (1988)     | 3-5, 7, 1, 4, 6, 2  |
|                                 | 3.40  | 2.23              | <i>Pseudorasbora parva</i>     | Kanazawa (1981)                | 4, 1, 6-7, 2, 3-5   |
| Toluene diamine                 | 3.16  | 1.96              | <i>Pimephales promelas</i>     | Veith <i>et al.</i> (1979)     | 1, 4-7, 3-5, 6, 2   |
| 2,4,6-Tribromoanisole           | 4.48  | 2.94              | <i>Pimephales promelas</i>     | Veith <i>et al.</i> (1979)     | 4, 1, 5, 3, 2, 6, 7 |
| 1,3,5-Tribromobenzene           | 4.51  | 3.23 <sup>§</sup> | <i>Poecilia reticulata</i>     | Gobas <i>et al.</i> (1989)     | 2, 3, 5, 6, 1, 7, 4 |
|                                 | 4.51  | 3.70              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1984)        | 5, 7, 6, 2, 3, 1, 4 |
|                                 | 4.51  | 3.97              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 7, 5, 6, 2, 3, 1, 4 |
|                                 | 4.51  | 4.08              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 7, 5, 6, 2, 3, 1, 4 |
| 2,4,6-Tribromobiphenyl          | 6.03  | 3.88 <sup>§</sup> | <i>Poecilia reticulata</i>     | Gobas <i>et al.</i> (1989)     | 6, 4, 2, 1, 7, 3, 5 |
| 2,4,6-Tribromophenol            | 4.23  | 2.71              | <i>Brachydanio rerio</i>       | Butte <i>et al.</i> (1987)     | 5, 4, 1, 3, 2, 6-7  |
| 2,3,4-Trichloroaniline          | 3.68  | 2.00              | <i>Poecilia reticulata</i>     | de Wolf <i>et al.</i> (1993)   | 3, 4, 1, 7, 6, 2, 5 |
| 2,4,5-Trichloroaniline          | 3.69  | 2.33              | <i>Poecilia reticulata</i>     | de Wolf <i>et al.</i> (1993)   | 3, 4, 1, 7, 2-6, 5  |
| 2,4,6-Trichloroaniline          | 3.69  | 2.33              | <i>Poecilia reticulata</i>     | de Wolf <i>et al.</i> (1993)   | 3, 4, 1, 7, 2-6, 5  |
| 3,4,5-Trichloroaniline          | 3.32  | 2.36              | <i>Poecilia reticulata</i>     | de Wolf <i>et al.</i> (1993)   | 2, 5, 6, 4, 1, 7, 3 |
| 1,2,3-Trichlorobenzene          | 4.14  | 2.90              | <i>Poecilia reticulata</i>     | de Wolf <i>et al.</i> (1993)   | 2, 7, 6, 1-3, 4, 5  |
|                                 | 4.14  | 3.08              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)        | 5, 6, 7, 2, 1-3, 4  |
|                                 | 4.14  | 3.28              | <i>Poecilia reticulata</i>     | Schrap and Opperhuizen (1990)  | 6, 7, 2, 1-3, 5, 4  |
|                                 | 4.14  | 3.41              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)        | 5, 6, 7, 2, 1-3, 4  |
|                                 | 4.05  | 2.95              | <i>Oncorhynchus mykiss</i>     | Veith <i>et al.</i> (1979)     | 6, 2-7, 1, 3, 4-5   |
|                                 | 4.05  | 3.11              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)        | 5, 6, 2-7, 1, 3, 4  |
|                                 | 4.05  | 3.31              | <i>Jordanella floridae</i>     | Smith <i>et al.</i> (1990)     | 5, 6, 2-7, 1, 3, 4  |
|                                 | 4.05  | 3.32              | <i>Pimephales promelas</i>     | Veith <i>et al.</i> (1979)     | 6, 2-7, 1, 3, 4-5   |
|                                 | 4.05  | 3.36              | <i>Lepomis cyanellus</i>       | Veith <i>et al.</i> (1979)     | 6, 2-7, 1, 3, 4-5   |
|                                 | 4.05  | 3.36              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 6, 5, 2-7, 1, 3, 4  |
| 1,3,5-Trichlorobenzene          | 4.05  | 3.51              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)        | 5, 6, 2-7, 1, 3, 4  |
|                                 | 4.05  | 3.57              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 6, 5, 2-7, 1, 3, 4  |
|                                 | 4.19  | 3.26              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)        | 5, 6, 7, 2, 3, 1, 4 |
|                                 | 4.19  | 3.48              | <i>Poecilia reticulata</i>     | Schrap and Opperhuizen (1990)  | 6, 7, 2, 3, 1-5, 4  |
|                                 | 4.19  | 3.61              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1983)        | 5, 6, 7, 2, 3, 1, 4 |
| 2,2',5-Trichlorobiphenyl        | 5.60  | 4.30 <sup>§</sup> | <i>Carassius auratus</i>       | Bruggeman <i>et al.</i> (1981) | 7, 3, 6, 5, 1, 2, 4 |
|                                 | 5.60  | 4.91              | <i>Oncorhynchus mykiss</i>     | Oliver and Niimi (1985)        | 5, 7, 3, 6, 1, 2, 4 |
| 2,4,5-Trichlorobiphenyl         | 5.90  | 3.78 <sup>§</sup> | <i>Poecilia reticulata</i>     | Gobas <i>et al.</i> (1989)     | 4, 6, 2, 1, 7, 3, 5 |
|                                 | 5.90  | 4.15              | <i>Poecilia reticulata</i>     | Gobas and Schrap (1990)        | 4, 6, 2, 1, 7, 3, 5 |
| 2,4',5-Trichlorobiphenyl        | 5.79  | 4.63 <sup>§</sup> | <i>Carassius auratus</i>       | Bruggeman <i>et al.</i> (1981) | 3, 7, 5, 1, 6, 2, 4 |
| 1,1,1-Trichloroethane           | 2.47  | 0.95              | <i>Lepomis macrochirus</i>     | Veith <i>et al.</i> (1980)     | 3, 5, 7, 1, 4-6, 2  |
| 1,1,2-Trichloroethylene         | 2.42  | 1.23              | <i>Lepomis macrochirus</i>     | Veith <i>et al.</i> (1980)     | 5, 1-3, 7, 4-6, 2   |
| 3,4,5-Trichloroguaiacol         | 4.11  | 2.41              | <i>Oncorhynchus mykiss</i>     | Niimi <i>et al.</i> (1990)     | 4, 1-3, 2, 7, 6, 5  |
| 4,5,6-Trichloroguaiacol         | 3.74  | 1.97              | <i>Oncorhynchus mykiss</i>     | Niimi <i>et al.</i> (1990)     | 3, 4, 1, 7, 2-6, 5  |
| 2,3,4-Trichloronitrobenzene     | 3.61  | 2.20              | <i>Oncorhynchus mykiss</i>     | Niimi <i>et al.</i> (1989)     | 3, 1-4-7, 6, 2-5    |
| 2,4,5-Trichloronitrobenzene     | 3.40  | 1.84              | <i>Oncorhynchus mykiss</i>     | Niimi <i>et al.</i> (1989)     | 3, 7, 1, 4, 5, 6, 2 |
| 2,4,6-Trichloronitrobenzene     | 3.69  | 2.88              | <i>Oncorhynchus mykiss</i>     | Niimi <i>et al.</i> (1989)     | 5, 2-6, 7, 1, 4, 3  |
| 2,4,6-Trichlorophenol           | 3.75  | 1.94              | <i>Jordanella floridae</i>     | Smith <i>et al.</i> (1990)     | 3, 4, 1, 7, 2-6, 5  |
|                                 | 3.75  | 2.37 <sup>§</sup> | <i>Oryzias latipes</i>         | Sugiura <i>et al.</i> (1984)   | 5, 3, 4, 1, 7, 2-6  |
| Tricresyl phosphate             | 5.11  | 2.22              | <i>Pimephales promelas</i>     | Veith <i>et al.</i> (1979)     | 4, 1, 2, 5, 3-6, 7  |
|                                 | 5.11  | 2.90              | <i>Alburnus alburnus</i>       | Bengtsson <i>et al.</i> (1986) | 4, 1, 2, 5, 3-6, 7  |
| Trifluralin                     | 5.34  | 3.50              | <i>Pseudorasbora parva</i>     | Kanazawa (1981)                | 4, 2, 1, 6, 5, 3, 7 |
| Triphenyl phosphate             | 4.59  | 2.60              | <i>Alburnus alburnus</i>       | Bengtsson <i>et al.</i> (1986) | 4, 1, 2-5, 3, 6, 7  |

<sup>1</sup>log BCF expressed on a lipid content basis; <sup>§</sup>log BCF estimated from the rate constants  $k_1$  and  $k_2$ ; \*Dichlorodibenzo-*p*-dioxin; <sup>§</sup>Hexa- and hepta-chlorodibenzo-*p*-dioxin; <sup>§</sup>Heptachlorodibenzofuran; \*\*Isomer not specified; <sup>§§</sup>Pentachlorodibenzo-*p*-dioxin; <sup>§§</sup>Pentachlorodibenzofuran; <sup>§</sup>Tetrachlorodibenzo-*p*-dioxin; <sup>§</sup>Tetrachlorodibenzofuran; <sup>o</sup>Classification of the models by increasing order of the absolute values of the residuals;  $\chi$ : model presenting a negative residual; x-y: models with identical residuals (absolute value).

**Table II** *n*-octanol/water partition coefficients (log P), bioconcentration data (log BCF) obtained under static and semi-static conditions and classification of the seven models.

| Chemical                                 | log P | log BCF           | Species                    | Reference                        | Classification <sup>o</sup> |
|--|-------|-------------------|----------------------------|----------------------------------|-----------------------------|
| Acridine                                 | 3.45  | 3.11              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 2, 5, 6, 4, 1, 7, 3         |
| Aniline                                  | 0.90  | 0.41              | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 2, 4, 1, 6, 5, 3, 7         |
| Anthracene                               | 4.45  | 2.95              | <i>Lepomis macrochirus</i> | Spacie <i>et al.</i> (1983)      | 4, 1, 5, 3, 2, 6, 7         |
|  | 4.45  | 3.86              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 7, 6, 2, 3, 1, 4         |
| Atrazine                                 | 2.61  | 0.78              | <i>Brachydanio rerio</i>   | Görge and Nagel (1990)           | 3, 5, 7, 1, 4, 6, 2         |
| Benzo[b]furan                            | 2.86  | 2.56              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 2, 6, 5, 4, 1, 7, 3         |
| Benzo[b]naphtho(2,3- <i>d</i> )thiophene | 5.07  | 4.17              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 7, 6, 3, 2, 1, 4         |
| Benzo[a]pyrene                           | 5.97  | 3.69              | <i>Lepomis macrochirus</i> | Spacie <i>et al.</i> (1983)      | 6, 4, 2, 1, 7, 5, 3         |
| Benzo[b]thiophene                        | 3.26  | 2.53              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 2, 5, 6, 4, 1, 7, 3         |
| 2,2'-Bithiophene                         | 3.75  | 3.55              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 2-6, 7, 1, 4, 3          |
| <i>p</i> - <i>sec</i> -Butylphenol       | 3.08  | 1.57              | <i>Salmo salar</i>         | McLeese <i>et al.</i> (1981)     | 3-5, 7, 1, 4, 6, 2          |
| <i>t</i> -Butylphenyldiphenyl phosphate  | 5.12  | 2.89              | <i>Pimephales promelas</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2, 5, 3-6, 7          |
|  | 5.12  | 3.04              | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2, 5, 3-6, 7          |
| Carbazole                                | 3.84  | 2.70              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 6, 2, 5, 7, 1, 4, 3         |
| 2-Chloroaniline                          | 1.93  | 1.18              | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 2, 7, 4, 6, 1, 5, 3         |
| 3-Chloroaniline                          | 1.91  | 1.06              | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 4, 6, 1, 2, 7, 5, 3         |
| 4-Chloroaniline                          | 1.88  | 0.91              | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 1-6, 4, 2, 5, 3, 7          |
| 2-Chloronaphthalene                      | 4.14  | 3.63              | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1985) | 6, 7, 2, 1-3, 5, 4          |
| 2-Chloronitrobenzene                     | 2.52  | 2.29 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 3, 5, 7, 1, 4, 6, 2         |
| 3-Chloronitrobenzene                     | 2.50  | 2.42 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 5, 7, 1, 3, 4, 6, 2         |
| 4-Chloronitrobenzene                     | 2.39  | 2.46 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 7, 1-4-6, 5, 2, 3           |
| 2-Chloro-6-nitrotoluene                  | 3.09  | 3.09 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 5, 4-6, 1, 2, 7, 3          |
| 4-Chloro-2-nitrotoluene                  | 3.05  | 3.02 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 4, 5, 1, 6, 7, 2, 3         |
| 2,8-DCDD*                                | 5.60  | 2.83              | <i>Carassius auratus</i>   | Sijm and Opperhuizen (1988)      | 5, 4, 2, 1, 6, 3, 7         |
| Dibenzo(1,4)dioxan                       | 4.19  | 3.85              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 6, 7, 2, 3, 1, 4         |
| Dibenzofuran                             | 4.21  | 3.54              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 6, 7, 2, 3, 1, 4         |
| Dibenzothiophene                         | 4.49  | 3.82              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 7, 6, 2, 3, 1, 4         |
| 1,2-Dibromobenzene                       | 3.64  | 2.70              | <i>Pimephales promelas</i> | Sijm <i>et al.</i> (1993)        | 2, 6, 7, 1-4, 3, 5          |
|  | 3.64  | 3.50              | <i>Poecilia reticulata</i> | Sijm <i>et al.</i> (1993)        | 5, 2, 6, 7, 1-4, 3          |
| 1,4-Dibromobenzene                       | 3.79  | 2.70              | <i>Poecilia reticulata</i> | Sijm <i>et al.</i> (1993)        | 6, 2, 7, 1, 4, 3, 5         |
|  | 3.79  | 3.40              | <i>Pimephales promelas</i> | Sijm <i>et al.</i> (1993)        | 6, 2, 7-5, 1, 4, 3          |
| 2,4-Dichloroaniline                      | 2.91  | 1.98              | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 2, 6, 4, 1, 7, 3-5          |
| 3,4-Dichloroaniline                      | 2.78  | 1.48              | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 3-5, 7, 1, 4, 6, 2          |
| 1,2-Dichlorobenzene                      | 3.43  | 2.40              | <i>Poecilia reticulata</i> | Sijm <i>et al.</i> (1993)        | 2, 6, 5, 1-4, 7, 3          |
|  | 3.43  | 2.70              | <i>Pimephales promelas</i> | Sijm <i>et al.</i> (1993)        | 2, 6, 1-4, 5, 7, 3          |
| 1,3-Dichlorobenzene                      | 3.53  | 3.78 <sup>†</sup> | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1988) | 2, 6, 1-4, 7, 3-5           |
| 1,4-Dichlorobenzene                      | 3.44  | 1.70              | <i>Poecilia reticulata</i> | Sijm <i>et al.</i> (1993)        | 5, 3, 7, 1, 4, 6, 2         |
|  | 3.44  | 2.40              | <i>Pimephales promelas</i> | Sijm <i>et al.</i> (1993)        | 2, 6, 4, 1-5, 7, 3          |
| 1,4-Dichloronaphthalene                  | 4.66  | 3.36              | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1985) | 3, 5, 2, 6, 1, 7, 4         |
| 1,8-Dichloronaphthalene                  | 4.19  | 3.79              | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1985) | 6, 7, 2, 3, 1-5, 4          |
| 2,3-Dichloronaphthalene                  | 4.51  | 4.04              | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1985) | 7, 6, 2, 3, 5, 1, 4         |
| 2,3-Dichloronitrobenzene                 | 3.05  | 3.01 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 4, 1, 5, 6, 7, 2-3          |
| 2,4-Dichloronitrobenzene                 | 3.05  | 3.02 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 4, 5, 1, 6, 7, 2, 3         |
| 2,5-Dichloronitrobenzene                 | 3.03  | 2.92 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 7, 1, 4, 5, 3, 6, 2         |
| 3,5-Dichloronitrobenzene                 | 3.09  | 3.01 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 1, 4, 7-5, 6, 3, 2          |
| 2,3-Dimethylnitrobenzene                 | 2.83  | 2.86 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 4, 6, 5, 1, 7, 2, 3         |
| 3,4-Dimethylnitrobenzene                 | 2.91  | 2.84 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 1, 7, 5, 4, 6, 3, 2         |
| 1,2-Dinitrobenzene                       | 1.69  | 1.02 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 3, 5, 1, 6, 4, 2, 7         |
| 1,3-Dinitrobenzene                       | 1.49  | 1.87 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 4, 2, 1-6, 5, 3, 7          |
| 2,4-Dinitrotoluene                       | 1.98  | 2.31 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 7, 2, 4, 6, 1, 5, 3         |
| 2,6-Dinitrotoluene                       | 2.10  | 2.44 <sup>†</sup> | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 2, 7, 4, 6, 1, 5, 3         |
| <i>p</i> -Dodecylphenol                  | 7.91  | 3.78              | <i>Salmo salar</i>         | McLeese <i>et al.</i> (1981)     | 7, 6, 5, 2, 4, 1, 3         |
| Fluorene                                 | 4.18  | 3.35              | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 6, 7, 2, 3, 1, 4         |
| Hexachlorobenzene                        | 5.73  | 5.62 <sup>†</sup> | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1988) | 7, 5, 3, 1, 6, 2, 4         |
| $\gamma$ -Hexachlorocyclohexane          | 3.72  | 2.34              | <i>Brachydanio rerio</i>   | Görge and Nagel (1990)           | 3-5, 1-4, 7, 2-6            |
|  | 3.72  | 2.41              | <i>Salmo salar</i>         | Carlberg <i>et al.</i> (1986)    | 3-5, 1-4, 7, 2-6            |
|  | 3.72  | 2.84              | <i>Salmo salar</i>         | Carlberg <i>et al.</i> (1986)    | 2-6, 7, 1-4, 3-5            |

**Table II (continued)**

| Chemical                           | log P | log BCF <sup>Species</sup> | Reference                  | Classification <sup>o</sup>      |                     |
|------------------------------------|-------|----------------------------|----------------------------|----------------------------------|---------------------|
| 2-Nitroaniline                     | 1.85  | 0.91                       | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 6, 1, 4, 2, 7, 5, 3 |
| 3-Nitroaniline                     | 1.37  | 0.92                       | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 2, 4, 1-6, 7, 5, 3  |
| 4-Nitroaniline                     | 1.39  | 0.64                       | <i>Brachydanio rerio</i>   | Kalsch <i>et al.</i> (1991)      | 4, 1-6, 2, 5, 3, 7  |
| Nitrobenzene                       | 1.85  | 1.47 <sup>†</sup>          | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 3, 5, 1, 6, 4, 2, 7 |
| 2-Nitrotoluene                     | 2.30  | 2.28 <sup>†</sup>          | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 5, 1, 6, 7, 4, 3, 2 |
| 3-Nitrotoluene                     | 2.42  | 2.21 <sup>†</sup>          | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 5, 3, 1, 7, 4, 6, 2 |
| 4-Nitrotoluene                     | 2.37  | 2.37 <sup>†</sup>          | <i>Poecilia reticulata</i> | Deneer <i>et al.</i> (1987)      | 5, 1, 7, 6, 4, 3, 2 |
| <i>p</i> -Nonylphenol              | 5.76  | 2.45                       | <i>Salmo salar</i>         | McLeese <i>et al.</i> (1981)     | 4, 2, 6, 1, 7, 5, 3 |
| Pentachloroaniline                 | 5.08  | 3.78                       | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1992)     | 6, 3, 7, 2, 1, 5, 4 |
| Pentachlorobenzene                 | 5.18  | 3.45                       | <i>Carassius auratus</i>   | Sijm and Opperhuizen (1988)      | 4, 1, 2, 6, 3, 5, 7 |
|                                    | 5.18  | 5.18 <sup>†</sup>          | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1988) | 3, 5-6, 7, 2, 1, 4  |
|                                    | 5.18  | 4.23                       | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1985) | 7, 3, 5-6, 2, 1, 4  |
| Pyrene                             | 5.18  | 3.68                       | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 1, 2, 4, 6, 3, 7, 5 |
| 2,3,7,8-TCDD <sup>‡</sup>          | 6.42  | 3.90                       | <i>Pimephales promelas</i> | Adams <i>et al.</i> (1986)       | 6, 7, 4, 2, 1, 5, 3 |
| 2,3,4,5-Tetrachloroaniline         | 4.57  | 3.28                       | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1992)     | 3, 2, 6, 1, 7, 4, 5 |
| 2,3,5,6-Tetrachloroaniline         | 4.46  | 3.59                       | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1992)     | 5, 7, 6, 2, 3, 1, 4 |
| 1,2,3,4-Tetrachlorobenzene         | 4.64  | 4.74 <sup>†</sup>          | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1988) | 6, 7, 3, 2-5, 1, 4  |
| 3,3',4,4'-Tetrachlorobiphenyl      | 6.63  | 4.59                       | <i>Poecilia reticulata</i> | Opperhuizen and Voors (1987)     | 7, 4, 2, 1, 6, 5, 3 |
| 3,3',4,4'-Tetrachlorodiphenylether | 5.78  | 4.51                       | <i>Poecilia reticulata</i> | Opperhuizen and Voors (1987)     | 3, 5, 7, 1, 6, 2, 4 |
| Tetrachloroguaiacol                | 4.45  | 3.15                       | <i>Salmo salar</i>         | Carlberg <i>et al.</i> (1986)    | 2, 3, 5, 1, 6, 4, 7 |
| Thianthrene                        | 4.47  | 3.56                       | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 7, 6, 2, 3, 1, 4 |
| 2,3,4-Trichloroaniline             | 3.68  | 2.61                       | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1992)     | 5, 2, 6, 7, 1, 4, 3 |
| 2,4,5-Trichloroaniline             | 3.69  | 2.88                       | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1992)     | 5, 2-6, 7, 1, 4, 3  |
| 2,4,6-Trichloroaniline             | 3.69  | 3.13                       | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1992)     | 5, 2-6, 7, 1, 4, 3  |
| 3,4,5-Trichloroaniline             | 3.32  | 3.04                       | <i>Poecilia reticulata</i> | de Wolf <i>et al.</i> (1992)     | 2, 5-6, 4, 1, 7, 3  |
| 1,3,5-Trichlorobenzene             | 4.19  | 4.35 <sup>†</sup>          | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1988) | 6, 7, 2, 3, 1-5, 4  |
| 2,4,5-Trichlorobiphenyl            | 5.90  | 4.26                       | <i>Poecilia reticulata</i> | Opperhuizen and Voors (1987)     | 2, 6, 1-4, 7, 5, 3  |
| 2,4,5-Trichlorodiphenylether       | 5.44  | 4.18                       | <i>Poecilia reticulata</i> | Opperhuizen and Voors (1987)     | 7, 3, 5, 6, 1, 2, 4 |
| 1,3,7-Trichloronaphthalene         | 5.35  | 4.43                       | <i>Poecilia reticulata</i> | Opperhuizen <i>et al.</i> (1985) | 7, 3, 5, 6, 1, 2, 4 |
| 2,4,6-Trichlorophenol              | 3.75  | 2.84                       | <i>Salmo salar</i>         | Carlberg <i>et al.</i> (1986)    | 2-6, 7, 1, 4, 3, 5  |
| Tri- <i>m</i> -cresyl phosphate    | 5.11  | 2.78                       | <i>Pimephales promelas</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2, 5, 3-6, 7  |
|                                    | 5.11  | 2.89                       | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2, 5, 3-6, 7  |
| Tri- <i>p</i> -cresyl phosphate    | 5.11  | 2.97                       | <i>Pimephales promelas</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2, 5, 3-6, 7  |
|                                    | 5.11  | 3.15                       | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2, 5, 3-6, 7  |
| Triphenyl phosphate                | 4.59  | 2.75                       | <i>Pimephales promelas</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2-5, 3, 6, 7  |
|                                    | 4.59  | 2.76                       | <i>Oncorhynchus mykiss</i> | Muir <i>et al.</i> (1983)        | 4, 1, 2-5, 3, 6, 7  |
| Xanthene                           | 4.23  | 3.62                       | <i>Poecilia reticulata</i> | de Voogt <i>et al.</i> (1991)    | 5, 6-7, 2, 3, 1, 4  |

<sup>†</sup>log BCF expressed on a lipid content basis; <sup>‡</sup>log BCF estimated from the rate constants  $k_1$  and  $k_2$ ; \*Dichlorodibenzo-*p*-dioxin; <sup>‡</sup>Tetrachlorodibenzo-*p*-dioxin; <sup>o</sup>Classification of the models by increasing order of the absolute values of the residuals; x: model presenting a negative residual; x-y: models with identical residuals (absolute value).

**Table III RMS values for the seven selected models.**

|                              | Eq. (1)* | Eq. (2) | Eq. (3) | Eq. (4) | Eq. (5) | Eq. (6) | Eq. (7) |
|------------------------------|----------|---------|---------|---------|---------|---------|---------|
| <b>Flow-through</b>          |          |         |         |         |         |         |         |
| log P ≤ 6 (n = 267)          | 0.58     | 0.59    | 0.60    | 0.58    | 0.60    | 0.59    | 0.64    |
| log P > 6 (n = 75)           | 1.93     | 1.78    | 2.32    | 1.79    | 1.83    | 0.86    | 1.13    |
| Total (n = 342)              | 1.04     | 0.99    | 1.21    | 0.98    | 1.01    | 0.66    | 0.77    |
| <b>Static or semi-static</b> |          |         |         |         |         |         |         |
| Total (n = 94)               | 0.58     | 0.55    | 0.67    | 0.57    | 0.55    | 0.51    | 0.60    |

\* (1) Veith *et al.* (1979); (2) Veith *et al.* (1980); (3) Mackay (1982); (4) Isnard and Lambert (1988); (5) Nendza (1991); (6) Bintein *et al.* (1993); (7) Connell and Hawker (1988).

Residuals




Figure 1 Residuals of Eq. (1) vs log P for BCF data obtained under flow-through conditions.

Residuals




Figure 2 Residuals of Eq. (1) vs log P for BCF data obtained under static and semi-static conditions.

Residuals




Figure 3 Residuals of Eq. (2) vs log P for BCF data obtained under flow-through conditions.

Residuals




Figure 4 Residuals of Eq. (2) vs log P for BCF data obtained under static and semi-static conditions.

Residuals




Figure 5 Residuals of Eq. (3) vs log P for BCF data obtained under flow-through conditions.

Residuals




Figure 6 Residuals of Eq. (3) vs log P for BCF data obtained under static and semi-static conditions.




Figure 7 Residuals of Eq. (4) vs log P for BCF data obtained under flow-through conditions.




Figure 8 Residuals of Eq. (4) vs log P for BCF data obtained under static and semi-static conditions.




Figure 9 Residuals of Eq. (5) vs log P for BCF data obtained under flow-through conditions.




Figure 10 Residuals of Eq. (5) vs log P for BCF data obtained under static and semi-static conditions.




Figure 11 Residuals of Eq. (6) vs log P for BCF data obtained under flow-through conditions.




Figure 12 Residuals of Eq. (6) vs log P for BCF data obtained under static and semi-static conditions.




Figure 13 Residuals of Eq. (7) vs log P for BCF data obtained under flow-through conditions.




Figure 14 Residuals of Eq. (7) vs log P for BCF data obtained under static and semi-static conditions.

octachlorodibenzo-*p*-dioxin (Figure 11). This underlines the necessity to estimate the performances of a model from different values for a same chemical.

Last, to obtain a general approximation of the predictive performances of the studied models, the RMS (i.e., root mean square error) values (Eq. (8)) for the seven studied models were calculated (Table III).

$$\text{RMS} = ((\sum (\log \text{BCF}_{\text{obs.}} - \log \text{BCF}_{\text{calc.}})^2) / \text{number of observations})^{1/2} \quad (8)$$

The results reported in Table III confirm that for compounds with log P values inferior to 6, the different models yield equivalent results. At the opposite, for molecules presenting log P values superior to 6, the bilinear model of Bintein *et al.* (1993) is largely superior to the other studied models. From our results, it appears that the different models under study do not present the same predictive power and domain of application. This can be problematic if they are incorporated in different computer programs designed to support decision-making in the evaluation of the hazards and risks of chemicals. For highly hydrophobic chemicals, the linear models (i.e., Eqs. (1) to (4)) tend to considerably overestimate the BCF values and Eq. (5) often gives unacceptable predictions. Eq. (7) provides better simulations than Eq. (5) but the statistical validity of this model is questionable. From a risk assessment point of view, highly conservative models could be judged interesting. However, from a modeling point of view, it is obvious that the first requirement for a model is to simulate with accuracy the reality. Therefore, among the seven models, we can consider that Eq. (6) (Bintein *et al.*, 1993) meets at best this objective. Indeed, it presents the best predictive power and the widest domain of application. However, it is important to stress that we do not claim that this model allows to simulate the bioconcentration behavior of all the organic chemicals susceptible to contaminate the aquatic ecosystems. Indeed, we simply assume that on the basis of BCF QSAR models using the *n*-octanol/water partition coefficient, it is the most efficient. It is obvious that all these models do not take into account the transformation of the chemicals in the fish. The lack of this information in the design of the models can explain some of the bad simulations obtained for the seven studied BCF models.

#### Acknowledgment

The work presented in this paper was supported by different contracts from the French Ministry of the Environment.

## References

- Adams, W.J., de Graeve, G.M., Sabourin, T.D., Cooney, J.D., and Mosher, G.M. (1986). Toxicity and bioconcentration of 2,3,7,8-TCDD to fathead minnows (*Pimephales promelas*). *Chemosphere* **15**, 1503-1511.
- Barron, M.G. (1990). Bioconcentration. *Environ. Sci. Technol.* **24**, 1612-1618.
- Bengtsson, B.E., Tarkpea, M., Sletten, T., Carlberg, G.E., Kringstad, A., and Renberg, L. (1986). Bioaccumulation and effects of some technical triaryl phosphate products in fish and *Nitocra spinipes*. *Environ. Toxicol. Chem.* **5**, 853-861.
- Bintein, S., Devillers, J., and Karcher, W. (1993). Nonlinear dependence of fish bioconcentration on *n*-octanol/water partition coefficient. *SAR QSAR Environ. Res.* **1**, 29-39.
- Bouraly, M. and Millischer, R.J. (1989). Bioaccumulation and elimination of tetrachlorobenzyltoluene (TCBT) by the rat and by fish. *Chemosphere* **18**, 2051-2063.
- Bruggeman, W.A., Martron, L.B.J.M., Kooiman, D., and Hutzinger, O. (1981). Accumulation and elimination kinetics of di-, tri- and tetra chlorobiphenyls by goldfish after dietary and aqueous exposure. *Chemosphere* **10**, 811-832.
- Bruggeman, W.A., Opperhuizen, A., Wijbenga, A., and Hutzinger, O. (1984). Bioaccumulation of super-lipophilic chemicals in fish. *Toxicol. Environ. Chem.* **7**, 173-189.
- Burke, A.B., Millburn, P., Huckle, K.R., and Hutson, D.H. (1991). Uptake and elimination of radioactivity from rainbow trout (*Oncorhynchus mykiss*) following exposure to [<sup>14</sup>C-ring]-2-phenyldodecane. *Environ. Toxicol. Chem.* **10**, 1467-1477.
- Butte, W., Fox, K., and Zauke, G.P. (1991). Kinetics of bioaccumulation and clearance of isomeric hexachlorocyclohexanes. *Sci. Total Environ.* **109/110**, 377-382.
- Butte, W., Willig, A., and Zauke, G.P. (1987). Bioaccumulation of phenols in zebrafish determined by a dynamic flow through test. In, *QSAR in Environmental Toxicology-II* (K.L.E. Kaiser, Ed.). D. Reidel Publishing Company, Dordrecht, pp. 43-53.
- Call, D.J., Brooke, L.T., Kent, R.J., Knuth, M.L., Poirier, S.H., Huot, J.M., and Lima, A.R. (1987). Bromacil and diuron herbicides: toxicity, uptake, and elimination in freshwater fish. *Arch. Environ. Contam. Toxicol.* **16**, 607-613.
- Carlberg, G.E., Martinsen, K., Kringstad, A., Gjessing, E., Grande, M., Källqvist, T., and Skare, J.U. (1986). Influence of aquatic humus on the bioavailability of chlorinated micropollutants in Atlantic salmon. *Arch. Environ. Contam. Toxicol.* **15**, 543-548.
- Connell, D.W. and Hawker, D.W. (1988). Use of polynomial expressions to describe the bioconcentration of hydrophobic chemicals by fish. *Ecotoxicol. Environ. Safety* **16**, 242-257.
- de Bruijn, J. and Hermens, J. (1991). Uptake and elimination kinetics of organophosphorous pesticides in the guppy (*Poecilia reticulata*): correlations with the octanol/water partition coefficient. *Environ. Toxicol. Chem.* **10**, 791-804.
- de Voogt, P., van Hattum, B., Leonards, P., Klamer, J.C., and Govers, H. (1991). Bioconcentration of polycyclic heteroaromatic hydrocarbons in the guppy (*Poecilia reticulata*). *Aquat. Toxicol.* **20**, 169-194.
- de Wolf, W., de Bruijn, J.H.M., Seinen, W., and Hermens, J.L.M. (1992). Influence of biotransformation on the relationship between bioconcentration factors and octanol-water partition coefficients. *Environ. Sci. Technol.* **26**, 1197-1201.

- de Wolf, W., Seinen, W., and Hermens, J.L.M. (1993). Biotransformation and toxicokinetics of trichloroanilines in fish in relation to their hydrophobicity. *Arch. Environ. Contam. Toxicol.* **25**, 110-117.
- de Wolf, W., Yedema, E.S.E., Seinen, W., and Hermens, J.L.M. (1994). Bioconcentration kinetics of chlorinated anilines in guppy, *Poecilia reticulata*. *Chemosphere* **28**, 159-167.
- Deneer, J.W., Sinnige, T.L., Seinen, W., and Hermens, J.L.M. (1987). Quantitative structure-activity relationships for the toxicity and bioconcentration factor of nitrobenzene derivatives towards the guppy (*Poecilia reticulata*). *Aquat. Toxicol.* **10**, 115-129.
- ECETOC (1995). *The Role of Bioaccumulation in Environmental Risk Assessment: The Aquatic Environment and Related Food Webs*. Technical Report no. 67, p. 112.
- EEC (1995). *Technical Guidance Documents in Support of the Commission Regulation (EC) n° 1488/94 on Risk Assessment for Existing Substances in Accordance With Council Regulation (EEC) n° 793/93*.
- Fujiwara, Y., Kinoshita, T., Sato, H., and Kojima, I. (1984). Biodegradation and bioconcentration of alkyl ethers. *Yukagaku* **33**, 111-114.
- Gobas, F.A.P.C., Clark, K.E., Shiu, W.Y., and Mackay, D. (1989). Bioconcentration of polybrominated benzenes and biphenyls and related superhydrophobic chemicals in fish: role of bioavailability and elimination into the feces. *Environ. Toxicol. Chem.* **8**, 231-245.
- Gobas, F.A.P.C. and Schrap, S.M. (1990). Bioaccumulation of some polychlorinated dibenzo-*p*-dioxins and octachlorodibenzofuran in the guppy (*Poecilia reticulata*). *Chemosphere* **20**, 495-512.
- Görge, G. and Nagel, R. (1990). Kinetics and metabolism of <sup>14</sup>C-lindane and <sup>14</sup>C-atrazine in early life stages of zebrafish (*Brachydanio rerio*). *Chemosphere* **21**, 1125-1137.
- Isnard, P. and Lambert, S. (1988). Estimating bioconcentration factors from octanol-water partition coefficient and aqueous solubility. *Chemosphere* **17**, 21-34.
- Jimenez, B.D., Cirno, C.P., and McCarthy, J.F. (1987). Effects of feeding and temperature on uptake, elimination and metabolism of benzo(a)pyrene in the bluegill sunfish (*Lepomis macrochirus*). *Aquat. Toxicol.* **10**, 41-57.
- Kalsch, W., Nagel, R., and Urich, K. (1991). Uptake, elimination, and bioconcentration of ten anilines in zebrafish (*Brachydanio rerio*). *Chemosphere* **22**, 351-363.
- Kanazawa, J. (1978). Bioconcentration ratio of diazinon by freshwater fish and snail. *Bull. Environ. Contam. Toxicol.* **20**, 613-617.
- Kanazawa, J. (1981). Measurement of the bioconcentration factors of pesticides by freshwater fish and their correlation with physicochemical properties or acute toxicities. *Pestic. Sci.* **12**, 417-424.
- Loonen, H., Tonkes, M., Parsons, J.R., and Govers, H.A.J. (1994). Bioconcentration of polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans in guppies after aqueous exposure to a complex PCDD/PCDF mixture: relationship with molecular structure. *Aquat. Toxicol.* **30**, 153-169.
- Mackay, D. (1982). Correlation of bioconcentration factors. *Environ. Sci. Technol.* **16**, 274-278.
- Mayer, F.L. (1976). Residue dynamics of di-2-ethylhexyl phthalate in fathead minnows (*Pimephales promelas*). *J. Fish. Res. Board Can.* **33**, 2610-2613.
- McCarthy, J.F. and Jimenez, B.D. (1985). Reduction in bioavailability to bluegills of polycyclic aromatic hydrocarbons bound to dissolved humic material. *Environ. Toxicol. Chem.* **4**, 511-521.
- McLeese, D.W., Zitko, V., Sergeant, D.B., Burrige, L., and Metcalfe, C.D. (1981). Lethality and accumulation of alkylphenols in aquatic fauna. *Chemosphere* **10**, 723-730.



- Mehrle, P.M., Buckler, D.R., Little, E.E., Smith, L.M., Petty, J.D., Peterman, P.H., Stalling, D.L., de Graeve, G.M., Coyle, J.J., and Adams, W.J. (1988). Toxicity and bioconcentration of 2,3,7,8-tetrachlorodibenzodioxin and 2,3,7,8-tetrachlorodibenzofuran in rainbow trout. *Environ. Toxicol. Chem.* **7**, 47-62.
- Muir, D.C.G., Hobden, B.R., and Servos, M.R. (1994). Bioconcentration of pyrethroid insecticides and DDT by rainbow trout: uptake, depuration, and effect of dissolved organic carbon. *Aquat. Toxicol.* **29**, 223-240.
- Muir, D.C.G., Marshall, W.K., and Webster, G.R.B. (1985). Bioconcentration of PCDDs by fish: effects of molecular structure and water chemistry. *Chemosphere* **14**, 829-833.
- Muir, D.C.G., Yarechewski, A.L., and Grift, N.P. (1983). Environmental dynamics of phosphate esters. III. Comparison of the bioconcentration of four triaryl phosphates by fish. *Chemosphere* **12**, 155-166.
- Nendza, M. (1991). QSARs of bioconcentration: validity assessment of log Pow/log BCF correlations. In, *Bioaccumulation in Aquatic Systems* (R. Nagel and R. Loskill, Eds.). VCH, Weinheim, pp. 43-66.
- Niimi, A.J., Lee, H.B., and Kisson, G.P. (1989). Octanol/water partition coefficients and bioconcentration factors of chloronitrobenzenes in rainbow trout (*Salmo gairdneri*). *Environ. Toxicol. Chem.* **8**, 817-823.
- Niimi, A.J., Lee, H.B., and Kisson, G.P. (1990). Kinetics of chloroguaiacols and other chlorinated phenolic derivatives in rainbow trout (*Salmo gairdneri*). *Environ. Toxicol. Chem.* **9**, 649-653.
- Oliver, B.G. and Niimi, A.J. (1983). Bioconcentration of chlorobenzenes from water by rainbow trout: correlations with partition coefficients and environmental residues. *Environ. Sci. Technol.* **17**, 287-291.
- Oliver, B.G. and Niimi, A.J. (1984). Rainbow trout bioconcentration of some halogenated aromatics from water at environmental concentrations. *Environ. Toxicol. Chem.* **3**, 271-277.
- Oliver, B.G. and Niimi, A.J. (1985). Bioconcentration factors of some halogenated organics for rainbow trout: limitations in their use for prediction of environmental residues. *Environ. Sci. Technol.* **19**, 842-849.
- Opperhuizen, A. and Schrap, S.M. (1987). Relationships between aqueous oxygen concentration and uptake and elimination rates during bioconcentration of hydrophobic chemicals in fish. *Environ. Toxicol. Chem.* **6**, 335-342.
- Opperhuizen, A., Serné, P., and van der Steen, J.M.D. (1988). Thermodynamics of fish/water and octan-1-ol/water partitioning of some chlorinated benzenes. *Environ. Sci. Technol.* **22**, 286-292.
- Opperhuizen, A., van der Velde, E.W., Gobas, F.A.P.C., Liem, D.A.K., and van der Steen, J.M.D. (1985). Relationship between bioconcentration in fish and steric factors of hydrophobic chemicals. *Chemosphere* **14**, 1871-1896.
- Opperhuizen, A. and Voors, P.I. (1987). Bioconcentration kinetics of 2,4,5-tri- and 3,3',4,4'-tetrachlorobiphenyl and 2,4,5-tri- and 3,3',4,4'-tetrachlorodiphenylether in fish. *Chemosphere* **16**, 2379-2388.
- Schrap, S.M. and Opperhuizen, A. (1990). Relationship between bioavailability and hydrophobicity: reduction of the uptake of organic chemicals by fish due to the sorption on particles. *Environ. Toxicol. Chem.* **9**, 715-724.
- Seguchi, K. and Asaka, S. (1981). Intake and excretion of diazinon in freshwater fishes. *Bull. Environ. Contam. Toxicol.* **27**, 244-249.
- Servos, M.R., Muir, D.C.G., and Webster, G.R.B. (1989). The effect of dissolved organic matter on the bioavailability of polychlorinated dibenzo-*p*-dioxins. *Aquat. Toxicol.* **14**, 169-184.
- Sijm, D.T.H.M. and Opperhuizen, A. (1988). Biotransformation, bioaccumulation and lethality of 2,8-dichlorodibenzo-*p*-dioxin: a proposal to explain the biotic fate and toxicity of PCDD's and PCDF's. *Chemosphere* **17**, 83-99.

- Sijm, D.T.H.M., Schipper, M., and Opperhuizen, A. (1993). Toxicokinetics of halogenated benzenes in fish: lethal body burden as a toxicological end point. *Environ. Toxicol. Chem.* **12**, 1117-1127.
- Smith, A.D., Bharath, A., Mallard, C., Orr, D., McCarty, L.S., and Ozburn, G.W. (1990). Bioconcentration kinetics of some chlorinated benzenes and chlorinated phenols in american flagfish, *Jordanella floridae* (Goode and Bean). *Chemosphere* **20**, 379-386.
- Southworth, G.R., Parkhurst, B.R., and Beauchamp, J.J. (1979). Accumulation of acridine from water, food, and sediment by the fathead minnow, *Pimephales promelas*. *Water Air Soil Pollut.* **12**, 331-341.
- Spacie, A., Landrum, P.F., and Laversee, G.J. (1983). Uptake, depuration, and biotransformation of anthracene and benzo[a]pyrene in bluegill sunfish. *Ecotoxicol. Environ. Safety* **7**, 330-341.
- Spehar, R.L., Tanner, D.K., and Nordling, B.R. (1983). Toxicity of the synthetic pyrethroids, permethrin and AC 222, 705 and their accumulation in early life stages of fathead minnows and snails. *Aquat. Toxicol.* **3**, 171-182.
- Sugiura, K., Aoki, M., Kaneko, S., Daisaku, I., Komatsu, Y., Shibuya, H., Suzuki, H., and Goto, M. (1984). Fate of 2,4,6-trichlorophenol, pentachlorophenol, *p*-chlorobiphenyl, and hexachlorobenzene in an outdoor experimental pond: comparison between observations and predictions based on laboratory data. *Arch. Environ. Contam. Toxicol.* **13**, 745-758.
- Tachikawa, M., Sawamura, R., Okada, S., and Hamada, A. (1991). Differences between freshwater and seawater killifish (*Oryzias latipes*) in the accumulation and elimination of pentachlorophenol. *Arch. Environ. Contam. Toxicol.* **21**, 146-151.
- Takimoto, Y., Ohshima, M., and Miyamoto, J. (1987). Comparative metabolism of fenitrothion in aquatic organisms. I. Metabolism in the euryhaline fish, *Oryzias latipes* and *Mugil cephalus*. *Ecotoxicol. Environ. Safety* **13**, 104-117.
- Takimoto, Y., Ohshima, M., Yamada, H., and Miyamoto, J. (1984). Fate of fenitrothion in several developmental stages of the killifish (*Oryzias latipes*). *Arch. Environ. Contam. Toxicol.* **13**, 579-587.
- Tsuda, T., Aoki, S., Inoue, T., and Kojima, M. (1995). Accumulation and excretion of diazinon, fenthion and fenitrothion by killifish: comparison of individual and mixed pesticides. *Water Res.* **29**, 455-458.
- Tsuda, T., Aoki, S., Kojima, M., and Fujita, T. (1992). Accumulation and excretion of organophosphorous pesticides by willow shiner. *Chemosphere* **25**, 1945-1951.
- Tsuda, T., Aoki, S., Kojima, M., and Fujita, T. (1993). Accumulation and excretion of chloroanilines by carp. *Chemosphere* **26**, 2301-2306.
- Tsuda, T., Aoki, S., Kojima, M., and Harada, H. (1988). Bioconcentration and excretion of benthocarb and simetryne by willow shiner. *Toxicol. Environ. Chem.* **18**, 31-36.
- Veith, G.D., DeFoe, D.L., and Bergstedt, B.V. (1979). Measuring and estimating the bioconcentration factor of chemicals in fish. *J. Fish. Res. Board Can.* **36**, 1040-1048.
- Veith, G.D., Macek, K.J., Petrocelli, S.R., and Carroll, J. (1980). An evaluation of using partition coefficients and water solubility to estimate bioconcentration factors for organic chemicals in fish. In, *Aquatic Toxicology, ASTM STP 707* (J.G. Eaton, P.R. Parrish and A.C. Hendricks, Eds.). pp. 116-129.
- Vigano, L., Galassi, S., and Gatto, M. (1992). Factors affecting the bioconcentration of hexachlorocyclohexanes in early life stages of *Oncorhynchus mykiss*. *Environ. Toxicol. Chem.* **11**, 535-540.

## RESUME


L'élaboration d'une équation bilinéaire reliant le coefficient de partage *n*-octanol/eau (log P) à la bioconcentration des molécules organiques chez les poissons a permis de mieux simuler le comportement bioaccumulatif des substances très lipophiles. L'adsorption par les sols et les sédiments des substances ionisées et non-ionisées a pu être modélisée par une équation régressive prenant en compte le log P, le potentiel d'ionisation des molécules, le pourcentage en carbone organique et le pH des substrats. L'intégration de ces deux relations dans CHEMFRANCE a élargi le domaine d'application de ce modèle régional de fugacité niveau III appliqué à la France. Des études menées sur le lindane et l'atrazine ont démontré le haut potentiel de simulation de CHEMFRANCE pour ces pesticides. Enfin, une analyse comparative des principales relations structure-bioconcentration utilisant le log P comme descripteur moléculaire a été réalisée afin de tester l'intérêt du modèle que nous proposons.

**Mots clés:** CHEMFRANCE; bioconcentration; adsorption; QSAR; validation; lindane; atrazine.

## SUMMARY

The elaboration of a bilinear relationship between the *n*-octanol/water partition coefficient (log P) and the accumulation of chemicals in fish allowed to simulate more accurately the bioconcentration potential of very lipophilic molecules. The soil and sediment adsorption of ionized and non-ionized chemicals has been modeled by a regression equation integrating log P, the ionization potential of compounds, the percentage of organic carbon and the pH of the substrates. The incorporation of these two equations in CHEMFRANCE allowed to increase the domain of application of this regional level III fugacity model applied to France. Studies performed on lindane and atrazine underlined the high simulation potential of this model for these pesticides. Last, a comparative analyse of structure-bioconcentration relationships using log P as molecular descriptor was performed in order to test the usefulness of the model we proposed.

**Key words:** CHEMFRANCE; bioconcentration; adsorption; QSAR; validation; lindane; atrazine.



UNIVERSITÄT BAYREUTH

PROF. DR. O. HUTZINGER

CHAIR OF ECOLOGICAL CHEMISTRY  
AND GEOCHEMISTRY, UNIVERSITY BAYREUTH

Universitätsstraße 30 · D-95440 Bayreuth  
Germany

---

Dr. J. Devillers  
Centre de Traitement de l'Informatin  
Scientifique  
21 rue de la Banniere  
69003 L y o n

France

May 28, 1996

AH/G

**RE: COMPARISON OF BCF MODELS BASED ON LOG P**

Dear Dr. Devillers:

We are pleased to advise you that the revised manuscript has been accepted for publication in the journal CHEMOSPHERE. The paper is being forwarded to Elsevier Science together with your request for 100 reprints.

Enclosed is a copyright agreement form. Kindly fill it in and mail it to the Production Editor, Chemosphere, Elsevier Science, The Boulevard, Langford Lane, Kidlington, Oxford OX5 1GB, England.

Sincerely yours,

Alfreda Hutzinger  
Assistant Editor



## COMPARISON OF BCF MODELS BASED ON LOG P

J. Devillers,\* S. Bintein, and D. Domine

CTIS, 21 rue de la Bannière, 69003 Lyon (France)

(Received in Germany 9 January 1996; accepted 24 May 1996)

### ABSTRACT

Seven linear and nonlinear BCF models based on log P have been compared in order to estimate their accuracy, predictive power, and domain of application. This comparison was performed from a data set constituted of 436 experimental BCF values recorded for 227 chemicals. We showed that for chemicals with  $\log P < 6$ , the different models yielded equivalent results. At the opposite, for highly hydrophobic chemicals ( $\log P > 6$ ), the bilinear model  $\log \text{BCF} = 0.91 \log P - 1.975 \log (6.8 \cdot 10^{-7} P + 1) - 0.786$  ( $n = 154$ ;  $r = 0.95$ ;  $s = 0.347$ ;  $F = 463.5$ ) was superior to the other studied models. Copyright © 1996 Elsevier Science Ltd

### INTRODUCTION

Numerous attempts have been made for modeling the accumulation of organic chemicals by fish since the experimental determination of this important biological activity is a time-consuming and costly exercise. The most widely used method consists in correlating laboratory bioconcentration factors (BCFs) determined in fish and *n*-octanol/water partition coefficients ( $\log P$ ). Bioconcentration can be defined as the result of the uptake, distribution, and elimination of a chemical in a fish due to water-borne exposure (Barron, 1990), whereas bioaccumulation includes all routes of exposure, and biomagnification deals with accumulation via food chains. Numerous test guidelines for the experimental determination of BCF in fish are available (e.g., OECD, ASTM). BCF is determined as the concentration of the chemical in whole fish at steady state divided by the concentration of the chemical in water during the exposure period and/or as the ratio between the uptake ( $k_1$ ) and depuration ( $k_2$ ) rate constants assuming first order kinetics. The accumulation of an organic compound in a fish basically depends on its hydrophobicity. Therefore,  $\log P$  is the physicochemical descriptor of choice for modeling bioconcentration. As a result, numerous BCF QSAR equations derived from  $\log P$  are available in the literature and there is a need for the evaluation of their domain of application. Under these conditions, the aim of this study was to compare different BCF/ $\log P$  models in order to estimate their accuracy, predictive power, and domain of application.

\* Author to whom all correspondence should be addressed.